

Measurement of the Deeply Virtual Neutral Pion Electroproduction Cross Section at the Thomas Jefferson National Accelerator Facility at 10.6 GeV

by

Robert Johnston

Submitted to the Department of Physics
in partial fulfillment of the requirements for the degree of

Interdisciplinary PhD in Physics and Statistics

at the

MASSACHUSETTS INSTITUTE OF TECHNOLOGY

June 2023

© Massachusetts Institute of Technology 2023. All rights reserved.

Author

Department of Physics

May 5, 2023

Certified by

Richard Milner

Professor of Physics

Thesis Supervisor

Accepted by

Lindley Winslow

Associate Department Head of Physics

**Measurement of the Deeply Virtual Neutral Pion
Electroproduction Cross Section at the Thomas Jefferson
National Accelerator Facility at 10.6 GeV**

by

Robert Johnston

Submitted to the Department of Physics
on May 5, 2023, in partial fulfillment of the
requirements for the degree of
Interdisciplinary PhD in Physics and Statistics

Abstract

Deeply virtual exclusive reactions provide unique channels to study both transverse and longitudinal properties of the nucleon simultaneously, allowing for a 3D image of nucleon substructure. This presentation will discuss work towards extracting an absolute cross section for one such exclusive process, deeply virtual neutral pion production, using 10.6 GeV electron scattering data off a proton target from the CLAS12 experiment in Jefferson Lab Hall B . This measurement is important as exclusive meson production has unique access to the chiral odd GPDs, and is also a background for other exclusive processes such as DVCS, making the determination of this cross section crucial for other exclusive analyses.

Thesis Supervisor: Richard Milner

Title: Professor of Physics

Acknowledgments

To Be Completed.

Contents

Contents	5
List of Figures	7
List of Tables	8
List of Equations	9
1 Introduction	11
1.1 Exploring Structure through Scattering	12
1.1.1 Discovery of the Proton	13
1.1.2 Electron as a Nucleon Probe	13
1.1.3 Structure of our World	14
1.2 Theoretical Background	14
1.2.1 GPDs and Deeply Virtual Scattering	14
1.2.2 The Handbag Approach	16
1.2.3 The Goloskokov-Kroll Model	17
1.3 Overview of Experimental Status	17
1.3.1 Previous Experimental Results	17
1.3.2 Overview of this Analysis	17

Bibliography	19
A	21
A.1 Full Cross Section Data	21
B	23
B.1 Cross check between Andrey Kim and Bobby Johnston	23

List of Figures

1-1	J.J. Thomson's Plum Pudding Model of the atom [6]	12
B-1	BSA Cross Check Results	24

List of Tables

1.1	GPDs Across Nucleon and Quark Polarizations	15
-----	---	----

List of Equations

1.1 DV π P Cross Section Decomposition	14
--	----

Chapter 1

Introduction

People have tried to understand the nature of the world around us for millennia, with discerning the structure of matter being a central effort in this quest. Famously, the Greek philosophers Leucippus and Democritus (\sim 5th century BCE) are credited with the concept of “atomism” - the belief that matter is composed of tiny indivisible particles called atoms (from the Greek *ατομος*, roughly translating to “uncuttable” [1]. Even further back, there are Indian records from as early as the 8th century BCE conceptualizing the world as being built from tiny fundamental particles [2].

Scientific progress on this front stalled until the early 1800s, when chemists explored how different elements combined in to form compounds in specific, repeatable, small integer ratios. John Dalton formulated this idea as the Law of Multiple Proportions, which paved the way for early scientific atomic theory [3]. In 1897, J.J. Thomson discovered the first subatomic particle, the electron, by studying cathode rays[4]. Accordingly, he devised a model of the atom which had electrons embedded in a ball of positively charged material, called the Thomson, or Plum Pudding, Model[5].




Figure 1-1: J.J. Thomson's Plum Pudding Model of the atom [6]

With the idea that the atom was a composite object, scientist began experimentation to study its exact structure. This was the starting point for the past 120 years of using particle scattering to probe atomic, and then nuclear, structure.

1.1 Exploring Structure through Scattering

Humanity interested in the structure of reality, atomic theory, subatomic, subnuclear, structure of this thesis. The rest of this chapter discusses these advancements, and describes the theoretical background for this experiment. Chapter 2 describes the experiment, CHapter 5 the analysis, etc. Chapter 8 summarizes this work and discusses how this analysis will proceed in the near future and a path for future experiments. The appendix includes numerous technical details and upplemental

plots.

As we increase the resolution (resolve features over smaller spatial distances), what we see is dependent on what resolution scale we are at. An exception to this case is if we are investigating point-like particles, which would have an identical response across all resolution scales

GPDs combine the kinematics of both elastic form factors and parton distributions. Andrey references 5, 6, and 7

1.1.1 Discovery of the Proton

Yimin has good references for this part

1.1.2 Electron as a Nucleon Probe

Include discussion of wavelength and momentum transfer (but in practice, limiting factor to resolution is lens system)

1961 hofstadter nobel prize (Andrey reference 1) Andrey reference 2 Friedman Kendall Tayleur scaling

Andrey thesis good write ups

Sangbaek: proton has properties Describe with QFT, words about QCD role of experiment in studying proton structure - proton discovery - neutron discovery - pointlike constituents - development of quark model - scaling behavior and asymptotic freedom - details about scattering experiments from first principles - words about elastic scattering (mott scattering) - Plot of elastic form factors, discussion of G_E and G_M , Rosenbluth formula - Mention TPEX - discussion of inelastic scattering - Structure functions - Discussion of spin and sum rules

Now discussion of DVEP: GPDs, Wigner functions relationships all the way

around some annoying math Handbag diagram Lepton hadron plane Status of experiments and future (EIC mapping)

invariant mass is W

Understanding the structure of matter has been a fundamental research pursuit for centuries.

Proton not a point mass - it has structure

1.1.3 Structure of our World

- atomic theory - discovery of proton / nucleon - proton structure measurements - lepton scattering experiments, HERA, etc

1.2 Theoretical Background

Discussion of Form Factors and generalized proton structure, Wigner Functions

1.2.1 GPDs and Deeply Virtual Scattering

The cross section for $DV\pi^0P$ has been theoretically linked to GPDs, which describe the 3D structure of the nucleon.

$$\frac{d^4\sigma_{\gamma^*p \rightarrow p'\pi^0}}{dQ^2 dx_B dt d\phi_\pi} = \Gamma(Q^2, x_B, E) \frac{1}{2\pi} \left\{ \left(\frac{d\sigma_T}{dt} + \epsilon \frac{d\sigma_L}{dt} \right) + \epsilon \cos(2\phi) \frac{d\sigma_{TT}}{dt} + \sqrt{2\epsilon(1+\epsilon)} \cos(\phi) \frac{d\sigma_{LT}}{dt} \right\} \quad (1.1)$$

There are 8 GPDs, 4 correspond to helicity conserving (chiral even) processes and 4 correspond to are helicity flipping (chiral odd) processes: H , E , \tilde{H} , and \tilde{E}

for chiral even, and H_T , E_T , \tilde{H}_T , and \tilde{E}_T

$\tilde{E}_T = 2^* \tilde{H}_T + E_T$ is commonly used

Nucleon Polarization	Quark Polarization		
	U	L	T
U	H		\tilde{E}_T
L		\tilde{H}	
T	E		H_T, \tilde{H}_T

Table 1.1: GPDs Across Nucleon and Quark Polarizations

Why do the structure combine in the way they do with the coefficients of cos phi terms and epsilons?

And the structure functions can be written as:

$$\frac{d\sigma_L}{dt} = \frac{4\pi\alpha}{kQ^2} \left\{ (1 - \xi^2) |\langle \tilde{H} \rangle|^2 - 2\xi^2 \Re [\langle \tilde{H} \rangle^* \langle \tilde{E} \rangle] - \frac{t'}{4m^2} \xi^2 |\langle \tilde{E} \rangle|^2 \right\} \quad (1.2)$$

$$\frac{d\sigma_T}{dt} = \frac{2\pi\alpha\mu_\pi^2}{kQ^4} \left\{ (1 - \xi^2) |\langle H_T \rangle|^2 - \frac{t'}{8m^2} |\langle \tilde{E}_T \rangle|^2 \right\} \quad (1.3)$$

$$\frac{d\sigma_{LT}}{dt} = \frac{4\pi\alpha\mu_\pi}{\sqrt{2}kQ^3} \xi \sqrt{1 - \xi^2} \frac{\sqrt{-t'}}{2m} \Re \left\{ \langle H_T \rangle^* \langle \tilde{E} \rangle \right\} \quad (1.4)$$

$$\frac{d\sigma_{TT}}{dt} = \frac{4\pi\alpha\mu_\pi^2}{kQ^4} \frac{-t'}{16m^2} \langle \tilde{E}_T \rangle^2 \quad (1.5)$$

and epsilon is...

and t' stands for.. $t - t_0$ where $t_0 = \frac{-4m^2\xi^2}{1-\xi^2}$

Where the skewness parameter is $\xi = \frac{x_B}{2-x_B}$ or whatever

The bracket $\langle \tilde{F} \rangle$ is the convolution of a GPD and an appropriate subprocess amplitude: $\langle \tilde{F} \rangle = \Sigma_\lambda \int_{-1}^1 d\bar{x} H_{0\lambda,0\lambda}(\bar{x}, \xi, Q^2, t=0) \tilde{F}(\bar{x}, \xi, Q^2, t)$

Where λ is the unobserved helicities of the partons participating in the subprocess

And k is the phase space factor given as $k = 16\pi (W^2 - m^2) \sqrt{\Lambda(W^2, -Q^2, m^2)}$

Where $\Lambda(W^2, -Q^2, m^2)$ is the Källén function and μ_{pi} is the reduced pion mass given as $\mu_{\pi^0} = \frac{m_{\pi^0}^2}{m_u + m_d}$ m_u and m_d are respective masses of up and down quarks.

Include proton pressure distribution plot

$$\Gamma(Q^2, x_B, E) = \frac{\alpha}{8\pi} \frac{Q^2}{m_p^2 E^2} \frac{1 - x_B}{x_B^3} \frac{1}{1 - \epsilon} \quad (1.6)$$

In addition to collinear momentum distribution of partons inside the nucleon, GPDs also encode the distribution of partons in the plane transverse to the nucleon's momentum in the infinite momentum frame [58]. Moreover, their relation to energy-momentum tensor (EMT) form factors allow us to access the EMT densities, the distribution of energy, angular momentum, pressure, and shear forces inside the nucleon [15].

Only valence quarks contribute electroproduction of uncharged pions.

1.2.2 The Handbag Approach

DVMP is sensitive to chiral odd GPDs, distinguishing it from DVCS as a GPD probe because why? Because something involving photon helicity and pion helicity, I forget exactly though

1.2.3 The Goloskokov-Kroll Model

1.3 Overview of Experimental Status

1.3.1 Previous Experimental Results

1.3.2 Overview of this Analysis

Hi [\[7\]](#) see more in section [1.1](#) just a test

Bibliography

- [1] C.C.W. Taylor, *The atomists, Leucippus and Democritus: fragments, a text and translation with a commentary* (University of Toronto Press, 1999), ISBN 0-8020-4390-9.
- [2] Thomas McEvilley, *The Shape of Ancient Thought: Comparative Studies in Greek and Indian Philosophies* (Allworth Press, New York, 2002), ISBN 978-0-8020-4390-0.
- [3] T. E. o. E. Britannica, *law of multiple proportions* (2010), URL <https://www.britannica.com/science/law-of-multiple-proportions>.
- [4] J. Thomson, *The Poular Science Monthly* pp. 323–325 (1901), URL <https://books.google.com/books?id=3CMDAAAAMBAJ&pg=PA323#v=onepage&q&f=false>.
- [5] Jaume Navarro, *A History of the Electron: J.J. and G.P. Thomson* (Cambridge University Press, 1995), ISBN 978-1-107-00522-8.
- [6] T. E. o. E. Britannica, *"Thomson atomic model"* (2023), URL <https://www.britannica.com/science/Thomson-atomic-model>.

- [7] I. Bedlinskiy, V. Kubarovsky, S. Niccolai, P. Stoler, K. P. Adhikari, M. D. Anderson, S. Anefalos Pereira, H. Avakian, J. Ball, N. A. Baltzell, et al., Physical Review C - Nuclear Physics **90** (2014), ISSN 1089490X, URL <https://journals.aps.org/prc/abstract/10.1103/PhysRevC.90.025205><https://arxiv.org/pdf/1405.0988.pdf>.

Appendix A

A.1 Full Cross Section Data

To be completed

Appendix B

B.1 Cross check between Andrey Kim and Bobby Johnston

As an additional cross check, Bobby calculated a $DV\pi^0P$ beam spin asymmetry and compared to Andrey Kim's results. This check will not comment on any acceptance, luminosity, or virtual photon flux factor calculations, but does validate exclusive event selection criteria. By examining figure [B-1](#) we can see that agreement is reasonable, especially considering Bobby's calculation does not have sideband subtraction included.

Fig [B-1](#) shows an overlay comparison of Andrey Kim's results (black datapoints, red fit line) and Bobby's results (red datapoints, orange fit line)




Figure B-1: BSA Cross Check Results