University of Cambridge Mathematical Tripos Part III

Functional Analysis

Lecturer:
András Zsák
a.zsak@dpmms.cam.ac.uk

Notes by:
Robert Frederik Uy
rfu20@cam.ac.uk

Course description

This course covers many of the major theorems of abstract Functional Analysis. It is intended to provide a foundation for several areas of pure and applied mathematics. We will cover as many of the following topics as time permits.

- Hahn–Banach theorems on the extension of linear functionals. Locally convex spaces. Discussion of the duals of the spaces $L_p(\mu)$ and $C(K)$.
- Weak and weak-star topologies. Hahn–Banach theorems on separation of convex sets. Theorems of Mazur, Goldstine and Banach–Alaoglu. Reflexivity and local reflexivity. Uniform convexity. The Milman–Pettis theorem. Duals of the spaces $L_p(\mu)$.
- Extreme points and the Krein–Milman theorem. Partial converse and the Banach–Stone theorem.
- Schauder bases in Banach spaces. Basis techniques and applications: compact and strictly singular operators; Pelczyński's decomposition method; the Eberlein–Šmulian theorem.
- The Bochner integral. The Krein–Šmulian theorem.
- Non-linear theory of Banach spaces. Lipschitz functions and the Lipschitz classification problem. Spaces with the Radon–Nikodym property. Linearization via differentiability and Rademacher's theorem. Linearization via invariant means. Uniform classification and Ribe's theorem.
- Banach algebras, elementary spectral theory. Commutative Banach algebras and the Gelfand representation theorem. Holomorphic functional calculus.
- Hilbert space operators, C^*-algebras. The Gelfand–Naimark theorem. Spectral theorem for commutative C^*-algebras. Spectral theorem and Borel functional calculus for normal operators.

Prerequisites

Thorough grounding in basic topology and analysis. Some knowledge of basic functional analysis and basic measure theory. In spectral theory we will make use of basic complex analysis. For example, Cauchy's theorem, Cauchy's integral formula and the maximum modulus principle. Much of the background material will be recalled either in lectures or via handouts.

Contents

1 The Hahn–Banach extension theorems **3**

1.1 Dual spaces 3

1.2 Biduals 5

1.3 Dual operators 6

1.4 Quotient spaces 7

1.5 Locally convex spaces 8

1.6 Dual space of $L_p(\mu)$ 11

1.7 Dual space of $C(K)$ 12

2 Weak topologies **13**

2.1 Weak topologies on vector spaces 14

2.2 Hahn–Banach separation theorems 17

1 The Hahn–Banach extension theorems

1.1 Dual spaces

Let X be a (real or complex) normed space. Recall from II Linear Analysis that the *dual space* X^* is defined as the space of all bounded linear functions on X, which is a Banach space with respect to the operator norm

$$||f|| := \sup\{|f(x)| \colon x \in B_X\}$$

Above, we denote by B_X the closed unit ball in X

$$B_X := \{x \in X \colon ||x|| \leq 1\}$$

Similarly, we denote by S_X the unit sphere

$$S_X := \{x \in X \colon ||x|| = 1\}$$

Examples *Normed spaces and their dual*

- $\ell_p^* \cong \ell_q$ for $1 \leq p < \infty, 1 < q \leq \infty$ with $p^{-1} + q^{-1} = 1$
- $c_0^* \cong \ell_1$
- If $\mathcal{H}$ is a Hilbert space, then $\mathcal{H}^* \cong \mathcal{H}$ (conjugate-linear in the complex case)

Notational remark. Let X, Y be normed spaces.

- We write $X \sim Y$ if X and Y are *isomorphic*, i.e. there exists a linear bijection $T \colon X \to Y$ such that T and T^{-1} are continuous.
- We write $X \cong Y$ if X, Y are *isometrically isomorphic*, i.e. there exists a linear surjective map $T \colon X \to Y$ such that, for every $x \in X$, $||Tx|| = ||x||$.
- For $x \in X$ and $f \in X^*$, we write $\langle x, f \rangle$ for the *action of f on x*, i.e. $f(x)$.

Definition Positive homogeneous and subadditive

Let X be a real vector space. We say that a functional $p \colon X \to \mathbb{R}$ is

- *positive homogeneous if $p(tx) = tp(x)$ for all $x \in X$ and $t \geq 0$*
- *subadditive if $p(x + y) \leq p(x) + p(y)$ for all $x, y \in X$*

Theorem 1.1 Hahn–Banach I

Let X be a real vector space and $p \colon X \to \mathbb{R}$ a positive homogeneous and subadditive functional. Let $Y \subset X$ be a subspace and $g \colon Y \to \mathbb{R}$ a linear map such that $g(y) \leq p(y)$ for every $y \in Y$. Then there exists a linear map $f \colon X \to \mathbb{R}$ such that

(i) $f|_Y = g$

(ii) $f(x) \leq p(x)$ *for every* $x \in X$

Proof. Consider the set

$$\mathcal{P} := \left\{ (Z, h) \colon \begin{array}{l} Y \subset Z \subset X, Z \text{ subspace}, \\ h \colon Z \to \mathbb{R} \text{ linear}, \\ h|_Y = g, \\ h(z) \leq p(z) \ \forall z \in Z \end{array} \right\}$$

Note that $\mathcal{P}$ is partially ordered by extension, i.e. $(Z_1, h_1) \preceq (Z_2, h_2)$ iff $Z_1 \subset Z_2$ and $h_2|_{Z_1} = h_1$. By Zorn's lemma, $\mathcal{P}$ has a maximal element (Z, h).

Suppose, on the contrary, that $Z \neq X$. Then we can fix $x \in X \backslash Z$ and $\alpha \in \mathbb{R}$. Define $h_1 \colon Z_1 := Z + \mathbb{R}x \to \mathbb{R}$ by $h_1(z + \lambda x) = h(z) + \lambda\alpha$. By construction, h_1 is linear and $h_1|_Z = h$.

It then suffices to find α such that $h_1(z+\lambda x) \leq p(z+\lambda x)$ for every $z \in Z$ and $\lambda \in \mathbb{R}$, since this would imply that $(Z_1, h_1) \in \mathcal{P}$ — a contradiction! To this end, note that, since p is positive homogeneous, we need

$$h(z) + \alpha \leq p(z+x)$$
$$h(z) - \alpha \leq p(z-x)$$

for all $z \in Z$, i.e. for every $y, z \in Z$,

$$h(y) - p(y-x) \leq \alpha \leq p(z+x) - h(z)$$

Of course, it suffices to show that, for every $y, z \in Z$,

$$h(y) - p(y-x) \leq p(z+x) - h(z)$$

Indeed, by hypothesis, we have

$$h(y) + h(z) = h(y+z) \leq p(y+z) \leq p(y-x+x+z) \leq p(y-x) + p(x+z)$$

as required. ■

Definition Seminorm

Let X be a real or complex normed space. A seminorm on X is a function $p\colon X \to \mathbb{R}$ such that

(i) $p(x) \geq 0$ *for all* $x \in X$

(ii) $p(\lambda x) = |\lambda| p(x)$ *for all* $x \in X$ *and scalars* λ

(iii) $p(x+y) \leq p(x) + p(y)$ *for all* $x, y \in X$

Remark. A norm is a seminorm, which is in turn a positive homogeneous and subadditive functional.

Theorem 1.2 Hahn–Banach II

Let X be a real or complex normed space and p a seminorm on X. Let $Y \subset X$ be a subspace and g a linear functional on Y such that $|g(y)| \leq p(y)$ for all $y \in Y$. Then there exists a linear functional f on X such that $f|_Y = g$ and $|f(x)| \leq p(x)$ for all $x \in X$.

Proof. We shall first prove the real case and then extend it to the complex case.

Case 1: Real case

Note that p is a positive homogeneous and subadditive functional satisfying $g(y) \leq |g(y)| \leq p(y)$ for all $y \in Y$. By Theorem 1.1, there exists a linear functional $f\colon X \to \mathbb{R}$ such that $f|_Y = g$ and $f(x) \leq p(x)$ for all $x \in X$. For $x \in X$, $-f(x) = f(-x) \leq p(-x) = p(x)$, so $|f(x)| \leq p(x)$.

Case 2: Complex case

Let $g_1 = \mathrm{Re}(g)$. Then $g_1\colon Y \to \mathbb{R}$ is $\mathbb{R}$-linear and, for every $y \in Y$, $|g_1(y)| \leq |g(y)|$. By the real case, there exists an $\mathbb{R}$-linear $f_1\colon X \to \mathbb{R}$ such that $f_1|_Y = g_1$ and, for every $x \in X$, $|f_1(x)| \leq p(x)$.

▶ **Claim.** There exists a unique linear map $f\colon X \to \mathbb{C}$ such that $\mathrm{Re}\, f = f_1$.

Proof. We simply take $f(x) = f_1(x) - i f_1(ix)$.

Note that $\mathrm{Re}(f|_Y) = f_1|_Y = g_1 = \mathrm{Re}(g)$, so $f|_Y = g$ by uniqueness. For $x \in X$, we have $|f(x)| = \lambda f(x) = f(\lambda x) = f_1(\lambda x) \leq p(\lambda x) = |\lambda| p(x)$ for some $\lambda \in \mathbb{C}, |\lambda| = 1$. ■

Remark. For a complex vector space V, we shall denote by $V_{\mathbb{R}}$ for V viewed as a real vector space. Then the proof of Theorem 1.2 shows that, for a complex normed space X, the map

$$\begin{aligned}(X^*)_{\mathbb{R}} &\to (X_{\mathbb{R}})^* \\ f &\mapsto \operatorname{Re} f\end{aligned}$$

is an isometric isomorphism.

Corollary 1.3
Let X be a real or complex vector space and p a seminorm on X. For any $x_0 \in X$, there exists a linear functional f on X such that

(i) *for every $x \in X$, $|f(x)| \leq p(x)$*
(ii) *$f(x_0) = p(x_0)$*

Proof. Let $Y = \operatorname{span}\{x_0\}$. Define a linear functional g on Y by setting $g(\lambda x_0) = \lambda p(x_0)$. Then for every scalar λ, $|g(\lambda x_0)| = |\lambda| p(x_0) = p(\lambda x_0)$. By Theorem 1.2, there exists a linear functional f on X such that $f|_Y = g$ and, for every $x \in X$, $|f(x)| \leq p(x)$. Of course, we also have $f(x_0) = g(x_0) = p(x_0)$. ∎

Theorem 1.4 Hahn–Banach III
Let X be a real or complex normed space.

(i) *Given a subspace Y of X and $g \in Y^*$, there exists $f \in X^*$ such that $f|_Y = g$ and $||f|| = ||g||$.*
(ii) *Given $x_0 \in X \backslash \{0\}$, there exists $f \in S_{X^*}$ such that $f(x_0) = ||x_0||$.*

Proof. To prove (i), we define $p\colon X \to \mathbb{R}$ by $p(x) := ||g||\,||x||$. This is a seminorm on X and, for every $y \in Y$, $|g(y)| \leq ||g||\,||y|| = p(y)$. We then proceed by applying Theorem 1.2 to obtain some $f \in X^*$ with $f|_Y = g$ and $||f|| \leq ||g||$. Since $f|_Y = g$, we immediately have $||f|| \geq ||g||$. For (ii), we take $p = ||\cdot||$. Apply the previous corollary. ∎

Remarks

- Theorem 1.4(i) is a linear version of Tietze's extension theorem (cf. II Linear Analysis).
- Theorem 1.4(ii) implies that X^* separates points of X, i.e. given $x \neq y$ in X, there exists $f \in X^*$ such that $f(x) \neq f(y)$. (Indeed, simply take $x_0 = x - y$.)
- The f in Theorem 1.4(ii) is sometimes called a *norming functional at* x_0:

  $$||x_0|| = \max\{|\langle x_0, g\rangle| \colon g \in B_{X^*}\}$$

  Alternatively, f is also called a *support functional at* x_0 (cf. separation theorems in II Analysis of Functions).

1.2 Biduals

Let X be a normed space. We call $X^{**} := (X^*)^*$ the *bidual* or *second dual* of X. Recall from II Linear Analysis that it is a Banach space in the operator norm. For each $x \in X$, define the map

$$\begin{aligned}\hat{x}\colon X^* &\to \mathbb{R} \\ f &\mapsto f(x)\end{aligned}$$

Note that $\hat{x}$ is linear and bounded: $|\hat{x}(f)| = |f(x)| \leq ||f||\,||x||$. So $\hat{x} \in X^{**}$ with $||\hat{x}|| \leq ||x||$. Thus, we can then define the *canonical embedding* of X into X^{**}

$$\hat{\ }\colon X \to X^{**}$$

$$x \mapsto \hat{x}$$

Theorem 1.5
The map $\hat{}$ is an isometric isomorphism of X into X^{**}.

Proof. The map is well-defined and norm-decreasing by above. It is easy to check linearity. The map is isometric since

$$||\hat{x}|| = \sup\{|\hat{x}(f)| \colon f \in B_{X^*}\} = ||x||$$

by Theorem 1.4(ii). ∎

Remarks

- For $x \in X$ and $f \in X^*$, we have $\langle f, \hat{x} \rangle = \langle x, f \rangle$.
- $X \cong \hat{X} := \hat{}(X) \subset X^{**}$, so $\hat{X}$ is closed in X^{**} iff X is complete.
- In general, $\overline{\hat{X}}$ (closure in X^{**}) is a Banach space containing an isometric copy of X as a dense subspace.

Definition Reflexive
We say that X is reflexive if $\hat{X} = X^{**}$.

Examples

- Reflexive spaces: ℓ_p for any $p \in (1, \infty)$, any Hilbert space, any finite-dimensional space (see also $L_p(\mu)$ later on)
- Not reflexive spaces: $c_0, \ell_1, L_1[0,1]$

Remark. If X is reflexive, then $X \cong X^{**}$. But there exist Banach spaces X such that $X \cong X^{**}$ but X is not reflexive.

1.3 Dual operators

Let X, Y be normed spaces. We denote by $\mathcal{B}(X,Y) := \{T \colon X \to Y \mid T \text{ is linear and bdd}\}$. Recall from II Linear Analysis that this is a normed space in the operator norm $||T|| := \sup\{||Tx|| \colon x \in B_X\}$ and that $\mathcal{B}(X,Y)$ is complete if Y is complete.

For $T \in \mathcal{B}(X,Y)$, the *dual operator* of T is the map

$$T^* \colon Y^* \to X^*$$
$$g \mapsto g \circ T$$

It is easy to see that T is a well-defined linear map and it is bounded with $||T^*|| = ||T||$:

$$||T^*|| = \sup_{g \in B_{Y^*}} ||T^* g|| = \sup_{g \in B_{Y^*}} \sup_{x \in B_X} |\langle x, T^*(g) \rangle| = \sup_{x \in B_X} \sup_{g \in B_{Y^*}} |\langle Tx, g \rangle| = \sup_{x \in B_X} ||Tx|| = ||T||$$

where the penultimate equality follows from Theorem 1.4(ii).

Remark. When X, Y are Hilbert spaces, the dual operator T^* 'is' the adjoint of T.

Example. For $1 < p, q < \infty$ with $p^{-1} + q^{-1} = 1$, the dual of the right-shift operator $R \colon \ell_p \to \ell_p$ is simply the left-shift operator $L \colon \ell_q \to \ell_q$.

Now, let's note some properties of dual operators:

(1) $(\mathrm{Id}_X)^* = \mathrm{Id}_{X^*}$
(2) $(\lambda S + \mu T)^* = \lambda S^* + \mu T^*$ for all $S, T \in \mathcal{B}(X,Y)$ and all scalars λ, μ

(3) $(ST)^* = T^*S^*$ for all $T \in \mathcal{B}(X,Y)$ and all $S \in \mathcal{B}(Y,Z)$

(4) The map

$$\begin{aligned} \mathcal{B}(X,Y) &\to \mathcal{B}(Y^*, X^*) \\ T &\mapsto T^* \end{aligned}$$

is an into isometric isomorphism.

(5) Let $T \in \mathcal{B}(X,Y)$. The diagram

$$\begin{array}{ccc} X & \xrightarrow{T} & Y \\ {\scriptstyle\wedge}\downarrow & & {\scriptstyle\wedge}\downarrow \\ X^{**} & \xrightarrow[T^{**}]{} & Y^{**} \end{array}$$

commutes, i.e., $\widehat{Tx} = T^{**}\hat{x}$ for all $x \in X$. For $x \in X, g \in Y^*$,

$$\langle g, T^{**}\hat{x}\rangle = \langle T^*g, \hat{x}\rangle = \langle x, T^*g\rangle = \langle Tx, g\rangle = \langle g, \widehat{Tx}\rangle$$

Remark. Properties (1) and (3) imply that $X \sim Y \implies X^* \sim Y^*$.

1.4 Quotient spaces

Let X be a normed space and Y a closed subspace. Then X/Y is a normed space in the quotient norm defined by

$$||x + Y|| := d(X,Y) := \inf\{||x+y|| \colon y \in Y\}$$

The quotient map $q\colon X \to X/Y$ is linear, surjective and bounded with $||q|| = 1$. Indeed, we have $||q(x)|| \le ||x||$ so $||q|| \le 1$. Note also that $q(D_X) = D_{X/Y}$ (for $x \in D_X$, we have $||q(x)|| \le ||x|| < 1$, so $q(x) \in D_{X/Y}$) and that, given $z \in D_{X/Y}$, we can pick some $x \in X$ such that $q(x) = z$ and, since $||z|| < 1$, we have $||x + y|| < 1$ for some $y \in Y$ and thus $x + y \in D_X$ with $q(x+y) = z$. It then follows that $||q|| = 1$ and q is an open map.

Given another normed space Z and a bounded linear map $T\colon X \to Z$ such that $Y \subset \ker T$, there exists a unique map $\tilde{T}$ such that

$$\begin{array}{ccc} X & \xrightarrow{\quad T \quad} & Z \\ & {\scriptstyle q}\searrow \quad \nearrow{\scriptstyle \tilde{T}} & \\ & X/Y & \end{array}$$

commutes, i.e. $T = \tilde{T} \circ q$. Moreover, $\tilde{T}$ is linear and bounded with $||\tilde{T}|| = ||T||$ since

$$\tilde{T}(D_{X/Y}) = \tilde{T}(q(D_X)) = T(D_X)$$

Theorem 1.6
Let X be a normed space. If X^* is separable, then so is X.

Proof. Pick a countable set $\{f_n \colon n \in \mathbb{N}\}$ dense in S_{X^*}. For every $n \in \mathbb{N}$, we can pick some $x_n \in B_X$ such that $f_n(x_n) > \frac{1}{2}$. Let $Y = \overline{\operatorname{span}}\{x_n \colon n \in \mathbb{N}\}$. It suffices to show that $Y = X$. Suppose, on the contrary, that $Y \neq X$. Then we can pick $g \in S_{X^*}$ such that $g|_Y = 0$. This is possible since we can, for instance, pick some $h \in (X/Y)^*$ such that $h \neq 0$ [e.g. Theorem 1.4(ii)], and we can then take $g = (h \circ q)/||h \circ q||$. Then there exists some $n \in \mathbb{N}$ such that $||f_n - g|| < \frac{1}{10}$. Then $||g(x_n)|| \ge \frac{1}{2} - \frac{1}{10} > 0$, which is a contradiction. ∎

Remark. The converse is false in general. For instance, the dual of $X = \ell_1$ is $X^* = \ell_\infty$, which is not separable.

Theorem 1.7
Any separable normed space X embeds isometrically into ℓ_∞.

Proof. Let $\{x_n : n \in \mathbb{N}\}$ be dense in X. For each $n \in \mathbb{N}$, let f_n be a norming functional at x_n, cf. Theorem 1.4(ii). We then define $T : X \to \ell_\infty$ by $Tx = (f_n(x))_{n=1}^\infty$. This is a bounded linear map since, for every $n \in \mathbb{N}$, we have $|f_n(x)| \leq ||x||$, so $Tx \in \ell_\infty$ and $||Tx||_\infty \leq ||x||$. For every $n \in \mathbb{N}$, we also have $||Tx_n||_\infty \geq |f_n(x_n)| = ||x_n||$ and thus $||Tx_n||_\infty = ||x_n||$. Hence, by density and continuity, it follows that $||Tx||_\infty = ||x||$. ∎

Remarks

- ℓ_∞ is isometrically universal for the class $\mathcal{SB}$ of separable Banach spaces. In fact, we will see later that there exists $Z \in \mathcal{SB}$ isometrically universal for $\mathcal{SB}$.
- A dual version: every separable Banach space is a quotient of ℓ_1.

Theorem 1.8 Vector-valued Liouville
Let X be a complex Banach space and $f : \mathbb{C} \to X$ holomorphic and bounded. Then f is constant.

Proof. Pick some $M > 0$ such that $||f(z)|| \leq M$ for all $z \in \mathbb{C}$. Since f is holomorphic, we have that, for every $z \in \mathbb{C}$, the limit

$$\lim_{w \to z} \frac{f(w) - f(z)}{w - z}$$

exists and shall be denoted by $f'(z)$. Now, fix $\varphi \in X^*$. Then $\varphi \circ f : \mathbb{C} \to \mathbb{C}$ is bounded

$$|\varphi(f(z))| \leq ||\varphi||||f(z)|| \leq M||\varphi||$$

and holomorphic

$$\lim_{w \to z} \frac{\varphi(f(w)) - \varphi(f(z))}{w - z} = \varphi\left(\lim_{w \to z} \frac{f(w) - f(z)}{w - z}\right)$$

where we have used the fact that φ is linear and continuous. By scalar Liouville, we have that $\varphi(f(z)) = \varphi(f(0))$ for every $z \in \mathbb{C}$.

Fix $z \in \mathbb{C}$. By above, we have that, for every $\varphi \in X^*$, $\varphi(f(z)) = \varphi(f(0))$. By Theorem 1.4(ii), it follows that $f(z) = f(0)$. ∎

1.5 Locally convex spaces

Definition Locally convex space (LCS)
A locally convex space is a pair $(X, \mathcal{P})$, where X is a (real or complex) vector space and $\mathcal{P}$ is a family of seminorms on X that separates points of X in the sense that, for every nonzero $x \in X$, there exists $p \in \mathcal{P}$ such that $p(x) \neq 0$.

Note that $\mathcal{P}$ induces a topology on X: $U \subset X$ is open iff

$$\forall\, x \in U\ \exists\, n \in \mathbb{N}\ \exists\, p_1, \cdots, p_n \in \mathcal{P}\ \exists\, \varepsilon > 0 \text{ s.t. } \{y \in X : p_k(y - x) < \varepsilon, 1 \leq k \leq n\} \subset U$$

Some basic properties of this topology include

- Addition and scalar multiplication are continuous
- It is Hausdorff
- $x_n \to x$ iff $p(x_n - x) \to 0$ for all $p \in \mathcal{P}$

- Let Y be a subspace of X. Let $\mathcal{P}_Y := \{p|_Y : p \in \mathcal{P}\}$. Then $(Y, \mathcal{P}_Y)$ is a LCS and its topology is the subspace topology induced by X on Y.
- Given two families $\mathcal{P}, \mathcal{Q}$ on X, they are equivalent (denoted by $\mathcal{P} \sim \mathcal{Q}$) if they induce the same topology on X. The topology of $(X, \mathcal{P})$ is metrisable iff there exists a countable $\mathcal{Q} \sim \mathcal{P}$.

Definition Fréchet space
A Fréchet space is a complete metrisable locally convex space.

Examples

- Any normed space $(X, ||\cdot||)$ is a LCS with $\mathcal{P} = \{||\cdot||\}$.
- Fix a nonempty open $U \subset \mathbb{C}$. Define

$$\mathcal{O}(U) := \{f : U \to \mathbb{C} \mid f \text{ is holomorphic}\}$$

  For a compact $K \subset U$, let $p_K(f) := \sup_{z \in K} |f(z)|$ for every $f \in \mathcal{O}(U)$. Let $\mathcal{P} = \{p_K : \text{ compact } K \subset U\}$. Then $(\mathcal{O}(U), \mathcal{P})$ is a LCS and its topology is the topology of local uniform convergence. It is also a Fréchet space and not normable, i.e. the topology is not induced by a norm.
- Fix $d \in \mathbb{N}$ and a nonempty open $\Omega \subset \mathbb{R}^d$. We define

$$C^\infty(\Omega) := \{f : \Omega \to \mathbb{R} \mid f \text{ is smooth}\}$$

  For a multi-index $\alpha \in \mathbb{Z}^d_{\geq 0}$, we can define

$$D^\alpha := \left(\frac{\partial}{\partial x_1}\right)^{\alpha_1} \cdots \left(\frac{\partial}{\partial x_d}\right)^{\alpha_d}$$

  For compact $K \subset \Omega$, we define

$$p_{K,\alpha}(f) := \sup_{w \in K} |(D^\alpha f)(w)|$$

  We then take $\mathcal{P} = \{p_{K,\alpha} : \text{ compact } K \subset \Omega, \alpha \in \mathbb{Z}^d_{\geq 0}\}$. Then $(C^\infty(\Omega), \mathcal{P})$ is a LCS. It is also Fréchet space and not normable.

Lemma 1.9
Let $(X, \mathcal{P})$ and $(Y, \mathcal{Q})$ be locally convex spaces and $T : X \to Y$ a linear map. The following are equivalent:

(i) *T is continuous*
(ii) *T is continuous at 0*
(iii) *for every $q \in \mathcal{Q}$, there exist $n \in \mathbb{N}$, $p_1, \cdots, p_n \in \mathcal{P}$ and $C \geq 0$ such that, for every $x \in X$,*

$$q(Tx) \leq C \max_{1 \leq k \leq n} p_k(x)$$

Proof. (i) $\implies$ (ii): Trivial.
(ii) $\implies$ (i): Translation is continuous.
(ii) $\implies$ (iii): Fix $q \in \mathcal{Q}$. Define $V = \{y \in Y : q(y) \leq 1\}$, which is a neighbourhood of 0 in Y. Then there exists a neighbourhood U of 0 in X such that $T(U) \subset V$. WLOG, $U = \{x \in X : p_k(x) \leq \varepsilon \ \forall 1 \leq k \leq n\}$ for some $n \in \mathbb{N}$, $p_1, \cdots, p_n \in \mathcal{P}$, $\varepsilon > 0$. It then remains to show that, for every $x \in X$, we have $q(Tx) \leq \frac{1}{\varepsilon} \max_{1 \leq k \leq n} p_k(x)$. Fix $x \in X$. If $p(x) \neq 0$, then $\varepsilon x / p(x) \in U$ and thus $T(\varepsilon x / p(x)) \in V$. It then follows that $q(T(\varepsilon x / p(x))) \leq 1$, i.e. $q(Tx) \leq p(x)/\varepsilon$. If $p(x) = 0$, then $\lambda x \in U$ for every scalar λ, so $q(T(\lambda x)) \leq 1$ for every scalar λ. Thus, $q(Tx) = 0 \leq p(x)/\varepsilon$.

(iii) $\Longrightarrow$ (ii): Given a neighbourhood V of 0 in Y, we seek a neighbourhood of U of 0 in X such that $T(U) \subset V$. WLOG, $V = \{y \in Y : q_k(y) \leq \varepsilon \ \forall 1 \leq k \leq n\}$ for some $n \in \mathbb{N}$, $q_1, \cdots, q_n \in \mathcal{Q}$. By (iii), for each $1 \leq k \leq n$, there exist $m_k \in \mathbb{N}$, $p_{k,1}, \cdots, p_{k,m_k} \in \mathcal{P}$, $C_k \geq 0$ such that

$$q_k(Tx) \leq C_k \max_{1 \leq j \leq n} p_{k,j}(x)$$

for every $x \in X$. Then $U = \{x \in X : p_{k,j}(x) \leq \frac{\varepsilon}{C_k+1}, 1 \leq j \leq m_k, 1 \leq k \leq n\}$ is a neighbourhood of 0 in X with the property that $T(U) \subset V$. ∎

Definition Dual space
Let $(X, \mathcal{P})$ be a locally convex space. The dual space X^* of X is the space of continuous linear functionals on X.

Lemma 1.10
Let X be a locally convex space and f a linear functional on X. Then f is continuous iff $\ker f$ *is closed.*

Proof. ($\Longrightarrow$) $\ker f = f^{-1}(\{0\})$
($\Longleftarrow$) The case $\ker f = X$ is trivial, so it remains to carefully consider the case where $\ker f \neq X$. Fix $x_0 \in X \setminus \ker f$. Then there exists an open neighbourhood U of 0 in X such that $(x_0 + U) \cap \ker f = \varnothing$. WLOG, we may assume that U is convex and balanced (i.e. $x \in U, |\lambda| = 1 \Longrightarrow \lambda x \in U$). Since f is linear, $f(U)$ is also convex and balanced. Note that $f(U)$ cannot be the whole scalar field since, in that case, we have $f(x_0 + U)$ — a contradiction since $0 \notin f(x_0 + U)$! Thus, $f(U)$ must be bounded, i.e. $f(U) \subset \{\lambda : |\lambda| \leq M\}$ for some $M \geq 0$. Then for every $\varepsilon > 0$, $\frac{\varepsilon}{M+1}U$ is a neighbourhood of 0 with $f(\frac{\varepsilon}{M+1}U) \subset \{\lambda : |\lambda| \leq \varepsilon\}$. Hence, f is continuous at 0 and thus $f \in X^*$. ∎

Theorem 1.11 Hahn–Banach theorem IV
Let $(X, \mathcal{P})$ be a locally convex space. Then

(i) *Given a subspace Y of X and $g \in Y^*$, there exists $f \in X^*$ such that $f|_Y = g$.*
(ii) *Given a closed subspace Y of X and $x_0 \in X \setminus Y$, there exists $f \in X^*$ such that $f|_Y = 0$ and $f(x_0) \neq 0$.*

Remark. X^* separates points of X: given $x \neq y$ in X, we can apply Theorem 1.11(ii) with $Y = \{0\}$ and $x_0 = x - y$.

Proof. (i): By Lemma 1.9, there exist $n \in \mathbb{N}$, $p_1, \cdots, p_n \in \mathcal{P}$ and $C \geq 0$ such that, for every $y \in Y$,

$$|g(y)| \leq C \max_{1 \leq k \leq n} p_k(y)$$

Set $p(x) := C \max_{1 \leq k \leq n} p_k(x)$ for every $x \in X$. It is easy to check that p is a seminorm on X. Moreover, we have $|g(y)| \leq p(y)$ for every $y \in Y$. By Theorem 1.2, there exists a linear functional f on X such that $f|_Y = g$ and, for every $x \in X$, $|f(x)| \leq p(x)$. By Lemma 1.9, f is continuous.
(ii): Let $Z = \operatorname{span}(Y \cup \{x_0\})$ and define the linear functional g on Z by $g(y + \lambda x_0) = \lambda$. Of course, $\ker g = Y$, which is closed and thus continuous by Lemma 1.10. By (i), there exists $f \in X^*$ such that $f|_Z = g$. Then $f|_Y = 0$ and $f(x_0) = g(x_0) = 1 \neq 0$. ∎

1.6 Dual space of $L_p(\mu)$

Fix a measure space $(\Omega, \mathcal{F}, \mu)$. Let $1 \leq p < \infty$ and $1 < q \leq \infty$ such that $p^{-1} + q^{-1} = 1$. For $g \in L_q$, we can define

$$\begin{aligned} \varphi_g \colon L_p &\to \{\text{scalars}\} \\ f &\mapsto \int_\Omega fg \, d\mu \end{aligned}$$

It is easy to check that φ_g is a well-defined linear map satisfying $|\varphi_g(f)| \leq ||f||_p ||g||_q$ (by Hölder). Then $\varphi_g \in L_p^*$ and $||\varphi_g|| \leq ||g||_q$. Then we can define a map

$$\begin{aligned} \varphi \colon L_q &\to L_p^* \\ g &\mapsto \varphi_g \end{aligned}$$

This is a bounded linear map with $||\varphi|| \leq 1$.

Theorem 1.12
Let $(\Omega, \mathcal{F}, \mu)$, p, q, φ be as above. Then

(i) *If $1 < p < \infty$, then φ is an isometric isomorphism, i.e. $L_p^* \cong L_q$.*

(ii) *If $p = 1$ and μ is σ-finite, then φ is an isometric isomorphism, i.e. $L_1^* \cong L_\infty$.*

Remark. Recall from II Probability and Measure that σ-finite means that there exist (A_n) in $\mathcal{F}$ such that $\Omega = \cup_n A_n$ and $\mu(A_n) < \infty$ for all n.

Proof. We first show that φ is isometric. Fix $g \in L_q$. Fix a measurable function λ such that $|\lambda| = 1$ a.e. and $\lambda g = |g|$ a.e.

(i) Let $f = \lambda |g|^{q-1}$. Then

$$\int_\Omega |f|^p \, d\mu = \int_\Omega |g|^{pq-p} \, d\mu = ||g||_q^q$$

Thus, $f \in L_p$ and $||f||_p^p = ||g||_q^q$. It then follows that

$$||f||_p ||\varphi_g|| \geq |\varphi_g(f)| = \int_\Omega |g|^q \, d\mu = ||g||_q^q$$

so $||\varphi_g|| \geq ||g||_q^{q-q/p} = ||g||_q$, as required.

(ii) The case $||g||_\infty = 0$ is trivial, so it remains to carefully consider the case where $||g||_\infty > 0$. Fix $0 \leq s < ||g||_\infty$. Then $A = \{u \in \Omega \colon |g(u)| \geq s\}$ has $\mu(A) > 0$. Since μ is σ-finite, there exists $B \subset A$ with $B \in \mathcal{F}$ such that $0 < \mu(B) < \infty$. Then $f = \lambda \mathbb{1}_B \in L_1$ and

$$||f||_1 = \int_B |\lambda| \, d\mu = \mu(B)$$

Then

$$\mu(B) ||\varphi_g|| \geq |\varphi_g(f)| = \int_B |g| \, d\mu \geq s\mu(B)$$

Hence, $||\varphi_g|| \geq s$ and thus $||\varphi_g|| \geq ||g||_\infty$.

Secondly, we show that φ is surjective. One approach is via the Radon–Nikodym theorem, but we will do it using uniform convexity. This will be covered in Chapter 4, so we defer the proof until then. ∎

1.7 Dual space of $C(K)$

Fix a compact Hausdorff space K. We then define the spaces

$$\begin{aligned} C(K) &:= \{f\colon K \to \mathbb{C} \mid f \text{ continuous}\} \\ C^{\mathbb{R}}(K) &:= \{f\colon K \to \mathbb{R} \mid f \text{ continuous}\} \\ C^{+}(\mathbb{K}) &:= \{f\colon K \to \mathbb{R}_{\geq 0} \mid f \text{ continuous}\} \\ M(K) &:= C(K)^* = \mathcal{B}(C(K), \mathbb{C}) \\ M^{\mathbb{R}}(K) &:= \{\varphi \in M(K) \mid \varphi(f) \in \mathbb{R}\ \forall f \in C^{\mathbb{R}}(K)\} \\ M^{+}(K) &:= \{\varphi\colon C(K) \to \mathbb{C} \mid \varphi \text{ linear, positive}\} \end{aligned}$$

Above, we say that φ is *positive* if $\varphi(f) \in \mathbb{R}_{\geq 0}$ for all $f \in C^+(K)$.

Remarks

- By considering the map $\varphi \mapsto \varphi|_{C^{\mathbb{R}}(K)}$, it is easy to see that $M^{\mathbb{R}}(K) \cong C^{\mathbb{R}}(K)^*$.
- $M^+(K) \subset M^{\mathbb{R}}(K)$ and, in fact, we have

$$M^+(K) = \{\varphi \in M(K)\colon ||\varphi|| = \varepsilon(\mathbb{1}_K)\}$$

Let ν be a *complex Borel measure* on K, i.e. $\nu\colon \mathcal{B} \to \mathbb{C}$ is a countably additive set-function on the Borel σ-field $\mathcal{B}$ on K. For $f \in C(K)$,

$$\int_K |f|\, d|\nu| \leq ||f||_\infty \cdot |\nu|(K)$$

where $|\nu|$ is the *total variation measure* of ν (cf. II Analysis of Functions). Thus, f is ν-integrable. We can then define a map

$$\begin{aligned} \varphi\colon C(K) &\to \mathbb{C} \\ f &\mapsto \int_K f\, d\nu \end{aligned}$$

Note that φ is linear and

$$|\varphi(f)| \leq \int_K |f|\, d|\nu| \leq ||f||_\infty \cdot |\nu|(K)$$

so $\varphi \in M(K) = C(K)^*$ with $||\varphi|| \leq |\nu|(K) =: ||\nu||_1$ (called the *total variation of* ν). Moreover, if ν is a signed measure (i.e. a countably additive set-function $\mathcal{B} \to \mathbb{R}$), then $\varphi \in M^{\mathbb{R}}(K)$. Similarly, if ν is a (positive) measure, then $\varphi \in M^+(K)$.

Theorem 1.13 Riesz representation theorem
For any $\varphi \in M(K)$, there exists a unique regular complex Borel measure ν on K that represents φ in the sense that

$$\varphi(f) = \int_K f\, d\nu$$

for all $f \in C(K)$. Moreover, $||\varphi|| = ||\nu||_1$. Furthermore, we have

(i) *if $\varphi \in M^{\mathbb{R}}(K)$, then ν is a signed measure.*
(ii) *if $\varphi \in M^+(L)$, then ν is a (positive) measure.*

Remark. Above, *regular* means that, for every $A \in \mathcal{B}$ and every $\varepsilon > 0$, there exists a closed set E and an open set U such that $E \subset A \subset U$ and $|\nu|(U \backslash E) < \varepsilon$.

Corollary 1.14

(i) *The space of regular complex Borel measures on K is a complex Banach space in $||\cdot||$. Moreover, it is isometrically isomorphic to $M(K) := C(K)^*$.*

(ii) *The space of regular signed Borel measures on K is a real Banach space in $||\cdot||$. Moreover, it is isometrically isomorphic to $M^{\mathbb{R}}(K) \cong C^{\mathbb{R}}(K)^*$.*

2 Weak topologies

Let X be a set and $\mathcal{F}$ a family of functions such that, every $f \in \mathcal{F}$ is a map $X \to Y_f$ where Y_f is a topological space. Then the *weak topology* $\sigma(X, \mathcal{F})$ on X is the smallest topology on X such that every $f \in \mathcal{F}$ is continuous.

Remarks

- The family $\mathcal{S} = \{f^{-1}(U) \colon f \in \mathcal{F}, \text{ open } U \subset Y_f\}$ is a sub-base of $\sigma(X, \mathcal{F})$, i.e. $\sigma(X, \mathcal{F})$ is generated by $\mathcal{S}$. In other words, the family of all finite intersections of elements of $\mathcal{S}$ is a base for $\sigma(X, \mathcal{F})$. Thus, $V \subset X$ is open iff

$$\forall\, x \in V\ \exists\, n \in \mathbb{N}\ \exists\, f_1, \cdots, f_n \in \mathcal{F}\ \exists \text{ open } U_f \subset Y_{f_j} \text{ s.t. } x \in \bigcap_{j=1}^{n} f_j^{-1}(U_j) \subset V$$

  or, equivalently,

$$\forall\, x \in V\ \exists\, n \in \mathbb{N}\ \exists\, f_1, \cdots, f_n \in \mathcal{F}\ \exists \text{ open nhoods } U_j \text{ of } f_j(x) \text{ in } Y_{f_j} \text{ s.t. } \bigcap_{j=1}^{n} f_j^{-1}(U_j) \subset V$$

- If, for every $f \in \mathcal{F}$, $\mathcal{S}_f$ is a sub-base for the topology on Y_f, then

$$\{f^{-1}(U) \colon f \in \mathcal{F}, U \in \mathcal{S}_f\}$$

  is a sub-base for $\sigma(X, \mathcal{F})$.
- If Y_f is Hausdorff for every $f \in \mathcal{F}$ and $\mathcal{F}$ separates points of X (i.e. for distinct $x, y \in X$, there exists $f \in \mathcal{F}$ such that $f(x) \neq f(y)$), then $\sigma(X, \mathcal{F})$ is Hausdorff.
- Let $Y \subset X$ and $\mathcal{F}_Y = \{f|_Y \colon f \in \mathcal{F}\}$. Then $\sigma(X, \mathcal{F})|_Y = \sigma(Y, \mathcal{F}_Y)$
- *Universal property*: Let Z be a topological space and $g \colon Z \to X$ a function. Then g is continuous iff, for every $f \in \mathcal{F}$, $f \circ g \colon Z \to Y_f$ is continuous.
  ($\Longrightarrow$) Trivial
  ($\Longleftarrow$) For every $f \in \mathcal{F}$ and open $U \subset Y_f$, we have that $g^{-1}(f^{-1}(U)) = (f \circ g)^{-1}(U)$ is open in Z. Thus, from the first remark above, we have that, for every $V \in \mathcal{S}$, $g^{-1}(V)$ is open in Z.

Examples

- Let X be a topological space, $Y \subset X$, and $\iota \colon Y \to X$ the inclusion map. Then the weak topology $\sigma(Y, \{\iota\})$ is the subspace topology on Y induced by X.
- Let Γ be an index set. Suppose X_γ is a topological space for each $\gamma \in \Gamma$, and define

$$X = \prod_{\gamma \in \Gamma} X_\gamma = \{x \colon x \text{ is a function with domain } \Gamma \text{ and } x(\gamma) \in X_\gamma\ \forall\, \gamma \in \Gamma\}$$

  For $x \in X$ and $\gamma \in \Gamma$, we write $x_\gamma = x(\gamma)$ and then think of X as the Γ-tuple $(x_\gamma)_{\gamma \in \Gamma}$. For each $\gamma \in \Gamma$, we define $\pi_\gamma \colon X \to X_\gamma$ by $\pi_\gamma(x) = x(\gamma)$ (evaluation at γ or projection onto X_γ). The *product topology* on X is the weak topology $\sigma(X, \{\pi_\gamma \colon \gamma \in \Gamma\})$. Note that $V \subset X$ is open iff

$$\forall\, x \in V\ \exists n \in \mathbb{N}\ \exists \gamma_1, \cdots, \gamma_n \in \Gamma\ \exists \text{ open nhood } U_j \text{ of } x_{\gamma_j} \text{ in } X_{\gamma_j} \text{ s.t. } \{(y_\gamma)_{\gamma \in \Gamma} \colon y_{\gamma_j} \in U_j\} \subset V$$

Proposition 2.1
Let X be a set. Suppose that, for each $n \in \mathbb{N}$, (Y_n, d_n) is a metric space and $f_n \colon X \to Y_n$ be a function. Further assume that $\mathcal{F} = \{f_n \colon n \in \mathbb{N}\}$ separates points of X (i.e. for distinct x, y in X, there exists $n \in \mathbb{N}$ such that $f_n(x) \neq f_n(y)$). Then the weak topology $\sigma(X, \mathcal{F})$ on X is metrisable.

Proof. WLOG, we may assume that $d_n \leq 1$ for all n since we can otherwise replace any such d_n such $\min\{1, d_n\}$. Define

$$d(x, y) := \sum_{n=1}^{\infty} 2^{-n} d_n(f_n(x), f_n(y))$$

It is easy to see that this is a metric on X and that, for every $n \in \mathbb{N}$ and $x, y \in X$,

$$d_n(f_n(x), f_n(y)) \leq 2^n d(x, y)$$

and thus f_n is d-continuous. Hence, $\sigma := \sigma(X, \mathcal{F}) \subset \tau$, where τ is the metric topology generated by d.

Note also that, by the Weierstrass M-test, the map $d \colon (X, \sigma) \times (X, \sigma) \to \mathbb{R}$ is continuous wrt the product topology, so for each $x \in X$ and $\varepsilon > 0$, we have

$$\{y \in X \colon d(x, y) < \varepsilon\} \in \sigma$$

Hence, $\tau \subset \sigma$. ■

Theorem 2.2 Tychonov's theorem
Let (X_γ, τ_γ) be a topological space for each $\gamma \in \Gamma$. Let $X = \prod_{\gamma \in \Gamma} X_\gamma$ and τ the product topology on X. If X_γ is compact for all $\gamma \in \Gamma$, then X is compact.

Proof. Let $\mathcal{F}$ be a non-empty family of closed subsets of X with the *finite intersection property*, i.e. for every $n \in \mathbb{N}$ and every $F_1, \cdots, F_n \in \mathcal{F}$, we have $\cap_{j=1}^n F_j \neq \varnothing$. We would then want to show that $\cap \mathcal{F} \neq \varnothing$.

Let $\mathcal{A}$ be a maximal (wrt inclusion) family of (not necessarily closed) subsets of X with the finite intersection property such that $\mathcal{A} \supset \mathcal{F}$. This can be shown to exist by Zorn's lemma. It then suffices to show that $\cap_{A \in \mathcal{A}} \overline{A} \neq \varnothing$. Indeed, by maximality, we have that

(i) for every $A, B \in \mathcal{A}$, we have $A \cap B \in \mathcal{A}$
(ii) if $A \subset X$ with the property that, for every $B \in \mathcal{A}$, we have $A \cap B \neq \varnothing$, then $A \in \mathcal{A}$

For $\gamma \in \Gamma$, the family $\{\pi_\gamma(A) \colon A \in \mathcal{A}\}$ has the finite intersection property and thus

$$\bigcap_{A \in \mathcal{A}} \overline{\pi_\gamma(A)} \neq \varnothing$$

since X_γ is compact. Then we can pick $x_\gamma \in \cap_{A \in \mathcal{A}} \overline{\pi_\gamma(A)}$ and then set $x = (x_\gamma)_{\gamma \in \Gamma}$.

We now show that $x \in \cup_{A \in \mathcal{A}} \overline{\pi_\gamma(A)}$. To do this, we fix $A \in \mathcal{A}$ and let V be a neighbourhood of x. WLOG, we have $V = \cap_{j=1}^n \pi_{\gamma_j}^{-1}(V_j)$ for some $n \in \mathbb{N}$ and $\gamma_1, \cdots, \gamma_n \in \Gamma$ and neighbourhoods V_j of x_{γ_j} in X_{γ_j}. We have that $V_j \cap \pi_{\gamma_j}(A) \neq \varnothing$ for all $1 \leq \jmath \leq n$ and thus $\pi_{\gamma_j}^{-1}(V_j) \in \mathcal{A}$. By (i), $V \in \mathcal{A}$ and $\mathcal{A}$ has the finite intersection property, so $A \cap V \neq \varnothing$. ■

2.1 Weak topologies on vector spaces

Let E be a real or complex vector space, and let F be a subspace of the space of all linear functionals on E. Suppose F separates points of E, i.e. for every nonzero $x \in E$, there exists

$f \in F$ such that $f(x) \neq 0$. We consider the weak topology $\sigma(E, F)$ on E. Recall that $U \subset E$ is open iff

$$\forall\, x \in U\ \exists\, n \in \mathbb{N}\ \exists\, f_1, \cdots, f_n \in F\ \exists\, \varepsilon > 0 \text{ s.t. } \{y \in E \colon |f_j(y - x)| < \varepsilon, 1 \leq j \leq n\} \subset U$$


For $f \in F$, we define $p_f \colon E \to \mathbb{R}$ by $p_f(x) := |f(x)|$. Let $\mathcal{P} := \{p_f \colon f \in F\}$. Then $(E, \mathcal{P})$ is a locally convex space whose topology is exactly $\sigma(E, F)$. In particular, $\sigma(E, F)$ is Hausdorff, and addition and scalar multiplication are continuous.

Lemma 2.3
Let E be as above. Let $f, g_1, \cdots, g_n$ be linear functionals on E such that

$$\bigcap_{j=1}^{n} \ker g_j \subset \ker f$$

Then $f \in \operatorname{span}\{g_1, \cdots, g_n\}$.

Proof. Define $T \colon E \to \mathbb{R}^n$ (or $\mathbb{C}^n$) by $Tx = (g_j(x))_{j=1}^n$. Then $\ker T = \cap_{j=1}^n \ker g_j \subset \ker f$. Hence, there exists a linear functional $h \colon \operatorname{im} T \to \mathbb{R}$ (or $\mathbb{C}$) such that $f = h \circ T$. Extend h to $\mathbb{R}^n$ to obtain the commutative diagram



Thus, for every $x \in E$, $f(x) = h(Tx) = \sum_{j=1}^n a_j g_j(x)$. ∎

Proposition 2.4
Let E, F be as above. Let f be a linear functional on E. Then f is continuous with respect to $\sigma(E, F)$ iff $f \in F$.

Remark. Equivalently, the proposition tells us that $(E, \sigma(E, F))^* = F$.

Proof. ($\Longleftarrow$) This is by definition of $\sigma(E, F)$.
($\Longrightarrow$) Let $V = \{x \in E \colon |f(x)| \leq 1\}$. This is a neighbourhood of 0 in $\sigma(E, F)$, so there exists $n \in \mathbb{N}$, $g_1, \cdots, g_n \in F$, $\varepsilon > 0$ such that

$$U = \{x \in E \colon |g_j(x)| < \varepsilon, 1 \leq j \leq n\} \subset V$$

If $x \in \cap_{j=1}^n \ker g_j$, then $\lambda x \in U$ for all scalars λ, so $|f(\lambda x)| \leq 1$ for all λ. Thus, $f(x) = 0$. By Lemma 2.3, $f \in \operatorname{span}\{g_1, \cdots, g_n\} \subset F$.

∎

Examples
The following two examples of weak topologies on vector spaces are central objects of study in this chapter. Below, X is a normed space.

- The weak topology on X is $\sigma(X, X^*)$. Note that X^* separates points of X by Hahn–Banach [Theorem 1.4(ii)]. We will sometimes denote by (X, w) the space $(X, \sigma(X, X^*))$ and refer to open sets as weakly open or w-open. Note that $U \subset X$ is w-open iff

$$\forall\, x \in U\ \exists\, n \in \mathbb{N}\ \exists\, f_1, \cdots, f_n \in X^*\ \exists\, \varepsilon > 0 \text{ s.t. } \{y \in X \colon |f_j(y - x)| < \varepsilon, 1 \leq j \leq n\} \subset U$$

- The weak-$*$ topology on X^* is $\sigma(X^*, X)$. Here, we think of X as the subspace of X^{**} under the canonical embedding. We will sometimes denote by $(X^*.w^*)$ the space $(X^*, \sigma(X^*, X))$ and refer to open sets as w^*-open. Note that $U \subset X^*$ is w^*-open iff

$$\forall\, x \in U\ \exists\, n \in \mathbb{N}\ \exists\, x_1, \cdots, x_n \in X\ \exists\, \varepsilon > 0 \text{ s.t. } \{g \in X^* \colon |(g \cdot f)(x_j)| < \varepsilon, 1 \leq j \leq n\} \subset U$$

Remarks

Let us briefly mention some properties of these topologies.

- (X, w) and (X^*, w^*) are locally convex spaces. In particular, they are Hausdorff, and addition and scalar multiplication are continuous.
- $\sigma(X, X^*)$ is contained in the norm topology $\tau_{||\cdot||}$, with equality iff $\dim X < \infty$
- $\sigma(X^*, X) \subset \sigma(X^*, X^{**}) \subset \tau_{||\cdot||}$, with equality in the first inclusion iff X is reflexive and in the second inclusion iff $\dim X < \infty$
- Given a subspace Y of X, we have $\sigma(X, X^*)|_Y = \sigma(Y, Y^*)$. Similarly, $\sigma(X^{**}, X^*)|_X = \sigma(X, X^*)$, so the canonical embedding $X \to X^{**}$ is a w-to-w^* homeomorphism of X onto $\hat{X} \subset X^{**}$, besides being an isometric isomorphism as previously established.

Proposition 2.5

Let X be a normed space. Then

(i) *A linear functional f on X is w-continuous iff $f \in X^*$. In other words, $(X, w)^* = X^*$.*

(ii) *A linear functional φ on X^* is w^*-continuous iff $\varphi \in X$, i.e. there exists $x \in X$ such that $\varphi = \hat{x}$. In other words, $(X^*, w^*)^* = X$.*

Moreover, it follows that $\sigma(X^*, X^{**}) = \sigma(X^*, X)$ iff X is reflexive.

Proof. (i) and (ii) follow from Proposition 2.4. For the last statement, ($\Longleftarrow$) is immediate whereas ($\Longrightarrow$) follows from the fact that

$$X^{**} \overset{(ii)}{=} (X^*, \sigma(X^*, X^{**}))^* = (X^*, \sigma(X^*, X))^* \overset{(i)}{=} X$$

■

Definition Weakly and weak-$*$ bounded

Let X be a normed space.

(i) *We say that a set $A \subset X$ is weakly bounded if $\{f(x) \colon x \in A\}$ is bounded for every $f \in X^*$.*

(ii) *We say that a set $B \subset X^*$ is weak-$*$ bounded if $\{f(x) \colon f \in B\}$ is bounded for every $x \in X$*

Remark. Equivalent definitions:

- $A \subset X$ is weakly bounded iff, for every w-neighbourhoods U of 0 in X, there exists $\lambda > 0$ such that $A \subset \lambda U$.
- $B \subset X$ is weak-$*$ bounded iff, for every w^*-neighbourhoods V of 0 in X^*, there exists $\lambda > 0$ such that $B \subset \lambda V$.

Recall from II Linear Analysis the *principle of uniform boundedness* (PUB): Let X be a Banach space, Y a normed space and $\mathcal{T} \subset \mathcal{B}(X, Y)$. If T is pointwise bounded (i.e., $\forall\, x \in X\ \sup_{T \in \mathcal{T}} ||Tx|| < \infty$), then T is uniformly bounded (i.e., $\sup_{T \in \mathcal{T}} ||T|| < \infty$).

Proposition 2.6

Let X be a normed space.

(i) *If $A \subset X$ is weakly bounded, then A is bounded in the norm topology.*

(ii) *If $B \subset X^*$ is w^*-bounded and X is complete, then B is bounded in the norm topology.*

Proof. For (i), note that $\hat{A} = \{\hat{x} \colon X \in A\} \subset X^{**}$, so if A is w-bounded, then $\hat{A}$ is pointwise bounded. Since X^* is complete, the PUB gives us

$$\sup_{x \in A} ||x|| = \sup_{x \in A} ||\hat{x}|| < \infty$$

To prove (ii), note that, if B is w^*-bounded, then B is pointwise bounded. Since X is complete, the PUB gives us

$$\sup_{f \in B} ||f|| < \infty$$

as required. ∎

Notation. It will be useful to introduce some notation.

- We write $x_n \overset{w}{\to} x$ and say that x_n weakly converges or w-converges to x if $x_n \to x$ in the weak topology. This is equivalent to
$$\langle x_n, f \rangle \to \langle x, f \rangle \qquad \forall\, f \in X^*$$
- We write $f_n \overset{w^*}{\to} f$ and say that f_n weak-$*$ converges or w^*-converges to f if $f_n \to f$ in the weak-$*$ topology. This is equivalent to
$$\langle x, f_n \rangle \to \langle x, f \rangle \qquad \forall\, x \in X$$

Recall from II Linear Analysis the following consequence of the PUB: If X is a Banach space, Y a normed space, and (T_n) a sequence in $\mathcal{B}(X, Y)$ that converges pointwise to a map $T \colon X \to Y$, then $T \in \mathcal{B}(X, Y)$, $\sup_{n \in \mathbb{N}} ||T_n|| < \infty$, and $||T|| \leq \liminf_{n \to \infty} ||T_n||$.

Proposition 2.7
Let X be a normed space.

(i) *If $x_n \overset{w}{\to} x$ in X, then $\sup_n ||x_n|| < \infty$ and $||x|| \leq \liminf_{n \to \infty} ||x_n||$*

(ii) *If $f_n \overset{w^*}{\to} f$ in X^* and X is complete, then $\sup_n ||f_n|| < \infty$ and $||f|| \leq \liminf_{n \to \infty} ||f_n||$.*

Proof. For (i), we have $\hat{x}_n \to \hat{x}$ pointwise in $\mathcal{B}(X^*, \text{scalars})$, so the result follows from above. For (ii), we have $f_n \to f$ pointwise in $\mathcal{B}(X, \text{scalars})$, so the result follows from above. ∎

2.2 Hahn–Banach separation theorems

Let $(X, \mathcal{P})$ be a locally convex space and C be a convex subset of X with $0 \in \operatorname{int} C$. Define

$$\begin{aligned} \mu_C \colon X &\to \mathbb{R} \\ x &\mapsto \inf\{t > 0 \colon x \in tC\} \end{aligned}$$

This is well-defined since the fact that $x/n \to 0$ as $n \to \infty$ implies the existence of some $n \in \mathbb{N}$ such that $x/n \in C$. We call μ_C the *Minkowski functional* or *gauge functional* of C.

Example. If X is a normed space and $C = B_X$, then $\mu_C = ||\cdot||$.

Lemma 2.8
μ_C is positive homogeneous and subadditive. Moreover,

$$\{x \in X \colon \mu_C(x) < 1\} \subset C \subset \{x \in X \colon \mu_C(x) \leq 1\}$$

with equality in the first inclusion if C is open and equality in the second inclusion if C is closed.

Proof. Positive homogeneity: Since $0 \in tC$ for every $t > 0$, we have that $\mu_C(0) = 0$ and thus $\mu_C(0 \cdot x) = 0 \cdot \mu_C(x)$. for any $x \in X$. For $s > 0$, we have $\mu_C(s \cdot x) = s \cdot \mu_C(x)$ since $x \in tC \ \forall \ t > 0 \iff sx \in stC$.

Subadditivity: Observe that $\mu_C(x) < s \iff x \in sC$. Indeed, there exists $t < s$ such that $x \in tC$ and thus

$$\frac{x}{s} = \frac{t}{s}\frac{x}{t} + \left(1 - \frac{t}{s}\right) 0 \in C$$

by convexity of C. Given $x, y \in X$, fix $s \in \mu_C(x)$ and $t > \mu_C(y)$. Then $x \in sC$ and $y \in tC$. By convexity, we have

$$\frac{x+y}{s+t} = \frac{s}{s+t}\frac{x}{s} + \frac{t}{s+t}\frac{y}{t} \in C$$

and thus $\mu_C(x+y) \leq s + t$. Hence, $\mu_C(x+y) \leq \mu_C(x) + \mu_C(y)$.

Inclusions: If $\mu_C(x) < 1$, then $x \in C$ by taking $s = 1$ above. If $x \in C$, then $\mu_C(x) \leq 1$ by definition. If C is open and $x \in C$, then $(1 + \frac{1}{n})x \to x$ so there exists some $n \in \mathbb{N}$ such that $(1 + \frac{1}{n})x \in C$ and thus $\mu_C(x) \leq \frac{1}{1+\frac{1}{n}} < 1$. If C is closed and $\mu_C(x) \leq 1$, then

$$\mu_C\left(\left(1 - \frac{1}{n}\right)x\right) = \left(1 - \frac{1}{n}\right)\mu_C(x) < 1$$

and thus $(1 - \frac{1}{n})x \in C$. Since $(1 - \frac{1}{n})x \to x$ and C is closed, it follows that $x \in C$. ∎

Remark. If C is balanced (i.e. $x \in C, |\lambda| = 1 \implies \lambda x \in C$), then μ_C is a seminorm. If, in addition, C is bounded (i.e. for every neighbourhood U of 0, there exists $\lambda > 0$ such that $C \subset \lambda U$), then μ_C is a norm.

Theorem 2.9 Hahn–Banach separation theorem I
Let $(X, \mathcal{P})$ be a locally convex space and C an open convex subset of X with $0 \in C$. For $x_0 \in X \backslash C$, there exists $f \in X^*$ such that $f(x) < f(x_0)$ for every $x \in C$ (if the scalar field is $\mathbb{R}$) or $\operatorname{Re} f(x) < \operatorname{Re} f(x_0)$ for every $x \in C$ (if the scalar field is $\mathbb{C}$).

Proof. We shall first prove the real case and then extend it to the complex case.

Case 1: Real case

By Lemma 2.8, $C = \{x \in X : \mu_C(x) < 1\}$, so $\mu_C(x_0) \geq 1$. Let $Y = \operatorname{span}\{x_0\}$ and define $g : Y \to \mathbb{R}$ by $g(\lambda x_0) = \lambda \mu_C(x_0)$. Note that

$$g(\lambda x_0) \begin{cases} = \mu_C(\lambda x_0), & \lambda \geq 0 \\ \leq 0 < \mu_C(\lambda x_0), & \lambda < 0 \end{cases}$$

so μ_C dominates g on Y in the sense of Theorem 1.1. Thus, there exists a linear map $f : X \to \mathbb{R}$ such that $f|_Y = g$ and, for every $x \in X$, $f(x) \leq \mu_C(x)$. it then follows that, for every $x \in C$,

$$f(x) \leq \mu_C(x) < 1 \leq \mu_C(x_0) = g(x_0) = f(x_0)$$

Moreover, on the open neighbourhood $\varepsilon C \cap (-\varepsilon C)$ of 0, $|f| < \varepsilon$ for any $\varepsilon > 0$, so f is continuous at 0. Hence, $f \in X^*$ by Lemma 1.9.

Case 2: Complex case

By the real case, there exists $g \in (X_{\mathbb{R}})^*$ such that, for every $x \in C$, $g(x) < g(x_0)$. Then $f(x) := g(x) - ig(ix)$ defines $f \in X^*$ such that $\operatorname{Re} f = g$. ∎

Remark. From now on, we only treat the real case; the complex case follows by the above argument.

Theorem 2.10 Hahn–Banach separation theorem II
Let $(X, \mathcal{P})$ be a locally convex space and let A, B be non-empty, disjoint, convex sets in X.

(i) *If A is open and B is closed, then there exists $f \in X^*$ and $\alpha \in \mathbb{R}$ such that $f(a) \leq \alpha \leq f(b)$ for all $a \in A$ and $b \in B$*

(ii) *If A is compact and B is closed, then there exists $f \in X^*$ such that $\sup_A f < inf_B f$.*

Proof. (i): Fix $a_0 \in A$ and $b_0 \in B$. Let $C = A - B + b_0 - a_0$ and $x_0 = b_0 - a_0$. Then C is convex and open as $C = \cup_{b \in B}(A - b + b_0 - a_0)$ and $0 \in C, x_0 \notin C$ as $A \cap B = \varnothing$. By Theorem 2.9, there exists $f \in X^*$ such that, for every $x \in C$, $f(x) < f(x_0)$. Then $f(a) < f(b)$ for every $a \in A$ and $b \in B$. In particular, $f \neq 0$. Set $\alpha = \inf_B f$. Then $f(a) \leq \alpha \leq f(b)$ for all $a \in A$ and $b \in B$. To upgrade $\leq$ to $<$, we fix $u \in X$ such that $f(u) > 0$. For $a \in A$, we have $a + u/n \to a$ as $n \to \infty$. Since A is open, $a + u/n \in A$ for some $n \in \mathbb{N}$ and thus $f(a) < f(a + u/n) \leq \alpha$.

(ii): For each $x \in A$, pick a neighbourhood W_x of 0 in X such that $(x + W_x) \cap B = \varnothing$. Next, choose an open neighbourhood V_x of 0 such that $V_x + V_x \subset W_x$. WLOG, we may take V_x to be convex and symmetric. Observe that $\{x + V_x \colon x \in A\}$ form an open cover for A, so we can pick $x_1, \cdots, x_n \in A$ such that

$$A \subset \bigcup_{k=1}^{n} (x_k + V_{x_k})$$

Set $U = \cap_{k=1}^n V_{x_k}$. Then U is an open convex neighbourhood of 0, so $A + U$ is open and convex. We shall now show that $(A + U) \cap B = \varnothing$: given $a \in A, u \in U$, there exists k such that

$$a + u \in x_k + V_{x_k} + V_{x_k} \subset x_k + W_{x_k}$$

which is disjoint from B. By (i), there exist $f \in X^*$ and $\alpha \in \mathbb{R}$ such that, for every $a \in A, u \in U, b \in B$, $f(a + u) < \alpha \leq f(b)$. In particular, we have

$$\sup_A f = \max_A f < \alpha \leq \inf_B f$$

as required. ∎

Theorem 2.11 Mazur's theorem
Let C be a convex set in a normed space X. Then $\overline{C}^{||\cdot||} = \overline{C}^w$. In particular, C is $||\cdot||$-closed iff C is w-closed.

Proof. We have that $\overline{C}^{||\cdot||} \subset \overline{C}^w$ since norm topology is finer than the weak topology. To prove $\supset$, we fix $x \in X \backslash \overline{C}^{||\cdot||}$. By Theorem 2.10(ii) applied to X, $A = \{x\}$ and $B = \overline{C}^{||\cdot||}$, there exists $f \in X^*$ such that $f(x) < \inf_B f$. Then $\{y \in X \colon f(y) < \inf_B f\}$ is a w-neighbourhood of x that is disjoint from B. Thus, $X \backslash \overline{C}^{||\cdot||}$ is w-open and thus $\overline{C}^{||\cdot||}$ is w-closed. Hence, $\overline{C}^{||\cdot||} = \overline{C}^w$. ∎

Corollary 2.12
Suppose $x_n \xrightarrow{w} 0$ in some normed space X. Then for every $\varepsilon > 0$, there exists $x \in \text{conv}\{x_n \colon n \in \mathbb{N}\}$ such that $||x|| < \varepsilon$.

Proof. $0 \in \overline{\{x_n \colon n \in \mathbb{N}\}}^w \subset \overline{\text{conv}\{x_n \colon n \in \mathbb{N}\}}^w = \overline{\text{conv}\{x_n \colon n \in \mathbb{N}\}}^{||\cdot||}$. ∎

Remark. It follows that there exist $p_1 < q_1 < p_2 < q_2 < \cdots$ and convex combinations $\sum_{k=p_n}^{q_n} t_k x_k$ that converge to 0 as $n \to \infty$.

Theorem 2.13 Banach–Alaoglu
For any normed space X, (B_{X^*}, w^*) is compact.

Proof. For $x \in X$, let $K_x = \{\lambda \colon \lambda \text{ scalar}, |\lambda| \leq |x|\}$. Let

$$K = \prod_{x \in X} K_x = \{f \colon X \to \{\text{scalars}\} \mid \forall\, x \in X, |f(x)| \leq ||x||\}$$

so $B_{X^*} \subset K$. The product topology on K is compact by Theorem 2.2 and is given by the evaluation maps $\varepsilon_x \colon K \to K_x, f \mapsto f(x)$. Note that $\varepsilon_x|_{B_{X^*}} = \hat{x}|_{B_{X^*}}$, so the subspace topology on B_{X^*} is the weak-$*$ topology. Note also that

$$\begin{aligned} B_{X^*} &= \{f \in K \colon f(\lambda x + \mu y) = \lambda f(x) + \mu f(y)\ \forall\, x, y \in X\ \forall \text{scalars} \lambda, \mu\} \\ &= \bigcap_{\substack{x,y \in X \\ \lambda,\mu \text{scalars}}} \{f \in K \colon (\varepsilon_{\lambda x + \mu y} - \lambda \varepsilon_x - \mu \varepsilon_y)(f) = 0\} \end{aligned}$$

is closed in K by continuity of the ε_x. Hence, it follows that (B_{X^*}, w^*) is compact. ∎

Proposition 2.14
Let X be a normed space and K a compact Hausdorff space. Then
(i) *X is separable iff (B_{X^*}, w^*) is metrisable.*
(ii) *$C(K)$ is separable iff K is metrisable.*

Proof.
$\overset{\text{(i)}}{\Longrightarrow}$ Pick a countable dense subset $\{x_n \colon n \in \mathbb{N}\} \subset X$. Let $\mathcal{F} = \{\hat{x}_n|_{B_{X^*}} \colon n \in \mathbb{N}\}$, which is countable and, by density, separates the points of B_{X^*}. By Proposition 2.1, the weak topology $\sigma(B_{X^*}, \mathcal{F})$ is metrisable. Moreover, $\sigma(B_{X^*}, \mathcal{F})$ is weaker than the weak-$*$ topology on B_{X^*}, which is $\sigma(B_{X^*}, \{\hat{x}|_{B_{X^*}} \colon x \in X\})$. Thus, $\mathrm{id} \colon (B_{X^*}, w^*) \to (B_{X^*}, \sigma(B_{X^*}, \mathcal{F}))$ is a continuous bijection from a compact space (by Theorem 2.13) to a Hausdorff space. Thus, id is a homeomorphism and the two topologies coincide. Hence, the weak-$*$ topology on B_{X^*} is metrisable.

$\overset{\text{(ii)}}{\Longrightarrow}$ By $\overset{\text{(i)}}{\Longrightarrow}$, $(B_{C(K)^*}, w^*)$ is metrisable. We then define $\delta \colon K \to (B_{C(K)^*}, w^*), k \mapsto \delta_k$, where $\delta_k(f) = f(k)$. Note that δ is well-defined since δ_k is linear and $|\delta_k(f)| \leq ||f||_\infty$. Moreover, δ is continuous by the universal property: for every $k \in K$,

$$(\hat{f} \circ \delta)(k) = \hat{f}(\delta_k) = \delta_k(f) = f(k)$$

Furthermore, δ is surjective: given distinct $k, l \in K$, Urysohn's lemma gives us $f \in C(K)$ such that $f(k) \neq f(l)$, and thus $\delta_k \neq \delta_l$ (different when evaluated at f). Putting everything together, $\delta \colon K \to \operatorname{im} \delta$ is a continuous bijection from a compact space to a Hausdorff space, so K is homeomorphic to a subset of $(B_{C(K)^*}, w^*)$ and hence K is metrisable.

$\overset{\text{(ii)}}{\Longleftarrow}$ Let d be a metric on K metrising K. As a compact metric space, (K, d) is separable. We then pick a countable dense subset $\{k_n \colon n \in \mathbb{N}\} \subset K$ and define $f_n \colon K \to \mathbb{R}, x \mapsto d(x, k_n)$. Let A be the algebra generated by $\{f_n \colon n \in \mathbb{N}\} \cup \{\mathbb{1}_K\}$. Note that A is a separable sublagebra of $C(K)$ that separates points of K (by density of $\{k_n \colon n \in \mathbb{N}\}$ and contains $\mathbb{1}_K$ (and, in the complex case, is closed under complex conjugation). By Stone–Weierstrass, $\overline{A} = C(K)$ and thus $C(K)$ is separable.

$\overset{\text{(i)}}{\Longleftarrow}$ Let $K = (B_{X^*}, w^*)$. By $\overset{\text{(ii)}}{\Longleftarrow}$, $C(K)$ is separable. Define $T \colon X \to C(K), x \mapsto \hat{x}|_{B_{X^*}}$. Note that T is well-defined, linear, and satisfies $||Tx|| = ||\hat{x}|_{B_{X^*}}||_\infty = ||\hat{x}|| = ||x||$. Thus, X isometrically embeds into $C(K)$. Hence, X is separable. ∎

Remarks

- If X is separable, then (B_{X^*}, w^*) is a compact metric space, so B_{X^*} is w^*-sequentially compact.
- If X is separable, then $X^* = \bigcup_{n\in\mathbb{N}} nB_{X^*}$ is w^*-separable. By Mazur, X is separable iff X is w-separable. Thus, X is w-separable implies that X^* is w^*-separable. However, the converse is false in general, e.g. $X = \ell_\infty$ (cf. Goldstein's theorem).
- We proved that K is a topological subspace of $(B_{C(K)^*}, w^*)$.
- Any normed space X isometrically embeds into $C(K)$ for some compact Hausdorff K, e.g. $K = (B_{X^*}, w^*)$.

Proposition 2.15
Let X be a normed space. Then (B_X, w) is metrisable iff X^* is separable.

Proof. ($\Longleftarrow$) By Proposition 2.14, $(B_{X^{**}}, w^*)$ is metrisable. Recall that $(B_{X^{**}}, w^*)|_{B_X} = (B_X, w)$ so (B_X, w) is metrisable.

($\Longrightarrow$) Let d be a metric on B_X metrising the weak topology on B_X. For $n \in \mathbb{N}$, $\{x \in B_X : d(x, 0) < \frac{1}{n}\}$ is a neighbourhood of 0 and hence contains a set $U_n = \{x \in B_X : \forall f \in F_n, |f(x)| < \varepsilon_n\}$, where $F_n \subset X^*$, F_n finite and $\varepsilon_n > 0$. For any weak neighbourhood V of 0 in B_X, there exists $n \in \mathbb{N}$ such that $U_n \subset V$. Let $Z = \overline{\operatorname{span}} \bigcup_{n\in\mathbb{N}} F_n$. We then want to show that $Z = X^*$.

Indeed, fix $f \in X^*$ and $\varepsilon > 0$. Consider the w-neighbourhood $V = \{x \in B_X : |f(x)| < \varepsilon\}$ of 0. Then there exists $n \in \mathbb{N}$ such that $U_n \subset V$. Let $Y = \bigcap_{g\in F_n} \ker g$. Then $||f|_Y|| \leq \varepsilon$. By Hahn–Banach, there exists $h \in X^*$ such that $h|_Y = f|_Y$ and $||h|| = ||f|_Y|| \leq \varepsilon$. Then $f - h|_Y = 0$, so by Lemma 2.3, $f - h \in \operatorname{span} F_n \subset Z$. Thus, $d(f, Z) \leq \varepsilon$. Hence $f \in Z$. ∎

Theorem 2.16 Goldstein's theorem
Let X be a normed space. Then $\overline{B_X}^{w^*} = B_{X^{**}}$, i.e. the closure of B_X in $X^*{*}$ is $B_{X^{**}}$.

Proof. Note that $B_X \subset B_{X^{**}}$ and $B_{X^{**}}$ is w^*-closed, so $\overline{B_X}^{w^*} \subset B_{X^{**}}$. To prove $\supset$, we fix $\varphi \in X^{**} \backslash \overline{B_X}^{w^*}$. It suffices to show that $||\varphi|| > 1$. Applying Theorem 2.10 to the locally convex space (X^{**}, w^*) to $A = \{\varphi\}$ and $B = \overline{B_X}^{w^*}$ to obtain $f \in X^*$ with $|\varphi(f)| > \sup_{\psi\in B} |\psi(f)| \geq \sup_{x\in B_X} |f(x)| = ||f||$. Thus, $||\varphi|| > 1$. ∎

Theorem 2.17
Let X be a Banach space. The following are equivalent:

(i) *X is reflexive.*
(ii) *(B_X, w) is compact.*
(iii) *X^* is reflexive.*

Proof. (i) $\Longrightarrow$ (ii): Since X is reflexive, $(B_X, w) = (B_{X^{**}}, w^*)$, which is compact by Banach–Alaoglu.
(ii) $\Longrightarrow$ (i): $(B_{X^{**}}, w^*)|_{B_X} = (B_X, w)$ is compact, so B_X is w^*-closed in $X^*{*}$. By Goldstein, $B_{X^{**}} = \overline{B_X}^{w^*} = B_X$.
(i) $\Longrightarrow$ (iii): Note that $(B_{X^*}, w) = (B_{X^*}, w^*)$ is compact by Banach–Alaoglu. By (ii) $\Longrightarrow$ (i), X^* is reflexive.
(iii) $\Longrightarrow$ (i): By (i) $\Longrightarrow$ (iii), we have that X^{**} is reflexive. Then by (i) $\Longrightarrow$ (ii), we have that $B_{X^{**}}, w)$ is compact. Since X is complete, B_X is closed and thus w-closed (by Mazur) in X^{**}. Thus, B_X is w-compact, so by (ii) $\Longrightarrow$ (i), X is reflexive. ∎

Remark. If X is a separable reflexive space, then (B_X, w) is a compact metric space.

Lemma 2.18
Let K be a non-empty compact metric space. Then there exists a continuous surjective map $\varphi\colon \Delta \to K$, where $\Delta = \{0,1\}^{\mathbb{N}}$ equipped with the product topology.

Remark. Note that Δ is homeomorphic to the Cantor set $(\varepsilon_i) \mapsto \sum_i (2\varepsilon_i)3^{-i}$.

Proof. ■