

Geometric Foundations of SU(3): Deriving Gauge Structure, Spacetime Dimension, and Spatial Extension from a Single Polyhedron

Robert Massman¹

¹*Rochester Institute of Technology**

(Dated: March 30, 2026)

We present a self-contained derivation of the geometric foundations underlying Chiral Geometrogenesis (CG), a framework in which gauge symmetry, spacetime dimension, and spatial structure emerge from a single polyhedral object. Starting from three irreducible inputs—observer existence, the requirement that the pre-geometric substrate have finite information content, and the restriction to compact simple gauge groups—we establish a chain of theorems: (1) Spacetime dimension $D = 4$ is uniquely compatible with stable bound-state observers. (2) The gauge group SU(3) is derived (not selected) via two complementary paths—topological ($\mathbb{Z}_3$ center from stella geometry plus rank constraint) and information-theoretic (given quantum interference structure, Fisher metric non-degeneracy eliminates $N \leq 2$; a parsimony criterion then selects $N = 3$ as the smallest stable prime)—and verified by categorical consistency (Tannaka reconstruction from polyhedral data). (3) The stella octangula—the compound of two interpenetrating tetrahedra—is the unique minimal geometric realization of SU(3) among all topological spaces satisfying weight correspondence, Weyl symmetry, and charge conjugation conditions. (4) The Euclidean metric on $\mathbb{R}^3$ emerges from the SU(3) Killing form. (5) Extended 3D space is derived as the unique FCC lattice from SU(3) representation theory, with a continuum limit recovering standard Yang-Mills theory. (6) The Standard Model gauge group $SU(3)_C \times SU(2)_L \times U(1)_Y$ is the minimal phenomenologically viable completion of the D_4 root system geometrically encoded by the stella’s polytope embedding chain via $D_4 \rightarrow \mathfrak{so}(10) \rightarrow \mathfrak{su}(5)$.

The framework rests on 8 independent inputs (3 irreducible, 5 supporting/redundant) and is subject to 5 explicit falsification conditions stated in Section I. This paper establishes the kinematic geometric foundations; the temporal dimension ($t = 1$, derived from phase evolution in Paper 2) is the sole forward dependency, while all spatial results are self-contained. Dynamical consequences (mass generation, confinement mechanism, gravity emergence) are deferred to Paper 2.

I. INTRODUCTION

A. Falsification Conditions

The following framework derives spacetime dimension, gauge symmetry, and spatial structure from a polyhedral encoding of SU(3). We state its falsification conditions first, then develop the theory. The geometric foundations of Paper 1 are refuted if *any* of the following is demonstrated:

- FC1:** $D \neq 4$. If spatiotemporal dimension differs from 4—e.g., large extra dimensions detected at colliders or in gravitational experiments—then Theorem III.2 fails. (Note: 4 independent dynamical mechanisms also select $D = 4$; all 5 must simultaneously fail.)
- FC2:** SU(3) is not the confining gauge group. If the strong interaction exhibits a different gauge structure, both derivation paths (topological, information-theoretic) and the categorical consistency check are refuted simultaneously.
- FC3:** Another polyhedron satisfies GR1–GR3. If a competing polyhedral or topological structure is

found that satisfies the geometric realization conditions with fewer or equal vertices, Theorem V.10 falls.

FC4: FCC is not the unique SU(3)-compatible lattice. If an alternative vertex-transitive tiling satisfying the four constraints derived in Theorem VI.3 (12-regularity, triangle prohibition, 4-squares-per-edge, O_h symmetry) is demonstrated, Theorem VI.4 is refuted.

FC5: The continuum limit does not recover SU(3) Yang-Mills. If the discrete-to-continuous procedure fails to reproduce standard gauge theory with the correct topological properties ($\pi_3 = \mathbb{Z}$, $\mathbb{Z}_3$ center), the bridge from geometry to physics is broken.

These conditions are sharp: each is a mathematical statement admitting definitive proof or disproof. We return to them in Section X.

B. Problem Statement

The Standard Model of particle physics requires approximately 30 free parameters, multiple postulated symmetries, and leaves fundamental questions unanswered—the flavor puzzle, the Strong CP problem, the origin of gravity, and the matter-antimatter asymmetry. This paper asks whether a subset of these structures can be *derived* rather than postulated.

* robert@robertmassman.com

Specifically, we pursue the question: *given that observers exist, what geometric structures are forced?* The answer, developed through Sections II–VII, is that space-time dimension, gauge symmetry, the specific polyhedron encoding that symmetry, spatial structure, and the Standard Model gauge group all follow from a small set of transparent assumptions.

a. The core narrative. The logic of this paper follows a single thread. Observer existence forces $D = 4$ (Section III). In four dimensions, the requirement that gauge structure be geometrically realized—encoded in a discrete object rather than presupposed on a smooth manifold—selects $SU(3)$ as the unique compact simple gauge group (Section IV). Once $SU(3)$ is fixed, the question becomes: *what geometric structure encodes it?* The answer is the stella octangula—two interpenetrating tetrahedra whose vertices biject with $SU(3)$ weights, whose symmetry group contains the Weyl group, and whose point-reflection symmetry realizes charge conjugation (Section V). No other polyhedron works. The Euclidean metric on $\mathbb{R}^3$ then emerges from the $SU(3)$ Killing form rather than being assumed. Tiling space with stellae produces the FCC lattice as the unique structure preserving color-phase coherence (Section VI), and the polytope embedding chain extending the stella encodes the D_4 root system from which the full Standard Model gauge group follows (Section VIII). The entire derivation is a sequence of *forced moves*: at each step, alternatives are eliminated by theorem rather than excluded by fiat.

b. A clarification on ontological status. The stella octangula is not a physical object embedded in a pre-existing space—no such space yet exists at this stage of the derivation. It is the unique pattern of algebraic constraints forced by gauge symmetry, charge conjugation, and confinement. Space, fields, and dynamics emerge from this pattern (Sections IX–X and Paper 2); the stella is prior to all of them.

C. Independent Inputs

The framework rests on 8 independent inputs. Of these, 3 are irreducible and the remaining 5 are derivable or redundant. Crucially, the geometric realization postulate (F1) is not assumed but *derived* from a more primitive principle: the pre-geometric substrate must be specifiable with finite information (Axiom FI).

a. Deriving F1 from finite information. Axiom FI states that the pre-geometric substrate $\mathcal{S}$ is specifiable by a finite quantity of information (finite Kolmogorov complexity). This is justified by the Bekenstein bound [1], the holographic principle [2], and the requirement that foundational structures be constructively definable [3]. Combined with gauge invariance (A1), CPT symmetry (A2), confinement (A3), and representation faithfulness (A4), Axiom FI forces the substrate to be a finite polyhedral complex satisfying (GR1)–(GR3) — which is precisely

TABLE I: Complete independent input structure for Paper 1. Inputs marked “Core” are irreducible; “Derived” inputs follow from core inputs via the theorems indicated.

#	Input	Class	Core?
I1	Observer existence $\Rightarrow D=4$	Anthropic	✓
FI	Finite information content of substrate	Info-theoretic	✓
F5	Compact simple gauge group	Framework	✓
F1	Gauge group geom. realized	Derived (FI)	
I3	Fisher metric on config. space (Sec. IV B)	Framework	
F2	$\mathbf{3} \oplus \mathbf{\bar{3}}$ rep. (Sec. V A)	Derived (F3)	
F3	Chirality geom. encoded	Derived ($SU(3)$)	
F4	Minimal vertex count	Redundant	
F6	Vertex-transitivity for tiling (Sec. VI)	Redundant	

the content of F1. The derivation chain is:

$$\begin{aligned} \text{FI} &\xrightarrow{\text{finite}} \text{discrete } \mathcal{S} \xrightarrow{A1, A4} \text{labeled vertices} \\ &\xrightarrow{\text{roots}} \mathcal{P} \xrightarrow{A1-A4} \text{GR1-GR3} \end{aligned} \quad (1)$$

(see Theorem 0.0.0b and Theorem 0.0.0c in the supplementary proof chain). F2 is derivable from F3 + CPT (Proposition 0.0.0h). F4 is a parsimony principle whose justification is *a posteriori*. F6 is required for $SU(3)$ phase coherence (non-vertex-transitive tilings fail the color singlet condition). The core set $\{I1, FI, F5\}$ is irreducible: removing any one collapses the derivation chain.

The framework’s primary strength is *explanatory economy*: from 8 transparent inputs (3 irreducible, of which the most specific — FI — asserts only that the substrate has finite information content), it derives spacetime dimension, gauge group, spatial metric, lattice structure, and field content. GR1–GR3 + MIN1–MIN3 collectively function as a selection device for the stella octangula; alternative axiom sets (e.g., adjoint representation content, non-minimal choices, simplicial constraints) could in principle select different geometric objects. The axiom package’s role as a definition space is made explicit throughout.

D. Derivation Chain

The logical structure forms a directed acyclic graph (Figure 1 shows the full dependency DAG). The *main*




FIG. 1: Input dependency graph for Paper 1. Blue boxes: irreducible inputs. Green: derived intermediate results. Orange: final geometric structures. Each arrow is a theorem proven in this paper.

thread reads:

$$\begin{aligned}
 & \text{FI} \xrightarrow{\text{Thm 0.0.0b}} \text{F1 (GR1-GR3)} \\
 & \text{Observers} \xrightarrow{\text{Thm III.2}} D=4 \\
 & D=4 + \text{F1} \xrightarrow{\text{Thm IV.3}} \text{SU}(3) \\
 & D=4 + \text{SU}(3) \xrightarrow{\text{Thm V.6}} \mathbb{R}^3 \\
 & \mathbb{R}^3 + \text{SU}(3) \xrightarrow{\text{Thm V.10}} \text{Stella} \\
 & \text{Stella} \xrightarrow{\text{Thm VI.4}} \text{FCC lattice} \\
 & \text{Stella} \xrightarrow{\text{Thm VIII.1}} D_4 \xrightarrow{\text{min.+pheno.}} G_{\text{SM}}
 \end{aligned} \tag{2}$$

Steps through D_4 are geometrically necessary and require no dynamical content from Papers 2–6; the final completion to the full SM gauge group invokes minimality and Standard Model compatibility as physical input (Remark VIII.4).

E. Roadmap

Section II establishes the geometric realization framework and $\text{SU}(3)$ weight system. Section III derives $D = 4$ from observer stability. Section IV presents two complementary derivations of $\text{SU}(3)$ (sharing the $D = 4$ dimensional input) and a categorical consistency check. Section V proves the stella octangula’s uniqueness. Section VI derives extended 3D space as the FCC lattice. Section VII establishes three continuum limits. Section VIII traces the polytope embedding chain from the stella to the D_4 root system and its minimal completion to the SM gauge group. Section IX derives color field existence from distinguishability and establishes the pressure function framework, bridging to the dynamics of Paper 2. Section X summarizes, addresses limitations, and revisits falsification.

II. DEFINITIONS AND FRAMEWORK

a. Conventions. Throughout this paper we adopt the following conventions:

- **Spacetime signature:** Mostly-plus (“East Coast”) convention $\eta_{\mu\nu} = \text{diag}(-1, +1, +1, +1)$.
- **Weight space metric:** The Killing form on the $\text{SU}(3)$ Cartan subalgebra induces a Euclidean metric on the 2D weight space, extending to $ds^2 = dx_1^2 + dx_2^2 + dx_3^2$ in the 3D stella embedding (Theorem V.6).
- **Natural units:** $\hbar = c = 1$ unless explicitly restored.
- **Index conventions:** Greek indices μ, ν, ρ, σ run over spacetime 0, 1, 2, 3; Latin i, j, k over spatial 1, 2, 3; uppercase A, B, C denote color indices.

A. Minimal Geometric Realization

a. Why polyhedral realization? We offer four motivations for seeking a polyhedral realization of gauge symmetry:

(i) *Discreteness from confinement.* QCD confines color charge into discrete hadrons. A polyhedron naturally encodes this: vertices represent localized charges, edges represent connections.

(ii) *Minimal encoding.* The weight diagram of $\text{SU}(3)$ is a discrete set of points. A polyhedral realization asks: what is the minimal geometric structure that encodes these relations while preserving algebraic symmetries?

(iii) *CPT as geometry.* In polyhedral realization, charge conjugation is geometric opposition—the point reflection exchanging matter tetrahedron T_+ with antimatter tetrahedron T_- .

(iv) *Pre-geometric coordinates.* Fiber bundles presuppose the manifold structure they cannot derive. For emergent spacetime, gauge structure must be encoded discretely.

Definition II.1 (Geometric Realization). *A geometric realization of a Lie group G is a polyhedral complex $\mathcal{P}$ embedded in $\mathbb{R}^n$ satisfying:*

(GR1) **Weight Correspondence:** *Vertices of $\mathcal{P}$ are in bijection with weights of the fundamental representation.*

(GR2) **Symmetry Preservation:** *The automorphism group $\text{Aut}(\mathcal{P})$ contains a subgroup isomorphic to the Weyl group $W(G)$.*

(GR3) **Conjugation Compatibility:** *Charge conjugation is encoded as a geometric involution.*

Remark II.2 (Combinatorial, Not Metric, Embedding). *The phrase “embedded in $\mathbb{R}^n$ ” in Definition II.1 refers to the combinatorial cell complex structure (vertices, edges,*

faces and their incidence relations), not to a metric space. No distances, angles, or inner products are assumed at this stage— $\mathbb{R}^n$ serves only as topological scaffolding supplying a sufficient number of coordinates to realize the incidence relations. The Euclidean metric on physical 3-space is derived from the $SU(3)$ Killing form in Theorem V.6; it is not presupposed here.

b. Necessity of these conditions. The conditions (GR1)–(GR3) follow from physical requirements:

(GR1): Informational minimality: For any discrete encoding to be faithful, its vertices must biject with weights—the eigenvalues of the Cartan generators that uniquely identify each color state.

(GR2): Gauge invariance: The Weyl group $W(SU(3)) \cong S_3$ permutes color charges; any faithful encoding must realize these permutations as automorphisms.

(GR3): CPT symmetry: The Lüders-Pauli theorem requires CPT invariance. For $SU(3)$, charge conjugation maps $\mathbf{3} \rightarrow \bar{\mathbf{3}}$, sending each weight to its negative.

Remark II.3 (GR1–GR3 as Necessary Conditions). *The derivation hierarchy:*

Layer	Content
1. Irreducible Observers exist ($\Rightarrow D = 4$, Thm. III.2)	
2. Physical	A1: Gauge invariance; A2: CPT symmetry; A3: Confinement; A4: Faithfulness
3. Derived	GR1 $\leftarrow A1+A4$; GR2 $\leftarrow A1+GR1$; GR3 $\leftarrow A1+A2+GR1$
4. Theorem	Stella uniqueness (Thm. V.10)

c. Necessity for emergent spacetime. Four requirements distinguish polyhedral encoding from alternatives: (a) No presupposed spacetime; (b) discrete charge classification via $\mathbb{Z}_3$ center; (c) pre-geometric integer coordinates; (d) combinatorial phase coherence.

TABLE II: Framework comparison for emergent spacetime requirements.

Framework	(a)	(b)	(c)	(d)
Fiber bundle	✗	✓	✗	✗
Lattice gauge	✗	✓	✓	✓
Spin foam	✓	✗	✓	✓
Causal set	✓	✗	✓	✗
Polyhedral	✓	✓	✓	✓

Remark II.4 (Complementarity with Standard Gauge Theory). *The polyhedral framework does not replace fiber bundle descriptions—it precedes them. The QCD gauge bundle remains the correct description of gluon dynamics on an already-existing Minkowski space; what fiber bundles*

cannot do is serve as the substrate from which spacetime emerges. The two frameworks address different questions: polyhedral encoding answers “from what does spacetime emerge?” while fiber bundles answer “how do gauge fields behave on spacetime?”

Theorem II.5 (Polyhedral Necessity). *Within the geometric realization framework (GR1–GR3), polyhedral (discrete) encoding is necessary for emergent spacetime. Specifically, the four alternative frameworks—fiber bundles, smooth manifolds, category-theoretic abstractions, and continuous Lie group actions on $\mathbb{R}^n$ —each fail at least one of the requirements (a)–(d) stated above. Alternative pre-geometric programs (simplicial complexes, causal sets) satisfy subsets of (a)–(d) and represent viable research directions with different foundational choices; polyhedral necessity is a consequence of the specific requirements adopted here, not a universal physical prohibition.*

Proof. We show each alternative fails at least one emergent-spacetime requirement.

(i) *Fiber bundles presuppose their base.* A principal G -bundle $P \xrightarrow{\pi} M$ requires a base manifold M as input data [4]. If spacetime is M , then M cannot emerge from P —requirement (a) fails.

(ii) *Smooth manifolds lack discrete charge classification.* A smooth manifold $\mathcal{M}$ carries continuous coordinates. The $\mathbb{Z}_3$ center of $SU(3)$ —which classifies color charges into exactly three triality sectors—has no natural encoding on a smooth manifold without additional discrete structure. Requirement (b) fails.

(iii) *Category-theoretic abstractions lack spatial content.* A monoidal category $\mathcal{C}$ with objects $\text{Rep}(G)$ encodes representation-theoretic data but contains no spatial information: no distances, no coordinates, no notion of “neighbor.” To extract spatial structure requires a geometric realization functor $F : \mathcal{C} \rightarrow \text{Top}$ —which outputs a discrete topological space (i.e., a polyhedral complex). Requirement (c) fails without such a functor; with it, one has returned to polyhedral encoding.

(iv) *Continuous group actions presuppose the continuum.* An $SU(3)$ action on $\mathbb{R}^n$ uses the continuous manifold structure of both the group and the ambient space, presupposing the metric geometry that is to be derived. Requirement (d) fails: phase coherence between neighboring cells requires shared combinatorial boundaries (faces), not a background metric.

Each alternative fails ≥ 1 requirement; polyhedral encoding satisfies all four (Table II). The requirements (a)–(d) are framework-specific choices motivated by the geometric realization program, not universal physical necessities. $\square$

Remark II.6 (Finite Information Foundation). *The requirements (a)–(d) that motivate polyhedral necessity themselves follow from a single, more primitive principle: the pre-geometric substrate must be specifiable with finite information (Axiom FI). A smooth manifold requires infinite information (uncountably many real coordinates) to specify a single point; a finite polyhedral complex is*

fully determined by a finite vertex set, edge set, and face set. This observation, combined with gauge invariance and CPT symmetry, yields the geometric realization conditions (GR1)–(GR3) as theorems rather than axioms (Theorem 0.0.0b). Empirical support comes from lattice QCD, which encodes SU(3) on a discrete structure and successfully reproduces hadron masses, the confinement transition, and the glueball spectrum from first principles [5]—demonstrating that discrete gauge encoding is not merely a regularization convenience but captures the full non-perturbative content of the theory.

Remark II.7 (Smooth Manifold Realizations of SU(3)). The flag manifold $SU(3)/T^2$ carries a transitive SU(3) action with 6 fixed points matching the non-zero weights of $\mathbf{3} \oplus \bar{\mathbf{3}}$, so polyhedral structure might seem unnecessary. The objection is correct about gauge theory on spacetime, but conflates this with whether smooth manifolds can serve as the pre-geometric substrate from which spacetime emerges. They cannot: (i) $SU(3)/T^2$ presupposes the continuum (local charts, C^∞ transition functions) that the emergence program aims to derive; (ii) its coordinates are real-valued, not the integer lattice labels available pre-geometrically; (iii) gauge parallel transport requires differential structure, whereas polyhedra enforce phase coherence by boundary identification alone. The algebraic content is encoding-independent; the emergence application requires discrete structure (see [supplementary derivation](#)).

Definition II.8 (Pre-Geometric Substrate). The pre-geometric substrate $\mathcal{S}$ is the stella octangula (hereafter “the stella” in prose, $\mathcal{S}$ in equations) equipped with:

- (i) **Combinatorial structure:** A cell complex with 8 vertices, 12 edges, and 8 faces organized as two interpenetrating tetrahedra T_+ and T_- . Of these 8 vertices, 6 carry the non-zero SU(3) weights of $\mathbf{3} \oplus \bar{\mathbf{3}}$, while 2 apex vertices encode the radial direction perpendicular to the weight plane. These apex vertices are not demanded by (GR1)–(GR3) alone; they arise from the embedding dimension constraint (Proposition II.10), which requires $d_{\text{embed}} = 3$ for a confining SU(3) theory. The 2D weight plane accounts for 2 of 3 dimensions; each tetrahedron therefore requires one additional vertex along the orthogonal (color-singlet) direction, giving exactly 2 apex vertices.
- (ii) **T_d -equivariant measure:** A measure μ on $\partial\mathcal{S}$ invariant under the tetrahedral symmetry group T_d .
- (iii) **No metric at this stage:** Distances between points are undefined. Coordinates are topological labels, not measurements.

The metric emerges later from the SU(3) Killing form (Theorem V.6).

Lemma II.9 (Confinement Dimension Lower Bound). Within the geometric realization framework, a faithful polyhedral realization of SU(N) requires spatial dimension $d_{\text{space}} \geq N - 1$.

Proof. The Weyl group $W(SU(N)) \cong S_N$ must act faithfully on the vertices of $\mathcal{P}$ (from GR2). The fundamental weights $\{\mu_1, \dots, \mu_N\}$ are N points in weight space $\mathfrak{h}^* \cong \mathbb{R}^{N-1}$, subject to the constraint $\sum_c \mu_c = 0$. These N points are affinely independent: they span an $(N-1)$ -dimensional affine subspace. A faithful S_N -action on N affinely independent points requires ambient dimension $\geq N - 1$. For SU(3): $d_{\text{space}} \geq 2$. $\square$

Proposition II.10 (Embedding Dimension from Confinement). For a confining SU(N) gauge group geometrically realized via a polyhedral complex satisfying (GR1)–(GR3), the embedding dimension is $d_{\text{embed}} = N$. For SU(3): $d_{\text{embed}} = 3$.

Proof. The proof is a three-part squeeze, with each step classified by its epistemic status: (E) established mathematics or physics, (F) framework-specific axiom.

Part A: Lower bound ($d_{\text{embed}} \geq N-1$). [E] From Lemma II.9, the N fundamental weights are affinely independent in $\mathfrak{h}^* \cong \mathbb{R}^{N-1}$, so faithful Weyl group action requires ambient dimension $\geq N - 1$.

Part B: Strict improvement ($d_{\text{embed}} \geq N$). [E+F] Confinement introduces a physical length scale $R_{\text{conf}} = \hbar c / \sqrt{\sigma}$, the characteristic size of bound states ($\sqrt{\sigma} = 440 \pm 30$ MeV [5]). Weight-space coordinates are kinematic labels classifying *which* color state; the confinement radius encodes *how strongly* states are bound—a dynamical degree of freedom orthogonal to weight space. Within the geometric realization framework, this dynamical separation must be encoded as a geometric direction in $\mathcal{P}$. If $d_{\text{embed}} = N-1$, the weight-space directions exhaust all embedding dimensions, leaving no coordinate to parameterize quark–antiquark separation. Hence $d_{\text{embed}} \geq N$.

Part C: Upper bound ($d_{\text{embed}} \leq N$). [E+F] Pure SU(N) Yang-Mills has a single gauge coupling g_s [E]. The β -function is a single ODE, and dimensional transmutation produces a single dynamical scale Λ_{QCD} [E]. We invoke the geometric realization principle that each independent gauge coupling contributes at most one embedding dimension beyond weight space [**F: framework axiom**]. Since SU(N) has one coupling, the radial contribution is ≤ 1 , giving $d_{\text{embed}} \leq (N-1) + 1 = N$.

Combining: $d_{\text{embed}} = N$. For SU(3): $d_{\text{embed}} = 3$. $\square$

Remark II.11 (Epistemic Status of the Upper Bound). Parts A and B rest on established mathematics and experimental physics. Part C invokes one irreducible framework axiom: the coupling-to-dimension correspondence, which posits that each independent gauge coupling maps to at most one radial embedding direction. This axiom is physically motivated—a second radial direction would require a second independent confinement scale, which is absent in SU(N) (the θ -angle is topological and does not undergo dimensional transmutation; quark masses are matter-sector parameters that do not introduce independent confining directions)—but it is not derivable from more primitive principles. The full classification table and objection analysis are in the [supplementary proof document](#).




FIG. 2: Geometric realization of the radial direction (Proposition II.10). (a) The SU(3) weight space is a 2D plane (horizontal); within the geometric realization framework, the single gauge coupling's RG flow curve $\alpha_s(\mu)$ maps to one radial dimension orthogonal to it: $2 + 1 = 3$. (b) The running coupling $\alpha_s(\mu)$ is a single function of one variable (the β -function), motivating the framework axiom that confinement contributes at most one radial direction. The confinement scale $\sqrt{\sigma} = \hbar c/R_{\text{stella}} \approx 440$ MeV and $\Lambda_{\overline{MS}}^{(5)} = 210$ MeV are marked.

Remark II.12 (Scope: Geometric Realization, Not Confinement). The result $d_{\text{embed}} = N$ applies to faithful geometric realizations satisfying (GR1)–(GR3)—it does not claim that SU(N) confinement requires N spatial dimensions. Lattice QCD conclusively demonstrates that SU(3) confines in 2+1 dimensions, and SU(N) for $N > 3$ confines in 3+1 dimensions. The distinction is that lattice gauge theory does not impose (GR1)–(GR3): gauge groups enter through link variables $U_\mu(x) \in G$, not through the lattice's spatial structure. The framework's claim is that if gauge group structure is geometrically realized—the core premise (F1)—then the embedding dimension is constrained by the gauge group.

Definition II.13 (Minimality). A geometric realization is minimal if:

- (M1) The vertex count equals the dimension of $\mathbf{3} \oplus \bar{\mathbf{3}}$.
- (M2) The embedding dimension is the smallest compatible with (GR1)–(GR3).

For SU(3), the minimal embedding dimension is 3 (Proposition II.10): two dimensions from weight space, one from the confinement radius that geometrically separates T_+ from T_- .

B. The SU(3) Weight System

The Lie algebra $\mathfrak{su}(3)$ has rank 2, with Cartan subalgebra spanned by $\{H_1, H_2\}$. The fundamental representation $\mathbf{3}$ has weights:

$$\mu_R = \left(\frac{1}{2}, \frac{1}{2\sqrt{3}}\right), \mu_G = \left(-\frac{1}{2}, \frac{1}{2\sqrt{3}}\right), \mu_B = \left(0, -\frac{1}{\sqrt{3}}\right) \quad (3)$$

a. Normalization convention. We choose the standard Dynkin normalization where fundamental weights form an equilateral triangle with unit side length and longest roots have squared length 2. This is a convention choice; the structural correspondence is convention-independent.

The Weyl group $\mathcal{W}(\text{SU}(3)) \cong S_3$ acts by permuting colors and by reflection (charge conjugation):

- Cyclic permutations $R \rightarrow G \rightarrow B \rightarrow R$ generate $\mathbb{Z}_3 \subset S_3$
- Pairwise exchanges generate transpositions
- Charge conjugation $C : \mu \mapsto -\mu$ exchanges $\mathbf{3} \leftrightarrow \bar{\mathbf{3}}$

The simple roots are:

$$\alpha_1 = (1, -1/\sqrt{3}), \quad \alpha_2 = (0, 2/\sqrt{3}) \quad (4)$$




FIG. 3: The SU(3) weight diagram for $\mathbf{3} \oplus \bar{\mathbf{3}}$, forming a hexagram pattern. Blue triangle: fundamental weights (circles). Red triangle: antifundamental weights (squares). Purple arrows: charge conjugation ($\vec{w} \rightarrow -\vec{w}$), corresponding to GR3. Gold star: origin where apex vertices project onto the weight plane.

III. OBSERVER-COMPATIBLE SPACETIME DIMENSIONALITY

With the geometric realization framework established, we must determine the spacetime dimension in which this realization is embedded. The dimension is not a free parameter: the following theorem shows that $D = 4$ is uniquely selected by the requirement that stable bound-state observers exist.

a. Nature of this argument. The following theorem is a *selection* argument, not a dynamical derivation. It identifies which spacetime dimensions are compatible with the existence of observers, not why our universe has observers or any particular dimension.

Definition III.1 (Stable Bound-State Observer). *A stable bound-state observer is a spatially localized, temporally persistent physical system capable of storing and processing information. Operationally, this requires: (i) stable gravitational bound states, (ii) stable atomic bound states, (iii) causal signal propagation, and (iv) sufficient topological complexity. Condition (v) establishes the temporal signature independently.*

Theorem III.2 (Unique Dimensionality). *Under general relativity and quantum mechanics, the spacetime dimension $D = 4$ is uniquely compatible with stable bound-state observers (Definition III.1).*

Proof. The proof proceeds by elimination of all $D \neq 4$ via five physical requirements.

(P1) *Gravitational Stability:* $D \leq 4$. In D -dimensional spacetime with $n = D - 1$ spatial dimensions, the gravitational potential scales as $V(r) \propto r^{-(n-2)}$. Stability of circular orbits requires $d^2V_{\text{eff}}/dr^2 > 0$, which fails for $n \geq 4$. This is the Ehrenfest instability argument [6].

(P2) *Atomic Stability:* $D = 4$ uniquely. In $D = 2 + 1$, the hydrogen atom has degeneracy $(2k + 1)$, not the k^2 degeneracy of 3D [7]. The k^2 degeneracy enables orbital hybridization (sp, sp^2 , sp^3) essential for tetrahedral carbon bonding. In $D = 4 + 1$ (Coulomb $\propto 1/r^2$), the Hamiltonian is unbounded below and atoms collapse. Thus $D = 4$ is uniquely compatible with stable atoms and chemistry-enabling spectra.

(P3) *Causal Wave Propagation.* Huygens' principle (sharp wavefronts) holds exactly only for odd spatial dimensions $n \geq 3$. Combined with (P1)–(P2), this selects $n = 3$.

(P4) *Topological Complexity.* Non-trivial knots exist only in $n = 3$ spatial dimensions. In $n \geq 4$, all knots can be untied [8]. The biological relevance of 3D knot theory includes DNA supercoiling [9], knotted proteins [10], and molecular machines [11].

(P5) *Temporal Uniqueness:* $t = 1$. For $t = 0$, no dynamics exist. For $t \geq 2$, the wave equation becomes ultrahyperbolic, violating causality. Thus $t = 1$ uniquely admits deterministic physics.

The intersection uniquely selects $n = 3$, $t = 1$:

$$\{n \leq 3\} \cap \{n = 3\} \cap \{n \geq 3, \text{ odd}\} \cap \{n = 3\} = \{3\} \quad (5)$$

□

b. Independent dynamical mechanisms. Theorem III.2 is a selection argument: it identifies $D = 4$ as the unique dimension compatible with observers. Critically, four independent *dynamical* mechanisms also yield $D = 4$ without any anthropic reasoning:




FIG. 4: Dimensional selection via observer stability constraints. (a) Effective potential: only $D = 4$ has a stable minimum. (b) Constraint intersection: $D = 4$ uniquely satisfies all requirements (P1)–(P5).

1. Causal dynamical triangulations recover $D = 4$ from a non-perturbative gravitational path integral [12, 13].
2. The Brandenberger–Vafa mechanism selects 3 large spatial dimensions via string winding mode annihilation [14].
3. Feng’s entropic argument shows that $D = 4$ maximizes gravitational entropy production [15].
4. Carlip’s “spontaneous dimensional reduction” demonstrates that short-distance dynamics in several quantum gravity approaches flow to $D = 2$, with $D = 4$ as the unique stable macroscopic fixed point [16].

The selection and dynamical streams are complementary: even if one rejects anthropic reasoning entirely, four independent physical mechanisms independently select $D = 4$. For the combined case to fail, all five arguments must simultaneously be wrong.

Remark III.3 (Constraint Hierarchy). *Constraints (P1) and (P2) alone uniquely select $D = 4$. Constraints (P3)–(P5) serve as consistency checks. The selection is robust: it does not depend on assumptions about signal cleanliness or knotted structures.*

c. Selection vs. measure problem. $D \neq 4$ yields zero observers, not fewer. These are absolute prohibitions, making $D = 4$ a sharp prediction rather than a soft selection effect.

An immediate objection arises: Theorem III.2 uses general relativity and quantum mechanics as inputs. If this framework claims to be more fundamental than both, is the derivation circular? The following theorem shows that GR and QM are *compatible with* and naturally produced by the geometric realization conditions, establishing self-consistency. The full dynamical derivations of GR and QM are completed in Papers 2–3; here we trace the structural chain that closes the logical loop.

Theorem III.4 (Framework-Internal $D = 4$ Consistency). *The geometric realization conditions (GR1)–(GR3) are compatible with and naturally lead to the standard physics (general relativity and quantum mechanics) used as input to Theorem III.2. This is a self-consistency*

verification, *not an independent derivation of $D = 4$: the physics assumed is recoverable from the geometry obtained, confirming that the logical loop closes without contradiction.*

Proof. We show that (GR1)–(GR3) naturally produce both GR and QM through a four-step chain. Each step is classified as established physics [E] or involving framework content [E+F].

Step 1: Non-abelian gauge structure. [E+F] (GR2) requires $\text{Aut}(\mathcal{P}) \supseteq W(G)$ [F]. For $\text{SU}(3)$, $W \cong S_3$ is non-abelian, forcing a non-abelian gauge group with self-interacting spin-1 mediators (Yang-Mills theory) [E].

Step 2: Spin-2 gravity. [E] By Weinberg’s soft graviton theorem [17], any Lorentz-invariant theory with massless spin-1 particles coupling to a conserved stress-energy tensor necessarily contains massless spin-2 exchange [E]. Lorentz invariance is itself derived within the framework: discrete O_h rotational symmetry of the FCC lattice extends to continuous $\text{SO}(3)$ in the continuum limit, and boosts emerge from the dynamical sector (the full derivation is in Paper 2; the supplementary [proof document](#) traces the complete chain through Theorems 0.0.8 and 0.0.11). The Einstein equations are recovered via the thermodynamic route (Theorem 5.2.3 in Paper 3).

Step 3: Quantum kinematics. [E+F] (GR1) requires vertices to biject with representation weights—the discrete eigenvalues of Cartan generators [F]. Discrete eigenvalue spectra are a characteristic feature of quantum mechanics [E]. The A_2 weight lattice structure, combined with the S_3 Weyl action, implies the full $\text{SU}(3)$ representation theory, from which quantum-mechanical operator algebra follows. Full quantum dynamics (Schrödinger equation, Born rule) requires additional development beyond this paper (Theorem 0.0.10 in the supplementary proofs; see Paper 2).

Step 4: Closure. Steps 1–3 establish that (GR1)–(GR3) are compatible with and naturally lead to non-abelian gauge theory, gravity, and quantum kinematics—the inputs to Theorem III.2. The logical chain closes:

$$\begin{aligned} (\text{GR1}–\text{GR3}) &\longrightarrow \text{GR} + \text{QM} \xrightarrow{\text{Thm III.2}} D=4 \\ &\longrightarrow \text{SU}(3) \longrightarrow \text{Stella} \longrightarrow (\text{GR1}–\text{GR3}) \checkmark \end{aligned} \quad (6)$$

The loop is self-consistent, not circular, because the entry point (observer existence, I1) is external to the loop. This theorem reproduces the same Ehrenfest physics used in Theorem III.2 and therefore does not constitute an independent confirmation of $D = 4$; its role is to verify that the framework’s outputs are compatible with its inputs. We use $\longrightarrow$ rather than $\Rightarrow$ for Steps 2–3 to emphasize that the full GR and QM derivations require results from Papers 2–3; this paper establishes only the kinematic compatibility and the structural chain. $\square$

A. Scope, Evidence, and Alternatives

a. Experimental confirmation. Three classes of experiments confirm $D = 4$: (1) LIGO/Virgo detect exactly 2 gravitational wave polarizations ($D(D-3)/2 = 2 \Rightarrow D = 4$); (2) torsion balance experiments test gravity to $52 \mu\text{m}$ with no deviation; (3) LHC constrains $M_D > 5 \text{ TeV}$ for 2 extra dimensions.

b. Scope conditions. The result assumes standard physics: general relativity (tensor gravity), gauge-invariant electromagnetism, and quantum mechanics. Scargill [18] showed 2+1D supports neural complexity if gravity is scalar; Burgbacher et al. [19] showed stable atoms in $D \geq 4$ with modified electromagnetism. These constitute modified physics, not counterexamples within the stated framework.

c. Compactified extra dimensions. Compactification scenarios (e.g., string theory) begin with $D = 10$ or $D = 26$ and reduce to $D = 4$ by compactifying extra dimensions on a Calabi-Yau or orbifold manifold. The present framework instead begins with observer existence and arrives at $D = 4$ directly. The two approaches ask different questions: compactification asks “how to reduce $D > 4$ to $D = 4$?” while Theorem III.2 asks “why $D = 4$ at all?” Unlike Kaluza-Klein theory, which posits continuous extra spatial dimensions whose isometries generate gauge symmetry (and faces the chiral fermion problem), the present framework operates in exactly $D = 4$ with discrete pre-geometric structure and no compactified dimensions. Detection of large extra dimensions (e.g., Kaluza-Klein excitations at colliders, deviations from $1/r^2$ gravity at sub-millimeter scales) remains falsification condition FC1.

IV. THREE PATHS TO $\text{SU}(3)$

Having established $D = 4$ from observer existence, the next step is identifying which gauge group is geometrically realized. The dimension constrains but does not uniquely determine this: $D = 4$ restricts $\text{SU}(N)$ candidates to rank ≤ 2 via the embedding dimension $d_{\text{embed}} = N \leq 3$ (Proposition II.10), but additional geometric structure is required to single out one group. We derive $\text{SU}(3)$ via two complementary paths—topological and information-theoretic—and verify the result with a third categorical consistency check. Both derivation paths share the $D = 4$ dimensional constraint (I1), making them convergent rather than fully independent; they employ different mathematical machinery and different subsets of the remaining inputs.

Remark IV.1 (Coleman–Mandula and the Pre-Geometric Regime). *The Coleman–Mandula theorem [20] states that the symmetry group of any QFT satisfying certain assumptions is necessarily a direct product $\text{Poincaré} \times G_{\text{internal}}$. Since this paper identifies gauge structure with spatial geometry, the apparent conflict must be addressed.*

The theorem requires five hypotheses, none of which hold pre-geometrically:

CM Hypothesis	Pre-geometric status
<i>S</i> -matrix exists	No scattering without spacetime
Poincaré invariance	No spacetime, no Poincaré
Mass gap	No Hamiltonian on spacetime
Two-particle scattering	No particles
Analytic amplitudes	No amplitudes

The gauge–geometry identification occurs before spacetime exists. Emergence is “complete”—and the CM hypotheses become satisfied—once the continuum limit (Theorem ??) produces a Lorentzian manifold with a well-defined *S*-matrix, a mass gap from confinement (Paper 2), and particle states carrying Poincaré quantum numbers. After that transition, the standard direct-product structure $\text{Poincaré} \otimes \text{SU}(3)_{\text{color}}$ obtains: weight space becomes the internal fiber, physical space becomes the base manifold, and the *S*-matrix is well-defined.

It is important to distinguish the two claims involved:

Standard:: $\text{SU}(3)$ weight vertices form an equilateral triangle in weight space $\mathfrak{h}^*$ (textbook Lie algebra).

Novel:: Weight space $\mathfrak{h}^*$ is a subspace of physical $\mathbb{R}^3$ in the pre-geometric phase (the load-bearing premise F1).

The first requires no defense. The second is precisely where the pre-geometric loophole is invoked: before spacetime emerges, the distinction between “internal” and “external” has not been established. This loophole is shared by all emergent-spacetime programs (LQG, causal sets, CDT; see Section X E).

The framework provides bounds on the transition scale: the discrete stella structure is manifest at $E \sim E_P$ (lattice spacing $a \approx 2.25 \ell_P$), while Lorentz violation is negligible at $E \ll E_P$ (Theorem VII.2: suppression $(E/E_P)^2$). The direct-product structure therefore obtains at all scales where the *S*-matrix is well-defined. The remaining ingredient—the mass gap scale—requires dynamical content from Paper 2.

A. Path A: Topological Derivation

Theorem IV.2 (Gauge Group Selection). *Among simple compact Lie groups of rank ≤ 4 , $\text{SU}(3)$ is uniquely compatible with 3D polyhedral realization satisfying (GR1)–(GR3).*

Proof. The simple compact Lie groups of rank ≤ 2 are: $\text{SU}(2)$ (rank 1), $\text{SU}(3)$ (A_2 , rank 2), $\text{SO}(5) \cong \text{Sp}(4)/\mathbb{Z}_2$ (B_2 , rank 2), and G_2 (rank 2). $\text{SU}(2)$ has a 2-weight fundamental (line segment), failing (GR2). $\text{SO}(5)$ has a square weight diagram with no 3-fold symmetry. G_2 has a 7-dimensional fundamental preventing vertex–weight bijection. Only $\text{SU}(3)$ admits a polyhedral realization satisfying (GR1)–(GR3). $\square$

Theorem IV.3 (Topological Derivation of $\text{SU}(3)$). *The stella octangula uniquely determines $\text{SU}(3)$ via its intrinsic $\mathbb{Z}_3$ rotational symmetry, independent of the weight-diagram argument.*

Proof. Step 1: $\mathbb{Z}_3$ from stella geometry. [E] The stella octangula has 3-fold rotational symmetry about each body diagonal. The rotations $\{I, R_{2\pi/3}, R_{4\pi/3}\}$ form $\mathbb{Z}_3$, derived from polyhedral geometry alone. (This mechanism— $\mathbb{Z}_3$ discrete symmetry from geometric frustration—has been experimentally observed in kagome-lattice metals, where three geometrically frustrated sublattices produce $\mathbb{Z}_3$ nematic order [21].)

Step 2: $\mathbb{Z}_3 \subseteq Z(G)$. [E] Color charges (triality) are gauge-invariant quantum numbers that must commute with all gauge transformations, placing them in the center $Z(G)$. This is established gauge theory [E].

Step 3: Classification by center. [E] Compact simple Lie groups with $\mathbb{Z}_3 \subseteq Z(G)$ are: $\text{SU}(3k)$ for $k \geq 1$ and E_6 (center exactly $\mathbb{Z}_3$). This is a standard classification result [E].

Step 4: Rank constraint from embedding dimension. [E+F] Proposition II.10 establishes $d_{\text{embed}} = N$ for $\text{SU}(N)$: the weight space contributes $N-1 = \text{rank}(G)$ dimensions, and the confinement radius contributes one additional dimension. Since Theorem III.2 gives $D_{\text{space}} = 3$, we require $\text{rank}(G) + 1 \leq D_{\text{space}}$, hence $\text{rank}(G) \leq 2$. The upper bound inherits the framework axiom from Proposition II.10 (Remark II.11).

Step 5: Unique intersection. [E]

Group	Rank	$\mathbb{Z}_3 \subseteq Z?$	Result
$\text{SU}(2)$	1	$\times$	Excluded
$\text{SU}(3)$	2	$\checkmark$	Unique
$\text{SO}(5)$	2	$\times$	Excluded
G_2	2	$\times$	Excluded
$\text{SU}(6)$	5	$\checkmark$	Rank fails
E_6	6	$\checkmark$	Rank fails

Therefore $G = \text{SU}(3)$ is uniquely determined. $\square$

Remark IV.4 ($\mathbb{Z}_4$ Exclusion: Independent of Rank Bound). *The rank constraint (Step 4) depends on the framework axiom in Proposition II.10. However, $\text{SU}(4)$ —the next candidate after $\text{SU}(3)$ in the $\text{SU}(3k)$ family—is excluded by an independent, purely algebraic argument that requires no framework input: $Z(\text{SU}(4)) = \mathbb{Z}_4$, and $\mathbb{Z}_4$ does not contain $\mathbb{Z}_3$ as a subgroup since $3 \nmid 4$. The element $\omega = e^{2\pi i/3}$ is not in $\mathbb{Z}_4 = \{e^{2\pi i k/4} : k = 0, 1, 2, 3\}$. Thus $\text{SU}(4)$ fails at Step 3, independently of any rank bound. More broadly, $N = 3$ is distinguished by the algebraic status of $\mathbb{Z}_3$ in the Weyl group $W(\text{SU}(N)) \cong S_N$. For $N = 3$, $\mathbb{Z}_3 = A_3 \trianglelefteq S_3$ is the unique normal cyclic-3 subgroup (the alternating group of index 2). For $N \geq 4$, $\mathbb{Z}_3$ embeds in S_N but is not normal—there are $\binom{N}{3}$ distinct 3-cycles, none distinguished. The stella’s $\mathbb{Z}_3$ is the maximal color symmetry, not a subgroup of a larger color rotation, matching the $N = 3$ algebraic structure exactly.*

Remark IV.5 (What Survives if Rank ≤ 2 Is Relaxed). *If the rank constraint (Step 4, which inherits one framework axiom from Proposition II.10) is dropped, the $\mathbb{Z}_3$ center condition alone leaves additional candidates:*

Group	Rank	dim(8)	Fund dim	Experimental conflict
SU(3)	2	8	3	None (QCD)
SU(6)	5	35	6	35 gluons, DIS rules out
SU(9)	8	80	9	80 gauge bosons
E_6	6	78	27	78 gauge bosons, 27-dim fund

All higher-rank candidates fail experimentally: deep inelastic scattering confirms exactly 8 gluonic degrees of freedom contributing to the proton momentum sum rule ($\langle x \rangle_g \approx 0.41$, consistent with 8 gluons). The rank constraint provides a geometric reason for what is otherwise an empirical fact—the geometric realization postulate (Definition II.1) requires the weight diagram to embed in the stella’s spatial structure, limiting the rank to $D_{\text{space}} - 1 = 2$.

Remark IV.6 (Homotopy Clarification: π_1 vs. π_3). *The $\mathbb{Z}_3$ color cycle $R \rightarrow G \rightarrow B \rightarrow R$ on the stella is a generator of $\pi_1(\text{PSU}(3)) = \mathbb{Z}_3$ (center symmetry, governing confinement and N -ality), not of $\pi_3(\text{SU}(3)) = \mathbb{Z}$ (instantons). The stella directly encodes the former—the $\mathbb{Z}_3$ phase structure of the center. The latter is a mathematical consequence: $\pi_3(\text{SU}(N)) = \mathbb{Z}$ for all $N \geq 2$, so once SU(3) is determined, the instanton classification follows automatically.*

B. Path B: Information-Theoretic Derivation

Path A depends on the geometric realization postulate (F1) and the stella’s $\mathbb{Z}_3$ symmetry. To verify that SU(3) is not an artifact of these specific assumptions, we present a second, complementary derivation from information geometry that does not invoke F1. Path B derives $N = 3$ purely from the structure of quantum interference, requiring three explicit assumptions beyond established mathematics:

Assumption IV.7 (Quantum Interference Form (A-IF)). *Distinguishable configurations produce probability distributions of the interference form $p_\phi(x) = |\sum_c A_c(x) e^{i\phi_c}|^2$. This presupposes complex amplitudes, coherent superposition, and the Born rule.*

Assumption IV.8 (Color Democracy (A-SN)). *All N field components are physically equivalent—the probability distribution $p_\phi(x)$ is invariant under the permutation group S_N acting on component labels.*

Assumption IV.9 (Irreducible Information Density (A-IID)). *Among Fisher-stable, algebraically irreducible configurations, nature realizes the one that maximizes per-DOF Fisher information $I_{\text{DOF}}(N) = 1/(2N)$.*

These three assumptions constitute Path B’s complete input set. A-IF is the load-bearing premise: without quantum interference structure, the $N = 2$ elimination fails. A-IID provides the selection principle that singles out $N = 3$ from the infinite set of Fisher-stable primes.

The Fisher information metric is evaluated on the *interference configuration space* T^{N-1} —the torus of relative phases among N components—not on the raw probability simplex. For N components with amplitudes $A_c(x)$ and phases ϕ_c , the interference pattern is

$$p_\phi(x) = \left| \sum_{c=1}^N A_c(x) e^{i\phi_c} \right|^2 \quad (7)$$

with the Fisher metric $g_{ij}^F(\phi) = \int p_\phi(x) \partial_i \log p_\phi \partial_j \log p_\phi dx$. Define the stability function $S(N) = 1$ if g^F is positive-definite at the S_N -symmetric equilibrium $\phi_c = 2\pi c/N$, and $S(N) = 0$ otherwise.

An additional physical constraint enters the lower bound: *color neutrality* requires the net phase to vanish, $\sum_{c=1}^N e^{i\phi_c} = 0$. This condition ensures that a color-singlet state exists—a physical necessity for confinement.

Proposition IV.10 (First Stable Principle). *Among Fisher-stable ($S(N) = 1$), algebraically irreducible (N prime) configurations, $N = 3$ uniquely maximizes the per-degree-of-freedom Fisher information $I_{\text{DOF}}(N) = \text{Tr}(g_N^F)/(N - 1) = 1/(2N)$.*

Proof. The proof has three components.

(i) *Fisher stability analysis.* For $N = 1$, the configuration space is zero-dimensional; no metric exists, so $S(1) = 0$.

For $N = 2$, two independent arguments exclude this case:

- (a) *Fisher degeneracy.* The equilibrium phases $(\phi_0, \phi_1) = (0, \pi)$ produce maximally destructive interference: $p_\phi(x) = |A_0(x) - A_1(x)|^2$, with $\partial p_\phi / \partial \phi|_{\text{eq}} = 0$ since $\sin(-\pi) = 0$. The Fisher metric vanishes: $g^F = 0$, so $S(2) = 0$.
- (b) *Color neutrality kills all DOF.* The constraint $e^{i\phi_1} + e^{i\phi_2} = 0$ forces $\phi_2 = \phi_1 + \pi$. Counting degrees of freedom: 2 phases $- 1$ neutrality $- 1$ overall gauge $= 0$. The configuration space is a single point.

Two further confirmations (Hessian degeneracy at the constrained equilibrium; complete destructive cancellation for S_2 -symmetric amplitudes) are given in the supplementary proof document (Proposition: SU(3) from Distinguishability Constraints, in preparation). This exclusion is purely information-theoretic—it does not invoke geometric realization (F1).

For $N = 3$, the equilibrium phases $(0, 2\pi/3, 4\pi/3)$ produce a non-degenerate 2×2 Fisher matrix with eigenvalues $\lambda_1 \approx 0.736$ and $\lambda_2 \approx 0.245$, both strictly positive. Hence $S(3) = 1$. The interference pattern evaluates to

$p = \frac{1}{2}[(A_R - A_G)^2 + (A_G - A_B)^2 + (A_B - A_R)^2]$, which is strictly positive for generic amplitudes (the zero set is a union of three perpendicular bisector planes, of Lebesgue measure zero). For all $N \geq 3$, $S(N) = 1$ (the Fisher metric is non-degenerate at S_N -symmetric equilibrium).

(ii) *Algebraic irreducibility.* For composite $N = a \cdot b$ ($a, b \geq 2$), the interference pattern decomposes into a sub-interference terms via the coset structure $\mathbb{Z}_N/\mathbb{Z}_a \cong \mathbb{Z}_b$. For prime N , no such decomposition exists (Lagrange’s theorem: $\mathbb{Z}_N$ has no proper non-trivial subgroups). The irreducible Fisher-stable systems are therefore $\{N \text{ prime} : N \geq 3\} = \{3, 5, 7, 11, \dots\}$.

(iii) *Per-DOF information maximization.* Among irreducible systems, the per-DOF Fisher information $I_{\text{DOF}}(N) = 1/(2N)$ is strictly decreasing in N . Applying Assumption IV.9 (A-IID): $N^* = \underset{N \text{ prime}, N \geq 3}{\operatorname{argmax}} \frac{1}{2N} = 3$. $\square$

Lemma IV.11 (Fisher-Killing Equivalence). *For a statistical manifold with S_N permutation symmetry: $g_{ij}^F = c_N \cdot g_{ij}^K$ where g^F is the Fisher metric and g^K the Killing form of $\text{SU}(N)$.*

Proof sketch. (1) The space of S_N -invariant symmetric 2-tensors on T^{N-1} is 1-dimensional. (2) Both g^F and g^K are S_N -invariant. (3) Since both are positive-definite and the space is 1-dimensional, they are proportional. $\square$

Corollary IV.12 ($\text{SU}(3)$ from First Stable Principle). *The derivation chain: First Stable $\Rightarrow N = 3 \Rightarrow W \cong S_3 \Rightarrow$ root system $A_2 \Rightarrow \mathfrak{su}(3) \Rightarrow \text{SU}(3)$.*

Remark IV.13 (Scope and Honesty). *The mathematical content— $S(2) = 0$ from destructive interference, $S(N) = 1$ for $N \geq 3$, composite- N decomposition, prime- N irreducibility—is rigorously proven. The residual selection step (Assumption IV.9) is a methodological criterion, not a derivable physical law. This selects N for the color sector specifically; $\text{U}(1)_Y$ and $\text{SU}(2)_L$ enter via the polytope embedding chain (Section VIII). Path B is best characterized as a retrodiction: given A-IF—which presupposes complex amplitudes, coherent superposition, and the Born rule—Path B confirms that $N = 3$ is uniquely selected. Since A-IF itself encodes quantum-mechanical structure, the argument is a consistency verification of $\text{SU}(3)$ from information-theoretic principles, not a derivation from pre-quantum first principles.*

Remark IV.14 (Three-Bound Convergence). *The $N = 3$ conclusion can be understood as the unique intersection of three independent constraints. From Theorem III.2 ($D_{\text{space}} = 3$) and Lemma II.9 ($d_{\text{space}} \geq N - 1$) one obtains a geometric upper bound $N \leq 4$. Color neutrality imposes $3 \mid N$. Together with the information-theoretic lower bound:*

$$\underbrace{N \geq 3}_{\text{Fisher stability}} \cap \underbrace{N \leq 4}_{\text{affine independence}} \cap \underbrace{3 \mid N}_{\text{color neutrality}} = \{3\}. \quad (8)$$

This reveals two independent routes to $N = 3$ within Path B (Table III):

- (a) A-IID route: *Fisher stability + irreducibility + per-DOF maximization (Assumption IV.9). Uses only A-IF and A-IID; no geometric input.*
- (b) Geometric route: *Eq. (8)—Fisher stability + affine independence (I1) + $\mathbb{Z}_3$ center. Uses I1 but not A-IID.*

The overdetermination—two routes with non-overlapping selection mechanisms—strengthens the conclusion beyond what either provides alone. A purely information-theoretic upper bound on N (without I1) remains an open problem: the Fisher metric has full rank for all $N \geq 3$ (see supplementary proof document: $\text{SU}(3)$ from Distinguishability Constraints, §6.1.1, in preparation).

TABLE III: Constraints on the color number N within Path B. The first three rows (above the rule) suffice for the geometric route; the last two rows constitute the A-IID route. Both routes share the Fisher stability lower bound.

Constraint	Source	Range
Fisher stability	$S(2) = 0$ (A-IF)	$N \geq 3$
Affine independence	Lem. II.9 + I1	$N \leq 4$
$\mathbb{Z}_3$ center	Color neutrality	$3 \mid N$
Irreducibility	Lagrange (prime N)	$\{3, 5, 7, \dots\}$
Info density	A-IID (max $1/(2N)$)	$N = 3$

C. Path C: Categorical Consistency Check

Path C does not independently derive $\text{SU}(3)$ —that work is done by Paths A and B. Instead, it verifies that the stella $\leftrightarrow \text{SU}(3)$ identification obtained from those paths is *categorically sound*: the polyhedral data are precisely the data required by Tannaka–Krein reconstruction, and no additional algebraic information is needed or lost.

Theorem IV.15 (Categorical Equivalence). *The category $\mathcal{C}_{\text{poly}}$ of A_2 -decorated polyhedral complexes satisfying (GR1)–(GR3) is equivalent to the category $\mathcal{C}_{\text{Weyl}}$ of S_3 -sets with A_2 weight structure.*

Remark IV.16 (Scope of the Equivalence). *Theorem IV.15 operates at the Cartan data level: root system $\Phi(A_2)$, weight lattice, and Weyl group $W \cong S_3$. It does not directly encode the full 8-parameter continuous group $\text{SU}(3)$ or the tensor product structure of $\text{Rep}(\text{SU}(3))$. The passage from Cartan data to the Lie algebra is supplied by Serre’s reconstruction theorem (a classical result); the further passage to the Lie group is Corollary IV.17. Thus the claim “ $\text{SU}(3)$ IS the stella” is established in two stages: Theorem IV.15 identifies the discrete Cartan data, and Corollary IV.17 recovers the continuous group.*




FIG. 5: Fisher information stability and the uniqueness of $N = 3$. (a) Stability transition: configurations with $N \leq 2$ fields (gray) are Fisher-unstable; $N \geq 3$ (green) are stable, with $N = 3$ (red) highlighted as the unique intersection $\{N \geq 3\} \cap \{N \leq 4\} \cap \{3 \mid N\}$. (b) Per-DOF Fisher information $I_{\text{DOF}} = 1/(2N)$ is maximized at the smallest stable value $N = 3$. (c) Angular coverage patterns: $N = 2$ (dashed blue) has deep valleys near zero, while $N = 3$ (solid red) maintains positive-definite coverage at all angles.

Together, these two results make precise the claim “SU(3) IS the stella.” The stella is the initial object in the category of A_2 -decorated polyhedral complexes.

Corollary IV.17 (Tannaka Reconstruction). *The full Lie group SU(3) can be reconstructed from the stella via Tannaka-Krein duality: $\text{SU}(3) \cong \text{Aut}^\otimes(\omega)$ where $\omega : \text{Rep}(\text{SU}(3)) \rightarrow \text{Vec}$ is the forgetful functor.*

Remark IV.18 (The Continuous from the Discrete). *Tannaka reconstruction recovers a continuous group (SU(3), 8 real parameters) from discrete categorical data (8 vertices, 12 edges, 8 faces)—showing how continuous symmetries can emerge from fundamentally discrete structure. (The equality $8_{\text{parameters}} = 8_{\text{vertices}}$ is a numerical coincidence, not a structural correspondence; the reconstruction depends on the full categorical data, not the vertex count alone.)*

Remark IV.19 (Tensor Products from Stella Geometry). *The stella encodes the tensor product decompositions of SU(3) concretely:*

- $\mathbf{3} \otimes \bar{\mathbf{3}} = \mathbf{8} \oplus \mathbf{1}$: The 6 root edges connecting weight vertices correspond to 6 charged gluons; the 2 apex vertices (zero weight, Cartan generators T_3 and T_8) correspond to 2 neutral gluons; vertex-antivertex pairing yields the singlet $\mathbf{1}$. The explicit 8 gluon color states are $\{r\bar{g}, r\bar{b}, g\bar{r}, g\bar{b}, b\bar{r}, b\bar{g}, (r\bar{r} - g\bar{g})/\sqrt{2}, (r\bar{r} + g\bar{g} - 2b\bar{b})/\sqrt{6}\}$; the would-be ninth combination $(r\bar{r} + g\bar{g} + b\bar{b})/\sqrt{3}$ is absent due to SU(3) tracelessness.
- $\mathbf{3} \otimes \mathbf{3} = \mathbf{6} \oplus \bar{\mathbf{3}}$: Each triangular face $F = \{R, G, B\}$ of T_+ is an oriented 2-simplex whose cyclic orientation $(R \rightarrow G \rightarrow B \rightarrow R)$ encodes the Levi-Civita symbol ϵ^{abc} . The antisymmetric combination $|a \wedge b\rangle$ has weights equal to $-w_{\bar{c}}$, reproducing $\bar{\mathbf{3}}$. The 3 diagonal states (cc) plus 3 symmetric off-diagonal states span the sextet $\mathbf{6}$.

The hadron phase structure follows immediately: in mesons, $\psi_{\text{meson}} \propto \sum_c \chi_c \bar{\chi}_c = \sum_c |A_c|^2$ (phases cancel pairwise: $e^{i\phi_c} e^{-i\phi_c} = 1$); in baryons, $\psi_{\text{baryon}} \propto \epsilon^{abc} \chi_a \chi_b \chi_c$ (the antisymmetric tensor ensures the color singlet). Both constructions are geometrically visible in the stella: mesons as vertex-antivertex pairs, baryons as oriented triangular faces. These decompositions make the claim “SU(3) IS the stella” concrete: not only the root system and Weyl group, but also the full tensor product structure is geometrically visible.

Remark IV.20 (Why Path C Is a Consistency Result). *The logical chain makes clear why Corollary IV.17 is a verification, not an independent derivation:*

1. $D = 4$ established (Theorem III.2), independent of SU(3).
2. SU(3) selected as the unique SU(N) compatible with $D = 4$ (Theorem IV.3).
3. Stella constructed as the unique minimal geometric realization (Theorem V.10).
4. The fiber functor ω is defined using the vertex $\leftrightarrow$ weight correspondence from step 3—this already uses knowledge that the stella encodes SU(3).
5. Tannaka reconstruction recovers SU(3) from stella + ω , confirming the identification.

If the stella harbored a hidden incompatibility with SU(3) representation theory (e.g., inconsistent tensor products), step 5 would expose it. The value of Path C is therefore not derivation but consistency certification: it guarantees that stella geometry and SU(3) algebra can be used interchangeably without loss of information.




FIG. 6: Confinement gradient on the stella octangula. T_+ vertex colors (R, G, B) gradient toward white at the singlet vertex W , visualizing how color charges confine into color singlets along the radial direction (Proposition II.10). The T_- tetrahedron (faint red) carries the anti-fundamental representation. White markers at the six T_+/T_- edge crossings (inner octahedron) show mesonic confinement points where vertex–antivertex pairs $q + \bar{q} \rightarrow \mathbf{1}$ form color singlets. The dashed hexagon marks the 2D weight plane $\mathfrak{h}^*$; the green axis is the 1D singlet (radial) direction.

D. Path Convergence

The two derivation paths and the consistency check use different mathematical machinery and different subsets of the framework’s inputs:




FIG. 7: Two complementary derivation paths (A, B) converge on $SU(3)$; Path C provides categorical consistency verification. The convergence of the derivation paths, using different mathematical machinery, ensures the result is not an artifact of any single approach.

Both paths share the $D = 4$ dimensional constraint (I1) and ultimately rest on the gauge–geometry correspondence encoded in GR1–GR3 (Def. 0.0.0). They are complementary—employing different mathematical ma-

TABLE IV: Three paths to $SU(3)$ and their input dependencies. Paths A and B provide complementary derivations sharing the $D = 4$ dimensional input; Path C provides categorical consistency verification of the result. The convergence of the two derivation paths, using different mathematical machinery, strengthens the conclusion.

	Path A	Path B	Path C [†]
	Topological	Info-theoretic	Categorical
Method	$\mathbb{Z}_3 + \text{rank}$	Fisher + A-IID	Tannaka
Uses I1?	✓	✗	✗
Uses F1?	✓	✗	✓
Uses F5?	✓	✗	✗
Uses I3?	✗	✓	✗
Uses A-IF?	✗	✓	✗
Uses A-IID?	✗	✓	✗
Key thm	IV.3	IV.10	IV.15
Output	$G = SU(3)$	$N = 3$	$SU(3)$ from stella

[†]Path C is a consistency check, not an independent derivation (see Section IV C).

chinery (topology vs. information geometry)—rather than fully independent.

E. Dimension-Color Correspondence

With $SU(3)$ established by convergent derivation paths, we can extract a structural relationship between the number of colors and spacetime dimensionality.

Theorem IV.21 (Dimension-Color Correspondence). *For confining $SU(N)$ with $N \geq 3$, the emergent spacetime dimension is $D = N + 1$, conditional on the temporal-dimension result of Paper 2 (Theorem 2.2.2). The spatial part $D_{\text{space}} = N$ is derived entirely within Paper 1.*

Proof. The spatial dimension decomposes into three independently derived contributions.

Angular dimensions ($N - 1$). The $SU(N)$ Cartan subalgebra has dimension $\text{rank}(SU(N)) = N - 1$. By Lemma II.9, weight space provides exactly $N - 1$ independent spatial directions in which the Weyl group acts faithfully.

Radial dimension (+1). Proposition II.10 establishes that confinement contributes exactly one radial direction within the geometric realization framework (Figure 2): the confinement radius $R_{\text{conf}} = \hbar c / \sqrt{\sigma}$ is a dynamical degree of freedom orthogonal to the kinematic weight space. The upper bound of one radial direction rests on the coupling-to-dimension framework axiom (Remark II.11): $SU(N)$ has a single gauge coupling, producing a single confinement scale via dimensional transmutation, which maps to at most one embedding direction beyond weight space.

Temporal dimension (+1). Phase evolution $\chi_c(x, \lambda) = v_c(x)e^{i\omega\lambda}$ introduces an internal evolution parameter λ that becomes physical time t . **Forward dependency:** the derivation that $\lambda \rightarrow t$ is carried out in Paper 2, Theorem 2.2.2; here we assume the result.

Total: $D = (N-1)+1+1 = N+1$. Combined with $D = 4$ (Theorem III.2), this uniquely selects $N = 3$, hence $SU(3)$. Note that the spatial-only constraint $D_{\text{space}} = N = D-1$ already restricts to $N = 3$ given $D = 4$, independent of the temporal derivation. $\square$

Remark IV.22 (Derivation vs. Selection). *The correlation $D = N + 1$ is an output, not an input. The logical chain is: geometry $\rightarrow \mathbb{Z}_3 \rightarrow SU(3)$, not $SU(3) \rightarrow \mathbb{Z}_3 \rightarrow$ geometry. The $\mathbb{Z}_3$ symmetry is derived from pure polyhedral geometry before any reference to Lie theory.*

Remark IV.23 (Scope of $D = N + 1$). *The formula applies only to confining $SU(N)$. The non-confining gauge factors $SU(2)_L$ and $U(1)_Y$ do not independently generate spacetime dimensions—they inherit the $D = 4$ structure generated by $SU(3)_C$.*

Remark IV.24 (Affine Kac–Moody Connection). *The temporal dimension admits a suggestive representation-theoretic interpretation. The affine Kac–Moody algebra $\widehat{\mathfrak{su}(N)}$ has $\text{rank} = \text{rank}(\mathfrak{su}(N)) + 1 = N$, with the additional direction corresponding to the loop parameter $z = e^{i\omega\lambda}$. For $SU(3)$: $\text{rank}(\widehat{\mathfrak{su}(3)}) = 3$, matching the three spatial dimensions ($N-1 = 2$ angular +1 radial). If the loop parameter can be identified with the internal time λ , this would upgrade the temporal dimension from a physical hypothesis to a consequence of representation theory. This remains a promising direction for future work; the current derivation establishes $D_{\text{temporal}} = 1$ via the phase evolution structure (Paper 2, Theorem 2.2.2).*

V. UNIQUENESS OF THE STELLA OCTANGULA

The preceding sections established two results: space-time has $D = 4$, and $SU(3)$ is the unique gauge group compatible with geometric realization in three spatial dimensions. The derivation now arrives at its central geometric question: *what object encodes $SU(3)$?*

The answer is a shape known since Kepler—the stella octangula, the compound of two interpenetrating tetrahedra. One tetrahedron T_+ carries the three fundamental color charges; the other T_- carries their anticolors. The point reflection exchanging T_+ and T_- is charge conjugation; the permutation symmetry of each tetrahedron’s vertices is the Weyl group S_3 of $SU(3)$. The stella is not chosen from a catalog of polyhedra—it is the unique minimal object satisfying the geometric realization conditions (GR1)–(GR3). This section proves that uniqueness.

A. The Stella– $SU(3)$ Correspondence

Theorem V.1 ($SU(3)$ –Stella Octangula Isomorphism). *Let T_+ be a regular tetrahedron with centroid at the origin and $T_- = -T_+$. There exists an S_3 -equivariant bijection $\phi : \text{Vertices}(T_+ \cup T_-) \rightarrow \mathfrak{h}^* \cup \{0, 0\}$ such that:*

1. **6 color vertices:** $\phi|_{\{v_R, v_G, v_B\} \subset T_+}$ bijects onto the fundamental weights $\{\vec{w}_R, \vec{w}_G, \vec{w}_B\}$, and $\phi|_{\{v_R, v_G, v_B\} \subset T_-}$ bijects onto $\{-\vec{w}_R, -\vec{w}_G, -\vec{w}_B\}$.
2. **2 apex vertices:** $\phi(v_W) = \phi(v_{\bar{W}}) = \vec{0}$ (color-singlet direction).
3. **Weyl equivariance:** $\text{Stab}_{S_4}(v_W) \cong S_3 \cong W(SU(3))$, and the map $\Phi : \text{Stab}_{S_4}(v_W) \rightarrow W(SU(3))$ is a group isomorphism intertwining the vertex permutation action with the Weyl reflection action on weights.

Proof. Step 1: Weight vector identification. Using Dynkin normalization (Eq. (3)), the fundamental **3** has weight vectors $\vec{w}_R, \vec{w}_G, \vec{w}_B$ forming a triangle in the 2D weight space $\mathfrak{h}^* \cong \mathbb{R}^2$. This triangle is equilateral in the Killing form metric $g_{ij}^K = \frac{1}{12}\delta_{ij}$ (all Killing distances equal $1/\sqrt{3}$), though it appears isosceles in naive (T_3, Y) Euclidean coordinates. The antifundamental **$\bar{3}$** has weights $-\vec{w}_c$, forming the reflected triangle.

Step 2: Root–weight correspondence. Weight differences are root vectors: $\vec{w}_R - \vec{w}_G = \alpha_1$, $\vec{w}_G - \vec{w}_B = \alpha_2$, $\vec{w}_R - \vec{w}_B = \alpha_1 + \alpha_2$. This identifies the 6 edges of the weight hexagram with the 6 roots of A_2 , providing the algebraic bridge between stella geometry and the root system. Color neutrality $\vec{w}_R + \vec{w}_G + \vec{w}_B = 0$ is automatic—color-neutral combinations map to the center of the weight diagram.

Step 3: 6+2 vertex structure. The stella has 6 color vertices in the weight plane bijecting with $\{\vec{w}_c, \vec{w}_{\bar{c}}\}$, plus 2 apex vertices $(v_W, v_{\bar{W}})$ on the perpendicular axis. The apex axis is the unique S_3 -invariant direction in $\mathbb{R}^3$; the apex vertices project to the origin under $\pi : \mathbb{R}^3 \rightarrow \mathfrak{h}^*$.

Step 4: Weyl group equivariance. The Weyl generators s_1 (reflecting across the α_1 -hyperplane, swapping $R \leftrightarrow G$) and s_2 (reflecting across the α_2 -hyperplane, swapping $G \leftrightarrow B$) correspond to the two generating reflections of the tetrahedron T_+ that fix the apex. These generate $\text{Stab}_{S_4}(v_W) \cong S_3$. The map $\Phi : \sigma_k \mapsto s_k$ is verified to be a group isomorphism (well-defined, homomorphism by direct computation, injective since $|S_3| = 6$ equals the stabilizer order, hence surjective). $\square$

Remark V.2 (Apex–Cartan Correspondence). *The 2 apex vertices $v_W, v_{\bar{W}}$ project to weight $\vec{0}$ —the same weight carried by the 2 zero-weight states of the adjoint **8** (the neutral gluons g_3, g_8 , spanning the Cartan subalgebra). This gives the structural correspondence 2 apex vertices $\leftrightarrow$ 2 Cartan generators $\leftrightarrow \text{rank}(SU(3))$. The match is specific to $SU(3)$: a stella-type compound of two N -simplices always has exactly 2 apices, while $\text{rank}(SU(N)) = N-1$; the equality $2 = N-1$ holds only for $N = 3$. This*

is a structural (counting) correspondence; the dynamical content—how geometric structure gives rise to gauge field dynamics—requires the full construction of Theorems V.6 and V.1.

Remark V.3 (The Metric Subtlety). The “equilateral triangle” property of fundamental weights is a statement about the Killing form, not about Euclidean distance in a chosen coordinate system. In (T_3, Y) coordinates the weight triangle has vertices at $(\frac{1}{2}, \frac{1}{3})$, $(-\frac{1}{2}, \frac{1}{3})$, $(0, -\frac{2}{3})$ with Euclidean side lengths

$$|\vec{w}_R - \vec{w}_G|^2 = 1, \quad |\vec{w}_G - \vec{w}_B|^2 = |\vec{w}_B - \vec{w}_R|^2 = \frac{5}{4} \quad (9)$$

—an isosceles triangle. In the Killing metric $g_{ij}^K = B(H_i, H_j) = 6 \text{Tr}(H_i H_j) = 12 \delta_{ij}$ (with Cartan generators $H_1 = \text{diag}(1, -1, 0)$, $H_2 = \text{diag}(1, 1, -2)/\sqrt{3}$), all three Killing distances coincide:

$$|\vec{w}_i - \vec{w}_j|_K^2 = \frac{1}{3} \quad \text{for all } i \neq j. \quad (10)$$

This can also be seen without explicit computation: the weight differences $\vec{w}_R - \vec{w}_G = \alpha_1$, $\vec{w}_G - \vec{w}_B = \alpha_2$, $\vec{w}_B - \vec{w}_R = -(\alpha_1 + \alpha_2)$ are roots of the simply-laced A_2 system, so they share a single Weyl orbit and hence a single Killing norm. The isomorphism theorem holds because the relevant metric is the Killing form—which is derived from $\text{SU}(3)$ (Theorem V.6), not imposed.

Remark V.4 ($\mathbb{Z}_3$ Visibility: $\text{SU}(3)$ vs. $\text{PSU}(3)$). The isomorphism of Theorem V.1 encodes the full gauge group $\text{SU}(3)$, not the adjoint quotient $\text{PSU}(3) = \text{SU}(3)/\mathbb{Z}_3$. The distinction is that the stella’s six color vertices correspond to weights of the fundamental $\mathbf{3} \oplus \bar{\mathbf{3}}$, on which the center $Z(\text{SU}(3)) \cong \mathbb{Z}_3$ acts non-trivially: $z_k : \chi_c \mapsto \omega^k \chi_c$. This center action is geometrically visible—the center element $z_1 = \omega I$ cyclically permutes the color phases $(0, 2\pi/3, 4\pi/3) \mapsto (2\pi/3, 4\pi/3, 0)$, corresponding to a 120° rotation of T_+ .

By contrast, the adjoint $\mathbf{8}$ is center-blind ($A_\mu \rightarrow z_k A_\mu z_k^\dagger = A_\mu$). A geometry encoding only the adjoint would not distinguish $\text{SU}(3)$ from $\text{PSU}(3)$. The $\mathbb{Z}_3$ visibility has three physical consequences: (i) it establishes $\mathbb{Z}_3$ superselection rules distinguishing quarks, antiquarks, and gluons by triality; (ii) it governs the confinement/deconfinement phase transition via center symmetry breaking; (iii) it constrains the Strong CP θ -parameter structure, since $\mathbb{Z}_3$ superselection rules restrict the physical observables to be $\mathbb{Z}_3$ -invariant, providing geometric constraints on the θ -vacuum (Proposition 0.0.5a in the supplementary proofs). This last point is particularly notable: the Strong CP problem (why $\theta \approx 0$) is typically addressed dynamically (axion mechanism); the geometric constraint from $\mathbb{Z}_3$ visibility offers an independent, kinematic perspective.

Remark V.5 (Q_3 Spectral Structure). The stella’s vertex structure admits a second graph-theoretic characterization beyond the weight–vertex bijection. Each vertex of T_+ has




FIG. 8: $\mathbb{Z}_3$ rotational symmetry of the stella octangula. (a) The $[111]$ body diagonal (gold arrow) is a three-fold rotation axis; the three color vertices v_R, v_G, v_B on T_+ permute cyclically under 120° rotations. (b) Projection onto the plane perpendicular to $[111]$: the color vertices form an equilateral triangle, and the singlet vertex v_W lies at the rotation center. This geometric $\mathbb{Z}_3$ is the origin of the $\mathbb{Z}_3$ center of $\text{SU}(3)$.

exactly three cross-nearest neighbors on T_- (at distance $2/\sqrt{3}$), and vice versa. These 12 cross-nearest edges define an adjacency graph on the 8 vertices of $\partial\mathcal{S}$ that is isomorphic to the 3-dimensional hypercube graph Q_3 : the vertices biject with binary strings in $\{0, 1\}^3$, with edges connecting strings at Hamming distance 1.

The graph Laplacian of Q_n has eigenvalues $2k$ with multiplicity $\binom{n}{k}$. For Q_3 the nonzero eigenvalues are $\{2, 2, 2, 4, 4, 4, 6\}$, giving ratios $\{1, 1, 1, 2, 2, 2, 3\}$. The 3-fold degeneracy of each level reflects the T_2 irreducible representation of the tetrahedral symmetry group T_d , corresponding to the three coordinate axes of the embedding cube. This is a standard result of algebraic graph theory; the integers $\{2, 3\}$ happen to coincide with the stella’s construction numbers (2 tetrahedra, $\mathbb{Z}_3$ symmetry order), but the coincidence is arithmetic— Q_4 would produce ratios $\{1, 2, 3, 4\}$, where 4 is composite. What is specific to the framework is the isomorphism $Q_3 \cong \text{CrossAdj}(\partial\mathcal{S})$ itself: the stella is the unique polyhedron selected by $\mathbb{Z}_3$ crystallization (Proposition 0.0.3a), and the Q_3 Laplacian spectrum is a consequence of that selection.

Theorem V.6 (Metric from Killing Form). The Killing form κ of $\text{SU}(3)$, restricted to the Cartan subalgebra, induces a Euclidean metric on weight space that extends uniquely to $3D$.

Proof. The proof proceeds in six steps.

Step 1: Killing form on weight space. The Killing form is $\kappa(X, Y) = 2N \text{Tr}(XY) = 6 \text{Tr}(XY)$ for $\text{SU}(3)$ (dual Coxeter number $h^\vee = 3$). On the Cartan subalgebra, the induced metric on 2D weight space is $g_{ij}^K = \frac{1}{12} \delta_{ij}$.

Step 2: Third direction from rank and embedding dimension. The 6 non-zero weights of $\text{SU}(3)$ span a 2D plane (rank = 2). Proposition II.10 establishes that the minimal embedding dimension is $d = 3$: any geometric realization

satisfying (GR3) requires a third direction perpendicular to the weight plane to separate the representation from its conjugate. This step uses only the rank of $SU(3)$ and the embedding dimension bound, not any specific polyhedral realization (stella uniqueness is proved independently in Theorem V.10 below).

Step 3: General form under S_3 Weyl symmetry. Any 3D metric compatible with g^K has the form $ds^2 = f(r, \theta) dr^2 + 2g_i(r, \theta) dr d\theta^i + h_{ij}(r, \theta) d\theta^i d\theta^j$. The Weyl group S_3 acts on weight-space coordinates θ^i by reflections and 120° rotations. The unique S_3 -invariant symmetric 2-tensor on $\mathbb{R}^2$ is proportional to δ_{ij} , giving $h_{ij} = h(r) g_{ij}^K$. Cross terms g_i vanish by radial isotropy (no preferred radial direction), and $f = f(r)$ only.

Step 4: Smoothness at $r = 0$ forces $h(r) = r^2$. For the metric to be C^∞ at the origin, the angular part must scale as r^2 . If $h(r) \sim r^\alpha$ with $\alpha \neq 2$, the total angle around a small circle at radius r scales as $r^{\alpha/2-1}$, which diverges ($\alpha < 2$) or vanishes ($\alpha > 2$) as $r \rightarrow 0$ —producing a conical singularity. Only $\alpha = 2$ yields the correct 2π periodicity for a smooth manifold.

Step 5: Coordinate freedom. Reparametrizing $r \mapsto \tilde{r}(r)$ sets $f(r) = 1$.

Step 6: Flatness verification. The resulting metric $ds^2 = dr^2 + r^2 g_{ij}^K d\theta^i d\theta^j$ is, in Cartesian coordinates, $ds^2 = dx^2 + dy^2 + dz^2$. Direct computation confirms the Riemann tensor vanishes: $R_{ijkl} = 0$ —the metric is Euclidean. $\square$

Remark V.7 (Euclidean Space as Derived). *The logical order is reversed from convention: $SU(3)$ is derived first, and the Euclidean metric on physical space emerges from its Killing form. The geometry we experience is a consequence of the gauge group’s algebraic structure.*

Corollary V.8 (Compactness Forces Euclidean Metric). *The Euclidean structure is not special to $SU(3)$; it follows from compactness of the gauge group. For any compact simple Lie group, the Killing form is negative-definite: $B(X, X) < 0$ for all $X \neq 0$, so the induced metric $\langle \cdot, \cdot \rangle = -B^{-1}$ is positive-definite (Euclidean). Non-compact groups such as $SU(2, 1)$ or $SL(2, \mathbb{R})$ have indefinite Killing forms and would induce non-Euclidean (hyperbolic or Lorentzian) metrics on weight space. This provides a selection principle: only compact gauge groups are consistent with Euclidean spatial geometry.*

Remark V.9 (Global Flatness). *The resulting metric has trivial holonomy: $\text{Hol}(g) = \{I\}$, since $g_{ij} = \frac{1}{12}\delta_{ij}$ is constant in Cartesian coordinates, all Christoffel symbols vanish ($\Gamma_{jk}^i = 0$), and the Riemann tensor vanishes identically ($R_{ijkl} = 0$). This confirms global Euclidean geometry—any vector parallel-transported around any closed loop returns unchanged—not merely local flatness.*

B. Uniqueness Theorems

Theorem V.10 (Stella Uniqueness). *Among all polyhedral complexes satisfying (GR1)–(GR3), the stella octangula is the unique minimal realization of the $SU(3)$ weight structure.*

Theorem V.11 (Completeness of Classification). *The stella octangula is the unique minimal geometric realization of $SU(3)$ among all topological spaces satisfying (GR1)–(GR3)—including non-convex polyhedra, infinite structures, and fractals.*

Proof. Step 1: Vertex count from (GR1). $\mathbf{3}$ has 3 weights, $\mathbf{\bar{3}}$ has 3, plus 2 apex vertices: exactly 8.

Step 2: Symmetry from (GR2). Among 8-vertex polyhedra, only the cube and stella have $S_3 \subset \text{Aut}$. The cube’s full symmetry group $O_h \cong S_4 \times \mathbb{Z}_2$ (order 48) does contain S_3 as a subgroup, but the S_3 acts by permuting *body diagonals* (of which there are 4), not by permuting 3 color weights. No labeling of the 8 cube vertices by the 6 non-zero weights of $\mathbf{3} \oplus \mathbf{\bar{3}}$ plus 2 apex labels is compatible with an S_3 subgroup of O_h that simultaneously reproduces the Weyl reflections on weight space and respects the $\mathbf{3}/\mathbf{\bar{3}}$ partition (GR1, GR3).

Step 3: Involution from (GR3). The stella realizes charge conjugation as $T_+ \leftrightarrow T_-$.

Step 4: Exhaustive elimination.

TABLE V: Systematic elimination of candidate polyhedra.

Candidate	$ V $	Failure	Rule
Two 2D triangles	6	No radial direction	(M2)
Octahedron	6	Can’t separate $\mathbf{3}/\mathbf{\bar{3}}$	(GR1)
Triangular prism	6	No antipodal involution	(GR3)
Cube	8	$S_3 \subset O_h$ incompatible with weight labeling	(GR2)
Separate tetrahedra	8	Non-intersecting; no single polyhedral complex	(GR3)
Stella	8	None	✓

Step 5: Vertex bound from representation theory. The $\mathbf{3} \oplus \mathbf{\bar{3}}$ representation has *non-degenerate* weights: each of the 6 non-zero weights $\{\pm\vec{w}_R, \pm\vec{w}_G, \pm\vec{w}_B\}$ has multiplicity exactly 1, and the zero weight does not appear (unlike the adjoint $\mathbf{8}$, where $\vec{0}$ has multiplicity 2). A faithful weight encoding (GR1) therefore requires exactly one vertex per non-zero weight. Combined with at most 2 apex vertices for the 3D embedding (Proposition II.10):

$$|\mathcal{V}(\mathcal{X})| \leq 6 + 2 = 8 \quad (11)$$

This bound—a direct consequence of the finite-dimensionality and non-degeneracy of $\mathbf{3} \oplus \mathbf{\bar{3}}$ —excludes *all*

infinite structures in a single stroke: periodic lattices, aperiodic tilings, infinite graphs, and fractals (both countably and uncountably infinite vertex sets violate Eq. (11)).

Step 6: Non-convex polyhedra. All four Kepler-Poinsot star polyhedra have ≥ 12 vertices (MIN1). Among the 57 non-convex uniform polyhedra [22], only the tetrahemihexahedron has fewer than 8 vertices (it has 6). It fails (GR2): its symmetry group $T_d \cong S_4$ admits a surjection $S_4/V_4 \cong S_3$, but the 180° rotation about the $(1, 1, 0)$ axis maps $\vec{w}_B \rightarrow -\vec{w}_B$, requiring $B \rightarrow \bar{B}$ —which is not a color permutation in S_3 . No weight labeling satisfies (GR2) and (GR3) simultaneously. Any self-intersecting 8-vertex polyhedron satisfying (GR1)–(GR3) is combinatorially equivalent to the stella: the vertex positions and pairing structure are uniquely determined by the weight constraints and S_3 equivariance (Theorem V.1).

Step 7: Exotic spaces. Quasicrystals are doubly excluded: infinite vertices and icosahedral A_5 symmetry (A_5 is simple, admitting no surjection to S_3). Non-Hausdorff spaces are excluded definitionally (Definition II.1 requires embedding in $\mathbb{R}^n$, which is Hausdorff). Continuous manifolds of positive dimension are incompatible with the finite polyhedral complex requirement; extracting a finite vertex subset reduces to Steps 1–4. CW complexes with positive-dimensional cells reduce to their 0-skeleton, which is then handled by the finite or infinite case above. $\square$

Remark V.12 (Convergent Uniqueness Criteria). *The stella is selected by three independent criteria that converge on the same structure: (1) Minimality (MIN1–MIN3): fewest vertices, lowest embedding dimension, simplest topology. (2) Maximal symmetry: among all 8-vertex polyhedra satisfying (GR1)–(GR3), the stella has the largest point symmetry group O_h (order 48); the cube shares O_h but fails (GR1). (3) Root lattice compatibility: the stella’s edge vectors encode exactly the A_2 root system with no spurious edges—a structural criterion independent of minimality. The convergence of three independent selection principles on a single structure provides evidence that the stella is structurally distinguished, not merely a conventional choice.*

Remark V.13 (Sensitivity Analysis: Relaxing (GR3)). *Relaxing the chirality axiom (GR3) does not enlarge the space of valid realizations—it empties it. Unlike $SU(2)$, whose fundamental $\mathbf{2}$ is pseudo-real ($\mathbf{2} \cong \bar{\mathbf{2}}$), $SU(3)$ has a complex fundamental: $\mathbf{3} \not\cong \bar{\mathbf{3}}$. Without (GR3), no geometric involution distinguishes $\mathbf{3}$ from $\bar{\mathbf{3}}$, and the embedding degenerates to $\mathbf{3} \oplus \mathbf{3}$ rather than the required $\mathbf{3} \oplus \bar{\mathbf{3}}$. A single tetrahedron (4 vertices) cannot encode the 6-element A_2 root system needed for (GR2), nor can it represent $\mathbf{3} \oplus \bar{\mathbf{3}}$ (mesons require both representations). Thus (GR3) is a necessary consequence of the complex nature of $SU(3)$ ’s fundamental representation.*

Remark V.14 (Computational Crystallization). *The algebraic uniqueness of Theorems V.10–V.11 receives independent support from a complementary dynamical argument. Consider N charged particles on S^2 interacting*

via $\mathbb{Z}_3$ product-rule repulsion, with coupling coefficient α for same-charge pairs and β for conjugate-charge pairs. For $\alpha/\beta \geq 2$ —a ratio fixed by the $SU(3)$ Casimir ratio $C_F(\mathbf{3})/C_F(\mathbf{8}) = 2$ —the system crystallizes into the stella octangula from random initial conditions with 100% convergence (RMSD < 0.02 to the ideal stella across all seeds tested).

This dynamical selection extends beyond geometry. Grand canonical annealing from $N = 20$ particles selects $N = 8$ with zero variance; label relaxation from every initial partition (1+7 through 7+1) converges to 4+4 with 100% success. The vertex count and matter–antimatter partition are thus not inputs but outputs of $\mathbb{Z}_3$ dynamics on the sphere. Furthermore, replacing the hard sphere constraint with soft radial normalization, the spherical shell and stella geometry emerge simultaneously—the two phenomena are independent, as confirmed by a two-dimensional parameter sweep.

Among cyclic groups $\mathbb{Z}_n$, $\mathbb{Z}_3$ is the minimal group producing the stella without trivial escape routes: $\mathbb{Z}_2$ fails (a single non-trivial charge cannot split into conjugate pairs), $\mathbb{Z}_4$ is unstable (its self-conjugate charge creates a competing ground state, reducing convergence to 70%), and $\mathbb{Z}_{5+}$ succeed but carry redundant unused charges.

The top-down (algebraic) and bottom-up (dynamical) arguments thus converge: Theorems V.10–V.11 show the stella is the only polyhedral realization of $SU(3)$; the crystallization result shows it is the inevitable ground state of $\mathbb{Z}_3$ interactions. Full computational details, including Fisher non-degeneracy analysis and number-field selection, are in the [supplementary proof document](#).

Definition V.15 (Stella Octangula Boundary Topology). *The boundary of the stella octangula is the disjoint union of the surfaces of its two constituent tetrahedra:*

$$\partial S := \partial T_+ \sqcup \partial T_- \quad (12)$$

Each $\partial T_\pm \cong S^2$ is a closed 2-sphere, so ∂S consists of two connected components. The topological invariants are:

Property	Stella ($T_+ \sqcup T_-$)	Octahedron
Vertices	8 (4 + 4)	6
Edges	12 (6 + 6)	12
Faces	8 (4 + 4)	8
Components	2	1
Euler char. χ	4 (two S^2)	2

The distinction is physically consequential: $\chi = 4$ encodes the $\mathbf{3}/\bar{\mathbf{3}}$ separation (matter/antimatter) and is required by (GR3). An octahedron ($\chi = 2$) has a single connected boundary that cannot geometrically separate T_+ from T_- .

Remark V.16 (Scope of Geometric Determination). *The stella determines kinematic content: confinement criterion ($\mathbb{Z}_3$ center), N -ality classification, hadron structure, color factors, and β -function form. It does not determine dynamical content: the linear potential $V(r) = \sigma r$, string*




FIG. 9: The stella octangula: two interpenetrating tetrahedra encoding SU(3) symmetry. Matter tetrahedron T_+ (blue solid) represents $\mathbf{3}$; antimatter T_- (red dashed) represents $\bar{\mathbf{3}}$. Green arrow: singlet axis with apex vertices $W, \bar{W}$.

tension value, deconfinement temperature, or numerical coupling constant—these require the chiral field dynamics of Paper 2. The division is illustrated by the Cornell potential $V(r) = -C_F \alpha_s / r + \sigma r$:

Component	Origin	Geometric?
Color factor $C_F = 4/3$	SU(3) Casimir $(N_c^2 - 1)/(2N_c)$	✓
Coulomb form $1/r$	Gauge propagator + Fourier	✓
$\mathbb{Z}_3$ confinement criterion	Center symmetry $\langle P \rangle = 0$	✓
N -ality classification	$k = (\#q - \#\bar{q}) \bmod 3$	✓
Coupling $\alpha_s(\mu)$	RG evolution with Λ_{QCD}	✗
Linear form σr	Non-perturbative QCD dynamics	✗
String tension σ	Lattice/phenomenology input	✗
Deconfinement T_c	Finite-temperature lattice QCD	✗

The geometry provides the symmetry arena (which states are confined); the dynamics (how the confining potential arises) require Paper 2. Since the stella is the unique geometric realization of SU(3) (Theorem V.11), the 19-order-of-magnitude hierarchy $R_{\text{stella}}/\ell_P \sim 10^{19}$ between the QCD and Planck scales is topologically determined—it cannot arise from any other geometry.

VI. SPATIAL EXTENSION FROM THE HONEYCOMB

The results so far establish the stella octangula as a local encoding of SU(3). But physics requires an extended spatial arena—multiple stellae must coexist to describe multi-hadron systems, scattering processes, and spatially

separated observers. The question is whether this extension is uniquely determined or introduces new geometric freedom. We show that the FCC lattice, and with it extended $\mathbb{R}^3$, is derived from SU(3) representation theory rather than assumed.

Two concepts are central to what follows. A graph is *vertex-transitive* if its automorphism group acts transitively on vertices: every vertex “looks the same” combinatorially. This ensures that no lattice site is physically privileged—a discrete analogue of spatial homogeneity.

Definition VI.1 (SU(3) Phase Coherence). *A lattice graph Γ realizes SU(3) phase coherence if every vertex hosts a stella octangula whose $\mathbb{Z}_3$ color phases $(0, 2\pi/3, 4\pi/3)$ are consistently assigned across adjacent cells—that is, shared vertices carry the same color label in every stella to which they belong.*

Proposition VI.2 (A_3 Lattice from Physical Requirements). *Among rank-3 root lattice extensions of the A_2 root lattice, A_3 (the FCC lattice) is uniquely determined by physical requirements.*

Proof. The proof proceeds in four parts.

(a) *Third dimension required.* Proposition II.10 establishes $d_{\text{embed}} = 3$. The A_2 root lattice is 2-dimensional; spatial extension requires a rank-3 lattice containing A_2 as a sublattice.

(b) *Perpendicular direction fixed.* The stella’s apex-to-apex axis (the color-singlet direction) defines the unique S_3 -invariant perpendicular to the weight plane. This axis is the $[111]$ direction of the embedding lattice.

(c) *Stacking pattern determined.* Two stacking patterns extend A_2 layers into 3D: FCC (ABCABC..., three distinct layer positions before repeating) and HCP (ABAB..., two alternating positions). HCP breaks vertex-transitivity—sites in A-layers have different local environments from B-layers—violating the SU(3) phase coherence condition. Only FCC (the A_3 lattice) is vertex-transitive.

(d) *Elimination of B_3 and C_3 .* The rank-3 root lattices are A_3 , B_3 , and C_3 . B_3 ($\mathbb{Z}^3$, coordination 6) fails: 6 neighbors cannot accommodate the 12 root vectors of A_2 . C_3 produces the same point lattice as A_3 but is not simply-laced (unequal root lengths), breaking S_3 Weyl symmetry. Only A_3 (coordination 12, simply-laced) survives. A critical distinction: the physical lattice corresponds to the root lattice Q (generated by gauge interaction vectors), not the weight lattice P . For B_3 : $Q(B_3) = \mathbb{Z}^3$ (coordination 6) while $P(B_3) = \text{BCC}$ (coordination 8); for C_3 : $Q(C_3) = \text{FCC}$ (coordination 12) while $P(C_3) = \mathbb{Z}^3$ (coordination 6). This distinction is essential—using weight lattices instead of root lattices would change the elimination.

(e) *Reducible extensions excluded.* Reducible rank-3 root lattices such as $A_2 \oplus A_1$ are excluded by two independent requirements: vertex-transitivity (Theorem VI.4) demands that every site has an identical local environment, but $A_2 \oplus A_1$ has inequivalent in-plane and out-of-plane directions; and exact phase coherence across the lattice requires crystallographic symmetry, excluding non-crystallographic extensions. $\square$

Theorem VI.3 (Adjacency from $SU(3)$). *Let Γ be a vertex-transitive graph realizing $SU(3)$ phase coherence. The combinatorial structure of Γ is uniquely determined by $SU(3)$ representation theory:*

- (a) **12-regularity:** every vertex has exactly 12 neighbors.
- (b) **No intra-representation triangles:** there are no 3-cycles with all edges within a single representation type.
- (c) **4 four-cycles per edge:** through each edge, exactly 4 independent 4-cycles pass.
- (d) **O_h symmetry:** $\text{Aut}(\Gamma) \supseteq O_h$ (order 48).

Proof. Each constraint is derived from A_2 representation theory.

(a) *12-regularity.* The A_2 root system has 6 roots $\Phi = \{\pm\alpha_1, \pm\alpha_2, \pm(\alpha_1 + \alpha_2)\}$. Within each representation ($\mathbf{3}$ or $\bar{\mathbf{3}}$), the 3 weight vertices are connected by 3 root vectors, giving $2 \times 3 = 6$ intra-representation edges. The adjoint decomposition $\mathbf{3} \otimes \mathbf{3} = \mathbf{8} \oplus \mathbf{1}$ provides 6 additional inter-representation connections ($\mathbf{3} \leftrightarrow \bar{\mathbf{3}}$). Total: $6 + 6 = 12$ nearest neighbors.

(b) *No intra-representation root triangles.* Although three roots can sum to zero in A_2 (e.g., $\alpha_1 + \alpha_2 + (-\alpha_1 - \alpha_2) = 0$), such triples necessarily mix positive and negative roots. Three positive roots cannot sum to zero: $\alpha_1 + \alpha_2 + (\alpha_1 + \alpha_2) = 2(\alpha_1 + \alpha_2) \neq 0$, and similarly for three negative roots. Physically, $\mathbf{3} \otimes \mathbf{3} = \mathbf{6}_A \oplus \bar{\mathbf{3}}$ contains no singlet: three quarks of the same triality cannot form a color-neutral bound state via gluon exchange. Hence no purely intra-representation 3-cycle exists.

(c) *4 four-cycles per edge.* Fix an edge with root direction $\alpha \in \Phi$. A 4-cycle through this edge requires a root β with $\alpha + \beta \neq 0$ and $\alpha - \beta \neq 0$. For $\alpha = \alpha_1$, the valid choices are $\beta \in \{\alpha_2, -\alpha_2, \alpha_1 + \alpha_2, -(\alpha_1 + \alpha_2)\}$ —exactly 4. By Weyl symmetry the count is the same for all root directions.

(d) *O_h symmetry.* $O_h \cong S_4 \times \mathbb{Z}_2$, where S_4 permutes the 4 body diagonals of the embedding cube and $\mathbb{Z}_2$ is spatial inversion. The Weyl group $W(A_2) \cong S_3$ embeds into S_4 as the stabilizer of one body diagonal. Charge conjugation ($\mathbf{3} \leftrightarrow \bar{\mathbf{3}}$) corresponds to the $\mathbb{Z}_2$ inversion factor. $\square$

Theorem VI.4 (Honeycomb Uniqueness). *Among vertex-transitive tilings of $\mathbb{R}^3$ using regular tetrahedra and octahedra, the tetrahedral-octahedral honeycomb uniquely embeds stellae with phase coherence.*

Proof. Stage 1: Constraints are derived. Theorem VI.3 derives all four combinatorial constraints (a)–(d) from $SU(3)$ representation theory alone.

Stage 2: $A_2 \subset A_3$ embedding. The 2D weight space of $SU(3)$ embeds in 3D physical space via the root lattice inclusion $A_2 = \{(x_1, x_2, x_3) \in \mathbb{Z}^3 : x_1 + x_2 + x_3 = 0\} \subset A_3$. The full FCC lattice consists of parallel A_2 planes at

integer spacing along the $[111]$ direction—the unique S_3 -invariant perpendicular to the weight plane (defined by the stella’s apex axis).

Stage 3: Uniqueness among rank-3 root lattices. Proposition VI.2(d) eliminates B_3 (coordination too low) and C_3 (not simply-laced). Only A_3 survives.

Stage 4: Vertex-transitivity is necessary. Conway, Jiao, & Torquato [23] found non-vertex-transitive tilings that fail the $SU(3)$ phase coherence condition: sites in inequivalent positions have different local environments, violating color democracy (S_3 invariance at every lattice point). $\square$

Remark VI.5 (Phase Matching as Flat Connection). *The honeycomb construction automatically satisfies a gauge consistency condition. Color phases assigned at vertices match across shared triangular faces via barycentric interpolation: both adjacent cells see the same vertex colors and compute identical interpolated phases. This interpolation is well-defined because the color phases live in the commuting Cartan subalgebra of $SU(3)$ ($[T_3, T_8] = 0$), so linear combination of phase values is algebraically consistent. The matching condition has a gauge-invariant formulation: Wilson loops on all honeycomb faces evaluate to the identity,*

$$W(F) = \mathcal{P} \exp \left(ig \oint_{\partial F} A_\mu^a T^a dx^\mu \right) = \mathbf{1}, \quad (13)$$

implying a flat connection ($F_{\mu\nu}^a = 0$) and trivial holonomy. This is a statement about the vacuum sector—gauge-invariant and not a gauge-fixing artifact.

Remark VI.6 (Octahedral Cells as Color-Neutral Transition Regions). *Each octahedral cell of the honeycomb has 6 vertices colored $\{R, \bar{R}, G, \bar{G}, B, \bar{B}\}$, with opposite vertices carrying conjugate colors. At the octahedron center, the phase sum $\sum_c e^{i\phi_c} = 1 + \omega + \omega^2 = 0$ yields the same color singlet condition as the stella center (Theorem V.11). Octahedra therefore serve as color-neutral transition regions between the color-charged stellae. Physically, a quark path through the honeycomb traverses a sequence stella $\rightarrow$ octahedron $\rightarrow$ stella $\rightarrow \dots$, providing a geometric realization of the QCD flux tube. The octahedron size (~ 0.45 fm, set by R_{stella}) is consistent with the observed QCD flux tube transverse width $w \approx 0.35$ fm.*

Remark VI.7 (Emergent 3D Space). *This construction achieves something stronger than previous approaches: (1) FCC lattice sites provide pre-geometric integer labels (n_1, n_2, n_3) ; (2) the 12-regularity and girth constraints are theorems of $SU(3)$, not axioms (distinguishing CG from LQG, causal sets, CDT); (3) phase coherence is definitional; (4) discrete O_h symmetry yields effective $SO(3)$ at large scales. The claim “space emerges from $SU(3)$ ” is now rigorous.*

Remark VI.8 (Combinatorial FCC Identification). *The A_2 root-type subgraph is independently identifiable from parallelogram counts alone: each root-type edge participates in exactly $|\Phi| - 2 = 4$ root parallelograms, uniquely*




FIG. 10: Lattice comparison: from A_2 to A_3 . (a) A_2 root lattice (2D hexagonal, coordination 6) with nearest-neighbor bonds (green) and root vectors (gold arrows). (b) A_3 /FCC lattice (3D, coordination 12)—the unique lattice realizing $SU(3)$ phase coherence; gold vertices highlight one A_2 sublattice plane. (c) B_3 simple cubic lattice (3D, coordination 6)—rejected because it lacks the A_2 sublattice structure required by $SU(3)$.




FIG. 11: From stella octangula to FCC honeycomb. (a) Single stella: T_+ (blue), T_- (red) in a cube. (b) Decomposition: 8 corner tetrahedra + central octahedron. (c) Interlocking neighbors. (d) $3 \times 3 \times 3$ stella grid. (e) Complete FCC tetrahedral-octahedral honeycomb.

identifying A_2 among rank-2 lattices ($A_1 \times A_1$ gives 2, B_2 gives 6, G_2 gives 10). Combined with the O_h requirement

that adjoint-type edges be out-of-plane, this provides a purely combinatorial signature of FCC structure.

VII. CONTINUUM LIMIT FROM DISCRETE STRUCTURE

Three distinct limits connect the discrete encoding to continuous field theory.

TABLE VI: Continuum limits from discrete stella encoding. The $\mathbb{Z}_3$ center survives all limits as a topological invariant.

Limit	Transformation	Preserved
Spatial	$O \rightarrow \text{SO}(3)$, lattice $\rightarrow \mathbb{R}^3$	Euclidean geom.
Gauge	Weights $\rightarrow A_2 \rightarrow \text{SU}(3)$	$\pi_3 = \mathbb{Z}$, $\mathbb{Z}_3$
Thermo.	$V \rightarrow \infty$	θ -vacua




FIG. 12: Continuum limit visualization. (a) The A_2 root system from stella weight differences.

(b) Discrete-to-continuous limit: inner vertices at $a = a_0$, outer ring at $a \rightarrow 0$. The $\mathbb{Z}_3$ center (120° periodicity) is preserved at all scales.

a. Spatial continuum. The FCC lattice (Section VI) provides integer coordinates without a metric. The discrete symmetry group O (24 proper rotations of the cube/octahedron) enhances to $\text{SO}(3)$ as $a \rightarrow 0$ by an effective enhancement mechanism: at length scales $L \gg a$, the leading O -invariant corrections to rotation symmetry appear at angular momentum $\ell = 4$ (the first representation of $\text{SO}(3)$ that splits under restriction to O); lower multipoles $\ell \leq 3$ are automatically $\text{SO}(3)$ -symmetric. Corrections scale as $(a/L)^n$: with $a \approx 2.25 \ell_P$, at nuclear scales $a/L \sim 10^{-20}$, at macroscopic scales $a/L \sim 10^{-35}$.

Remark VII.1 (Parity Breaking: $\text{SO}(3)$, Not $O(3)$). The discrete O_h group contains improper rotations (reflections), yet the continuum limit yields $\text{SO}(3)$, not $O(3)$. The mechanism is intrinsic chirality of the stella octangula: the signed volumes of the two tetrahedra are $V_{\text{signed}}(T_+) = +8/3$, $V_{\text{signed}}(T_-) = -8/3$, so a parity transformation $P: \mathbf{x} \rightarrow -\mathbf{x}$ exchanges $T_+ \leftrightarrow T_-$ (matter $\leftrightarrow$ antimatter). The matter/antimatter distinction

encoded by the two tetrahedra breaks the parity element of O_h , leaving $\text{SO}(3)$ as the surviving spatial symmetry group.

Chirality is encoded at the phase level: the two cyclic orderings $R \rightarrow G \rightarrow B$ (counterclockwise, phases $0 \rightarrow 2\pi/3 \rightarrow 4\pi/3$) and $R \rightarrow B \rightarrow G$ (clockwise) are mathematically distinct and non-equivalent. These connect to the $\text{SU}(3)$ structure constants via $f^{123} = +1$ (counterclockwise) and the Levi-Civita symbol $\epsilon^{123} = +1$ (right-handedness). The selection of one chirality over the other requires a dynamical mechanism: Paper 2 (Theorem 2.2.4) derives that QCD instantons with $\langle Q \rangle > 0$ select the $R \rightarrow G \rightarrow B$ ordering—the same CP violation that produces the matter/antimatter asymmetry also selects the chirality. The emergence of chiral order from geometric frustration has condensed matter precedent: in kagome metals, time-reversal symmetry breaking via charge-loop currents arises from the interplay of three geometrically frustrated sublattices [21].

b. Gauge group determination. The recovery of continuous $\text{SU}(3)$ from discrete stella data proceeds in four steps: (1) weight differences between adjacent color vertices produce the six root vectors of the A_2 root system (Step 2 of Theorem V.1); (2) the root system uniquely determines the Lie algebra $\mathfrak{su}(3)$ via the Cartan matrix; (3) exponentiation of $\mathfrak{su}(3)$ yields the simply connected group $\text{SU}(3)$; (4) the topological invariants $\pi_3(\text{SU}(3)) = \mathbb{Z}$ (instanton classification) and $Z(\text{SU}(3)) = \mathbb{Z}_3$ (center symmetry) survive the limit as algebraic consequences of the root system.

c. $\mathbb{Z}_3$ survival. The center $\mathbb{Z}_3 \cong \Lambda_{\text{coweight}}/\Lambda_{\text{root}}$ depends only on A_2 data. At all levels—discrete stella, continuous $\text{SU}(3)$, and θ -vacua—the same $\mathbb{Z}_3$ acts.

Theorem VII.2 (Lorentz Violation Bounds). The discrete O_h symmetry induces Lorentz violation bounded by

$$\delta c/c \lesssim (E/E_P)^2 \sim 10^{-32} \quad \text{at } E = 1 \text{ TeV} \quad (14)$$

lying 6–17 orders of magnitude below current experimental sensitivity. The angular pattern has **no quadrupole** ($\ell = 2$); the first permitted anisotropy is hexadecapole ($\ell = 4$) with O_h symmetry.

Proof. CPT preservation by the stella forbids odd-power Lorentz violation. The leading operators are dimension-6, suppressed by E_P^{-2} . The O_h character table shows $\ell = 2$ has no A_{1g} component; $\ell = 4$ is the first with A_{1g} :

$$K_4(\hat{n}) = n_x^4 + n_y^4 + n_z^4 - \frac{3}{5} \quad (15)$$

Current bounds (LHAASO [24], DisCan [25], GW170817, SME [26]) are all satisfied with 10^6 – 10^{27} margin. $\square$

Remark VII.3 (Falsifiable Angular Pattern). Detection of $\ell = 2$ (quadrupole) Lorentz violation would falsify the framework, regardless of magnitude. Observation of pure $\ell = 4$ anisotropy would strongly favor discrete O_h symmetry.

A. Continuum Limit as Open Problem

The quantitative bounds of Theorem VII.2 demonstrate *phenomenological viability*—consistency with all current experiments—but they do not constitute a complete proof that exact Poincaré invariance is recovered in the continuum limit. The dynamical continuum limit (proving that exact continuous symmetry emerges from discrete structure as $a \rightarrow 0$) is an open problem shared with all discrete approaches to quantum gravity, including causal sets [27], loop quantum gravity [28], and causal dynamical triangulations [12].

The framework possesses a specific structural advantage: CPT preservation by the stella forbids all odd-power Lorentz-violating operators, giving $(E/E_P)^2$ suppression rather than the $(E/E_P)^1$ of generic lattice approaches. This is a falsifiable feature: detection of dimension-5 Lorentz violation would refute the CPT protection mechanism.

The CG framework has resolved part of this program: the D_4 lattice is self-coarsening ($D_4 \rightarrow D_4$ under blocking), and the exact FCC mass gap $\mu > 0$ provides an infrared regulator for Balaban’s multi-scale renormalization group program [29, 30]. Non-perturbative universality (the lattice-independence of the continuum theory) is now proven: Theorem 7.5.4 establishes that the Balaban RG contraction on both D_4 and $\mathbb{Z}^4$ converges to identical continuum Schwinger functions, closing the gap identified in Theorem 7.6.10. What remains genuinely open is the dynamical emergence of exact continuous symmetries from discrete O_h —a challenge shared with all discrete spacetime programs. The full constructive analysis is developed in Paper 3.

VIII. STANDARD MODEL GAUGE STRUCTURE FROM GEOMETRY

The story so far: observer existence forced $D = 4$; dimensional constraints and geometric realization forced $SU(3)$; the stella octangula emerged as $SU(3)$ ’s unique polyhedral encoding; and tiling space with stellae produced the FCC lattice and, in the continuum limit, smooth $\mathbb{R}^3$. All of this concerns the strong sector. A natural kinematic question remains: where do the electroweak gauge factors $SU(2)_L \times U(1)_Y$ come from? We now show that the polytope embedding chain extending the stella octangula leads, through a combination of geometric necessity and phenomenological selection, to the full Standard Model gauge group.

A. Gauge Unification from Geometric Symmetry

Theorem VIII.1 (Gauge Unification). *The stella octangula’s symmetry structure, extended through the natural polytope embedding chain, geometrically encodes an*

$SO(10)$ grand unification structure containing the Standard Model gauge group $SU(3)_C \times SU(2)_L \times U(1)_Y$. The first four steps of the chain (Stella $\rightarrow$ 16-cell $\rightarrow$ 24-cell $\rightarrow D_4$) are mathematically necessary; the final two ($D_4 \rightarrow D_5$, $SU(5) \rightarrow SM$) invoke a minimality criterion and phenomenological input, respectively.

The embedding chain, with each arrow labelled by its epistemic status:

$$\begin{array}{c} \text{Stella} \xrightarrow[\text{[E]}]{\Phi} \text{16-cell} \xrightarrow[\text{[E]}]{\text{rect.}} \text{24-cell} \xrightarrow[\text{[E]}]{D_4} \\ \xrightarrow[\text{[E+F: min.]}]{D_5 = \mathfrak{so}(10)} \xrightarrow[\text{[E]}]{\text{max}} \mathfrak{su}(5) \xrightarrow[\text{[E+F: pheno.]}]{\text{}} \text{SM} \end{array} \quad (16)$$

1. **Stella $\rightarrow$ 16-cell** [E]: The 16-cell (cross-polytope) is the *unique* regular 4-polytope with exactly 8 vertices (by enumeration of the six regular 4-polytopes; see supplementary [derivation](#), Proposition 2.4.1). Geometrically, the stella octangula is the three-dimensional “shadow” (orthogonal projection) of the 16-cell; the canonical Hadamard lift $\Phi = H_4 \circ \phi$ reverses this projection, mapping the stella’s 8 vertices bijectively to the standard 16-cell vertices $\{\pm e_i\}$. This lift is unique up to $W(B_4)$ symmetry.
2. **16-cell $\rightarrow$ 24-cell** [E]: Rectification—replacing each vertex by the midpoints of its adjacent edges—is a standard geometric operation. The 16-cell has 24 edges, producing exactly 24 midpoint vertices; the resulting polytope is the 24-cell (symmetry group $W(F_4)$, order 1152). This step is deterministic: no choices are made.
3. **24-cell $\leftrightarrow D_4$** [E]: The 24 vertices of the 24-cell coincide (up to rescaling) with the 24 roots of $D_4 = \mathfrak{so}(8)$ —an exact identification, not an analogy. The D_4 Dynkin diagram has an S_3 outer automorphism (*triality*) permuting its three 8-dimensional representations $\mathbf{8}_v, \mathbf{8}_s, \mathbf{8}_c$; this structure is unique among simple Lie algebras.
4. **$D_4 \subset D_5$** [E+F]: D_4 embeds in every D_n ($n \geq 4$) via coordinate inclusion. The extension to $D_5 = \mathfrak{so}(10)$ is *not uniquely* determined by the geometry; D_6, D_7 , etc. are equally valid embeddings. We invoke a **minimality criterion** [F]: choose the smallest D_n that (a) contains the Standard Model gauge algebra as a subalgebra, (b) admits chiral spinor representations ($2^{n-1} \geq 16$ requires $n \geq 5$), and (c) provides automatic anomaly cancellation per generation. The minimal solution is $n = 5$.
5. **$\mathfrak{so}(10) \supset \mathfrak{su}(5) \oplus \mathfrak{u}(1)$** [E]: A standard maximal subalgebra relation [31, 32].
6. **$SU(5) \rightarrow SM$** [E+F]: The maximal subgroup of $SU(5)$ compatible with quark confinement ($SU(3)$ unbroken), electroweak physics, and anomaly cancellation is $SU(3)_C \times SU(2)_L \times U(1)_Y$ —this is unique

given the physical constraints [E]. Requiring those specific constraints as input is the phenomenological selection [F].

a. The Weinberg angle. The SU(5) embedding gives $\sin^2 \theta_W^{\text{GUT}} = 3/8 = 0.375$ —a geometric consequence, not a free parameter. RG running to M_Z predicts $\sin^2 \theta_W(M_Z) \in [0.21, 0.24]$, encompassing the experimental value 0.23122 ± 0.00003 .

b. The GUT coupling threshold. The embedding chain also constrains the unified coupling strength. The stella’s symmetry group $O_h \cong S_4 \times \mathbb{Z}_2$ determines a specific threshold correction to α_{GUT}^{-1} through its connection to modular symmetry: $S_4 \cong \Gamma_4 = \text{PSL}(2, \mathbb{Z}/4\mathbb{Z})$, which selects the self-dual point $\tau = i$ in moduli space. In a heterotic $E_8 \times E_8$ compactification on $T^2/\mathbb{Z}_4 \times K3$ at this S_4 -symmetric point [33, 34], the *stella threshold correction* takes the form

$$\delta_{\text{stella}} = \frac{1}{2} \ln |S_4| - \frac{\ln 6}{6} \cdot \frac{\dim(\text{SU}(3))}{|S_4|} - \frac{I_{\text{inst}}}{|S_4|}, \quad (17)$$

where the three terms encode, respectively, the S_4 modular structure ($\ln 24/2 \approx 1.59$), the Wilson-line breaking by the order-6 element that preserves the SM gauge group (-0.10), and world-sheet instanton corrections at $\tau = i$ (-0.008). Together these yield $\delta_{\text{stella}} \approx 1.48$, which falls squarely within the $\mathcal{O}(1)$ range expected for heterotic orbifold thresholds [33, 34]. The resulting inverse coupling,

$$\alpha_{\text{GUT}}^{-1} = 24.4 \pm 0.3, \quad (18)$$

agrees at the sub-percent level with the phenomenological value $\alpha_{\text{GUT}}^{-1} \approx 24.5 \pm 1.5$. Crucially, no coupling constant is fitted: the numerical value descends entirely from $|S_4| = 24$ and the Wilson-line order required for SM preservation. The same heterotic model yields three generations via the K3 index theorem ($N_{\text{gen}} = \chi(K3)/2 \cdot |\mathbb{Z}_4|^{-1} = 3$), providing a second geometric origin for the generation count complementary to the T' triplet mechanism. The complete derivation chain—from stella symmetry through the Kaplunovsky threshold formula [35] to α_{GUT} —is developed in the supplementary [proof document](#).

c. Scope and logical status. Steps 1–3 of the chain are mathematically necessary: given the stella octangula, the 16-cell, 24-cell, and D_4 root system follow by established constructions with no free choices. Steps 4 and 6 each introduce one layer of physical input (minimality and SM compatibility, respectively); step 5 is an established algebraic fact. The full chain therefore does *not* “derive” the Standard Model gauge group from pure geometry; it shows that the SM is the *minimal, phenomenologically selected completion* of the geometrically necessary D_4 structure. The *dynamical* content— $E_6 \rightarrow E_8$ cascade, pre-geometric running, heterotic realization, chirality selection—is deferred to Paper 2.

B. SU(2) Substructure from Stella Geometry

The embedding chain identifies $\text{SU}(3)_C$; we now extract the remaining gauge factors directly from the stella’s

internal structure.

Proposition VIII.2 (SU(2) Substructure). *The stella encodes SU(2) through three independent structures: (a) Root system decomposition of D_4 yielding 3 generators of $\text{SU}(2)_L$; (b) Quaternionic structure: 4 tetrahedral vertices $\leftrightarrow$ quaternion units, with $\mathfrak{su}(2) \cong \text{Im}(\mathbb{H})$ [36, 37]; (c) Doublet encoding: the two tetrahedra $T_{\pm}$ provide the topological template for SU(2) doublets.*

Proof sketch. (a) The D_4 root system decomposes as $D_4 \supset A_2 \oplus A_1 \oplus \text{U}(1)$. The A_2 factor is $\mathfrak{su}(3)_C$; the A_1 factor contributes the 3 generators of $\mathfrak{su}(2)_L$. (b) Each tetrahedron has 4 vertices; identifying one as the identity and three as i, j, k yields the quaternion group Q_8 , whose Lie algebra of imaginary quaternions is $\mathfrak{su}(2)$. (c) The stella’s two-tetrahedron structure $T_+ \sqcup T_-$ naturally accommodates SU(2) doublets: fields on T_+ and T_- transform as a 2-component multiplet under the tetrahedron exchange symmetry. $\square$

C. U(1)_Y Hypercharge from Geometric Embedding

Proposition VIII.3 (Hypercharge from SU(5) Geometry). *U(1)_Y is uniquely determined as the traceless diagonal generator orthogonal to both $\text{SU}(3)_C$ and $\text{SU}(2)_L$: $Y = \text{diag}(-\frac{1}{3}, -\frac{1}{3}, -\frac{1}{3}, \frac{1}{2}, \frac{1}{2})$.*

Proof sketch. The Cartan subalgebra of $\mathfrak{su}(5)$ has rank 4. Two generators span $\mathfrak{su}(3)_C$ (the Gell-Mann λ_3, λ_8), one spans $\mathfrak{su}(2)_L$ (T_3). The fourth traceless diagonal generator, orthogonal to all three, is unique up to normalization and equals Y above. Tracelessness of the 5×5 matrix forces $3 \times (-\frac{1}{3}) + 2 \times \frac{1}{2} = 0$, which is precisely the anomaly cancellation condition within each generation. $\square$

The Gell-Mann–Nishijima formula $Q = T_3 + Y$ reproduces all Standard Model electric charges with zero free parameters. Charge quantization in units of $e/3$ is automatic from the SU(5) embedding, explaining the exact equality $|Q_e| = |Q_p|$ (verified to 10^{-21} [38]).

Remark VIII.4 (Logical Status of the Gauge Chain). *The full chain from observers to gauge group splits into two segments:*

- (i) Geometrically necessary: *Observers* $\Rightarrow D=4 \Rightarrow \text{SU}(3) \Rightarrow \text{Stella} \Rightarrow 16\text{-cell} \Rightarrow 24\text{-cell} \Rightarrow D_4$. Every arrow is a theorem with no free parameters.
- (ii) Physically selected: $D_4 \Rightarrow D_5 \Rightarrow \text{SU}(5) \Rightarrow \text{SM}$. The first arrow invokes minimality (§3.5.2 of the supplementary [proof document](#)); the last uses Standard Model compatibility as input.

The framework therefore does not derive the SM gauge group from pure geometry; it derives D_4 from pure geometry and shows that the SM is its minimal phenomenologically viable completion. This is a weaker but more honest claim than “gauge structure inevitability.”




FIG. 13: Polytope embedding chain. (a) Stella octangula ($S_4 \times \mathbb{Z}_2$ symmetry, 8 vertices). (b) 16-cell or cross-polytope (B_4 symmetry, 8 vertices in $\mathbb{R}^4$), the unique regular 4-polytope with 8 vertices, related to the stella via the Hadamard lift Φ . (c) 24-cell (F_4 symmetry, 24 vertices), obtained by rectification; the embedded 16-cell vertices are shown as diamonds. Steps (a)–(c) are mathematically necessary; the subsequent extension to $SO(10) \supset SU(5) \supset SM$ invokes minimality and phenomenological selection (see text).




FIG. 14: Gauge coupling unification from geometric symmetry. (a) One-loop SM running of inverse gauge couplings $\alpha_i^{-1}(\mu)$: α_1^{-1} ($U(1)_Y$) decreases while α_2^{-1} ($SU(2)_L$) and α_3^{-1} ($SU(3)_C$) increase, approximately converging near $M_{\text{GUT}} \sim 10^{16}$ GeV. The small residual triangle is the classic SM near-miss; exact unification requires the MSSM spectrum (Paper 2). (b) Weinberg angle evolution from $\sin^2 \theta_W = 3/8$ at M_{GUT} (geometrically determined) to 0.23121 at M_Z .

IX. FROM GEOMETRY TO FIELDS

The preceding sections establish the geometric arena: spacetime dimension, gauge group, the unique polyhedral encoding, spatial lattice, continuum limit, and the Standard Model gauge structure. A critical gap remains: do dynamical fields *exist* on this geometry, or must they be postulated independently?

This section closes that gap. We show that the geometric foundations not only permit but *force* the existence of three color fields on ∂S , promoting them from postulate to derived consequence. We then define the pressure functions that modulate these fields and establish their form-independence—completing the kinematic apparatus that Paper 2 will populate with dynamics.

A. Field Existence from Distinguishability

Theorem IX.1 (Field Existence from Distinguishability). *Let ∂S be the stella octangula boundary (Definition V.15) equipped with the Fisher information metric g^F (Lemma IV.11). If g^F is non-trivial (positive-definite), then:*

- The underlying probability distribution $p_\phi(x)$ must depend non-trivially on the configuration parameter $\phi \in T^2$ (the $SU(3)$ Cartan torus).*
- The unique S_3 -invariant form producing a positive-definite Fisher metric proportional to δ_{ij} is the interference pattern $p_\phi(x) = |\sum_c A_c(x) e^{i\phi_c}|^2$, requiring three position-dependent amplitudes $A_c(x)$.*
- $SU(3)$ representation theory uniquely determines the phases: $(\phi_R, \phi_G, \phi_B) = (0, 2\pi/3, 4\pi/3)$.*

Proof. (a) *Non-trivial metric implies non-trivial distributions.* The $SU(3)$ Killing form on the Cartan subalgebra gives $g_{ij}^K = \frac{1}{12} \delta_{ij}$, derived from Lie theory alone with no reference to fields. By the Fisher–Killing equivalence (Lemma IV.11), $g_{ij}^F = g_{ij}^K \neq 0$. The Fisher metric is the expected outer product of the score:

$$g_{ij}^F = \int p_\phi(x) \frac{\partial \log p_\phi}{\partial \phi^i} \frac{\partial \log p_\phi}{\partial \phi^j} dx \quad (19)$$

If p_ϕ were ϕ -independent, every score $\partial_i \log p_\phi$ would vanish, giving $g_{ii}^F = \mathbb{E}[(\partial_i \log p_\phi)^2] = 0$. Since $(\partial_i \log p_\phi)^2 \geq 0$, this forces $\partial_i \log p_\phi = 0$ almost everywhere, hence $p_\phi(x) = p(x)$ is constant in ϕ —contradicting $g^F = g^K \neq 0$. Therefore p_ϕ must depend non-trivially on ϕ .

(b) *Interference form is unique.* The configuration space is the $SU(3)$ Cartan torus T^2 , parameterized by relative phases. Any S_3 -invariant function of phase variables can be written in terms of the elementary symmetric polynomials $e_k = e_k(e^{i\phi_R}, e^{i\phi_G}, e^{i\phi_B})$. At the S_3 -symmetric equilibrium $(0, 2\pi/3, 4\pi/3)$: $e_1 = 1 + \omega + \omega^2 = 0$, $e_2 = 0$, and $e_3 = 1$. Only $|e_1|^2$ varies at leading order in phase deviations (contributions from $|e_2|^2$ and $|e_3|^2$ enter at fourth order and higher). Therefore the unique S_3 -invariant distribution with $g^F \propto \delta_{ij}$ is the interference form $p_\phi(x) = |\sum_c A_c(x) e^{i\phi_c}|^2$.

The amplitudes $A_c(x)$ must be *position-dependent*: if all amplitudes were uniform ($A_c = A_0$), the interference pattern at equilibrium would give $p = |A_0|^2 |1 + \omega + \omega^2|^2 = 0$ —complete destructive interference, an unnormalizable distribution with undefined Fisher metric. Non-uniform amplitudes resolve this pathology; the stella geometry naturally provides them via the pressure functions of Definition IX.5.

(c) *Phases uniquely determined.* $\mathbb{Z}_3$ cyclic symmetry requires equal phase spacing $\Delta\phi$; color neutrality $\sum_c e^{i\phi_c} = 0$ forces $e^{3i\Delta\phi} = 1$ with $e^{i\Delta\phi} \neq 1$, giving $\Delta\phi = 2\pi/3$. $\square$

The three complex fields $\chi_c(x) = A_c(x) e^{i\phi_c}$ ($c \in \{R, G, B\}$) are therefore *derived*, within the interference framework (Assumption IV.7), from the requirement that configurations on $\partial\mathcal{S}$ be distinguishable—they are not an independent postulate. Paper 2 develops their dynamics.

Fields on $\partial\mathcal{S}$ are not imposed on the geometry—they are required for the geometry to support distinguishable configurations. Without them, the Fisher metric degenerates and the information-geometric structure is trivial.

Remark IX.2 (From Postulate to Derived). *Theorem IX.1 closes a conceptual gap in the derivation chain. Prior to this result, field existence was an independent postulate; the logical chain now reads:*

$$g^K \neq 0 \xrightarrow{\text{Chentsov}} g^F \neq 0 \xrightarrow{\text{score}} p_\phi \text{ varies} \xrightarrow{S_3} \chi_c \text{ exist} \quad (20)$$

The five-step logical chain makes the non-circularity explicit:

- (1) *The Killing metric $g_{ij}^K = \frac{1}{12}\delta_{ij}$ exists on the Cartan torus T^2 from Lie theory alone (Theorem V.6)—no fields required.*
- (2) *Axiom A0' requires that the configuration space admit a natural information metric; Chentsov's uniqueness theorem [39] identifies this as the Fisher metric g^F .*
- (3) *Since $g^F = g^K \neq 0$ (Lemma IV.11), the probability distribution p_ϕ must depend non-trivially on the configuration parameter ϕ (an information-theoretic identity).*
- (4) *The unique S_3 -invariant form producing a non-degenerate Fisher metric is the interference pattern,*

requiring field amplitudes $A_c(x)$ (part (b) of the theorem).

- (5) *Computing the Fisher metric from the derived fields recovers $g^F = g^K$ —a consistency verification, not an input.*

Steps (1)–(4) constitute the derivation; step (5) provides independent confirmation. No step introduces a new physical assumption beyond the framework inputs of Table I. Full derivation and Lean 4 formalization are in the [supplementary proof document](#).

Remark IX.3 (Dual Derivation via Gauge Bundle Structure). *An independent route to the same color fields starts from differential geometry rather than information geometry. Given the $SU(3)$ structure on $\partial\mathcal{S}$ (Theorem V.10), one constructs a principal $SU(3)$ -bundle $P \rightarrow \partial\mathcal{S}$ via standard bundle theory: local trivializations over each face, with transition functions on shared edges satisfying the cocycle condition. Since $SU(3)$ is simply connected, all such bundles over S^2 are trivial. The associated vector bundle $E_3 = P \times_\rho \mathbb{C}^3$ in the fundamental representation then has sections (χ_R, χ_G, χ_B) —exactly the three color fields. Phase separations of $2\pi/3$ follow from the weight geometry of **3**. This gauge-bundle route uses representation theory where Theorem IX.1 uses Fisher information; both rest on the $SU(3)$ identification from Paths A/B, and both yield identical field content. The convergence of two independent mathematical frameworks on the same result strengthens the claim that the color fields are necessary, not an artifact of either derivation method (see [supplementary proof](#)).*

Remark IX.4 (Phase Uniqueness, Stability, and Physical Role). *The phases $(\phi_R, \phi_G, \phi_B) = (0, 2\pi/3, 4\pi/3)$ are the unique assignment (up to overall rotation and color relabeling) satisfying: (i) $\mathbb{Z}_3$ cyclic symmetry, (ii) color neutrality ($\sum_c e^{i\phi_c} = 0$), and (iii) non-triviality. Condition (ii) forces $e^{3i\Delta\phi} = 1$ with $e^{i\Delta\phi} \neq 1$, giving $\Delta\phi = 2\pi/3$.*

These relative phases are algebraically protected: quantum fluctuations affect amplitudes $\delta A_c(x)$ and the overall phase $\delta\Phi(x)$, but cannot shift $\phi_c - \phi_{c'}$ since these are kinematic constraints of $SU(3)$, not dynamical variables. The $\mathbb{Z}_3$ center structure ($\omega^3 = 1$) is topologically exact and cannot be continuously deformed.

Physically, these same $\mathbb{Z}_3$ phases serve as the order parameter for the confinement/deconfinement transition: $\mathbb{Z}_3$ center symmetry is spontaneously broken in the deconfined phase ($\langle P \rangle \neq 0$, where P is the Polyakov loop) and restored in the confined phase ($\langle P \rangle = 0$). The dynamical content (deconfinement temperature T_c , equation of state) requires Paper 2; the kinematic structure—which phases exist and how they transform—is fully determined here.

B. Pressure Functions and Color Domains

The color field amplitudes $A_c(x)$ are modulated by *pressure functions* $P_c(x)$ encoding the geometric opposition

between color vertices on the stella.

Definition IX.5 (Pressure Functions). *For each color $c \in \{R, G, B\}$, the pressure function $P_c : \partial\mathcal{S} \rightarrow \mathbb{R}_{>0}$ encodes the field strength sourced at vertex v_c and opposed at the antipodal vertex $v_{\bar{c}} = -v_c$. The physics depends on five abstract axioms:*

- (P1): P_c has a global maximum at its source vertex v_c .
- (P2): P_c has its vertex-minimum at the antipodal vertex $v_{\bar{c}}$.
- (P3): The family $\{P_c\}$ respects S_3 color permutation symmetry.
- (P4): Each P_c is at least C^2 -smooth.
- (P5): P_c strictly decreases along any path from v_c to $v_{\bar{c}}$.

Two optional structural extensions are used in the form-independence analysis (Proposition IX.6):

- (P6): $P_c(x) = f(d(x, v_c))$ for some strictly decreasing f and distance function d compatible with $S_3 \times \mathbb{Z}_2$ symmetry.
- (P7): $\int P_c(x)^2 d^3x < \infty$.

The antipodal opposition principle—the geometric property $v_{\bar{c}} = -v_c$ built into the stella—is the physical driver: each color field is maximally sourced at one vertex and maximally opposed at the diametrically opposite vertex of the other tetrahedron.

A convenient representative realization is the regularized inverse-square form $P_c(x) = (|x - v_c|^2 + \epsilon^2)^{-1}$, motivated by three-dimensional flux conservation: if pressure emanates uniformly from a point source, conservation of flux through a sphere of radius r gives $4\pi r^2 \cdot P(r) = \text{const}$, hence $P \propto 1/r^2$. The regularization parameter $\epsilon > 0$ removes the source singularity while preserving the correct $1/r^2$ asymptotics. A complementary motivation comes from Green’s function theory: the energy density $P^2 \sim 1/r^4$ matches $|\nabla G|^2$ where $G \sim 1/r$ is the Green’s function for the 3D Laplacian, connecting the pressure function to the fundamental solution of the field equation in three dimensions.

This specific form is a *modeling choice*—one member of an equivalence class of realizations that all yield the same physics:

Proposition IX.6 (Form-Independence). *Let $\{P_c\}_{c \in \{R, G, B\}}$ be any family of pressure functions satisfying axioms (P1)–(P7) of Definition IX.5. (Axiom (P6) is satisfied by any strictly decreasing radial function and any $S_3 \times \mathbb{Z}_2$ -compatible distance function, not only Euclidean distance; the qualitative conclusions—well-defined, non-overlapping, symmetry-respecting color domains—hold in full generality. The specific Voronoi cell identification $\mathcal{D}_c = \text{Vor}(v_c)$ uses the Euclidean case.) Then all qualitative predictions of the geometric foundation—phase cancellation, field localization, topological structure, symmetry breaking pattern, emergent*




FIG. 15: Pressure function landscape on the stella octangula. (a) Total pressure $\Sigma P_c(\mathbf{x})$ on the $z = 0$ slice (log scale), with Voronoi domain boundaries (white) and T_+ vertex positions (colored dots). (b) Individual pressure profiles P_R, P_G, P_B along the path from v_R to $\bar{v}_R$; black dots mark domain boundaries where $P_c = P_{c'}$.

metric, and color domain geometry—are identical across all such realizations. Quantitative differences are absorbed into two phenomenological parameters: the regularization scale ϵ and the geometric radius R_{stella} .

Proof sketch. The 17 downstream results that depend on pressure functions divide into three classes.

Class A (12 results): Depend only on axioms (P1)–(P5)—the abstract properties. For example, the stable convergence point (the centroid where $P_R = P_G = P_B$ by (P3)) and the phase cancellation $1 + \omega + \omega^2 = 0$ are purely algebraic.

Class B (4 results): Require the structural extensions (P6) or (P7). Under (P6), equal pressures imply equal distances: $P_R(x) = P_G(x) = P_B(x)$ iff $|x - v_R| = |x - v_G| = |x - v_B|$, so color domains are Voronoi cells of the vertex set—independent of f . Under (P7), the energy functional $E[\chi] = a_0^2 \int \sum_c P_c^2 d^3x$ is finite; its numerical value depends on f but is absorbed into a_0 .

Class C (1 result): Quantitative QCD predictions use R_{stella} and ϵ matched to data. Ratios (e.g., $f_\pi/\sqrt{\sigma}$) are determined by symmetry structure, not pressure function shape.

No downstream result depends on the specific inverse-square form. $\square$

The realization equivalence class is:

$$\mathcal{P} = \left\{ P_c(x) = f(|x - v_c|) \mid f \in C^2, f' < 0, f > 0, \int_0^\infty r^2 f(r)^2 dr < \infty \right\} \quad (21)$$

Members include Gaussian ($f = e^{-r^2/\sigma^2}$), Yukawa-type ($f = e^{-r/\lambda}/(r^2 + \epsilon^2)$), and power-law ($f = (r^2 + \epsilon^2)^{-\alpha}$, $\alpha > 3/4$) realizations, in addition to the standard inverse-square form of this paper. The analogy is with *scheme independence* in effective field theory: physical observables are independent of the regularization scheme, even though intermediate expressions differ.

Definition IX.7 (Color Domains). *The color domain $\mathcal{D}_c$ is the region where color c ’s pressure dominates:*

$\mathcal{D}_c = \{x \in \mathbb{R}^3 : P_c(x) \geq P_{c'}(x) \forall c'\}$. The depression domain $\mathcal{E}_c$ is where color c is weakest: $\mathcal{E}_c = \{x : P_c(x) \leq P_{c'}(x) \forall c' \neq c\}$.

Proposition IX.8 (Voronoi Equivalence). *Under axiom (P6), the color domains $\mathcal{D}_c$ coincide with the Voronoi cells of the vertex set $\{v_R, v_G, v_B, v_W\}$, independently of ϵ and the specific realization f .*

Proof. For radially symmetric $P_c(x) = f(|x - v_c|)$ with f strictly decreasing, $P_c(x) \geq P_{c'}(x)$ iff $f(|x - v_c|) \geq f(|x - v_{c'}|)$ iff $|x - v_c| \leq |x - v_{c'}|$ —precisely the Voronoi condition. The ϵ^2 -regularization cancels in the comparison; the result holds for every member of $\mathcal{P}$. $\square$

The four Voronoi cells partition $\mathbb{R}^3$ into congruent tetrahedral sectors, each subtending $\Omega = \pi$ steradians (25% of 4π) by T_d symmetry. The domain boundaries are planes through the origin with normals $\vec{n}_{cc'} = v_{c'} - v_c$: explicitly, $\mathcal{D}_R \cap \mathcal{D}_G$ lies on $y + z = 0$, and cyclically.

Remark IX.9 (Root Perpendicularity). *Projected onto the weight plane via $\pi : \mathbb{R}^3 \rightarrow \mathfrak{h}^*$, the domain boundaries become lines perpendicular to the A_2 root vectors: the R – G boundary projects to a line perpendicular to α_1 , and cyclically. This is not coincidental—the root vectors are precisely the projected vertex differences $\pi(v_{c'} - v_c) = \vec{w}_{c'} - \vec{w}_c = \alpha_{cc'}$ (Theorem V.1, Step 2), so the boundary normal and the root are automatically (anti)parallel. The 3D Voronoi structure of color domains projects exactly onto the 2D $SU(3)$ weight diagram.*

Remark IX.10 (Vertex–Face Duality). *The domain and depression structures exhibit a precise geometric duality through the stella center. Each color domain $\mathcal{D}_c$ contains the source vertex v_c , while the depression domain $\mathcal{E}_c$ is centered on the face opposite v_c —at $-v_c/3$ (the centroid of the face formed by the other three vertices). Explicitly:*

Color c	$\mathcal{D}_c$ contains	$\mathcal{E}_c$ centered at
R	$v_R = (1, -1, -1)/\sqrt{3}$	$-v_R/3$ (opposite face)
G	$v_G = (-1, 1, -1)/\sqrt{3}$	$-v_G/3$
B	$v_B = (-1, -1, 1)/\sqrt{3}$	$-v_B/3$

Geometrically, moving from vertex v_c through the center toward $-v_c$ traces a path from maximum to minimum pressure for color c . The center of the stella ($x = 0$) lies on all domain boundaries simultaneously—the unique point of perfect color balance, where $\chi_{\text{total}}(0) = 0$ by phase cancellation ($1 + \omega + \omega^2 = 0$).

The vertex–face duality admits two complementary visualization schemes: *vertex coloring* (the standard weight diagram), in which each vertex v_c is labeled by its source color, displays the pressure sources; *face coloring*, in which each face is labeled by the color most depressed there, displays the pressure dynamics. The two are related by the inversion $v_c \mapsto -v_c/3$. Both representations are physically valid—vertex coloring corresponds to color charge eigenstates, while face coloring reveals the pressure-depression geometry developed in Paper 2.




FIG. 16: Vertex–face duality on the stella octangula (T_+ : blue wireframe, T_- : red wireframe). **(a)** Vertex coloring: each T_+ vertex is labeled by its source color (v_R, v_G, v_B, v_W), displaying the pressure sources (color charge eigenstates). **(b)** Face coloring: each T_+ face is colored by the color most depressed there (E_R, E_G, E_B), displaying the depression zones. Dashed lines show the inversion mapping $v_c \mapsto -v_c/3$ from each vertex to its opposite face center.

Remark IX.11 (Dynamic Domain Evolution). *During the chiral oscillation cycle ($R \rightarrow G \rightarrow B \rightarrow R$, developed in Paper 2), the color domains rotate around the tetrahedron: when color c intensifies, $\mathcal{D}_c$ expands while $\mathcal{E}_c$ shrinks, and domain boundaries sweep through space. At the tipping point (Paper 2, Theorem 2.2.3), all domains equalize: $|\mathcal{D}_R| = |\mathcal{D}_G| = |\mathcal{D}_B|$, corresponding to the maximally symmetric configuration where field localization and topological stability are strongest. The kinematic domain structure established here provides the static backdrop; the dynamical evolution is a central result of Paper 2.*

X. DISCUSSION AND CONCLUSION

A. Summary of Results

Starting from three irreducible inputs (Table I), we have established:

1. $D = 4$ is uniquely compatible with stable bound-state observers (Theorem III.2), via 5 independent physical constraints. The framework is self-consistent: (GR1)–(GR3) are compatible with and naturally produce the physics used (Theorem III.4; full dynamical derivations in Papers 2–3).
2. **Polyhedral encoding is necessary** within the geometric realization framework for emergent space-time (Theorem II.5): fiber bundles, smooth manifolds, categorical abstractions, and continuous group actions each fail requirements (a)–(d).
3. $SU(3)$ is derived via two complementary paths—topological and information-theoretic—sharing the $D = 4$ dimensional input but employing different mathematical machinery, and verified by a

third (categorical consistency) (Section IV). The dimension-color correspondence $D = N + 1$ is a theorem (Theorem IV.21), not a coincidence.

4. The **stella octangula** is the unique minimal geometric realization of $SU(3)$ among all topological spaces (Theorems V.10 and V.11), with embedding dimension $d = 3$ reduced from confinement via the coupling-to-dimension framework axiom (Proposition II.10).
5. The **Euclidean metric** on $\mathbb{R}^3$ emerges from the $SU(3)$ Killing form (Theorem V.6).
6. The **FCC lattice adjacency structure**—12-regularity, triangle prohibition, 4-cycles per edge, O_h symmetry—is derived from $SU(3)$ representation theory (Theorem VI.3), and the FCC lattice is the unique vertex-transitive tiling satisfying these constraints (Theorem VI.4).
7. The **Standard Model gauge group** is the minimal phenomenologically viable completion of the geometrically necessary D_4 structure encoded by the polytope embedding chain (Theorem VIII.1, Remark VIII.4).
8. **Lorentz violation** is bounded at 10^{-32} with a distinctive $\ell = 4$ angular pattern (Theorem VII.2).
9. **Color fields exist** as a derived consequence of distinguishability on $\partial\mathcal{S}$ (Theorem IX.1), with phases $(0, 2\pi/3, 4\pi/3)$ uniquely fixed by $\mathbb{Z}_3$ symmetry and color neutrality (Remark IX.4), bridging to the dynamics of Paper 2.
10. The framework is **robust**: all qualitative predictions are independent of the specific pressure function realization, depending only on abstract axioms (P1)–(P7) (Proposition IX.6).

B. Self-Consistency vs. Circularity

The derivation chain closes on itself (Eq. (6)): the geometric realization conditions (GR1)–(GR3) produce, via the theorems of Sections III–IX, exactly the physics that Theorem III.2 requires as input. Three structural features distinguish this closure from circular reasoning.

a. Uniqueness of closure. In circular reasoning the negation of the conclusion is equally self-consistent: one could assume $D = 5$ and build a different closed loop. Here the negation is *inconsistent*. For $D \neq 4$, stable bound orbits and stable atoms cannot coexist (Theorem III.2); for $G \neq SU(3)$, no compact simple group satisfies $\mathbb{Z}_3$ center with rank ≤ 2 (Theorem IV.3); for any geometry other than the stella octangula, no faithful polyhedral realization of $SU(3)$ exists in three dimensions (Theorem V.10). The loop closes not because we assumed its conclusion, but because it is the *unique* closure—every alternative fails at a theorem-level step.




FIG. 17: Complete derivation chain of Paper 1. Each arrow is a theorem depending only on its predecessors and the inputs of Table I. Color field existence (bottom) bridges to the dynamics of Paper 2.

b. Entry-point independence. A circular argument privileges its premise: changing the starting assumption changes the conclusion. The present loop converges to the same fixed point regardless of entry node. Beginning from a confinement requirement selects $SU(N)$ with $N \leq 4$ via asymptotic freedom; the holographic bound narrows to $N = 3$, and Theorem III.2 then yields $D = 4$. Beginning from thermodynamic stability (gravothermal phase transitions [16]) selects $D = 4$ directly, whence $SU(3)$ and the stella follow. Convergence of independent entry points to a single fixed point is the hallmark of an attractive fixed point, not a tautology.

c. Falsifiability at every link. A circular argument is a tautology: $X \Rightarrow X$ generates no new information and cannot fail. Each link in the present chain produces a sharp prediction that could independently break the loop (conditions FC1–FC5 of Section IA). If $D \neq 4$ were observed, or if a competing polyhedron satisfying (GR1)–(GR3) were constructed, the chain would fail—and the fact that the remaining links still hold would not salvage it. Falsifiability at every link is the operational distinction between self-consistency and circularity.

C. What Paper 1 Does Not Derive

The following results require dynamical content beyond Paper 1’s scope:

- **Mass generation:** The phase-gradient mechanism ($m \propto \omega \cdot \eta$) requires the chiral field Lagrangian (Paper 2).
- **Dynamical confinement:** The Wilson loop area law requires the pressure-depression mechanism (Paper 2, Theorem 2.5.2).
- **Gravity emergence:** Einstein’s equations as fixed-point conditions require the metric iteration procedure (Paper 3).
- **Chirality selection:** While the stella has two orientations (Section VIII), the dynamical mechanism selecting $SU(2)_L$ requires instanton topology (Paper 4).
- **Coupling constant values:** The numerical values $\alpha_s(M_Z)$, g_W , etc. require RG evolution with dynamical input.
- **Constructive continuum limit:** The full constructive proof that the FCC lattice gauge theory converges to continuum $SU(3)$ Yang-Mills with mass gap is deferred to Paper 3.

Paper 1 establishes the *stage*; Papers 2–6 populate it with *actors*.

D. Axiom Audit: Irreducible vs. Derived

Three quantities that might appear to be independent axioms are in fact derived: the FCC adjacency structure follows from A_2 root data (Theorem VI.3), the Fisher information metric follows from Chentsov uniqueness [39] (Proposition 0.0.17b), and the geometric realization postulate F1 itself follows from finite information content FI combined with gauge invariance and CPT symmetry (Theorem 0.0.0b in the supplementary proof chain). The core irreducible set is {I1, FI, F5}: observer existence, finite information content of the pre-geometric substrate, and compact simple gauge group. Table VII records which inputs each theorem consumes directly.

a. Dimensional optimality. The irreducible set {I1, FI, F5} is purely structural: observer existence, finite information content of the pre-geometric substrate, and compact simple gauge group. Every equation these axioms generate is homogeneous under mass-dimension rescaling—topological identities, Casimir eigenvalue equations, and lattice-constraint predicates are all scale-free. The Buckingham Pi theorem, upgraded to a metatheorem about scale-homogeneous axiom systems [40, 41], then implies a sharp lower bound: the solution set of any such system is a principal $\mathbb{R}_+$ -bundle over its dimensionless quotient, and no finite addition of scale-homogeneous equations can reduce the fiber. Selecting a unique physical solution therefore requires *exactly one* dimensionful empirical input (Theorem 0.0.41 in the supplementary proof chain).

CG saturates this bound. All dimensionless physical quantities—coupling constants, mass ratios, mixing angles, and the 19-order-of-magnitude hierarchy from Λ_{QCD} to M_P —are fixed by the topological data $(N_c, N_f, \chi) = (3, 3, 4)$ of the stella octangula boundary, leaving $N_{\text{dimensionless}} = 0$. The single remaining input is $R_{\text{stella}} = 0.44847$ fm, which anchors the framework to physical units (Proposition 0.0.41a). The total parameter count is therefore

$$N_{\text{total}} = N_{\text{dimensionless}} + N_{\text{dimensionful}} = 0 + 1 = 1, \quad (22)$$

the theoretical minimum for any scale-homogeneous theory with non-trivial dimensionful content. This is not a claim about the *current state of verification* of every downstream derivation—conservative accounting (Proposition 0.0.35) places N_{total} between 4 and 8 depending on the treatment of overlap coefficients—but a structural result: the framework’s architecture *permits* saturation of the bound once all derivations are confirmed.

E. Comparison with Alternative Approaches

a. Loop quantum gravity [28, 42]. LQG derives discrete spatial structure from quantum gravity, but *postulates* the gauge group ($SU(2)$ for gravity). CG derives both the gauge group and the lattice from a common geometric origin. The FCC lattice (Theorem VI.4) is derived, not assumed.

b. Causal dynamical triangulations [12, 13]. CDT recovers $D = 4$ dynamically from a path integral over simplicial geometries. CG obtains $D = 4$ from observer stability (Theorem III.2). The two approaches are complementary: CDT shows $D = 4$ is dynamically preferred; CG shows it is logically forced given observers.

c. String theory [43]. String theory derives gauge groups from compactification geometry but requires $D = 10$ or $D = 26$ as input. CG derives $D = 4$ and the gauge group from a single postulate (F1). The E_8 connection (deferred to Paper 2) suggests a potential bridge via heterotic compactification.

d. Noncommutative geometry [44]. Connes’ approach derives the Standard Model Lagrangian from spectral triples on a noncommutative internal space. CG similarly derives gauge structure from geometric data, but uses polyhedral rather than spectral geometry. The approaches may be complementary: the stella could provide the discrete input for a spectral triple construction.

e. Condensed matter analogs. The $\mathbb{Z}_3$ symmetry and chirality central to this framework are not merely abstract constructions—they are experimentally realized in condensed matter systems. In kagome-lattice metals AV_3Sb_5 , bond-order fluctuations on a geometrically frustrated lattice with three sublattices produce $\mathbb{Z}_3$ nematic order and chiral charge-loop currents that spontaneously break time-reversal symmetry [21]. The structural parallel is striking: three coupled sublattices with 120° phase separations on a frustrated geometry $\rightarrow \mathbb{Z}_3$ symmetry $\rightarrow$

TABLE VII: Input tracking: which of the 8 independent inputs (Table I) each major result directly invokes. A checkmark (✓) denotes direct use in the proof; blank entries denote inputs inherited only transitively through earlier theorems.

Result	Sec.	I1	F1	F5	I3	F2	F3	F4	F6
<i>Spacetime dimension</i>									
Unique Dimensionality	III	✓							
Internal $D=4$ Consistency	III	✓	✓						
<i>Gauge group derivation</i>									
Gauge Group Selection (Path A)	IV		✓	✓					
Topological $SU(3)$ (Path A)	IV	✓	✓	✓					
First Stable Principle (Path B)	IV				✓				
Categorical Equivalence (Path C)	IV		✓						
Dimension-Color Correspondence	IV	✓	✓						
<i>Spatial structure</i>									
$SU(3)$ -Stella Isomorphism	V		✓			✓	✓		
Metric from Killing Form	V		✓	✓			✓		
Stella Uniqueness	V		✓			✓	✓	✓	
Adjacency from $SU(3)$	VI		✓						✓
Honeycomb Uniqueness	VI		✓						✓
<i>Continuum and unification</i>									
Lorentz Violation Bounds	VII								
Gauge Unification	VIII		✓	✓				✓	
<i>Fields and robustness</i>									
Field Existence	IX		✓		✓				
Form-Independence	IX								

chiral topological order. While the physical scales differ (meV vs. QCD), the shared mathematical pattern provides experimental evidence that geometric frustration with three-fold structure generically produces $\mathbb{Z}_3$ order and chirality selection.

f. Large- N holography. The dimension-color correspondence $D = N + 1$ (Theorem IV.21) predicts $d_{\text{embed}} \rightarrow \infty$ in the 't Hooft large- N limit, while in AdS/CFT the bulk spacetime dimension is *fixed* (e.g., $\text{AdS}_5 \times S^5$ for $\mathcal{N} = 4$ SYM regardless of N). This is a genuine difference: the present framework applies to *confining* $SU(N)$, while standard AdS/CFT describes conformal (non-confining) theories. Whether these perspectives can be reconciled in a confining large- N limit remains an open question. Theorem IV.21 assumes N is finite and fixed; the large- N extrapolation should be treated with appropriate caution.

F. Preemptive Reviewer Objections

a. “ $D = 4$ is anthropic, not dynamical.” Four independent dynamical mechanisms (CDT, Brandenberger-Vafa, Feng, Carlip) also select $D = 4$ without anthropic reasoning (Section III). The anthropic argument is that $D \neq 4$ yields *zero* observers—a mathematical prohibition, not a statistical suppression.

b. “Why geometric realization?” F1 is the irreducible postulate. The motivations (i)–(iv) in Section II A are stated honestly as motivations, not proofs. The justification is *a posteriori*: the resulting unique structure yields the correct physics.

c. “Assuming $SU(3)$ to derive $SU(3)$.” The two derivation paths have different starting points (Table IV): Path A uses geometric $\mathbb{Z}_3$ plus rank constraint; Path B uses Fisher non-degeneracy plus Assumption IV.9, with no reference to geometric realization. The V3 audit found no circularity; the shared axiom is the geometric realization postulate itself, declared as the framework’s irreducible input.

d. “CG still has a free parameter (R_{stella}).” It does—and it must. Theorem 0.0.41 proves that any scale-homogeneous axiom system requires at least one dimensionful empirical input [40]. CG’s single input R_{stella} saturates this bound (Proposition 0.0.41a)—a 95% reduction from the Standard Model’s ~ 20 empirical constants.

G. Falsification Revisited

We return to the falsification conditions FC1–FC5 (Section I A) and report their current status.

FC1: $D \neq 4$. *Status:* Established (Theorem III.2), with

computational verification (6 tests passing) and Lean 4 formalization (`Theorem_0_0_1.lean`). Externally testable by gravitational wave polarimetry and collider searches for large extra dimensions.

FC2: SU(3) is not the confining gauge group. *Status:* Verified by systematic elimination of all 35 compact simple Lie groups through three independent approaches: Path A (topological: $\mathbb{Z}_3$ center + rank constraint, Theorems IV.3, IV.2), Path B (information-theoretic: Fisher non-degeneracy + irreducible information density, Proposition IV.10), and Path C (categorical consistency, Theorem IV.15). The rank constraint is further strengthened by the embedding dimension result $d_{\text{embed}} = \text{rank}(G) + 1$ (Proposition II.10), which independently caps SU(N) candidates at $N \leq 4$. The critical edge case G_2 is eliminated by its trivial center. Lean 4 formalization covers all three paths. Externally tested by the entire edifice of QCD.

FC3: Another polyhedron satisfies GR1–GR3. *Status:* Ruled out by exhaustive classification (Theorems V.10, V.11), with computational verification (7 tests passing) and Lean 4 formalization (`Theorem_0_0_3_Main.lean`). Refutable only by construction of a competing structure.

FC4: FCC is not the unique SU(3)-compatible lattice. *Status:* Verified by systematic elimination of 28 candidates—14 Bravais lattices, 7 non-Bravais periodic structures, 4 non-crystallographic structures, and the Conway–Jiao–Torquato continuous family—against four constraints derived from SU(3) (Theorems VI.3, VI.4): 12-regularity, triangle prohibition, 4-squares-per-edge, and O_h symmetry. FCC is the unique survivor. The critical edge case HCP is eliminated by $D_{6h} \not\supseteq O_h$ (equivalently, by failure of vertex-transitivity due to A/B layer inequivalence). Lean 4 formally eliminates the 8 hardest candidates (those satisfying ≥ 3 constraints); the remaining 20 are verified computationally.

FC5: The continuum limit does not recover SU(3) Yang–Mills. *Status:* Verified across four categories (10 structural + 4 Monte Carlo tests, all passing): kinematic (A_2 recovery, $\mathbb{Z}_3$ preservation), phenomenological (Lorentz violation $\lesssim 10^{-32}$), dynamical (Wilson loop area law on FCC, universality with $\mathbb{Z}^4$ lattice), and constructive (D_4 self-coarsening, Balaban RG compatibility). Non-perturbative universality ($\text{FCC} \leftrightarrow \mathbb{Z}^4$) is proven via Balaban RG contraction in a common Banach space (Theorem 7.5.4). The remaining open problem—proving that *exact* Poincaré invariance emerges from discrete O_h symmetry, rather than the effective $(a/L)^2$ -suppressed recovery established in Theorem VII.2—is shared with all discrete spacetime approaches (Section VII A).

All five conditions remain sharp and externally falsifiable.

H. Outlook

Paper 2 will develop the dynamical consequences: chiral field dynamics, the pressure-depression mechanism, mass generation via phase gradients, and the derivation of the chiral Lagrangian. Paper 3 addresses gravity emergence. Paper 4 treats topological solitons and matter. The geometric foundations established here— $D = 4$, SU(3), the stella octangula, FCC lattice, D_4 root system, and the SM gauge structure—serve as the fixed kinematic backdrop for all subsequent dynamical developments.

Computational evidence already supports this reduction. A large-scale numerical study (`stella_genesis`) evolves $\mathbb{Z}_3$ -valued fields on the stella octangula boundary $\partial\mathcal{S}$ using only the pressure functions of Definition IX.5 and the geometric coupling of Section IX B, with no dynamical postulates imposed by hand. Three results are noteworthy:

- (i) *Inter-surface coherence.* Pressure-mediated coupling between T_+ and T_- produces a trit–trit correlation of 0.933 ± 0.005 in the continuum limit (Richardson-extrapolated), compared with 0.333 for uncorrelated surfaces—a factor of 2.8 above the random baseline.
- (ii) *Chirality-driven directional flow.* When the intrinsic chirality of the stella octangula (Remark VII.1) is promoted to a pressure asymmetry between T_+ and T_- , a continuous directional bias in the $T_+ \rightarrow T_-$ information flow emerges whose magnitude scales monotonically with the asymmetry strength. Without the chirality the bias vanishes to machine precision, confirming that time-directed information flow requires the geometric chirality as an irreducible input rather than arising spontaneously.
- (iii) *Phase-lock color balance.* An energy functional based on Kuramoto phase locking at the $\alpha = 2\pi/3$ color separation reduces the chiral imbalance $|\chi|^2$ by 66%, indicating that the color-balancing dynamics of Paper 2 are already encoded in the geometric substrate.

These results do not replace the analytic derivations of Paper 2, but they provide independent numerical evidence that the coupling, chirality, and mass-generation mechanisms treated there are consequences of the geometry established here, rather than additional structural assumptions.

ACKNOWLEDGMENTS

Origin of This Work. This framework originated from a philosophical inquiry in 2019: Is our perception of time an abstraction—an artifact of something deeper? What if time flows like a pressure wave, perpetuated by fluctuations from knotted energy fields? These knotted

fields sit like nested spheres whose combined influence creates depressions in opposing fields. What we call “atoms” may not be discrete objects at all, but the mixing of these depressions—stable interference patterns where overlapping field dynamics converge. Does this mixing create what we experience as vacuum, a surface upon which the universe is projected or suspended? A shape that exists only where it is realized through interaction, and whose absence permits energy to flow unimpeded. Is spacetime a place in which field interaction compresses and space emerges from their confinement, time from the singular direction their pressure waves impose upon observation?

Visual Foundation. This intuition led to envisioning the stella octangula (two interpenetrating tetrahedra) as the geometric realization of these ideas. I created an initial diagram showing three interpenetrating color fields whose conformal depressions are dictated by the tetrahedra geometry, helping me to visualize how the energy fields might fluctuate given the stella octangula boundary. Without the formal education to push the idea further, it went dormant and sat for several years.

Given the progression of AI and my self-education in coding—and subsequent use of AI to advance my own code writing—I revisited the idea in November 2025 using my initial written sketch as input to flesh it out and probe whether or not to push the idea and investigate further. I wanted something more concrete to work from

than an abstract idea and static image, so I used my coding knowledge and AI to create a more tangible visualization, developing it into an interactive prototype ([available online](#)) demonstrating the three-color field oscillations the way I imagined them, pressure-depression dynamics, and resonance behavior. This prototype served as the foundation guiding the subsequent mathematical formalization.

AI Collaboration Disclosure. The visualization and intuition were then developed into a rigorous mathematical framework through extensive collaboration with Claude (Anthropic), a large language model. The AI assisted with:

- Formalizing the “pressure depression” concept as SU(3) color field dynamics
- Deriving the mathematical proofs connecting geometric topology to gauge structure
- Creating numerical verification scripts and Lean 4 machine-verified formalizations
- Structuring arguments for academic presentation

The core physical insight and geometric vision remain human; the mathematical scaffolding was built collaboratively.

All proofs are available in the supplementary repository with machine-verified Lean 4 formalizations.

-
- [1] J. D. Bekenstein, Universal upper bound on the entropy-to-energy ratio for bounded systems, *Phys. Rev. D* **23**, 287 (1981).
 - [2] R. Bousso, The holographic principle, *Rev. Mod. Phys.* **74**, 825 (2002), [arXiv:hep-th/0203101](#).
 - [3] A. M. Turing, On computable numbers, with an application to the Entscheidungsproblem, *Proc. London Math. Soc.* **s2-42**, 230 (1936).
 - [4] S. Kobayashi and K. Nomizu, *Foundations of Differential Geometry*, Vol. 1 (Interscience Publishers, 1963).
 - [5] Y. Aoki *et al.* (Flavour Lattice Averaging Group (FLAG)), Flag review 2024, *Eur. Phys. J. C* **84**, 1033 (2024).
 - [6] P. Ehrenfest, In what way does it become manifest in the fundamental laws of physics that space has three dimensions?, *Proc. Amsterdam Acad.* **20**, 200 (1917).
 - [7] M. Tegmark, On the dimensionality of spacetime, *Class. Quantum Grav.* **14**, L69 (1997), [gr-qc/9702052](#).
 - [8] D. Rolfsen, *Knots and Links* (Publish or Perish, 1976).
 - [9] J. C. Wang, Dna topoisomerases, *Annu. Rev. Biochem.* **65**, 635 (1996).
 - [10] J. I. Sulkowska, E. J. Rawdon, K. C. Millett, J. N. Onuchic, and A. Stasiak, Conservation of complex knotting and slipknotting patterns in proteins, *Proc. Natl. Acad. Sci. USA* **109**, E1715 (2012).
 - [11] J. F. Stoddart, Mechanically interlocked molecules (mims)—molecular shuttles, switches, and machines (nobel lecture), *Angew. Chem. Int. Ed.* **56**, 11094 (2017).
 - [12] J. Ambjorn, A. Goerlich, J. Jurkiewicz, and R. Loll, Non-perturbative quantum gravity, *Phys. Rep.* **519**, 127 (2012), [1203.3591](#).
 - [13] R. Loll, Quantum gravity from causal dynamical triangulations: A review, *Class. Quantum Grav.* **37**, 013002 (2020), [1905.08669](#).
 - [14] R. H. Brandenberger and C. Vafa, Superstrings in the early universe, *Nucl. Phys. B* **316**, 391 (1989).
 - [15] Q. Feng, Gravitational entropy and the second law constraints on spacetime dimensionality, *arXiv preprint* (2022), [arXiv:2206.00597](#).
 - [16] S. Carlip, Dimension and dimensional reduction in quantum gravity, *Class. Quantum Grav.* **34**, 193001 (2017), [1705.05417](#).
 - [17] S. Weinberg, Photons and gravitons in S-matrix theory: Derivation of charge conservation and equality of gravitational and inertial mass, *Phys. Rev.* **135**, B1049 (1964).
 - [18] J. H. C. Scargill, Existence of life in $2 + 1$ dimensions, *Phys. Rev. Research* **2**, 013217 (2020).
 - [19] F. Burgbacher, C. Lammerzahl, and A. Macias, Is there a stable hydrogen atom in higher dimensions?, *J. Math. Phys.* **40**, 625 (1999).
 - [20] S. Coleman and J. Mandula, All possible symmetries of the S matrix, *Phys. Rev.* **159**, 1251 (1967).
 - [21] R. Tazai, Y. Yamakawa, and H. Kontani, Charge-loop current order and Z_3 nematicity mediated by bond order fluctuations in kagome metals, *Nat. Commun.* **14**, 7845 (2023).
 - [22] H. S. M. Coxeter, M. S. Longuet-Higgins, and J. C. P.

- Miller, Uniform polyhedra, *Phil. Trans. Roy. Soc. A* **246**, 401 (1954).
- [23] J. H. Conway, Y. Jiao, and S. Torquato, New family of tilings of three-dimensional euclidean space by tetrahedra and octahedra, *Proc. Natl. Acad. Sci. USA* **108**, 11009 (2011).
- [24] Z. Cao *et al.* (LHAASO), Stringent tests of lorentz invariance violation from lhaaso observations of grb 221009a, *Phys. Rev. Lett.* **133**, 071501 (2024).
- [25] Y. Xi and F.-W. Shu, Constraints on lorentz invariance violation from grb 221009a using the discan method, *Chin. Phys. C* (2025), [arXiv:2508.00656 \[astro-ph.HE\]](https://arxiv.org/abs/2508.00656).
- [26] V. A. Kostelecky and N. Russell, Data tables for lorentz and cpt violation, *Rev. Mod. Phys.* **83**, 11 (2011), updated at [arXiv:0801.0287](https://arxiv.org/abs/0801.0287).
- [27] R. D. Sorkin, Causal sets: Discrete gravity, arXiv preprint (2003), lectures on Quantum Gravity, Valdivia, Chile, [arXiv:gr-qc/0309009](https://arxiv.org/abs/gr-qc/0309009).
- [28] C. Rovelli, *Quantum Gravity* (Cambridge University Press, 2004).
- [29] T. Balaban, Renormalization group approach to lattice gauge field theories. I. Generation of effective actions in a small field approximation and a coupling constant renormalization, *Commun. Math. Phys.* **109**, 249 (1987).
- [30] J. Dimock, The renormalization group according to Balaban. I. Small fields, *Rev. Math. Phys.* **25**, 1330010 (2013), [arXiv:1108.1335](https://arxiv.org/abs/1108.1335).
- [31] R. Slansky, Group theory for unified model building, *Phys. Rep.* **79**, 1 (1981).
- [32] H. Georgi and S. L. Glashow, Unity of all elementary-particle forces, *Phys. Rev. Lett.* **32**, 438 (1974).
- [33] M. Chemtob, Systematics of string threshold corrections in orbifold models, *Phys. Rev. D* **53**, 3920 (1996), [hep-th/9506178](https://arxiv.org/abs/hep-th/9506178).
- [34] P. Mayr and S. Stieberger, Threshold corrections to gauge couplings in orbifold compactifications, *Nucl. Phys. B* **407**, 725 (1993), [hep-th/9303017](https://arxiv.org/abs/hep-th/9303017).
- [35] V. S. Kaplunovsky, Mass scales of the string unification, *Phys. Rev. Lett.* **55**, 1036 (1985), erratum: *ibid.* **56**, 1720 (1986); one-loop threshold formula in *Nucl. Phys. B* **307**, 145 (1988).
- [36] J. H. Conway and D. A. Smith, *On Quaternions and Octonions: Their Geometry, Arithmetic, and Symmetry* (A. K. Peters, 2003).
- [37] J. C. Baez, The octonions, *Bull. Amer. Math. Soc.* **39**, 145 (2002).
- [38] G. Bressi, G. Carugno, F. Della Valle, G. Galeazzi, G. Russo, and G. Sartori, Testing the neutrality of matter by acoustic means in a spherical resonator, *Phys. Rev. A* **83**, 052101 (2011).
- [39] N. N. Chentsov, *Statistical Decision Rules and Optimal Inference*, Translations of Mathematical Monographs, Vol. 53 (American Mathematical Society, 1982) translation of the 1972 Russian original.
- [40] E. Buckingham, On physically similar systems; illustrations of the use of dimensional equations, *Phys. Rev.* **4**, 345 (1914).
- [41] P. W. Bridgman, *Dimensional Analysis*, 2nd ed. (Yale University Press, 1931).
- [42] A. Ashtekar and J. Lewandowski, Background independent quantum gravity: a status report, *Class. Quantum Grav.* **21**, R53 (2004).
- [43] J. Polchinski, *String Theory* (Cambridge University Press, 1998).

- [44] A. Connes, Gravity coupled with matter and the foundation of non-commutative geometry, *Commun. Math. Phys.* **182**, 155 (1996).

Appendix A: Lean 4 Excerpts

The following excerpts illustrate key machine-verified proofs from the Lean 4 formalization.

a. Note on code presentation. These are pedagogical summaries; the complete, machine-verifiable proofs are in the supplementary repository (run `lake build` to verify).

1. Topological Derivation of SU(3) (Theorem 0.0.15)

```

/-
  Theorem 0.0.15: Topological Derivation of SU(3)
  Status: SORRY-FREE (704 lines)

  Key constraints:
  1. Z_3 subset Z(G) (from phase structure)
  2. rank(G) <= 2 (from D = 4 spacetime)
  Result: G = SU(3) is UNIQUE.
-/

inductive LieGroupSeries
| A (n : Nat) -- SU(n+1)
| B (n : Nat) -- SO(2n+1)
| C (n : Nat) -- Sp(2n)
| D (n : Nat) -- SO(2n)
| G2 | F4 | E6 | E7 | E8

def SU3 : LieGroupSeries := .A 2

theorem SU3_has_Z3_center :
  SU3.centerContainsZ3 = true := by decide

theorem SU3_satisfies_rank :
  SU3.rank <= 2 := by norm_num

-- MAIN THEOREM
theorem topological_uniqueness_SU3 :
  forall G : LieGroupSeries,
    G.centerContainsZ3 /\ G.rank <= 2
    -> G = SU3 := by
  intro G (hcenter, hrank)
  cases G with
  | A n => interval_cases n <|> simp_all
  | _ => simp_all

```

Appendix B: Supplementary Remarks

1. Explicit Embedding Matrix

The weight-space-to-physical-space map is uniquely determined by the requirement of preserving the Killing metric. The weight plane $\mathfrak{h}^* \cong \mathbb{R}^2$ embeds perpendicular to the color-singlet direction $\hat{n} = (1, 1, 1)/\sqrt{3}$ via the orthonormal basis

$$\vec{e}_1 = \frac{1}{\sqrt{2}}(1, -1, 0), \quad \vec{e}_2 = \frac{1}{\sqrt{6}}(1, 1, -2), \quad (\text{B1})$$

giving the embedding matrix $M = (\vec{e}_1 \ \vec{e}_2)^T \in \mathbb{R}^{3 \times 2}$ with $M^T M = \mathbb{I}_2$ —confirming an isometric embedding (distances are preserved). This explicit construction verifies

that the abstract weight-space geometry of Theorem V.1 has a concrete, metric-preserving realization in $\mathbb{R}^3$.

2. Immirzi Parameter from Stella Geometry

The stella geometry yields a natural Immirzi-like parameter: $\gamma_{\text{CG}} = \sqrt{3} \ln(3)/(4\pi) \approx 0.151$, where $\sqrt{3}/4$ is a geometric factor (triangle area in natural units), $\ln(3)$ is the entropy factor (3 color states), and $1/\pi$ is the angular normalization. This value is comparable to loop quantum gravity estimates:

Source	γ	Origin
Chiral Geometrogenesis	0.151	SU(3) weight geometry
BKLR (1998)	0.127	Black hole entropy counting
Dreyer (2003)	0.124	Quasinormal modes
Barbero-Immirzi	0.20–0.24	SU(2) spin networks

The appearance of $\ln(3)$ in both γ_{CG} and Dreyer’s quasinormal mode calculation is suggestive: in CG, $3 = \dim(\mathbf{3})$; in LQG, 3 is the number of asymptotic quasinormal mode families—hinting at a deep connection between SU(3) color structure and black hole horizon degrees of freedom.

Appendix C: Notation and Conventions

Symbol Meaning

S	Stella octangula (pre-geometric substate)
∂S	Boundary $\partial T_+ \sqcup \partial T_-$
T_+, T_-	Matter / antimatter tetrahedra
$\mathbf{3}, \mathbf{\bar{3}}, \mathbf{8}$	SU(3) representations $\mathbf{3}, \mathbf{\bar{3}}, \mathbf{8}$
$\mathcal{W}$	Weyl group $W(\text{SU}(3)) \cong S_3$
$\mathbb{Z}_3$	Center of SU(3)
μ_c	Weight vectors ($c \in \{R, G, B\}$)
α_1, α_2	Simple roots of A_2
g_{ij}^K	Killing metric on weight space
O_h	Full octahedral symmetry (order 48)
$K_\ell(\hat{n})$	O_h -invariant cubic harmonics

a. Metric signature. Mostly-plus convention $(-, +, +, +)$.

b. Natural units. $\hbar = c = 1$ unless stated otherwise.

c. Index conventions. Greek $\mu, \nu, \dots$: spacetime 0, 1, 2, 3. Latin $i, j, \dots$: spatial 1, 2, 3. Color $c, c', \dots$: R, G, B .

d. Weight normalization. Standard Dynkin convention: longest roots have squared length 2; fundamental weights form equilateral triangle with unit side length; generators normalized as $\text{Tr}[T^a T^b] = \frac{1}{2} \delta^{ab}$.