

# Contents

<b>1</b>	<b>Introduction</b>	<b>2</b>
<b>2</b>	<b>Description of Model</b>	<b>2</b>
2.1	Description of purpose of the network in the model . . . . .	2
2.2	Description of Model Parameters . . . . .	3
2.3	Compliance . . . . .	4
<b>3</b>	<b>Childcare Model and Benchmark</b>	<b>4</b>
<b>4</b>	<b>Non-Strategic Model</b>	<b>6</b>
<b>5</b>	<b>Analysis of Testing Compliance</b>	<b>8</b>
<b>6</b>	<b>Strategic Model</b>	<b>8</b>
<b>7</b>	<b>Results</b>	<b>9</b>
<b>8</b>	<b>Discussion</b>	<b>9</b>
<b>9</b>	<b>Conclusion</b>	<b>9</b>
<b>A</b>	<b>Appendix Section</b>	<b>10</b>

# Agent-based models of non-pharmaceutical interventions for epidemic control

Robert Brian Milligan and Supervised by Julian Garcia Gallego & Buser Say

7 July 2022

## 1 Introduction

This paper investigates the modification of an existing model disease spread produced by Ryan McGee and used a part of various peer reviewed research articles. I have looked at applying the model to the situation of COVID-19 in childcare and involves creating both a non strategic and strategic behavioural models. The Behaviour relates to compliance to various actions the agent can choose to do such as doing an additional COVID rapid antigen test if they have symptoms of the virus.

## 2 Description of Model

### 2.1 Description of purpose of the network in the model

The model has two structures agents can be arranged in , one which uses a one single connected network designed to simulate a workplace, the other has a group of age segregated connected networks as well as small networks each agent is a part of that represents the household they are in. These large networks are made up of a number of cohorts which are loosely connected and each of these has a number of subgroups which are highly connected.

The network is set up before the simulation begins and does not change throughout the simulations run. 80% of transmission spread between agents occurs along the edges connecting them, while 20% of transmission is at random

Each day agents will be asked to do a test if their day has come up on a surveillance testing schedule, they show symptoms or have been contacted that they are a close contact.

During the day agents can spread the contagion to each other and can progress through the stages of the disease if they have it. They additionally have the choice to participate in contact tracing.

Agents will also be asked to isolate for one of six reasons. They or a group member develop a symptomatic case, returns a positive test or is told they are a close contact through using contact tracing




Figure 1: An Example Netowrk

These systems can effectively be disabled by overriding the compliance for them to be a large negative number, for example compliance with contact tracing can be set to -10 to disable the use of that system.

Example of the network of 100 agents To simulate the childcare scenario they are split into 5 groups of about 20 agents each with high connectivity inside the group and low connectivity between the groups The agents current state is shown by the nodes colour and the edges are that agents close contacts which the disease can spread easiest though the population

## 2.2 Description of Model Parameters

The Current Modifications of the model relate to allowing 10 of the Model Parameters that relate to Compliance to be dynamically updated each day dependent on a given rule. The 5 most relevent of these being.

- TESTING\_COMPLIANCE\_RATE\_SYMPOMATIC (what proportion of agents take a test immediately as a result of having symptoms)
- TESTING\_COMPLIANCE\_RATE\_RANDOM (what proportion of agents will do surveillance)

- ISOLATION\_COMPLIANCE\_RATE\_SYMPTOMATIC\_INDIVIDUAL (what proportion of agents will isolate given they have a symptomatic case)
- ISOLATION\_COMPLIANCE\_RATE\_POSITIVE\_INDIVIDUAL (what proportion of agents will isolate given a positive result from a test)
- ISOLATION\_COMPLIANCE\_RATE\_POSITIVE\_GROUPMATE (what proportion of agents in a group isolate given one of them has a positive result from a test)

A simple model might only use a few of these compliance parameters such as and is how the parameters work in the base model

- TESTING\_COMPLIANCE\_RATE\_SYMPTOMATIC = 0.5
- TESTING\_COMPLIANCE\_RATE\_RANDOM = 0.5
- ISOLATION\_COMPLIANCE\_RATE\_POSITIVE\_INDIVIDUAL = 1

## 2.3 Compliance

## 3 Childcare Model and Benchmark

- Parameter justification
- test false negative rate 0.36 <https://www.abc.net.au/news/2022-02-05/staff-children-preschool-childcare-protected-covid-national-plan/100805610>
- R0 mean of Omicron is 9.5, delta is 5.4 <https://www1.racgp.org.au/news/clinical/what-to-expect-from-a-third-omicron-wave-in-australia> <https://academic.oup.com/jtm/article/29/3/taac037/654>
- base compliance for behaviour is set to 0.5
- 100 agents in model split into 5 groups with high interconnectedness within the group and low connections between groups
- compliance increases for 2 actions, symptomatic test and regular interval surveillance tests
- The cost of compliance is fixed and there is a reward based on if connecting agents are isolated, hospitalised or a fatality
- from looking at early results since only 2-3 tests a week there is a delay and the higher the R0 the fewer chances they can have to change their mind about compliance
- the R0 of 9.5 or 5.4 may be too high as it assumes children spread the disease at the same rate as the general population as well as all disease spread occurring in childcare which it probably is not. Therefore 2 other R0 cases of 3 and 2 are used




Figure 2: Benchmark Run with Unchanging Compliance at 50% for regular twice weekly surveillance testing and testing if symptomatic with the resulting final network. 92 of the 100 agents received the infection




Figure 3: An Example Simplified Network

## 4 Non-Strategic Model

We have a fixed cost of compliance to an action and varying factors that can raise it. A mix of global and local behavioural factors can be added to make an agent more compliant. In the current build these are the known positive cases in the network in the past 14 days. The proportion of contact agents (those which share an edge) which are a fatality, hospitalised or in isolation. These values are taken away from the base cost and if the result is lower from a specified value the agent will be compliant to that action, otherwise they are not

For a simple example we have a situation for this central agent

TESTING\_COMPLIANCE\_RATE\_SYMPOMATIC = 0.5 TESTING\_COMPLIANCE\_RATE\_RANDOM = 0.5

14 day positive test rate for network = 0.02 = 2This agent's base aptitude +0.30 which can range (-0.5,0.5) Base cost = 0.5 Compliance =  $0.5 - (5 * 1/10) - (4 * 0.02) + 0.3 = 0.22$  In this case the agent is compliant will test if they develop a symptomatic case immediately will do surveillance testing if the neighbouring agent that entered hospital returns a positive test, our agent will enter isolation

the reward for agents can be based on a mix of both a local and global network situation as in the example runs

Of the 10 possible compliance parameters built into the existing mode, 4 are used these are, • immediately testing if the agent has a symptomatic case • using regular surveillance testing • isolating if a positive test is returned




Figure 4: Test Results of a non strategic model

## 5 Analysis of Testing Compliance

On average, what percentage of the population caught the contagion On average, how many days did it take for the outbreak to stop with 0 active cases

We can analyse compliance by comparing the effect varying levels have on the length a contagion actively spreads and what proportion of the population becomes infected. Firstly a baseline can be set to explore the parameter space, then further tests done to see the effect having compliance change as a result of the current known spread of the contagion in the network.

Over 3 R0 virus base reproduction rate was the model run with the most important find being the non-strategic behavioural model providing the most improvement over a base static level of compliance as the R0 is reduced, from a -2% improvement with an R0 of 9.5 to a 28% improvement with an R0 of 2. This is probably due to one of the aspects the model relating to the 14 day known positive cases recorded. With a low R0 in the system there are more chances for agents to test themselves before they pass on the disease and these reported cases incentivise even more agents to do symptomatic and surveillance testing. With extremely high R0 numbers it is observed that the non-strategic behavioural model has little effect on the total number infected. Across all cases it can be seen that the amount of time the simulation runs for is tied with how many agents get infected. Both the static values of compliance and non-strategic behavioural model follow this trend.

## 6 Strategic Model

The view of benefit and cost can be grouped into 3 categories

- 1. Global Situation of the model e.g. changes over time but is identical across all agents e.g. number of active cases or rate of change in spread rate
- 2. Structural Layout of the model, identical over time but varies between each agents e.g. number of close contacts or number of agents one can reach in 2 steps
- 3. Structural Situation of the model, changes over time and varies for each agent e.g. number of close contacts in a particular state or group of states like hospitalised, fatality, isolated etc.

for the following tests these can be used to create a personal benefit for each agent

it could be possible to use of a mix of these situations and in real life that may be the case with peoples decision making process

Structural layout of the network, for example number of close contacts an agent has



global situation for example the number of reported cases in the last 2 weeks , the number of actual cases in the last 2 weeks, the growth of the disease spread rate

local situation for example the number of neighbouring agents that are a fatality, hospitalised or isolated

benefit depends on change on rate of disease , number of agents around you that are hospitalised, fatality or isolated

one model is exclusively global, one is exclusively local

number of infections of those connected to me, 1 or 2 degrees , that are exposed or infectious

number of infections globally

agents will know the total percent of agents that are compliant and use that to work out how compliant they are

one universal curve for compliance , would this mean it has to be based off global factors ?

do I actually need to mathematically work out a curve and its intercepts

every day am I updating this curve? , can agents move to exactly where they want, or is it a slow process, they can only move X amount every day?, a limited amount makes sense as that would match the curves in other papers like thomas? Or would you see where the maxima is on that day and instantly assign the end that value?

come up with a equation (payoff function), x parameter to produce a curve like in the papers parameters go in , map plot, make it a notebook

does it change based on rate of change or on number of cases - if its global every curve would be the same if its local every curve is different for each agent notebook

how do curves change with infection numbers/rate

keep adding to latex doc.

once its a good function integrate it

Redo table of content Childcare first introduces compliance Describe the model

When you comply you pay a cost

Growth of disease rate

## 7 Results

## 8 Discussion

## 9 Conclusion

## A Appendix Section

Removed parameter section

- `TRACING_COMPLIANCE_RATE` (what proportion of agents comply with contact tracing)
- `ISOLATION_COMPLIANCE_RATE_POSITIVE_CONTACT` (what proportion of agents isolate given they are a close contact)
- `TESTING_COMPLIANCE_RATE_TRACED` (what proportion of agents take a test immediately as a result of being informed they are a close contact)
- `ISOLATION_COMPLIANCE_RATE_SYMPTOMATIC_GROUPMATE` (what proportion of agents will isolate given one of their group mates has a symptomatic case)
- `ISOLATION_COMPLIANCE_RATE_POSITIVE_CONTACTGROUPMATE` (what proportion of agents in a group isolate given one of them is a close contact)