

MATH 222 (Lectures 1,2, and 4) Fall 2015  
Midterm 1

Name: \_\_\_\_\_

Student ID#: \_\_\_\_\_

Circle your TA's name from the following list.

Carolyn Abbott

Tejas Bhojraj

Zachary Carter

Mohamed Abou Dbai

Ed Dewey

Jale Dinler

Di Fang

Bingyang Hu

Canberk Irimagzi

Chris Janjigian

Tao Ju

Ahmet Kabakulak

Dima Kuzmenko

Ethan McCarthy

Tung Nguyen

Jaeun Park

Adrian Tovar Lopez

Polly Yu

	Problem 1	Problem 2	Problem 3	Problem 4	Problem 5	Problem 6	Problem 7
Score							

### Instructions

- Write neatly on this exam. If you need extra paper, let us know.
- On Problems 1, 2, and 3, only the answer will be graded.
- On Problems 4, 5, 6, and 7 you must show your work and we will grade the work and your justification, and not just the final answer.
- Each problem worth either 14 or 15 points.
- No calculators, books, or notes (except for those notes on your 3 inch by 5 inch notecard.)
- Please simplify any formula involving a trigonometric function and an inverse trigonometric function. For example, please write  $\cos(\arcsin x) = \sqrt{1 - x^2}$ . Note that we have provided some formulas on the next page to help with this.

## Formulas

You may freely quote any algebraic or trigonometric identity, as well as any of the following formulas or minor variants of those formulas.

- $\cos(\arcsin x) = \sqrt{1 - x^2}$
- $\sec(\arctan x) = \sqrt{1 + x^2}$ .
- $\tan(\operatorname{arcsec} x) = \sqrt{x^2 - 1}$ .
- $\int x^n dx = \begin{cases} \frac{x^{n+1}}{n+1} + C & \text{when } n \neq -1 \\ \ln |x| + C & \text{when } n = -1 \end{cases}$
- $\int e^x dx = e^x + C$
- $\int \cos x dx = \sin x + C$
- $\int \sin x dx = -\cos x + C$
- $\int \tan x dx = -\ln |\cos x| + C$
- $\int \cot x dx = \ln |\sin x| + C$
- $\int \sec x dx = \ln |\sec x + \tan x| + C$ .
- $\int \csc x dx = -\ln |\csc x + \cot x| + C$ .
- $\int \frac{1}{1+x^2} dx = \arctan(x) + C$ .

1. For each statement below, CIRCLE true or false.

(a)		(b)		(c)		(d)		(e)	
True	False	True	False	True	False	True	False	True	False

(a) If  $\frac{x}{7} = \cos \theta$  then  $\tan \theta = \frac{\sqrt{49-x^2}}{x}$ .

(b)  $\int 3 \sin^2(\theta) d\theta = \frac{\sin^3 \theta}{\cos \theta} + C$

(c)  $\frac{1+\sin(x)}{x^3} \geq \frac{1}{x^3}$  for all  $x \geq 1$ .

(d)  $\int_2^\infty \frac{1}{x^2-9} dx$  is a finite number.

(e)  $\int_3^\infty \frac{x-\sqrt{x}}{3x^3+11} dx$  is a finite number.

2. On this page, only the answer will be graded.

(a) Compute  $\int \sin^2(x) - \cos^2(x) dx$ .

(b) Compute  $\int \frac{4}{(x-1)(3x+1)} dx$ .

(c) Compute  $\int_{-3}^{\infty} \frac{1}{x^2+6x+10} dx$ .

3. On this page, only the answer will be graded.

(a) Find a positive number  $A$  such that  $\int_{100}^{\infty} \frac{1}{x^2+73x-5} dx < A$ .

(b) Compute  $\int x e^{7x+1} dx$ .

(c) Compute  $\int \frac{1}{\sqrt{2x-x^2}} dx$ .

4. Compute  $\int_1^\infty \frac{4x+3}{x(2x+1)(2x+3)} dx$  or explain why the integral does not exist. (You may freely use the formula  $\frac{4x+3}{x(2x+1)(2x+3)} = \frac{1}{x} - \frac{1}{2x+1} - \frac{1}{2x+3}$ .)

5. Compute  $\int (z + e^z) \sin(3z) dz$ .

6. Compute  $\int e^{-x}\sqrt{4-e^{2x}}dx$ .



7. (a) For  $n = 0, 1, \dots$  let  $I_n = \int x^n e^{13x+2} dx$ . Derive a reduction formula for  $I_n$ .
- (b) Let  $J_n = \int x^5 (\ln x)^n dx$  for  $n \geq 0$ . This satisfies the reduction formula  $J_n = (\ln x)^n \frac{x^6}{6} - \frac{n}{6} J_{n-1}$  for  $n \geq 1$ . Compute  $J_2$ .

This page left blank for additional work.

This page left blank for additional work.