

MATH 222 (Lectures 1,2, and 4) Fall 2015  
**Midterm 1 Solutions**

Student ID#: \_\_\_\_\_

Circle your TA's name from the following list.

Carolyn Abbott      Tejas Bhojraj      Zachary Carter      Mohamed Abou Dbai      Ed Dewey

Jale Dinler      Di Fang      Bingyang Hu      Canberk Irimagzi      Chris Janjigian

Tao Ju      Ahmet Kabakulak      Dima Kuzmenko      Ethan McCarthy      Tung Nguyen

Jaeun Park      Adrian Tovar Lopez      Polly Yu

Please inform your TA if you find any errors in the solutions.

	Problem 1	Problem 2	Problem 3	Problem 4	Problem 5	Problem 6	Problem 7
Score							

**Instructions**

- Write neatly on this exam. If you need extra paper, let us know.
- On Problems 1, 2, and 3, only the answer will be graded.
- On Problems 4, 5, 6, and 7 you must show your work and we will grade the work and your justification, and not just the final answer.
- Each problem worth either 14 or 15 points.
- No calculators, books, or notes (except for those notes on your 3 inch by 5 inch notecard.)
- Please simplify any formula involving a trigonometric function and an inverse trigonometric function. For example, please write  $\cos(\arcsin x) = \sqrt{1 - x^2}$ . Note that we have provided some formulas on the next page to help with this.

## Formulas

You may freely quote any algebraic or trigonometric identity, as well as any of the following formulas or minor variants of those formulas.

- $\cos(\arcsin x) = \sqrt{1 - x^2}$
- $\sec(\arctan x) = \sqrt{1 + x^2}$ .
- $\tan(\operatorname{arcsec} x) = \sqrt{x^2 - 1}$ .
- $\int x^n dx = \begin{cases} \frac{x^{n+1}}{n+1} + C & \text{when } n \neq -1 \\ \ln |x| + C & \text{when } n = -1 \end{cases}$
- $\int e^x dx = e^x + C$
- $\int \cos x dx = \sin x + C$
- $\int \sin x dx = -\cos x + C$
- $\int \tan x dx = -\ln |\cos x| + C$
- $\int \cot x dx = \ln |\sin x| + C$
- $\int \sec x dx = \ln |\sec x + \tan x| + C$ .
- $\int \csc x dx = -\ln |\csc x + \cot x| + C$ .
- $\int \frac{1}{1+x^2} dx = \arctan(x) + C$ .

1. For each statement below, CIRCLE true or false.

(a)		(b)		(c)		(d)		(e)	
True	False	True	False	True	False	True	False	True	False

(a) If  $\frac{x}{7} = \cos \theta$  then  $\tan \theta = \frac{\sqrt{49-x^2}}{x}$ .

(b)  $\int 3 \sin^2(\theta) d\theta = \frac{\sin^3 \theta}{\cos \theta} + C$

(c)  $\frac{1+\sin(x)}{x^3} \geq \frac{1}{x^3}$  for all  $x \geq 1$ .

(d)  $\int_2^\infty \frac{1}{x^2-9} dx$  is a finite number.

(e)  $\int_3^\infty \frac{x-\sqrt{x}}{3x^3+11} dx$  is a finite number.

**Solution:**

(a) True.

(b) False.

(c) False.

(d) False.

(e) True.

2. On this page, only the answer will be graded.

(a) Compute  $\int \sin^2(x) - \cos^2(x) dx$ .

**Solution:**

$$\begin{aligned}\int \sin^2(x) - \cos^2(x) dx &= \int \sin^2(x) - (1 - \sin^2(x)) dx \\ &= \int (2 \sin^2(x) - 1) dx \\ &= \int (1 - \cos(2x)) - 1 dx \\ &= - \int \cos(2x) dx \\ &= -\frac{1}{2} \sin(2x) + C\end{aligned}$$

(b) Compute  $\int \frac{4}{(x-1)(3x+1)} dx$ .

**Solution:** We rewrite this in the form:

$$\int \frac{4}{(x-1)(3x+1)} dx = \int \frac{1}{x-1} - \frac{3}{3x+1}$$

Solving using the method of equating coefficients yields  $A = 1$  and  $B = -3$ .

(c) Compute  $\int_{-3}^{\infty} \frac{1}{x^2+6x+10} dx$ .

**Solution:**

$$\begin{aligned}\int_{-3}^{\infty} \frac{1}{x^2+6x+10} dx &= \lim_{b \rightarrow \infty} \int_{-3}^b \frac{1}{1+(x+3)^2} dx \\ &= \lim_{b \rightarrow \infty} [\arctan(x+3)]_{-3}^b \\ &= \lim_{b \rightarrow \infty} (\arctan(b+3) - \arctan(0)) = \pi/2.\end{aligned}$$

3. On this page, only the answer will be graded.

(a) Find a positive number  $A$  such that  $\int_{100}^{\infty} \frac{1}{x^2+73x-5} dx < A$ .

**Solution:** Any  $A$  bigger than .0075 will work.

(b) Compute  $\int x e^{7x+1} dx$ .

**Solution:** Let  $f = x$  and  $g' = e^{7x+1}$  so that  $f' = 1$  and  $g = \frac{1}{7}e^{7x+1}$ . Then

$$\begin{aligned}\int x e^{7x+1} dx &= \int f g' \\ &= f g - \int f' g \\ &= \frac{x}{7} e^{7x+1} - \frac{1}{7} \int e^{7x+1} dx \\ &= \frac{x}{7} e^{7x+1} - \frac{1}{49} e^{7x+1} + C\end{aligned}$$

(c) Compute  $\int \frac{1}{\sqrt{2x-x^2}} dx$ .

**Solution:** Complete the square to get  $2x - x^2 = 1 - (x - 1)^2$ . Then we get:

$$\int \frac{1}{\sqrt{2x-x^2}} dx = \int \frac{1}{\sqrt{1-(x-1)^2}} dx$$

Using  $x - 1 = \sin \theta$  and  $dx = \cos \theta d\theta$  this yields:

$$\begin{aligned}&= \int \frac{\cos \theta d\theta}{\sqrt{1-\sin^2 \theta}} \\ &= \int 1 d\theta \\ &= \theta + C \\ &= \arcsin(x - 1) + C.\end{aligned}$$

4. Compute  $\int_1^\infty \frac{4x+3}{x(2x+1)(2x+3)} dx$  or explain why the integral does not exist. (You may freely use the formula  $\frac{4x+3}{x(2x+1)(2x+3)} = \frac{1}{x} - \frac{1}{2x+1} - \frac{1}{2x+3}$ .)

**Solution:** We compute:

$$\begin{aligned}
 \int_1^\infty \frac{4x+3}{x(2x+1)(2x+3)} dx &= \int_1^\infty \frac{1}{x} - \frac{1}{2x+1} - \frac{1}{2x+3} dx \\
 &= \lim_{b \rightarrow \infty} \left[ \ln|x| - \frac{1}{2} \ln|2x+1| - \frac{1}{2} \ln|2x+3| \right]_1^b \\
 &= \lim_{b \rightarrow \infty} \left[ \ln \frac{|x|}{\sqrt{(2x+1)(2x+3)}} \right]_1^b \\
 &= \lim_{b \rightarrow \infty} \left[ \ln \frac{|x|}{\sqrt{4x^2 + 8x + 3}} \right]_1^b \\
 &= \lim_{b \rightarrow \infty} \ln \frac{b}{\sqrt{4b^2 + 8b + 3}} - \ln \frac{1}{\sqrt{15}} \\
 &= \ln\left(\frac{1}{\sqrt{4}}\right) - \ln\left(\frac{1}{15}\right) = \ln\left(\frac{\sqrt{15}}{2}\right)
 \end{aligned}$$

There are other equivalent answers.

5. Compute  $\int (z + e^z) \sin(3z) dz$ .

**Solution:** We compute this as the sum of two integrals:

$$\int (z + e^z) \sin(3z) dz = \int z \sin(3z) dz + \int e^z \sin(3z) dz$$

For  $\int z \sin(3z) dz$  we “double back”. First do integration by parts with  $f = z$  so  $f' = 1$  and  $g' = \sin(3z)$  so  $g = -\frac{1}{3} \cos(3z)$ . And we get:

$$\begin{aligned} \int z \sin(3z) dz &= fg - \int f'g \\ &= -\frac{1}{3}z \cos(3z) + \frac{1}{3} \int \cos(3z) dz \\ &= -\frac{1}{3}z \cos(3z) + \frac{1}{9} \sin(3z) + C \end{aligned}$$

Then we let  $I = \int e^z \sin(3z) dz$ . We first integrate by parts with  $f = \sin(3z)$  and  $g' = e^z$ . Then  $f' = 3 \cos(3z)$  and  $g = e^z$ , yielding:

$$\begin{aligned} I &= \int e^z \sin(3z) dz = fg - \int f'g \\ &= \sin(3z)e^z - 3 \int \cos(3z)e^z dz \end{aligned}$$

We integrate by parts again, with  $h = \cos(3z)$  and  $k' = e^z$  so  $h' = -3 \sin(3z)$  and  $k = e^z$ :

$$\begin{aligned} &= \sin(3z)e^z - 3 \int h k' \\ &= \sin(3z)e^z - 3 \left( h k - \int h' k \right) \\ &= \sin(3z)e^z - 3 \left( \cos(3z)e^z - \int (-3 \sin(3z))e^z dz \right) \\ &= \sin(3z)e^z - 3 \cos(3z)e^z - 9 \int \sin(3z)e^z dz \\ &= \sin(3z)e^z - 3 \cos(3z)e^z - 9I \end{aligned}$$

We thus have the equation:

$$I = \sin(3z)e^z - 3 \cos(3z)e^z - 9I$$

which, after moving all of the  $I$  terms to the left side, yields:

$$(1 + 9)I = \sin(3z)e^z - 3 \cos(3z)e^z + C.$$

We thus obtain:

$$I = \frac{1}{10} (\sin(3z)e^z - 3 \cos(3z)e^z) + C$$

Putting this together yields:

$$\int (z + e^z) \sin(3z) dz = -\frac{1}{3}z \cos(3z) + \frac{1}{9} \sin(3z) + \frac{1}{10} (\sin(3z)e^z - 3 \cos(3z)e^z)$$

6. Compute  $\int e^{-x} \sqrt{4 - e^{2x}} dx$ .

**Solution:** Set  $z = e^x$  so that  $dz = e^x dx$  and  $dx = \frac{dz}{e^x} = \frac{dz}{z}$ . Then we have:

$$\int e^{-x} \sqrt{4 - e^{2x}} dx = \int z^{-2} \sqrt{4 - z^2} dz$$

Now let  $z = 2 \sin \theta$  so that  $dz = 2 \cos \theta d\theta$  and we get:

$$\begin{aligned} &= \int (2 \sin \theta)^{-2} \sqrt{4 - 4 \sin^2 \theta} 2 \cos \theta d\theta \\ &= \int \frac{4 \cos^2 \theta}{4 \sin^2 \theta} d\theta \\ &= \int \frac{\cos^2 \theta}{\sin^2 \theta} d\theta \\ &= \int (\cot^2 \theta - 1) d\theta \\ &= \int \csc^2 \theta - 1 d\theta \\ &= -\cot \theta - \theta + C \end{aligned}$$

Since  $\sin(\theta) = \frac{z}{2}$  we get  $\theta = \arcsin(\frac{z}{2})$  and  $\cot(\theta) = \frac{\sqrt{4-z^2}}{z}$  yielding:

$$\begin{aligned} &= -\frac{\sqrt{4 - z^2}}{z} - \arcsin\left(\frac{z}{2}\right) + C \\ &= -\frac{\sqrt{4 - e^{2x}}}{e^x} - \arcsin\left(\frac{e^x}{2}\right) + C \end{aligned}$$



7. (a) For  $n = 0, 1, \dots$  let  $I_n = \int x^n e^{13x+2} dx$ . Derive a reduction formula for  $I_n$ .
- (b) Let  $J_n = \int x^5 (\ln x)^n dx$  for  $n \geq 0$ . This satisfies the reduction formula  $J_n = (\ln x)^n \frac{x^6}{6} - \frac{n}{6} J_{n-1}$  for  $n \geq 1$ . Compute  $J_2$ .

**Solution:**

(a): Let  $f = x^n$  so  $f' = nx^{n-1}$  and let  $g' = e^{13x+2}$  so that  $g = \frac{1}{13}e^{13x+2}$ . Then:

$$\begin{aligned} I_n &= \int x^n e^{13x+2} dx = \int f g' \\ &= fg - \int f' g \\ &= \frac{1}{13} x^n e^{13x+2} - \frac{n}{13} \int x^{n-1} e^{13x+2} dx \\ &= \frac{1}{13} x^n e^{13x+2} - \frac{n}{13} I_{n-1} \end{aligned}$$

(b):  $J_0 = \int x^5 dx = \frac{x^6}{6} + C$ . Then

$$J_1 = (\ln x)^1 \frac{x^6}{6} - \frac{1}{6} J_0 = \frac{1}{6} x \ln x - \frac{x^6}{36} + C.$$

Then

$$J_2 = (\ln x)^2 \frac{x^6}{6} - \frac{2}{6} J_1 = (\ln x)^2 \frac{x^6}{6} - \frac{1}{3} \left( \frac{1}{6} x \ln x - \frac{x^6}{36} \right) + C$$

If you simplify, you might get other, equivalent, answers.