

# MASTER REFERENCE LIST

---

- C.M. Ablow and J.L. Brenner[1963], Roots and canonical forms for circulant matrices, *Trans. Amer. Math. Soc.*, 107, pp. 360–376.
- R. Ahlswede and D.E. Daykin, The number of values of combinatorial functions, *Bull. London Math. Soc.*, 11, pp. 49–51.
- A.V. Aho, J.E. Hopcroft and J.D. Ullman[1975], *The Design and Analysis of Computer Algorithms*, Addison-Wesley, Reading, Mass.
- N. Alon, R.A. Brualdi and B.L. Shader[1991], Multicolored trees in bipartite decompositions of graphs, *J. Combin. Theory, Ser. B*, to be published.
- S.A. Amitsur and J. Levitzki[1950], Minimal identities for algebras, *Proc. Amer. Math. Soc.*, 1, pp. 449–463.
- [1951], Remarks on minimal identities for algebras, *Proc. Amer. Math. Soc.*, 2, pp. 320–327.
- L.D. Andersen and A.J.W. Hilton[1983], Thanks Evans!, *Proc. London Math. Soc.* (3), 47, pp. 507–522.
- I. Anderson[1971], Perfect matchings of graphs, *J. Combin. Theory*, 10, pp. 183–186.
- R.P. Anstee[1982], Properties of a class of  $(0,1)$ -matrices covering a given matrix, *Canad. J. Math.*, 34, pp. 438–453.
- M. Aschbacher[1971], The non-existence of rank three permutation groups of degree 3250 and subdegree 57, *J. Algebra*, 19, pp. 538–540.
- M.J. Atallah[1982], Finding the cycle index of an irreducible, nonnegative matrix, *SIAM J. Computing*, 11, pp. 567–570.
- K.B. Athreya, C.B. Pranesachar and N.M. Singhi[1980], On the number of latin rectangles and chromatic polynomial of  $L(K_{r,s})$ , *Europ. J. Combinatorics*, 1, pp. 9–17.
- K. Balasubramanian[1990], On transversals in latin squares, *Linear Alg. Applics.*, 131, pp. 125–129.
- S.E. Bammel and J. Rothstein[1975], The number of  $9 \times 9$  latin squares, *Discrete Math.*, 11, pp. 93–95.
- E. Bannai and T. Ito[1973], On finite Moore graphs, *J. Fac. Sci. Univ. Tokyo*, 20, pp. 191–208.
- L. Bassett, J. Maybee and J. Quirk[1983], Qualitative economics and the scope of the correspondence principle, *Econometrica*, 26, pp. 544–563.

- V.J. Baston[1982], A Marica-Schönheim theorem for an infinite sequence of finite sets, *Math. Z.*, 179, pp. 531–533.
- C. Berge[1958], Sur le couplage maximum d'un graphe, *C.R. Acad. Sciences (Paris)*, 247, pp. 258–259.
- [1976], *Graphs and Hypergraphs*, North Holland, Amsterdam.
- S. Beslin and S. Ligh[1989], Greatest common divisor matrices, *Linear Alg. Appls.*, 118, pp. 69–76.
- T. Beth[1983], Eine Bemerkung zur Abschätzung der Anzahl orthogonaler lateinischer Quadrate mittels Siebverfahren, *Abh. Math. Sem. Hamburg*, 53, pp. 284–288.
- N. Biggs[1974], *Algebraic Graph Theory*, Cambridge Tracts in Mathematics No. 67, Cambridge University Press, Cambridge.
- G. Birkhoff[1946], Tres observaciones sobre el algebra lineal, *Univ. Nac. Tucumán Rev. Ser. A*, pp. 147–151.
- B. Bollobás[1979], *Graph Theory*, Springer-Verlag, New York.
- V.I. Bol'shakov[1986], Upper values of a permanent in  $\Lambda_n^k$ , *Combinatorial analysis, No. 7 (Russian)*, Moskov. Gos. Univ., Moscow, pp. 92–118 and 164–165.
- [1986], The spectrum of the permanent on  $\Lambda_n^k$ , *Proceedings of the All-Union seminar on discrete mathematics and its applications, Moscow (1984)*, (Russian), Moskov. Gos. Univ. Mekh.-Mat. Fak., Moscow, pp. 65–73.
- J.A. Bondy and U.S.R. Murty[1976], *Graph Theory with Applications*, North-Holland, New York.
- C.W. Borchardt[1860], Über eine der Interpolation entsprechende Darstellung der Eliminations-Resultante, *J. Reine Angew. Math.*, 57, pp. 111–121.
- R.C. Bose[1938], On the application of the properties of Galois fields to the problem of construction of hyper-Graeco-Latin squares, *Sankhyā*, 3, pp. 323–338.
- R.C. Bose and S.S. Shrikhande[1959], On the falsity of Euler's conjecture about the non-existence of two orthogonal Latin squares of order  $4t + 2$ , *Proc. Nat. Acad. Sci. U.S.A.*, 45, pp. 734–737.
- [1960], On the composite of balanced incomplete block designs, *Canad. J. Math.*, 12, pp. 177–188.
- [1960], On the construction of sets of mutually orthogonal Latin squares and the falsity of a conjecture of Euler, *Trans. Amer. Math. Soc.*, 95, pp. 191–209.
- R.C. Bose, S.S. Shrikhande and E.T. Parker[1960], Further results on the construction of mutually orthogonal Latin squares and the falsity of Euler's conjecture, *Canad. J. Math.*, 12, pp. 189–203.
- A. Brauer[1947], Limits for the characteristic roots of a matrix II, *Duke Math. J.*, 14, pp. 21–26.
- A. Brauer and I.C. Gentry[1968], On the characteristic roots of tournament matrices, *Bull. Amer. Math. Soc.*, 74, pp. 1133–1135.
- R.K. Brayton, D. Coppersmith and A.J. Hoffman[1974], Self-orthogonal latin squares of all orders  $n \neq 2, 3, 6$ , *Bull. Amer. Math. Soc.*, 80, pp. 116–118.
- L.M. Brégman[1973], Certain properties of nonnegative matrices and their permanents, *Dokl. Akad. Nauk SSSR*, 211, pp. 27–30. (*Soviet Math. Dokl.*, 14, pp. 945–949.)
- W.G. Bridges[1971], The polynomial of a non-regular digraph, *Pacific J. Math.*, 38, pp. 325–341.
- [1972], The regularity of  $x^3$ -graphs, *J. Combin. Theory, Ser. B*, 12, pp. 174–176.

- W.G. Bridges and R.A. Mena[1981],  $x^k$ -digraphs, *J. Combin. Theory, Ser. B*, 30, pp. 136–143.
- A. Brouwer, A. Cohen and A. Neumaier[1989], *Distance Regular Graphs*, Springer-Verlag, Berlin.
- A.E. Brouwer, A.J. de Vries and R.M.A. Wieringa[1978], A lower bound for the length of partial transversals in a latin square, *Nieuw Archief voor Wiskunde* (3), XXVI, pp. 330–332.
- R.A. Brualdi[1966], Term rank of the direct product of matrices, *Canad. J. Math.*, 18, pp. 126–138.
- [1966], Permanent of the direct product of matrices, *Pacific J. Math.*, 16, pp. 471–482.
- [1971], Matchings in arbitrary graphs, *Proc. Cambridge Phil. Soc.*, 69, pp. 401–407.
- [1976], Combinatorial properties of symmetric non-negative matrices, *Teorie Combinatoire*, Toma II, Accademia Nazionale dei Lincei, Roma, pp. 99–120.
- [1977], *Introductory Combinatorics*, Elsevier Science Publishers, New York.
- [1979], Matrices permutation equivalent to irreducible matrices and applications, *Linear Multilin. Alg.*, 7, pp. 1–12.
- [1979], The diagonal hypergraph of a matrix (bipartite graph), *Discrete Math.*, 27, pp. 127–147.
- [1980], On the diagonal hypergraph of a matrix, *Annals of Discrete Math.*, 8, pp. 261–264.
- [1982], Matrices, eigenvalues, and directed graphs, *Linear Multilin. Alg.*, 11, pp. 143–165.
- [1985], Combinatorial verification of the elementary divisors of tensor products, *Linear Alg. Applies.*, 71, pp. 31–47.
- [1987], The Jordan canonical form: an old proof, *Amer. Math. Monthly*, 94, pp. 257–267.
- [1987], Combinatorially determined elementary divisors, *Congressus Numerantium*, 58, pp. 193–216.
- [1988], Counting permutations with restricted positions: permanents of (0,1)-matrices. A tale in four parts. In The 1987 Utah State University Department of Mathematics Conference Report by L. Beasley and E.E. Underwood, *Linear Alg. Applies.*, 104, pp. 173–183.
- [1988], In Memoriam Herbert J. Ryser 1923–1985, *J. Combin. Theory, Ser. A*, 47, pp. 1–5.
- [1990], The many facets of combinatorial matrix theory, in *Matrix Theory and Applications*, C.R. Johnson ed., Proc. Symposia Pure and Applied Math. vol. 40, Amer. Math. Soc., Providence.
- [1992], The symbiotic relationship of combinatorics and matrix theory, *Linear Alg. Applies.*, to be published.
- R.A. Brualdi and K.L. Chavey[1991], Elementary divisors and ranked posets with application to matrix compounds, *Linear Multilin. Alg.*, to be published.
- R.A. Brualdi and J. Csima[1986], Extending subpermutation matrices in regular classes of matrices, *Discrete Math.*, 62, pp. 99–101.
- [1991], Butterfly embedding proof of a theorem of König, *Amer. Math. Monthly*, to be published.

- R.A. Brualdi and P.M. Gibson[1977], Convex polyhedra of doubly stochastic matrices I. Applications of the permanent function, *J. Combin. Theory, Ser. A*, 22, pp. 194–230.
- R.A. Brualdi, J.L. Goldwasser and T.S. Michael[1988], Maximum permanents of matrices of zeros and ones, *J. Combin. Theory, Ser. A*, 47, pp. 207–245.
- R.A. Brualdi, F. Harary and Z. Miller[1980], Bigraphs versus digraphs via matrices, *J. Graph Theory*, 4, pp. 51–73.
- R.A. Brualdi and M.B. Hedrick[1979], A unified treatment of nearly reducible and nearly decomposable matrices, *Linear Alg. Applies.*, 24, pp. 51–73.
- R.A. Brualdi and M. Lewin[1982], On powers of nonnegative matrices, *Linear Alg. Applies.*, 43, pp. 87–97.
- R.A. Brualdi and B. Liu[1991], Fully indecomposable exponents of primitive matrices, *Proc. Amer. Math. Soc.*, to be published.
- [1990], Hall exponents of Boolean matrices, *Czech. Math. J.*, 40 (115), pp. 663–674.
- [1990], Generalized exponents of primitive directed graphs, *J. Graph Theory*, 14, pp. 483–499.
- R.A. Brualdi and T.S. Michael[1989], The class of 2-multigraphs with a prescribed degree sequence, *Linear Multilin. Alg.*, 24, pp. 81–102.
- R.A. Brualdi and M. Newman[1965], Some theorems on the permanent, *J. Res. National Bur. Stands.*, 69B, pp. 159–163.
- R.A. Brualdi, S.V. Parter and H. Schneider[1966], The diagonal equivalence of a nonnegative matrix to a stochastic matrix, *J. Math. Anal. Applies.*, 16, pp. 31–50.
- R.A. Brualdi and J.A. Ross[1980], On the exponent of a primitive, nearly reducible matrix, *Math. Oper. Res.*, 5, pp. 229–241.
- [1981], Matrices with isomorphic diagonal hypergraphs, *Discrete Math.*, 33, pp. 123–138.
- R.A. Brualdi and B.L. Shader[1990], Matrix factorizations of determinants and permanents, *J. Combin. Theory, Ser. A*, 54, pp. 13–134.
- [1991], On converting the permanent into the determinant and sign-nonsingular matrices, *Applied Geometry and Discrete Mathematics* (P. Gritzmann and B. Sturmfels, eds.), to be published.
- R.H. Bruck[1951], Finite nets. I. Numerical invariants, *Canad. J. Math.*, 3, pp. 94–107.
- [1963], Finite nets. II. Uniqueness and imbedding, *Pacific J. Math.*, 13, pp. 421–457.
- T. Brylawski[1975], Modular constructions for combinatorial geometries, *Trans. Amer. Math. Soc.*, 203, pp. 1–44.
- D. de Caen and D. Gregory[1987], On the decomposition of a directed graph into complete bipartite subgraphs, *Ars Combinatoria*, 23B, pp. 139–146.
- D. de Caen and D.G. Hoffman[1989], Impossibility of decomposing the complete graph on  $n$  points into  $n - 1$  isomorphic complete bipartite graphs, *SIAM J. Disc. Math.*, 2, pp. 48–50.
- P.J. Cameron[1978], Strongly regular graphs, *Selected Topics in Graph Theory* (L.W. Beineke and R.J. Wilson, eds.), Academic Press, New York, pp. 337–360.
- P.J. Cameron, J.-M. Goethals and J.J. Seidel[1978], Strongly regular graphs

- having strongly regular subconstituents, *J. Algebra*, 55, pp. 257–280.
- P.J. Cameron, J.M. Goethals, J.J. Seidel and E.E. Shult[1976], Line graphs, root systems, and elliptic geometry, *J. Algebra*, 43, pp. 305–327.
- P.J. Cameron and J.H. van Lint[1975], *Graph Theory, Coding Theory and Block Designs*, London Math. Soc. Lecture Note Series No.19, Cambridge University Press, Cambridge.
- P. Camion[1965], Characterization of totally unimodular matrices, *Proc. Amer. Math. Soc.*, 16, pp. 1068–1073.
- P. Cartier and D. Foata[1969], *Problèmes Combinatoires de Commutation et Réarrangement*, Lec. Notes in Math. No. 85, Springer, Berlin.
- A. Cayley[1854], Sur les déterminantes gauches, *Crelle's J.*, 38, pp. 93–96.
- [1889], A theorem on trees, *Quarterly J. Math.*, 23, pp. 376–378.
- S. Chaiken[1982], A combinatorial proof of the all minors matrix tree theorem, *SIAM J. Alg. Disc. Meth.*, 3, pp. 319–329.
- C.Y. Chao[1977], On a conjecture of the semigroup of fully indecomposable relations, *Czech. Math. J.*, 27, pp. 591–597.
- C.Y. Chao and T. Wang[1987], On the matrix equation  $A^2 = J$ , *J. Math. Res. and Exposition*, 2, pp. 207–215.
- C.Y. Chao and M.C. Zhang[1983], On the semigroup of fully indecomposable relations, *Czech. Math. J.*, 33, pp. 314–319.
- G. Chaty and M. Chein[1976], A note on top down and bottom up analysis of strongly connected digraphs, *Discrete Math.*, 16, pp. 309–311.
- W.C. Chu[1985], Some algebraic identities concerning determinants and permanents, *Linear Alg. Applics.*, 116, pp. 35–40.
- F.R.K. Chung, P. Diaconis, R.L. Graham and C.L. Mallows[1981], On the permanents of complements of the direct sum of identity matrices, *Advances in Applied Math.*, 2, pp. 121–137.
- V. Chungphaisan[1974], Conditions for sequences to be  $r$ -graphic, *Discrete Math.*, 7, pp. 31–39.
- C.L. Coates[1959], Flow graph solutions of linear algebraic equations, *IEEE Trans. Circuit Theory*, CT-6, pp. 170–187.
- C.C. Colburn[1984], The complexity of completing partial latin squares, *Discrete Applied Math.*, 8, pp. 25–30.
- S.A. Cook[1971], The complexity of theorem proving procedures, *Proc. 3rd ACM Symp. on Theory of Computing*, pp. 151–158.
- A.B. Cruse[1974], On embedding incomplete symmetric latin squares, *J. Combin. Theory, Ser. A*, 16, pp. 18–22.
- J. Csima[1979], On the plane term rank of three dimensional matrices, *Discrete Math.*, 28, pp. 147–152.
- D.M. Cvetković[1975], The determinant concept defined by means of graph theory, *Mat. Vesnik*, 12, pp. 333–336.
- D.M. Cvetković, M. Doob and H. Sachs[1982], *Spectra of Graphs – Theory and Application*, 2nd ed., Deutscher Verlag der Wissenschaften, Berlin, Academic Press, New York.
- D. Cvetković, M. Doob, I. Gutman and A. Torgâsev[1988], *Recent Results in the Theory of Graph Spectra*, Annals of Discrete Mathematics No. 36, Elsevier Science Publishers, New York.
- R.H. Damerell[1973], On Moore graphs, *Proc. Cambridge Phil. Soc.*, 74, pp. 227–236.

- [1983], On Smetaniuk's construction for latin squares and the Andersen-Hilton theorem, *Proc. London Math. Soc.* (3), 47, pp. 523–526.
- P.J. Davis[1979], *Circulant Matrices*, John Wiley & Sons, New York.
- D.E. Daykin and R. Häggvist[1981], Problem No. 6347, *Amer. Math. Monthly*, 88, p. 446.
- D.E. Daykin and L. Lovász[1976], The number of values of a Boolean function, *J. London Math. Soc.*, 12 (2), pp. 225–230.
- E.V. Denardo[1977], Periods of connected networks, *Math. Oper. Res.*, 2, pp. 20–24.
- J. Dénes and A.D. Keedwell[1974], *Latin Squares and Their Applications*, Academic Press, New York.
- J. Donald, J. Elwin, R. Hager and P. Salamon[1984], A graph-theoretic upper bound on the permanent of a nonnegative matrix, *Linear Alg. Applics.*, 61, pp. 187–198.
- M. Doob[1984], Applications of graph theory in linear algebra, *Math. Mag.*, 57, pp. 67–76.
- A.L. Dulmage and N.S. Mendelsohn[1958], Coverings of bipartite graphs, *Canad. J. Math.*, 10, pp. 517–534.
- [1958], Some generalizations of the problem of distinct representatives, *Canad. J. Math.*, 10, pp. 230–241.
- [1959], A structure theory of bipartite graphs of finite exterior dimension, *Trans. Roy. Soc. Canad.*, Third Ser. Sec. III 53, pp. 1–3.
- [1962], The exponent of a primitive matrix, *Canad. Math. Bull.*, 5, pp. 642–656.
- [1963] The characteristic equation of an imprimitive matrix, *SIAM J. Appl. Math.*, 11, pp. 1034–1045.
- [1964], Gaps in the exponent set of primitive matrices, *Illinois J. Math.*, 8, pp. 642–656.
- [1964], The exponents of incidence matrices, *Duke Math. J.*, 31, pp. 575–584.
- [1967], Graphs and matrices, *Graph Theory and Theoretical Physics* (F. Harary, ed.), Academic Press, New York, pp. 167–227.
- J. Edmonds[1967], Systems of distinct representatives and linear algebra, *J. Res. Nat. Bur. Standards*, 71B, pp. 241–245.
- G.P. Egor'čev[1981], A solution of van der Waerden's permanent problem, *Dokl. Akad. Nauk SSSR*, 258, pp. 1041–1044. (*Soviet Math. Dokl.*, 23, pp. 619–622.)
- P. Erdős and T. Gallai[1960], Graphs with prescribed degrees of vertices, *Mat. Lapok*, 11, pp. 264–274 (in Hungarian).
- P. Erdős, D.R. Hickerson, D.A. Norton and S.K. Stein[1988], Has every latin square of order  $n$  a partial transversal of size  $n - 1$ ?, *Amer. Math. Monthly*, 95, pp. 428–430.
- P. Erdős, A. Rényi and V.T. Sós[1966], On a problem of graph theory, *Studia Sci. Math. Hungar.*, 1, pp. 215–235.
- R. Euler, R.E. Burkhard and R. Grommes[1986], On latin squares and the facial structure of related polytopes, *Discrete Math.*, 62, pp. 155–181.
- T. Evans[1960], Embedding incomplete latin squares, *Amer. Math. Monthly*, 67, pp. 958–961.
- S. Even[1979], *Graph Algorithms*, Computer Science Press, Potomac, Maryland.
- C.J. Everett and P.R. Stein[1973], The asymptotic number of  $(0,1)$ -matrices with zero permanent, *Discrete Math.*, 6, pp. 29–34.

- D.I. Falikman[1981], A proof of van der Waerden's conjecture on the permanent of a doubly stochastic matrix, *Mat. Zametki*, 29, pp. 931–938. (*Math. Notes*, 29, pp. 475–479.)
- M. Fiedler[1973], Algebraic connectivity of graphs, *Czech. Math. J.*, 23, pp. 298–305.
- [1975], A property of eigenvectors of nonnegative symmetric matrices and its application to graph theory, *Czech. Math. J.*, 25, pp. 619–633.
- [1990], Absolute connectivity of trees, *Linear Multilin. Alg.*, 26, pp. 86–106.
- S. Fiorini and R.J. Wilson[1977], *Edge-colorings of graphs*, Pitman, London.
- J. Folkman and D.R. Fulkerson[1969], Edge colorings in bipartite graphs, *Combinatorial Mathematics and Their Applications* (R.C. Bose and T. Dowling, eds.), University of North Carolina Press, Chapel Hill, pp. 561–577.
- L.R. Ford, Jr. and D.R. Fulkerson[1962], *Flows in Networks*, Princeton University Press, Princeton.
- T.H. Foregger[1975], An upper bound for the permanent of a fully indecomposable matrix, *Proc. Amer. Math. Soc.*, 49, pp. 319–324.
- G. Frobenius[1896], Über die Primfactoren der Gruppendeterminante, *Sitzungsber. der Königlich Preuss. Akad. Wiss. Berlin*, pp. 1343–1382.
- [1879], Theorie der linearen Formen mit ganzen Coefficienten, *J. für reine und angew. Math.*, 86, pp. 146–208.
- [1917], Über zerlegbare Determinanten, *Sitzungsber. Preuss. Akad. Wiss. Berlin*, pp. 274–277.
- [1912], Über Matrizen aus nicht negativen Elementen, *Sitzungsber. Preuss. Akad. Wiss. Berlin*, pp. 476–457.
- D.R. Fulkerson[1960], Zero-one matrices with zero trace, *Pacific J. Math.*, 10, pp. 831–836.
- [1964], The maximum number of disjoint permutations contained in a matrix of zeros and ones, *Canad. J. Math.*, 10, pp. 729–735.
- [1970], Blocking Polyhedra, *Graph Theory and Its Applications* (B. Harris, ed.), Academic Press, New York, pp. 93–111.
- D.R. Fulkerson, A.J. Hoffman and M.H. McAndrew[1965], Some properties of graphs with multiple edges, *Canad. J. Math.*, 17, pp. 166–177.
- D. Gale[1957], A theorem on flows in networks, *Pacific J. Math.*, 7, pp. 1073–1082.
- T. Gallai[1963], Neuer Beweis eines Tutte'schen Satzes, *Magyar Tud. Akad. Közl.*, 8, pp. 135–139.
- E.R. Gansner[1981], Acyclic digraphs, Young tableaux and nilpotent matrices, *SIAM J. Alg. Disc. Meth.*, 2, pp. 429–440.
- F.R. Gantmacher[1959], *The Theory of Matrices*, vol. 2, Chelsea, New York.
- A.M.H. Gerards[1989], A short proof of Tutte's characterization of totally unimodular matroids, *Linear Alg. Applies.*, 114/115, pp. 207–212.
- I. Gessel[1985], Counting three-line latin rectangles, *Proc. Colloque de Combinatoire Énumérative*, UQAM, pp. 106–111.
- [1987], Counting latin rectangles, *Bull. Amer. Math. Soc. (New Series)*, 16, pp. 79–82.
- P.M. Gibson[1966], A short proof of an inequality for the permanent function, *Proc. Amer. Math. Soc.*, 17, pp. 535–536.
- [1971], Conversion of the permanent into the determinant, *Proc. Amer. Math. Soc.*, 27, pp. 471–476.



- [1972], A lower bound for the permanent of a  $(0,1)$ -matrix, *Proc. Amer. Math. Soc.*, 33, pp. 245–246.
- [1972], The pfaffian and 1-factors of graphs, *Trans. N.Y. Acad. Sci.*, 34, pp. 52–57.
- [1972], The pfaffian and 1-factors of graphs II, *Graph Theory and Applications* (Y. Alavi, D.R. Lick and A.T. White, eds.), *Lec. Notes in Mathematics*, No. 303, pp. 89–98, Springer-Verlag, New York.
- J.-M. Goethals and J.J. Seidel[1967], Orthogonal matrices with zero diagonal, *Canad. J. Math.*, 19, pp. 1001–1010.
- [1970], Strongly regular graphs derived from combinatorial designs, *Canad. J. Math.*, 22, pp. 597–614.
- R.L. Graham and H.O. Pollak[1971], On the addressing problem for loop switching, *Bell System Tech. J.*, 50, pp. 2495–2519.
- [1972], On embedding graphs in squashed cubes, *Lecture Notes in Math*, vol. 303, Springer-Verlag, New York, pp. 99–110.
- R. Grone[1991], On the geometry and Laplacian of a graph, *Linear Alg. Applics.*, to be published.
- B. Grone and R. Merris[1987], Algebraic connectivity of trees, *Czech. Math. J.*, 37, pp. 660–670.
- R. Guérin[1968], Existence et propriétés des carrés latins orthogonaux II, *Publ. Inst. Statist. Univ. Paris*, 15, pp. 215–293.
- R.P. Gupta[1967], On basis diagraphs, *J. Comb. Theory*, 3, pp. 16–24.
- [1967], A decomposition theorem for bipartite graphs, *Théorie des Graphes Rome I.C.C.* (P. Rosenstiehl, ed.), Dunod, Paris, pp. 135–138.
- [1974], On decompositions of a multigraph with spanning subgraphs, *Bull. Amer. Math. Soc.*, 80, pp. 500–502.
- [1978], An edge-coloration theorem for bipartite graphs with applications, *Discrete Math.*, 23, pp. 229–233.
- H. Hadwiger, H. Debrunner and V. Klee[1964], *Combinatorial Geometry in the Plane*, Holt, Rinehart and Winston, New York.
- W. Haemers[1979], *Eigenvalue Techniques in Design and Graph Theory*, Mathematisch Centrum, Amsterdam.
- R. Häggvist[1978], A solution to the Evans conjecture for latin squares of large size, *Combinatorics*, Proc. Conf. on Combinatorics, Kesthely (Hungary) 1976, János Bolyai Math. Soc. and North Holland, pp. 495–513.
- S.L. Hakimi[1962], On realizability of a set of integers as degrees of the vertices of a linear graph I, *J. Soc. Indust. Appl. Math.*, 10, pp. 496–506.
- M. Hall, Jr.[1945], An existence theorem for latin squares, *Bull. Amer. Math. Soc.* 51, pp. 387–388.
- [1948], Distinct representatives of subsets, *Bull. Amer. Math. Soc.*, 54, pp. 922–926.
- [1952], A combinatorial problem on abelian groups, *Proc. Amer. Math. Soc.*, 3, pp. 584–587.
- [1986], *Combinatorial Theory*, 2nd edition, Wiley, New York,
- P. Hall[1935], On representatives of subsets, *J. London Math. Soc.*, 10, pp. 26–30.
- H. Hanani[1970], On the number of orthogonal latin squares, *J. Combin. Theory*, 8, pp. 247–271.



- F. Harary[1959], A graph theoretic method for the complete reduction of a matrix with a view toward finding its eigenvalues, *J. Math. and Physics*, 38, pp. 104–111.
- [1962], The determinant of the adjacency matrix of a graph, *SIAM Review*, 4, pp. 202–210.
- [1967], Graphs and matrices, *SIAM Review*, 9, pp. 83–90.
- [1969], *Graph Theory*, Addison-Wesley, Reading, Mass.
- F. Harary, R. Norman and D. Cartwright[1965], *Structural Models*, Wiley, New York.
- F. Harary and E.M. Palmer[1973], *Graphical Enumeration*, Academic Press, New York.
- D.J. Hartfiel[1970], A simplified form for nearly reducible and nearly decomposable matrices, *Proc. Amer. Math. Soc.*, 24, pp. 388–393.
- [1971], On constructing nearly decomposable matrices, *Proc. Amer. Math. Soc.*, 27, pp. 222–228.
- [1975], A canonical form for fully indecomposable (0,1)-matrices, *Canad. Math. Bull.*, 18, pp. 223–227.
- D.J. Hartfiel and R. Loewy[1984], A determinantal version of the Frobenius-König theorem, *Linear Multilin. Alg.*, 16, pp. 155–165.
- V. Havel[1955], A remark on the existence of finite graphs (in Hungarian), *Časopis Pěst. Mat.*, 80, pp. 477–480.
- B.R. Heap and M.S. Lynn[1964], The index of primitivity of a non-negative matrix, *Numer. Math.*, 6, pp. 120–141.
- M. Hedrick and R. Sinkhorn[1970], A special class of irreducible matrices—the nearly reducible matrices, *J. Algebra*, 16, pp. 143–150.
- J.R. Henderson[1975], Permanents of (0,1)-matrices having at most two 0's per line, *Canad. Math. Bull.*, 18, pp. 353–358.
- P. Heymans[1969], Pfaffians and skew-symmetric matrices, *Proc. London Math. Soc.* (3), 19, pp. 730–768.
- D.G. Higman[1971], Partial geometries, generalized quadrangles and strongly regular graphs, *Atti di Conv. Geometria Combinatoria e sue Applicazione* (A. Barlotti, ed.), Perugia, pp. 263–293.
- A.J. Hoffman[1960], On the uniqueness of the triangular association scheme, *Ann. Math. Statist.*, 31, pp. 492–497.
- [1960], Some recent applications of the theory of linear inequalities to extremal combinatorial analysis, *Proc. Symp. in Applied Mathematics*, vol. 10, Amer. Math. Soc., pp. 113–127.
- [1963], On the polynomial of a graph, *Amer. Math. Monthly*, 70, pp. 30–36.
- [1967], Research problem 2-11, *J. Combin. Theory*, 2, p. 393.
- [1970],  $-1 - \sqrt{2}$ ? *Combinatorial Structures and Their Applications*, Gordon and Breach, New York, pp. 173–176.
- [1977], On graphs whose least eigenvalue exceeds  $-1 - \sqrt{2}$ , *Linear Alg. Applics.*, 16, pp. 153–165.
- A.J. Hoffman and J.B. Kruskal[1956], Integral boundary points of convex polyhedra, *Annals of Math. Studies*, No. 38, Princeton University Press, Princeton, pp. 223–246.
- A.J. Hoffman and M.H. McAndrew[1965], The polynomial of a directed graph, *Proc. Amer. Math. Soc.*, 16, pp. 303–309.

- A.J. Hoffman and D.K. Ray-Chaudhuri, On a spectral characterization of regular line graphs, unpublished manuscript.
- J.C. Holladay and R.S. Varga[1958], On powers of non-negative matrices, *Proc. Amer. Math. Soc.*, 9, pp. 631–634.
- R. Horn and C.R. Johnson[1985], *Matrix Analysis*, Cambridge University Press, Cambridge.
- X.L. Hubaut[1975], Strongly regular graphs, *Discrete Math.*, 13, pp. 357–381.
- C.G. Jacobi[1841], De determinantibus functionalibus, *J. Reine Angew. Math.*, 22, pp. 319–359.
- K. Jones, J.R. Lundgren, N.J. Pullman and R. Rees[1988], A note on the covering numbers of  $K_n - K_m$  and complete  $t$ -partite graphs, *Congressus Num.*, 66, pp. 181–184.
- D.D. Joshi[1987], *Linear Estimation and Design of Experiments*, Wiley, New York.
- A.-A.A. Jucys[1976], The number of distinct latin squares as a group-theoretical constant, *J. Combin. Theory, Ser. A*, 20, pp. 265–272.
- W.B. Jurkat and H.J. Ryser[1966], Matrix factorizations of determinants and permanents, *J. Algebra*, 3, pp. 1–27.
- [1967], Term ranks and permanents of nonnegative matrices, *J. Algebra*, 5, pp. 342–357.
- [1968], Extremal configurations and decomposition theorems, *J. Algebra*, 8, pp. 194–222.
- R. Kallman[1982], A method for finding permanents of 0,1 matrices, *Maths. of Computation*, 38, pp. 167–170.
- I. Kaplansky[1939], A generalization of the ‘Problème des rencontres,’ *Amer. Math. Monthly*, 46, pp. 159–161.
- [1943], Solution of the “Problème des ménages,” *Bull. Amer. Math. Soc.*, 49, pp. 784–785.
- [1944], Symbolic solution of certain problems in permutations, *Bull. Amer. Math. Soc.*, 50, pp. 906–914.
- R.M. Karp[1972], Reducibility among combinatorial problems, In *Complexity of Computer Calculations* (R.E. Miller and J.W. Thatcher, eds.), Plenum, New York, pp. 85–104.
- P.W. Kasteleyn[1961], The statistics of dimers on a lattice, *Physica*, 27, pp. 1209–1225.
- [1967], Graph theory and crystal physics, *Graph Theory and Theoretical Physics* (F. Harary, ed.), Academic Press, New York, pp. 43–110.
- J.G. Kemeny and J.L. Snell[1960], *Finite Markov Chains*, Van Nostrand, Princeton.
- D.G. Kendall[1969], Incidence matrices, interval graphs and seriation in archaeology, *Pacific J. Math.*, 28, pp. 565–570.
- F. King and K. Wang[1985], On the  $g$ -circulant solutions to the matrix equation  $A^m = \lambda J$ , II, *J. Combin. Theory, Ser. A*, 38, pp. 182–186.
- V. Klee, R. Ladner and R. Manber[1983], Signsolvability revisited, *Linear Alg. Applies.*, 59, pp. 131–157.
- D.E. Knuth[1970], Notes on central groupoids, *J. Combin. Theory*, 8, pp. 376–390.
- [1974], Wheels within wheels, *J. Combin. Theory, Ser. B*, 16, pp. 42–46.
- K.K. Koksma[1969], A lower bound for the order of a partial transversal, *J. Combin. Theory*, 7, pp. 94–95.

- J. Komlós[1967], On the determinant of  $(0,1)$ -matrices, *Studia Sci. Math. Hung.*, 2, pp. 7–21.
- D. König[1916], Über Graphen und ihre Anwendung auf Determinantentheorie und Mengenlehre, *Math. Ann.*, 77, pp. 453–465.
- [1936], *Theorie der endlichen und unendlichen Graphen*, Leipzig, reprinted by Chelsea[1960], New York.
- J.P.S. Kung[1984], Jacobi's identity and the König-Egerváry theorem, *Discrete Math.*, 49, pp. 75–77.
- C.W.H. Lam[1975], A generalization of cyclic difference sets I, *J. Combin. Theory, Ser. A*, 19, pp. 51–65.
- [1975], A generalization of cyclic difference sets II, *J. Combin. Theory, Ser. A*, 19, pp. 177–191.
- [1977], On some solutions of  $A^k = dI + \lambda J$ , *J. Combin. Theory, Ser. A*, 23, pp. 140–147.
- C.W.H. Lam and J.H. van Lint[1978], Directed graphs with unique paths of fixed length, *J. Combin. Theory, Ser. B*, 24, pp. 331–337.
- C.W.H. Lam, L.H. Thiel and S. Swiercz[1989], The nonexistence of finite projective planes of order 10, *Canad. J. Math.*, XLI, pp. 1117–1123.
- H. Laue[1988], A graph theoretic proof of the fundamental trace identity, *Discrete Math.*, 69, pp. 197–198.
- J.S. Lew[1966], The generalized Cayley-Hamilton theorem in  $n$  dimensions, *Z. Angew. Math. Phys.*, 17, pp. 650–653.
- M. Lewin[1971], On exponents of primitive matrices, *Numer. Math.*, 18, pp. 154–161.
- [1971], On nonnegative matrices, *Pacific. J. Math.*, 36, pp. 753–759.
- [1974], Bounds for the exponents of doubly stochastic matrices, *Math. Zeit.*, 137, pp. 21–30.
- M. Lewin and Y. Vitek[1981], A system of gaps in the exponent set of primitive matrices, *Illinois J. Math.*, 25, pp. 87–98.
- Chang Li-Chien[1959], The uniqueness and non-uniqueness of the triangular association scheme, *Sci. Record. Peking Math.* (New Ser.), 3, pp. 604–613.
- [1960], Associations of partially balanced designs with parameters  $v = 28$ ,  $n_1 = 12$ ,  $n_2 = 15$ , and  $p_{11}^2 = 4$ , *Sci. Record. Peking Math.*, (New Ser.), 4, pp. 12–18.
- E.H. Lieb[1968], A theorem on Pfaffians, *J. Combin. Theory*, 5, pp. 313–368.
- C.C. Lindner[1970], On completing latin rectangles, *Canad. Math. Bull.*, 13, pp. 65–68.
- J.H. van Lint[1985],  $(0, 1, *)$  distance problems in Combinatorics, *Surveys in Combinatorics* (I. Anderson, ed.), London Math. Soc. Lecture Notes 103, Cambridge University Press, pp. 113–135.
- J.H. van Lint and J.J. Seidel[1966], Equilateral point sets in elliptic geometry, *Nederl. Akad. Wetensch. Proc, Ser. A*, 69 (=Indag. Math., 28), pp. 335–348.
- C.H.C. Little[1975], A characterization of convertible  $(0,1)$ -matrices, *J. Combin. Theory, Ser. B*, 18, pp. 187–208.
- C.H.C. Little and J.M. Pla[1972], Sur l'utilisation d'un pfaffien dans l'étude des couplages parfaits d'un graphe, *C. R. Acad. Scie. Paris*, 274, p. 447.
- B. Liu[1990], A note on the exponents of primitive  $(0,1)$ -matrices, *Linear Alg. Applies.*, 140, pp. 45–51.

- [1991], New results on the exponent set of primitive, nearly reducible matrices, *Linear Alg. Applies*, to be published.
- B. Liu, B. McKay, N. Wormwald and K.M. Zhang[1990], The exponent set of symmetric primitive  $(0,1)$ -matrices with zero trace, *Linear Alg. Applies*, 133, pp. 121–131.
- L. Lovász[1970], Subgraphs with prescribed valencies, *J. Combin. Theory*, 8, pp. 391–416.
- [1979], On determinants, matchings, and random algorithms, *Proceedings of Fundamentals of Computation Theory*, Akademie Verlag, Berlin, pp. 565–574.
- L. Lovász and M.D. Plummer[1977], On minimal elementary bipartite graphs, *J. Combin. Theory, Ser. A*, 23, pp. 127–138.
- R.D. Luce[1952], Two decomposition theorems for a class of finite oriented graphs, *Amer. J. Math.*, 74, pp. 701–722.
- S.L. Ma and W.C. Waterhouse[1987], The  $g$ -circulant solutions of  $A^m = \lambda J$ , *Linear Alg. Applies.*, 85, pp. 211–220.
- P.A. MacMahon[1915], *Combinatory Analysis*, vol. 1, Cambridge University Press, Cambridge.
- H.F. MacNeish[1922], Euler squares, *Ann. Math.*, 23, pp. 221–227.
- W. Mader[1973], Grad und lokaler Zusammenhang in endlichen Graphen, *Math. Ann.*, 205, pp. 9–11.
- H.B. Mann[1942], The construction of orthogonal latin squares, *Ann. Math. Statist.*, 13, pp. 418–423.
- M. Marcus and H. Minc[1961], On the relation between the determinant and the permanent, *Illinois J. Math.*, 5, pp. 376–381.
- [1963], Disjoint pairs of sets and incidence matrices, *Illinois J. Math*, 7, pp. 137–147.
- [1964], *A Survey of Matrix Theorems and Matrix Inequalities*, Allyn and Bacon, Boston.
- J. Marica and J. Schönheim[1969], Differences of sets and a problem of Graham, *Canad. Math. Bull.*, 12, pp. 635–637.
- J.S. Maybee[1981], Sign solvability, in *Computer assisted analysis and model simplification* (H. Greenberg and J. Maybee, eds.), Academic Press, New York.
- P.K. Menon[1961], Method of constructing two mutually orthogonal latin squares of order  $3n + 1$ , *Sankhyā A*, 23, pp. 281–282.
- D. Merriell[1980], The maximum permanent in  $\Lambda_n^k$ , *Linear Multilin. Alg.*, 9, pp. 81–91.
- H. Minc[1963], Upper bound for permanents of  $(0,1)$ -matrices, *Bull. Amer. Math. Soc.*, 69, pp. 789–791.
- [1967], A lower bound for permanents of  $(0,1)$ -matrices, *Proc. Amer. Math. Soc.*, 18, pp. 1128–1132.
- [1969], On lower bounds for permanents of  $(0,1)$ -matrices, *Proc. Amer. Math. Soc.*, 22, pp. 117–123.
- [1969], Nearly decomposable matrices, *Linear Alg. Applies.*, 5, pp. 181–187.
- [1973],  $(0,1)$ -matrices with minimal permanents, *Israel J. Math.*, 77, pp. 27–30.
- [1974], An unresolved conjecture on permanents of  $(0,1)$ -matrices, *Linear Multilin. Alg.*, 2, pp. 57–64.
- [1974], The structure of irreducible matrices, *Linear Multilin. Algebra*, 2, pp. 85–90.

- [1978], *Permanents*, Addison-Wesley, Reading, Mass.
- L. Mirsky[1968], Combinatorial theorems and integral matrices, *J. Combin. Theory*, 5, pp. 30–44.
- [1971], *Transversal Theory*, Academic Press, New York.
- B. Mohar[1988], The Laplacian spectrum of graphs, *Preprint Series Dept. Math. University E.K. Ljubljana*, 26, pp. 353–382.
- J.W. Moon and L. Moser[1966], Almost all (0,1)-matrices are primitive, *Studia Scient. Math. Hung.*, 1, pp. 153–156.
- J.W. Moon and N.J. Pullman[1967], On the powers of tournament matrices, *J. Combin. Theory*, 3, pp. 1–9.
- T. Muir[1882], On a class of permanent symmetric functions, *Proc. Roy. Soc. Edinburgh*, 11, pp. 409–418.
- [1897], A relation between permanents and determinants, *Proc. Roy. Soc. Edinburgh*, 22, pp. 134–136.
- K. Murota[1989], On the irreducibility of layered mixed matrices, *Linear Multilin. Alg.*, 24, pp. 273–288.
- J.R. Nechvatal[1981], Asymptotic enumeration of generalized latin rectangles, *Utilitas Math.*, 20, pp. 273–292.
- A. Nijenhuis[1976], On permanents and the zeros of rook polynomials, *J. Combin. Theory, Ser. A*, 21, pp. 240–244.
- A. Nijenhuis and H. Wilf[1970], On a conjecture of Ryser and Minc, *Nederl. Akad. Wetensch. Proc., Ser. A*, 73 (= *Indag. Math.*, 32), pp. 151–158.
- [1978], *Combinatorial Algorithms*, second ed., Academic Press, New York.
- G.N. de Oliveira[1974], Note on the characteristic roots of tournament matrices, *Linear Alg. Applics.*, 8, pp. 271–272.
- W.E. Opencomb[1984], On the intricacy of combinatorial problems, *Discrete Math.*, 50, pp. 71–97.
- J. Orlin[1977], Contentment in graph theory: covering graphs with cliques, *Indag. Math.*, 39, pp. 406–424.
- [1978], Line-digraphs, Arborescences, and theorems of Tutte and Knuth, *J. Combin. Theory, Ser. B*, 25, pp. 187–198.
- P.A. Ostrand[1970], Systems of distinct representatives II, *J. Math. Anal. Applics.*, 32, pp. 1–4.
- A. Ostrowski[1937], Über die Determinanten mit überwiegender Hauptdiagonale, *Comm. Math. Helv.*, 10, pp. 69–96.
- [1951], Über das Nichtverschwinden einer Klassen von Determinanten und die Lokalisierung der charakteristischen Wurzeln von Matrizen, *Compositio Math.*, 9, pp. 209–226.
- M.W. Padberg[1976], A note on the total unimodularity of matrices, *Discrete Math.*, 14, pp. 273–278.
- L.J. Paige[1947], A note on finite abelian groups, *Bull. Amer. Math. Soc.*, 53, pp. 590–593.
- E.T. Parker[1959], Orthogonal Latin squares, *Proc. Nat. Acad. Sci.*, 45, pp. 859–862.
- [1959], Construction of some sets of mutually orthogonal Latin squares, *Proc. Amer. Math. Soc.*, 10, pp. 946–949.
- [1990], private communication.
- G.W. Peck[1984], A new proof of a theorem of Graham and Pollak, *Discrete Math.*, 49, pp. 327–328.

- H. Perfect[1966], Symmetrized form of P. Hall's theorem on distinct representatives, *Quart. J. Math. Oxford* (2), 17, pp. 303–306.
- J.M. Pla[1965], Sur l'utilisation d'un pfaffien dans l'étude des couplages parfaits d'un graphe, *C. R. Acad. Sci. Paris*, 260, pp. 2967–2970.
- H. Poincaré[1901], Second complément à l'analysis situs, *Proc. London Math. Soc.*, 32, pp. 277–308.
- G. Pólya[1913], Aufgabe 424, *Arch. Math. Phys.* (3), 20, p. 271.
- C.R. Pranesachar[1981], Enumeration of latin rectangles via SDR's, *Combinatorics and Graph Theory* (S.B. Rao, ed.), Lecture Notes in Math., 885, Springer-Verlag, Berlin and New York, pp. 380–390.
- C. Procesi[1976], The invariant theory of  $n \times n$  matrices, *Advances in Math.*, 19, pp. 306–381.
- V. Pták[1958], On a combinatorial theorem and its applications to nonnegative matrices, *Czech. Math. J.*, 8, pp. 487–495.
- V. Pták and J. Sedláček[1958], On the index of imprimitivity of non-negative matrices, *Czech. Math. J.*, 8, pp. 496–501.
- R. Rado[1967], On the number of systems of distinct representatives, *J. London Math. Soc.*, 42, pp. 107–109.
- D. Raghavarao[1971], *Construction and combinatorial problems in design of experiments*, Wiley, New York (reprinted[1988] by Dover, Mineola, NY).
- Yu. P. Ramyslov[1974], Trace identities of full matrix algebras over a field of characteristic zero, *Math. USSR-Izv.*, 8, pp. 727–760.
- J. Riordan[1946], Three-line latin rectangles-II, *Amer. Math. Monthly*, 53, pp. 18–20.
- V. Romanovsky[1936], Recherches sur les Chains de Markoff, *Acta. Math.*, 66, pp. 147–251.
- D. Rosenblatt[1957], On the graphs and asymptotic forms of finite boolean relation matrices and stochastic matrices, *Naval Research Logistics Quarterly*, 4, pp. 151–167.
- J.A. Ross[1980], *Some problems in combinatorial matrix theory*, Ph.D. thesis, University of Wisconsin, Madison.
- [1982], On the exponent of a primitive, nearly reducible matrix II, *SIAM J. Alg. Disc. Meth.*, 3, pp. 385–410.
- L.R. Rowen[1980], *Polynomial Identities in Ring Theory*, Academic Press, New York.
- D.E. Rutherford[1964], The Cayley-Hamilton theorem for semirings, *Proc. Roy. Soc. Edinburgh Sec. A*, 66, pp. 211–215.
- H.J. Ryser[1951], A combinatorial theorem with an application to latin rectangles, *Proc. Amer. Math. Soc.*, 2, pp. 550–552.
- [1957], Combinatorial properties of matrices of 0's and 1's, *Canad. J. Math.*, 9, pp. 371–377.
- [1960], Matrices of zeros and ones, *Bull. Amer. Mat. Soc.*, 66, pp. 442–464.
- [1963], *Combinatorial Mathematics*, Carus Mathematical Monograph No. 14, Math. Assoc. of Amer., Washington, D.C.
- [1967], Neuere Probleme in der Kombinatorik (prepared by D.W. Miller), *Vorträge über Kombinatorik*, Oberwolfach, pp. 69–91.
- [1969], Combinatorial configurations, *SIAM J. Appl. Math.*, 17, pp. 593–602.
- [1970], A generalization of the matrix equation  $A^2 = J$ , *Linear Alg. Applics.*, 3, pp. 451–460.

- [1972], A fundamental matrix equation for finite sets, *Proc. Amer. Math. Soc.*, 34, pp. 332–336.
- [1973], Analogs of a theorem of Schur on matrix transformations, *J. Algebra*, 25, pp. 176–184.
- [1973], Indeterminates and incidence matrices, *Linear Multilin. Alg.*, 1, pp. 149–157.
- [1974], Indeterminates and incidence matrices, in *Combinatorics*, Math. Centre Tracts Vol. 55, Mathematisch Centrum, Amsterdam, pp. 3–17.
- [1975], The formal incidence matrix, *Linear Multilin. Alg.*, 3, pp. 99–104.
- [1982], Set intersection matrices, *J. Combin. Theory, Ser. A*, 32, pp. 162–177.
- [1984], Matrices and set differences, *Discrete Math.*, 49, pp. 169–173.
- H. Sachs[1967], Über Teiler, Faktoren und charakteristische Polynome von Graphen II, *Wiss. Z. Techn. Hochsch. Ilmenau*, 13, pp. 405–412.
- M. Saks[1986], Some sequences associated with combinatorial structures, *Discrete Math.*, 59, pp. 135–166.
- H. Schneider[1977], The concepts of irreducibility and full indecomposability of a matrix in the works of Frobenius, König and Markov, *Linear Alg. Applics.*, 18, pp. 139–162.
- [1986], The influence of the marked reduced graph of a nonnegative matrix on the Jordan form and on related properties: A survey, *Linear Alg. Applics.*, 84, pp. 161–189.
- A. Schrijver[1978], A short proof of Minc's conjecture, *J. Combin. Theory, Ser. A*, 25, pp. 80–83.
- [1983], Bounds on permanents, and the number of 1-factors and 1-factorizations of bipartite graphs, *Surveys in Combinatorics*, Cambridge University Press, Cambridge, pp. 107–134.
- [1986], *Theory of Linear and Integer Programming*, Wiley, New York.
- A. Schrijver and W.G. Valiant[1980], On lower bounds for permanents, *Indag. Math.*, 42, pp. 425–427.
- A. Schuchat[1984], Matrix and network models in archaeology, *Math. Magazine*, 57, pp. 3–14.
- S. Schwarz[1973], The semigroup of fully indecomposable relations and Hall relations, *Czech. Math. J.*, 23, pp. 151–163.
- A.J. Schwenk and R.J. Wilson[1978], On the eigenvalues of a graph, *Selected Topics in Graph Theory* (L.W. Beineke and R.J. Wilson, eds.), Academic Press, New York, pp. 307–336.
- J. Sedláček[1959], O incidenčních maticích orientovaných grafů, *Časop. Pěst. Mat.*, 84, pp. 303–316.
- J.J. Seidel[1968], Strongly regular graphs with  $(-1, 1, 0)$  adjacency matrix having eigenvalue 3, *Linear Alg. Applics.*, 1, pp. 281–298.
- [1969], Strongly regular graphs, *Recent Progress in Combinatorics* (W.T. Tutte, ed.), Academic Press, New York, pp. 185–198.
- [1974], Graphs and two-graphs, *Proceedings of the Fifth Southeastern Conference on Combinatorics, Graph Theory and Computing*, Congressus Numerantium X, *Utilitas Math.*, Winnipeg, pp. 125–143.
- [1976], A survey of two-graphs, *Teorie Combinatorie*, Tomo I, (B. Segre, ed.), Accademia Nazionale dei Lincei, Rome, pp. 481–511.
- [1979], Strongly regular graphs, *Surveys in Combinatorics, Proc. 7th British*



- Combinatorial Conference*, London Math. Soc. Lecture Note Ser. 38 (B. Bollobás, ed.), Cambridge University Press, Cambridge.
- P.D. Seymour[1980], Decomposition of regular matroids, *J. Combin. Theory, Ser. B*, 28, pp. 305–359.
- [1982], Applications of the regular matroid decomposition, *Colloquia Math. Soc. János Bolyai*, No. 40 Matroid Theory, Szeged (Hungary), pp. 345–357.
- P. Seymour and C. Thomassen[1987], Characterization of even directed graphs, *J. Combin. Theory, Ser. B*, 42, pp. 36–45.
- C.E. Shannon[1949], A theorem on coloring the lines of a network, *J. Math. Phys.*, 28, pp. 148–151.
- J.Y. Shao[1985], On a conjecture about the exponent set of primitive matrices, *Linear Alg. Applies.*, 65, pp. 91–123.
- [1985], On the exponent of a primitive digraph, *Linear Alg. Applies.*, 64, pp. 21–31.
- [1985], Matrices permutation equivalent to primitive matrices, *Linear Alg. Applies.*, 65, pp. 225–247.
- [1987], The exponent set of symmetric primitive matrices, *Scientia Sinica Ser. A*, vol. XXX, pp. 348–358.
- P.W. Shor[1982], A lower bound for the length of a partial transversal in a latin square, *J. Combin Theory, Ser. A*, 33, pp. 1–8.
- S.S. Shrikhande[1959], The uniqueness of the  $L_2$  association scheme, *Ann. Math. Statist.*, 30, pp. 781–798.
- [1961], A note on mutually orthogonal latin squares, *Sankhyā A*, 23, pp. 115–116.
- R. Sinkhorn and P. Knopp[1969], Problems involving diagonal products in non-negative matrices, *Trans. Amer. Math. Soc.*, 136, pp. 67–75.
- B. Smetaniuk[1981], A new construction for latin squares I. Proof of the Evans conjecture, *Ars Combinatoria*, 11, pp. 155–172.
- [1982], A new construction on latin squares-II: The number of latin squares is strictly increasing, *Ars Combinatoria*, 14, pp. 131–145.
- H.J.S. Smith[1876], On the value of a certain arithmetical determinant, *Proc. London Math. Soc.*, 7, pp. 208–212.
- J.H. Smith[1970], Some properties of the spectrum of a graph, *Combinatorial Structures and Their Applications* (R. Guy, N. Sauer and J. Schönheim, eds.), Gordon and Breach, New York, pp. 403–406.
- R.P. Stanley[1980], Weyl groups, the hard Lefschetz theorem, and the Sperner property, *SIAM J. Alg. Disc. Methods*, 1, pp. 168–184.
- [1986], *Enumerative Combinatorics*, vol. I, Wadsworth & Brooks/Cole, Monterey.
- H. Straubing[1983], A combinatorial proof of the Cayley-Hamilton theorem, *Discrete Math.*, 43, pp. 273–279.
- R.G. Swan[1963], An application of graph theory to algebra, *Proc. Amer. Math. Soc.*, 14, pp. 367–373.
- [1969], Correction to “An application of graph theory to algebra,” *Proc. Amer. Math. Soc.*, 21, pp. 379–380.
- G. Szegő[1913], Zu Aufgabe 424, *Arch. Math. Phys.* (3), 21, pp. 291–292.
- R.E. Tarjan[1972], Depth first search and linear graph algorithms, *SIAM J. Computing*, 1, pp. 146–160.

- G. Tarry[1900,1901], Le problème de 36 officiers, *Compte Rendu de l'Association Française pour l'Avancement de Science Naturel*, 1, pp. 122–123, and 2, pp. 170–203.
- O. Taussky[1949], A recurring theorem on determinants, *Amer. Math. Monthly*, 10, pp. 672–676.
- H.N.V. Temperley[1964], On the mutual cancellation of cluster integrals in Mayer's fugacity series, *Proc. Phys. Soc.*, 83, pp. 3–16.
- [1981], *Graph Theory and Applications*, Ellis Horwood, Chichester.
- C. Thomassen[1981], A remark on the factor theorems of Lovász and Tutte, *J. Graph Theory*, 5, pp. 441–442.
- [1986], Sign-nonsingular matrices and even cycles in directed graphs, *Linear Alg. Applics.*, 75, pp. 27–41.
- J. Touchard[1934], Sur un problème de permutations, *C. R. Acad. Sci. Paris*, 198, pp. 631–633.
- W.T. Tutte[1947], The factorization of linear graphs, *J. London Math. Soc.*, 22, pp. 107–111.
- [1948], The dissection of equilateral triangles into equilateral triangles, *Proc. Cambridge Philos. Soc.*, 44, pp. 463–482.
- [1952], The factors of graphs, *Canadian J. Math.*, 4, pp. 314–328.
- [1958], A homotopy theorem for matroids, I and II, *Trans. Amer. Math. Soc.*, 88, pp. 144–174.
- [1978], The subgraph problem, *Discrete Math.*, 3, pp. 289–295.
- [1981], Graph factors, *Combinatorica*, 1, pp. 79–97.
- [1984], Graph Theory, *Encyclopedia of Mathematics and Its Applications*, vol. 21, Addison-Wesley, Reading, Mass.
- H. Tverberg[1982], On the decomposition of  $K_n$  into complete bipartite graphs, *J. Graph Theory*, 6, pp. 493–494.
- L.G. Valiant[1979], Completeness classes in algebra, *Proc. 11th Ann. ACM Symp. Theory of Computing*, pp. 249–261.
- [1979], The complexity of computing the permanent, *Theoret. Comput. Sci.*, 8, pp. 189–201.
- R.S. Varga[1962], *Matrix Iterative Analysis*, Prentice-Hall, Englewood Cliffs, N.J.
- D. Vere-Jones[1984], An identity involving permanents, *Linear Alg. Applics.*, 63, pp. 267–270.
- [1988], A generalization of permanents and determinants, *Linear Alg. Applics.*, 111, pp. 119–124.
- V.G. Vizing[1964], On an estimate of the chromatic class of a p-graph (in Russian), *Diskret. Analiz.*, 3, pp. 25–30.
- [1965], The chromatic class of a multigraph (in Russian), *Cybernetics*, 3, pp. 32–41.
- W. Vogel[1963], Bemerkungen zur Theorie der Matrizen aus Nullen und Einsen, *Archiv der Math.*, 14, pp. 139–144.
- J. von zur Gathen[1987], Permanent and determinant, *Linear Alg. Applics.*, 96, pp. 87–100.
- [1987], Feasible arithmetic computations: Valiant's hypothesis, *J. Symbolic Comp.*, 4, pp. 137–172.
- M. Voorhoeve[1979], A lower bound for the permanents of certain (0,1)-matrices, *Indag. Math.*, 41, pp. 83–86.
- A.J. de Vries and R.M.A. Wieringa[1978], Een ondergrens voor de lengte van een partiele transversaal in een Latijns vierkant, preprint.

- K. Wang[1980], On the matrix equation  $A^m = \lambda J$ , *J. Combin. Theory, Ser. A*, 29, pp. 134–141.
- [1981], A generalization of group difference sets and the matrix equation  $A^m = dI + \lambda J$ , *Aequationes Math.*, 23, pp. 212–222.
- [1982], On the g-circulant solutions to the matrix equation  $A^m = \lambda J$ , *J. Combin. Theory, Ser. A*, 33, pp. 287–296.
- N. White (ed.)[1986], Theory of Matroids, *Encyclopedia of Maths. and Its Applics.*, Cambridge University Press, Cambridge.
- [1987], Unimodular matroids, Combinatorial Geometries (N. White, ed.), *Encyclopedia of Maths. and Its Applics.*, Cambridge University Press, Cambridge.
- H. Wielandt[1958], Unzerlegbare, nicht negative Matrizen, *Math. Zeit.*, 52, pp. 642–645.
- H.S. Wilf[1968], A mechanical counting method and combinatorial applications, *J. Combin. Theory*, 4, pp. 246–258.
- [1986], *Algorithms and Complexity*, Prentice-Hall, Englewood Cliffs, N.J.
- R.J. Wilson[1985], *Introduction to Graph Theory*, 3rd edition, Longman, Harlow, Essex.
- R.M. Wilson[1974], Concerning the number of mutually orthogonal latin squares, *Discrete Math.*, 9, pp. 181–198.
- [1974], A few more squares, *Congressus Numerantium*, No. X, pp. 675–680.
- D.E. Woolbright[1978], An  $n \times n$  latin square has a transversal with at least  $n - \sqrt{n}$  distinct symbols, *J. Combin. Theory, Ser. A*, 24, pp. 235–237.
- D.E. Woolbright and H.-L. Fu[1987], The rainbow theorem of 1-factorization, preprint.
- S. Yang and G.P. Barker[1988], On the exponent of a primitive, minimally strong digraph, *Linear Alg. Applics.*, 99, pp. 177–198.
- D. Zeilberger[1985], A combinatorial approach to matrix algebra, *Discrete Math.*, 56, pp. 61–72.
- K.M. Zhang[1987], On Lewin and Vitek's conjecture about the exponent set of primitive matrices, *Linear Alg. Applics.*, 96, pp. 101–108.
- L. Zhu[1982], A short disproof of Euler's conjecture concerning orthogonal latin squares, *Ars Combinatoria*, 14, pp. 47–55.