



ELSEVIER

Linear Algebra and its Applications 325 (2001) 109–139

---

---

LINEAR ALGEBRA  
AND ITS  
APPLICATIONS

---

---

[www.elsevier.com/locate/laa](http://www.elsevier.com/locate/laa)

# The inverse of a tridiagonal matrix

Ranjan K. Mallik\*

*Department of Electrical Engineering, Indian Institute of Technology, Hauz Khas,  
New Delhi 110016, India*

Received 6 March 1998; accepted 1 September 2000

Submitted by R.A. Brualdi

---

## Abstract

In this paper, explicit formulae for the elements of the inverse of a general tridiagonal matrix are presented by first extending results on the explicit solution of a second-order linear homogeneous difference equation with variable coefficients to the nonhomogeneous case, and then applying these extended results to a boundary value problem. A formula for the characteristic polynomial is obtained in the process. We also establish a connection between the matrix inverse and orthogonal polynomials. In addition, the case of a cyclic tridiagonal system is discussed. © 2001 Elsevier Science Inc. All rights reserved.

*AMS classification:* 15A09; 39A10

*Keywords:* Tridiagonal matrix; Inverse; Second-order linear difference equation; Variable coefficients; Explicit solution; Orthogonal polynomials

---

## 1. Introduction

Tridiagonal matrices [1–4] are connected with different areas of science and engineering, including telecommunication system analysis [5] and finite difference methods for solving partial differential equations [4,6,7]. In many of these areas, inversions of tridiagonal matrices are necessary. Efficient algorithms [8], indirect formulae [1,9–12], and direct expressions in some special cases [4,7] for such inversions are known. Bounds on the elements of the inverses of diagonally dominant tridiagonal matrices have also been obtained [13]. However, explicit formulae for the

---

\* Tel.: +91-11-659-1049; fax: +91-11-658-1606.

*E-mail address:* [rkmallik@ee.iitd.ernet.in](mailto:rkmallik@ee.iitd.ernet.in) (R.K. Mallik).

elements of a general tridiagonal matrix inverse, which can give a better analytical treatment to a problem, are not available in the open literature [1].

In this paper, we present explicit formulae for the elements of the inverse of a general tridiagonal matrix. The approach is based on linear difference equations [14,15], and is as follows. Results on the *explicit solution of a second-order linear homogeneous difference equation with variable coefficients* [16] are extended to the nonhomogeneous case. Then these extended results are applied to a boundary value problem to obtain the desired formulae. A formula for the characteristic polynomial is obtained in the process. We also establish a connection between the matrix inverse and orthogonal polynomials. In addition, the case of a cyclic tridiagonal system is discussed.

## 2. Results on the second-order linear homogeneous difference equation with variable coefficients

In this section, we will review some results on the second-order linear homogeneous difference equation with variable coefficients [16]. Let  $\mathbb{N}$  denote the set of natural numbers. A set  $S_q(L, U)$ , where  $q, L, U \in \mathbb{N}$ , has been defined in [16] as the set of all  $q$ -tuples with elements from  $\{L, L+1, \dots, U\}$  arranged in ascending order so that no two consecutive elements are present, that is,

$$S_q(L, U) \triangleq \{L, L+1, \dots, U\} \quad \text{if } U \geq L \text{ and } q = 1, \quad (1a)$$

$$\triangleq \{(k_1, \dots, k_q) : k_1, \dots, k_q \in \{L, L+1, \dots, U\}; k_l - k_{l-1} \geq 2 \text{ for } l = 2, \dots, q\} \\ \text{if } U \geq L+2 \text{ and } 2 \leq q \leq \left\lfloor \frac{U-L+2}{2} \right\rfloor, \quad (1b)$$

$$\triangleq \emptyset \text{ otherwise.} \quad (1c)$$

For example,

$$\begin{aligned} S_1(2, 6) &= \{2, 3, 4, 5, 6\}, \\ S_2(2, 6) &= \{(2, 4), (2, 5), (2, 6), (3, 5), (3, 6), (4, 6)\}, \\ S_3(2, 6) &= \{(2, 4, 6)\}, \\ S_4(2, 6) &= \emptyset. \end{aligned}$$

The following results have been proved in [16]:

- For  $U \geq L$ ,  $1 \leq q \leq \lfloor (U-L+2)/2 \rfloor$ ,

$$|S_q(L, U)| = \binom{U-L-q+2}{q} = \frac{(U-L-q+2)!}{q!(U-L-2q+2)!}. \quad (2)$$

- For  $U \geq L$ ,  $q \geq 2$ ,

$$S_q(L, U+1) = S_q(L, U) \cup \left\{ (k_1, \dots, k_{q-1}, k_q): k_q = U+1; \right. \\ \left. (k_1, \dots, k_{q-1}) \in S_{q-1}(L, U-1) \right\}. \quad (3)$$

We add a proposition which is similar to result (3).

**Proposition 1.** For  $U \geq L$ ,  $q \geq 2$ ,

$$S_q(L-1, U) = S_q(L, U) \cup \left\{ (k_1, k_2, \dots, k_q): k_1 = L-1; \right. \\ \left. (k_2, \dots, k_q) \in S_{q-1}(L+1, U) \right\}. \quad (4)$$

**Proof.** If either  $U = L$ ,  $q \geq 2$  or  $U \geq L+1$ ,  $q > \lfloor (U-L+3)/2 \rfloor$ , then Eq. (4) holds trivially.

For  $U \geq L+1$ ,  $2 \leq q \leq \lfloor (U-L+3)/2 \rfloor$ , we get, from (1a)–(1c),

$$S_q(L-1, U) = \left\{ (k_1, \dots, k_q): k_1, \dots, k_q \in \{L-1, L, \dots, U\}; \right. \\ \left. k_l - k_{l-1} \geq 2 \text{ for } l = 2, \dots, q \right\}. \quad (5)$$

The right-hand side of (5) can be expressed as the union of two disjoint sets, one containing  $q$ -tuples with  $k_1 \neq L-1$  and the other containing  $q$ -tuples with  $k_1 = L-1$ , which we denote as  $S_{q_0}(L-1, U)$  and  $S_{q_1}(L-1, U)$ , respectively. Therefore,

$$S_q(L-1, U) = S_{q_0}(L-1, U) \cup S_{q_1}(L-1, U), \quad (6)$$

where

$$S_{q_0}(L-1, U) = \left\{ (k_1, \dots, k_q): k_1, \dots, k_q \in \{L, L+1, \dots, U\}; \right. \\ \left. k_l - k_{l-1} \geq 2 \text{ for } l = 2, \dots, q \right\}, \\ S_{q_1}(L-1, U) = \left\{ (k_1, \dots, k_q): k_1 = L-1; \right. \\ \left. k_2, \dots, k_q \in \{L, L+1, \dots, U\}; \right. \\ \left. k_l - k_{l-1} \geq 2 \text{ for } l = 2, \dots, q \right\}. \quad (7)$$

It is clear from (1a)–(1c) that

$$S_{q_0}(L-1, U) = S_q(L, U). \quad (8)$$

In the set  $S_{q_1}(L-1, U)$ , since  $k_1 = L-1$ , we have  $k_2 - (L-1) \geq 2$ , which implies that  $k_2, \dots, k_q \in \{L+1, L+2, \dots, U\}$ . Therefore,

$$S_{q_1}(L-1, U) = \left\{ (k_1, k_2, \dots, k_q): k_1 = L-1; \right. \\ \left. (k_2, \dots, k_q) \in S_{q-1}(L+1, U) \right\}. \quad (9)$$

Combining Eqs. (6), (8) and (9), we get (4).  $\square$

Consider the *second-order linear homogeneous difference equation*

$$y_{n+2} = A_n y_{n+1} + B_n y_n, \quad n \geq 1, \quad (10)$$

with integral index  $n$ , variable complex coefficients  $A_n$  and  $B_n$ ,  $B_n \neq 0$ , and complex initial values  $y_1, y_2$ . Define, for  $k \geq 2$ ,

$$\sigma_k \triangleq \frac{B_k}{A_{k-1} A_k}. \quad (11)$$

It has been shown in [16] that the *explicit solution* of difference equation (10), which is an expression for  $y_{n+2}$  in terms of only coefficients  $A_1, \dots, A_n, B_1, \dots, B_n$ , and initial values  $y_1, y_2$ , is given by

$$y_{n+2} = C_n y_2 + D_n y_1, \quad n \geq 0, \quad (12)$$

where

$$C_0 = 1, \quad C_1 = A_1, \quad (13a)$$

$$D_0 = 0, \quad D_1 = B_1, \quad D_2 = B_1 A_2 \quad (13b)$$

and

$$C_n = (A_1 \cdots A_n) \left( 1 + \sum_{q=1}^{\lfloor n/2 \rfloor} \sum_{(k_1, \dots, k_q) \in S_q(2, n)} (\sigma_{k_1} \cdots \sigma_{k_q}) \right) \quad (14a)$$

for  $n \geq 2$ ,

$$D_n = B_1 (A_2 \cdots A_n) \left( 1 + \sum_{q=1}^{\lfloor (n-1)/2 \rfloor} \sum_{(k_1, \dots, k_q) \in S_q(3, n)} (\sigma_{k_1} \cdots \sigma_{k_q}) \right) \quad (14b)$$

for  $n \geq 3$ ,

where  $S_q(L, U)$  is defined by (1a)–(1c), and  $\sigma_k$  is defined by (11).

### 3. Solution of the second-order linear nonhomogeneous difference equation with variable coefficients

We now focus on the *second-order linear nonhomogeneous difference equation*

$$y_{n+2} = A_n y_{n+1} + B_n y_n + x_{n+2}, \quad n \geq 1, \quad (15)$$

with integral index  $n$ , variable complex coefficients  $A_n$  and  $B_n$ ,  $B_n \neq 0$ , complex forcing term  $x_{n+2}$ , and complex initial values  $y_1, y_2$ .

Let  $\sigma_k = B_k/(A_{k-1}A_k)$  for  $k \geq 2$  as in (11). For  $n \geq 0, i \geq 1$ , define a quantity  $E_n(i)$ , which is a function of the  $A_k$ s and the  $\sigma_k$ s, as

$$E_n(i) \triangleq (A_i \cdots A_n) \left( 1 + \sum_{q=1}^{\lfloor (n-i+1)/2 \rfloor} \sum_{(k_1, \dots, k_q) \in S_q(i+1, n)} (\sigma_{k_1} \cdots \sigma_{k_q}) \right) \\ \text{if } i = 1, \dots, n-1, n \geq 2, \\ \triangleq A_n \quad \text{if } i = n, n \geq 1, \\ \triangleq 1 \quad \text{if } i = n+1, n \geq 0, \\ \triangleq 0 \quad \text{otherwise.} \quad (16)$$

Therefore,  $E_n(i)$  is nontrivial only for  $n \geq i-1, i \geq 1$ . Note that

$$E_n(1) = C_n \quad \text{for } n \geq 0, \quad (17)$$

where  $C_n$  is given by (13a) and (14a).

Two recurrences for  $E_n(i)$  are established by the following two propositions.

**Proposition 2.** For  $n \geq i-1, i \geq 1$ ,

$$E_{n+2}(i) = A_{n+2}E_{n+1}(i) + B_{n+2}E_n(i). \quad (18)$$

**Proof.** Using the definition of  $E_n(i)$  in (16) and  $\sigma_i$  in (11), the right-hand side of (18) can be written for  $n = i-1, i \geq 1$  as

$$A_{i+1}E_i(i) + B_{i+1}E_{i-1}(i) = A_{i+1}A_i + B_{i+1} \\ = (A_iA_{i+1})(1 + \sigma_{i+1}) \\ = E_{i+1}(i) \quad (19)$$

and for  $n = i, i \geq 1$  as

$$A_{i+2}E_{i+1}(i) + B_{i+2}E_i(i) = (A_{i+2}A_{i+1}A_i)(1 + \sigma_{i+1}) + B_{i+2}A_i \\ = (A_iA_{i+1}A_{i+2})(1 + \sigma_{i+1} + \sigma_{i+2}) \\ = E_{i+2}(i), \quad (20)$$

which imply that (18) holds for  $n = i-1, i, i \geq 1$ .

Now consider the case when  $n \geq i+1, i \geq 1$ . Again, using (16) and noting the fact that  $B_{n+2} = A_{n+1}A_{n+2}\sigma_{n+2}$ , we get

$$A_{n+2}E_{n+1}(i) + B_{n+2}E_n(i) \\ = (A_i \cdots A_{n+2}) \left( 1 + \sum_{q=1}^{\lfloor (n-i+2)/2 \rfloor} \sum_{(k_1, \dots, k_q) \in S_q(i+1, n+1)} (\sigma_{k_1} \cdots \sigma_{k_q}) \right)$$

$$\begin{aligned}
& + \sigma_{n+2} + \sum_{q=2}^{\lfloor (n-i+3)/2 \rfloor} \sum_{(k_1, \dots, k_{q-1}) \in S_{q-1}(i+1, n)} (\sigma_{k_1} \cdots \sigma_{k_{q-1}}) \sigma_{n+2} \Big) \\
& = (A_i \cdots A_{n+2}) \left( 1 + \sum_{k_1=i+1}^{n+2} \sigma_{k_1} \right. \\
& \quad + \sum_{q=2}^{\lfloor (n-i+2)/2 \rfloor} \sum_{(k_1, \dots, k_q) \in S_q(i+1, n+1)} (\sigma_{k_1} \cdots \sigma_{k_q}) \\
& \quad \left. + \sum_{q=2}^{\lfloor (n-i+3)/2 \rfloor} \sum_{(k_1, \dots, k_{q-1}) \in S_{q-1}(i+1, n)} (\sigma_{k_1} \cdots \sigma_{k_{q-1}}) \sigma_{n+2} \right). \quad (21)
\end{aligned}$$

Now

$$\left\lfloor \frac{n-i+3}{2} \right\rfloor = \begin{cases} \left\lfloor \frac{n-i+2}{2} \right\rfloor & \text{if } (n-i+2) \text{ is even,} \\ \left\lfloor \frac{n-i+2}{2} \right\rfloor + 1 & \text{if } (n-i+2) \text{ is odd.} \end{cases} \quad (22)$$

If  $(n-i+2)$  is odd, by (1a)–(1c),

$$S_{\lfloor (n-i+3)/2 \rfloor}(i+1, n+1) = S_{\lfloor (n-i+2)/2 \rfloor+1}(i+1, n+1) = \emptyset.$$

Then, using (3), we get

$$\begin{aligned}
S_q(i+1, n+2) &= S_q(i+1, n+1) \cup \{(k_1, \dots, k_q): k_q = n+2; \\
&\quad (k_1, \dots, k_{q-1}) \in S_{q-1}(i+1, n)\}, \quad (23)
\end{aligned}$$

for  $q = 2, \dots, \lfloor (n-i+3)/2 \rfloor$ . Putting together the summation terms of (21) using (22) and (23), we obtain

$$\begin{aligned}
& A_{n+2}E_{n+1}(i) + B_{n+2}E_n(i) \\
& = (A_i \cdots A_{n+2}) \left( 1 + \sum_{q=1}^{\lfloor (n-i+3)/2 \rfloor} \sum_{(k_1, \dots, k_q) \in S_q(i+1, n+2)} (\sigma_{k_1} \cdots \sigma_{k_q}) \right) \\
& = E_{n+2}(i) \quad (24)
\end{aligned}$$

for  $n \geq i+1, i \geq 1$ . A combination of (19), (20) and (24) yields (18).  $\square$

**Proposition 3.** For  $1 \leq i \leq n-1, n \geq 2$ ,

$$E_n(i) = A_i E_n(i+1) + B_{i+1} E_n(i+2). \quad (25)$$

**Proof.** From the definition of  $E_n(i)$  in (16), the right-hand side of (25) can be written for  $i = n - 1$  as

$$\begin{aligned} A_{n-1}E_n(n) + B_nE_n(n+1) &= A_{n-1}A_n + B_n \\ &= (A_{n-1}A_n)(1 + \sigma_n) \\ &= E_n(n-1) \end{aligned} \quad (26)$$

and for  $i = n - 2$  as

$$\begin{aligned} A_{n-2}E_n(n-1) + B_{n-1}E_n(n) &= (A_{n-2}A_{n-1}A_n)(1 + \sigma_n) + B_{n-1}A_n \\ &= (A_{n-2}A_{n-1}A_n)(1 + \sigma_n + \sigma_{n-1}) \\ &= E_n(n-2), \end{aligned} \quad (27)$$

which imply that (25) holds for  $i = n - 2, n - 1$ .

Now consider the case when  $1 \leq i \leq n - 3$ . Using (16) and (11) we obtain

$$\begin{aligned} &A_iE_n(i+1) + B_{i+1}E_n(i+2) \\ &= (A_i \cdots A_n) \left( 1 + \sum_{q=1}^{\lfloor (n-i)/2 \rfloor} \sum_{(k_1, \dots, k_q) \in S_q(i+2, n)} (\sigma_{k_1} \cdots \sigma_{k_q}) + \sigma_{i+1} \right. \\ &\quad \left. + \sum_{q=2}^{\lfloor (n-i+1)/2 \rfloor} \sum_{(k_2, \dots, k_q) \in S_{q-1}(i+3, n)} \sigma_{i+1}(\sigma_{k_2} \cdots \sigma_{k_q}) \right) \\ &= (A_i \cdots A_n) \left( 1 + \sum_{k_1=i+1}^n \sigma_{k_1} + \sum_{q=2}^{\lfloor (n-i)/2 \rfloor} \sum_{(k_1, \dots, k_q) \in S_q(i+2, n)} (\sigma_{k_1} \cdots \sigma_{k_q}) \right. \\ &\quad \left. + \sum_{q=2}^{\lfloor (n-i+1)/2 \rfloor} \sum_{(k_2, \dots, k_q) \in S_{q-1}(i+3, n)} \sigma_{i+1}(\sigma_{k_2} \cdots \sigma_{k_q}) \right). \end{aligned} \quad (28)$$

Now

$$\left\lfloor \frac{n-i+1}{2} \right\rfloor = \begin{cases} \left\lfloor \frac{n-i}{2} \right\rfloor & \text{if } (n-i) \text{ is even,} \\ \left\lfloor \frac{n-i}{2} \right\rfloor + 1 & \text{if } (n-i) \text{ is odd.} \end{cases} \quad (29)$$

If  $(n-i)$  is odd, then by (1a)–(1c),

$$S_{\lfloor (n-i+1)/2 \rfloor}(i+2, n) = S_{\lfloor (n-i)/2 \rfloor+1}(i+2, n) = \emptyset.$$

Then, using Proposition 1, we get

$$\begin{aligned} S_q(i+1, n) &= S_q(i+2, n) \cup \{(k_1, \dots, k_q): k_1 = i+1; \\ &\quad (k_2, \dots, k_q) \in S_{q-1}(i+3, n)\}, \end{aligned} \quad (30)$$

for  $q = 2, \dots, \lfloor (n-i+1)/2 \rfloor$ . Putting together the summation terms of (28) using (29) and (30), we obtain (25).  $\square$

The following proposition provides the *explicit solution* of the difference equation (15), which consists of an expression for  $y_{n+2}$  in terms of only coefficients  $A_1, \dots, A_n, B_1, \dots, B_n$ , initial values  $y_1, y_2$ , and forcing terms  $x_3, \dots, x_{n+2}$ .

**Proposition 4.** *The solution of difference equation (15) with initial values  $y_1, y_2$  is given by*

$$y_{n+2} = C_n y_2 + D_n y_1 + \sum_{k=3}^{n+2} E_n(k-1)x_k, \quad n \geq 1, \quad (31)$$

where  $E_n(k)$  is defined by (16), and  $C_n$  and  $D_n$  by (13a), (13b), (14a), and (14b).

**Proof.** The solution of (15) with initial values  $y_1, y_2$  can be expressed as

$$y_{n+2} = H_n + P_n, \quad n \geq 1, \quad (32)$$

where  $H_n$  is the *solution of the homogeneous equation*, and  $P_n$  the *particular solution*. From the results of Section 2 (Eqs. (10) and (12)), we have

$$H_n = C_n y_2 + D_n y_1, \quad n \geq 1. \quad (33)$$

It is also known that the particular solution  $P_n$  satisfies the recurrence

$$P_{n+2} = A_{n+2} P_{n+1} + B_{n+2} P_n + x_{n+4}, \quad n \geq 1. \quad (34)$$

Using (15), (13a), (13b), (14a) and (14b), we get

$$\begin{aligned} y_3 &= C_1 y_2 + D_1 y_1 + x_3, \\ y_4 &= C_2 y_2 + D_2 y_1 + (x_4 + A_2 x_3), \end{aligned} \quad (35)$$

where

$$\begin{aligned} C_1 &= A_1, & C_2 &= A_1 A_2 + B_2, \\ D_1 &= B_1, & D_2 &= B_1 A_2, \end{aligned} \quad (36)$$

which in turn implies that

$$P_1 = x_3, \quad P_2 = x_4 + A_2 x_3. \quad (37)$$

We now proceed to prove the validity of the expression

$$P_n = \sum_{k=3}^{n+2} E_n(k-1)x_k \quad (38)$$

for  $n \geq 1$ .

It is clear from (37) and the definition of  $E_n(k)$  in (16) that expression (38) holds for  $n = 1, 2$ . Let (38) hold for  $P_n$  and  $P_{n+1}$  for  $n \geq 1$ . Then, from (34), Proposition 2 and (16), we get



$$\begin{aligned}
P_{n+2} &= A_{n+2} \sum_{k=3}^{n+3} E_{n+1}(k-1)x_k + B_{n+2} \sum_{k=3}^{n+2} E_n(k-1)x_k + x_{n+4} \\
&= \sum_{k=3}^{n+2} [A_{n+2}E_{n+1}(k-1) + B_{n+2}E_n(k-1)]x_k + A_{n+2}x_{n+3} + x_{n+4} \\
&= \sum_{k=3}^{n+2} E_{n+2}(k-1)x_k + E_{n+2}(n+2)x_{n+3} + E_{n+2}(n+3)x_{n+4}. \quad (39)
\end{aligned}$$

Thus,

$$P_{n+2} = \sum_{k=3}^{n+4} E_{n+2}(k-1)x_k, \quad (40)$$

which agrees with (38). By mathematical induction, expression (38) is valid for all  $n \geq 1$ .

From (32), (33) and (38), we obtain (31).  $\square$

### 3.1. An example

Consider the case when we need to find  $y_6$  in the difference equation (15) in terms of coefficients  $A_1, A_2, A_3, A_4, B_1, B_2, B_3, B_4$ , initial values  $y_1, y_2$ , and forcing terms  $x_3, x_4, x_5, x_6$ .

From Proposition 4, we have

$$y_6 = C_4 y_2 + D_4 y_1 + \sum_{k=3}^6 E_4(k-1)x_k.$$

Eqs. (14a) and (14b) imply that

$$\begin{aligned}
C_4 &= (A_1 A_2 A_3 A_4) \left( 1 + \sum_{k_1 \in S_1(2,4)} \sigma_{k_1} + \sum_{(k_1, k_2) \in S_2(2,4)} (\sigma_{k_1} \sigma_{k_2}) \right), \\
D_4 &= (B_1 A_2 A_3 A_4) \left( 1 + \sum_{k_1 \in S_1(3,4)} \sigma_{k_1} \right).
\end{aligned}$$

Now, from the definition of  $S_q(L, U)$  in (1a)–(1c), we get

$$S_1(2, 4) = \{2, 3, 4\},$$

$$S_2(2, 4) = \{(2, 4)\},$$

$$S_1(3, 4) = \{3, 4\}.$$

Therefore,

$$C_4 = (A_1 A_2 A_3 A_4)(1 + \sigma_2 + \sigma_3 + \sigma_4 + \sigma_2 \sigma_4),$$

$$D_4 = (B_1 A_2 A_3 A_4)(1 + \sigma_3 + \sigma_4),$$

where, from (11),

$$\sigma_2 = \frac{B_2}{A_1 A_2}, \quad \sigma_3 = \frac{B_3}{A_2 A_3}, \quad \sigma_4 = \frac{B_4}{A_3 A_4}.$$

In addition, from (16), we obtain

$$E_4(2) = (A_2 A_3 A_4)(1 + \sigma_3 + \sigma_4),$$

$$E_4(3) = (A_3 A_4)(1 + \sigma_4),$$

$$E_4(4) = A_4,$$

$$E_4(5) = 1.$$

Therefore, by substituting the expressions for  $\sigma_2$ ,  $\sigma_3$  and  $\sigma_4$  in  $C_4$ ,  $D_4$ ,  $E_4(2)$ ,  $E_4(3)$ ,  $E_4(4)$ ,  $E_4(5)$ , we finally have the desired solution, which is

$$\begin{aligned} y_6 = & (A_1 A_2 A_3 A_4 + B_2 A_3 A_4 + B_3 A_1 A_4 + B_4 A_1 A_2 + B_2 B_4) y_2 \\ & + (B_1 A_2 A_3 A_4 + B_1 B_3 A_4 + B_1 B_4 A_2) y_1 \\ & + (A_2 A_3 A_4 + B_3 A_4 + B_4 A_2) x_3 \\ & + (A_3 A_4 + B_4) x_4 + A_4 x_5 + x_6. \end{aligned}$$

#### 4. A boundary value problem and the tridiagonal matrix inverse

Consider the difference equation (15) over the finite index interval  $[1, K + 2]$ ,  $K \geq 2$ , with zero boundary values, that is, the recurrence

$$y_{n+2} = A_n y_{n+1} + B_n y_n + x_{n+2}, \quad 1 \leq n \leq K, \quad (41)$$

with boundary values  $y_1, y_{K+2}$  such that

$$y_1 = y_{K+2} = 0. \quad (42)$$

The  $K$  equations in (41) can be written in matrix form as

$$\begin{bmatrix} -B_1 & -A_1 & 1 & & & & \\ & -B_2 & -A_2 & 1 & & & \\ & & \ddots & \ddots & \ddots & & \\ & 0 & & -B_{K-1} & -A_{K-1} & 1 & \\ & & & -B_K & -A_K & 1 & \\ & & & & & & 1 \end{bmatrix} \begin{bmatrix} y_1 \\ y_2 \\ \vdots \\ y_{K+1} \\ y_{K+2} \end{bmatrix}$$

$$= \begin{bmatrix} x_3 \\ \vdots \\ x_{K+2} \end{bmatrix}. \quad (43)$$

Using boundary conditions (42), the solution of the difference equation (41) can be expressed as

$$\begin{bmatrix} y_2 \\ \vdots \\ y_{K+1} \end{bmatrix} = \begin{bmatrix} -A_1 & 1 & & & \\ -B_2 & -A_2 & 1 & & \\ & \ddots & \ddots & \ddots & \\ \mathbf{0} & & -B_{K-1} & -A_{K-1} & 1 \\ & & & -B_K & -A_K \end{bmatrix}^{-1} \begin{bmatrix} x_3 \\ \vdots \\ x_{K+2} \end{bmatrix}. \quad (44)$$

However, from Proposition 4, the quantities  $y_3, \dots, y_{K+2}$  of recurrence (41) can be expressed in terms of  $y_1, y_2$  as

$$y_{i+1} = C_{i-1}y_2 + D_{i-1}y_1 + \sum_{j=1}^{i-1} E_{i-1}(j+1)x_{j+2}, \quad 2 \leq i \leq K+1. \quad (45)$$

As a result of boundary conditions (42), substitution of  $i = K+1$  in (45) gives

$$y_{K+2} = C_K y_2 + \sum_{j=1}^K E_K(j+1)x_{j+2} = 0, \quad (46)$$

and therefore, when  $C_K \neq 0$ ,

$$y_2 = \sum_{j=1}^K \left[ -\frac{E_K(j+1)}{C_K} \right] x_{j+2}. \quad (47)$$

Substituting (47) in (45) and setting  $y_1 = 0$  we obtain

$$\begin{aligned} y_{i+1} = & - \sum_{j=1}^K \frac{C_{i-1}E_K(j+1)}{C_K} x_{j+2} \\ & + \sum_{j=1}^{i-1} E_{i-1}(j+1)x_{j+2}, \quad 2 \leq i \leq K, \quad C_K \neq 0. \end{aligned} \quad (48)$$

In other words,

$$\begin{aligned} y_{i+1} = & \sum_{j=1}^{i-1} \left[ E_{i-1}(j+1) - \frac{C_{i-1}E_K(j+1)}{C_K} \right] x_{j+2} \\ & + \sum_{j=i}^K \left[ -\frac{C_{i-1}E_K(j+1)}{C_K} \right] x_{j+2}, \quad 2 \leq i \leq K, \quad C_K \neq 0. \end{aligned} \quad (49)$$

Using (16) and (17), Eqs. (47) and (49) can be written together as

$$y_{i+1} = \sum_{j=1}^K \left[ E_{i-1}(j+1) - \frac{E_{i-1}(1)E_K(j+1)}{E_K(1)} \right] x_{j+2},$$

$$1 \leq i \leq K, \quad E_K(1) \neq 0. \quad (50)$$

Thus the *solution of boundary value problem* (41), (42) is given by (47) and (49) or alternatively by (50). This solution also gives the inverse of a tridiagonal matrix.

Denoting the  $K \times K$  tridiagonal matrix derived from the  $K \times (K+2)$  matrix on the left-hand side of (43) (by removing the first and  $(K+2)$ th columns) as

$$\mathbf{T}_K = \begin{bmatrix} -A_1 & 1 & & & \\ -B_2 & -A_2 & 1 & & \\ & \ddots & \ddots & \ddots & \\ \mathbf{0} & & -B_{K-1} & -A_{K-1} & 1 \\ & & & -B_K & -A_K \end{bmatrix}, \quad K \geq 2, \quad (51)$$

and its inverse as

$$\mathbf{R}_K = \mathbf{T}_K^{-1} = [R_{i,j}]_{i,j=1}^K, \quad (52)$$

we obtain, from (44), (47) and (49), an expression for  $R_{i,j}$ , the element in the  $i$ th row and  $j$ th column of  $\mathbf{T}_K^{-1}$ . It is given by

$$R_{i,j} = E_{i-1}(j+1) - \frac{C_{i-1}E_K(j+1)}{C_K}$$

$$\text{for } j = 1, \dots, i-1, \quad i = 2, \dots, K,$$

$$= -\frac{C_{i-1}E_K(j+1)}{C_K}$$

$$\text{for } j = i, \dots, K, \quad i = 1, \dots, K, \quad C_K \neq 0, \quad (53)$$

where  $E_i(j)$  is defined in (16), and  $C_i$  in (13a) and (14a). Alternatively, from (44) and (50), we get

$$R_{i,j} = E_{i-1}(j+1) - \frac{E_{i-1}(1)E_K(j+1)}{E_K(1)}$$

$$\text{for } i, j = 1, \dots, K, \quad E_K(1) \neq 0. \quad (54)$$

The following proposition gives an expression for the determinant of the tridiagonal matrix  $\mathbf{T}_K$ .

**Proposition 5.** For  $K \geq 2$ ,

$$\det(\mathbf{T}_K) = (-1)^K C_K, \quad (55)$$

where  $\mathbf{T}_K$  is given by (51) and  $C_K$  by (14a).

**Proof.** From (51), it is clear that if we find  $\det(\mathbf{T}_K)$  by using the  $K$ th row of  $\mathbf{T}_K$  followed by its  $K$ th column, we obtain

$$\det(\mathbf{T}_K) = -A_K \det(\mathbf{T}_{K-1}) + B_K \det(\mathbf{T}_{K-2}), \quad K \geq 4. \quad (56)$$

It is simple to show that

$$\det(\mathbf{T}_2) = \det \begin{pmatrix} -A_1 & 1 \\ -B_2 & -A_2 \end{pmatrix} = A_1 A_2 (1 + \sigma_2) = C_2, \quad (57a)$$

$$\begin{aligned} \det(\mathbf{T}_3) &= \det \begin{pmatrix} -A_1 & 1 & 0 \\ -B_2 & -A_2 & 1 \\ 0 & -B_3 & -A_3 \end{pmatrix} \\ &= -A_1 A_2 A_3 (1 + \sigma_2 + \sigma_3) \\ &= -C_3. \end{aligned} \quad (57b)$$

Since  $C_K = E_K(1)$ , it is governed by the second-order recursion

$$C_K = A_K C_{K-1} + B_K C_{K-2}, \quad K \geq 4 \quad (58)$$

by Proposition 2. Therefore,

$$(-1)^K C_K = -A_K (-1)^{K-1} C_{K-1} + B_K (-1)^{K-2} C_{K-2}, \quad K \geq 4. \quad (59)$$

Recursions (56) and (59) for  $\det(\mathbf{T}_K)$  and  $(-1)^K C_K$ , respectively, are identical, and from (57a) and (57b) we find that they have the same initial values  $(-1)^2 C_2$  and  $(-1)^3 C_3$ . Hence,  $\det(\mathbf{T}_K) = (-1)^K C_K$  for  $K \geq 2$ .  $\square$

If it so happens that  $\det(\mathbf{T}_K) = 0$ , then Proposition 5 implies that  $C_K = 0$ . In that case, the matrix  $\mathbf{T}_K$  is not invertible, and Eq. (53) for the elements of  $\mathbf{T}_K^{-1}$  does not hold.

Let  $\Delta_{j,i}$  denote the *minor* corresponding to the  $j$ th row and  $i$ th column of  $\mathbf{T}_K$ . Then, from (53) and Proposition 5, we have

$$\begin{aligned} \Delta_{j,i} &= (-1)^{(K-i-j)} [C_K E_{i-1}(j+1) - C_{i-1} E_K(j+1)] \\ &\quad \text{for } j = 1, \dots, i-1, \quad i = 2, \dots, K, \\ &= -(-1)^{(K-i-j)} C_{i-1} E_K(j+1) \\ &\quad \text{for } j = i, \dots, K, \quad i = 1, \dots, K, \end{aligned} \quad (60)$$

or, alternatively, from (54) and Proposition 5,

$$\begin{aligned} \Delta_{j,i} &= (-1)^{(K-i-j)} [E_K(1) E_{i-1}(j+1) - E_{i-1}(1) E_K(j+1)] \\ &\quad \text{for } i, j = 1, \dots, K. \end{aligned} \quad (61)$$

#### 4.1. The tridiagonal matrix inverse

We now consider a nonsingular  $K \times K$  tridiagonal matrix  $\Phi_K$  ( $K \geq 2$ ), with complex entries, given by

$$\Phi_K = \begin{bmatrix} \alpha_1 & \beta_1 & & & \\ \gamma_2 & \alpha_2 & \beta_2 & & \\ & \ddots & \ddots & \ddots & \\ \mathbf{0} & & \gamma_{K-1} & \alpha_{K-1} & \beta_{K-1} \\ & & & \gamma_K & \alpha_K \end{bmatrix},$$

$$\beta_1, \dots, \beta_{K-1} \neq 0, \gamma_2, \dots, \gamma_K \neq 0. \quad (62)$$

Without loss of generality, let  $\beta_K$  be defined as

$$\beta_K \triangleq 1, \quad (63a)$$

and

$$A_i \triangleq -\frac{\alpha_i}{\beta_i}, \quad 1 \leq i \leq K,$$

$$B_i \triangleq -\frac{\gamma_i}{\beta_i}, \quad 2 \leq i \leq K, \quad (63b)$$

such that  $A_1, \dots, A_K, B_2, \dots, B_K$  are entries of  $\mathbf{T}_K$  in (51). We can express  $\Phi_K$  as

$$\Phi_K = \text{diag}(\beta_1, \dots, \beta_K) \mathbf{T}_K,$$

and therefore, the inverse  $\Psi_K$  of  $\Phi_K$  can be expressed as

$$\Psi_K = \Phi_K^{-1} = [\psi_{i,j}]_{i,j=1}^K = \mathbf{T}_K^{-1} \text{diag}\left(\frac{1}{\beta_1}, \dots, \frac{1}{\beta_K}\right), \quad (64)$$

where, from (52) and (54),

$$\psi_{i,j} = \frac{R_{i,j}}{\beta_j}$$

$$= \frac{1}{\beta_j} \left[ E_{i-1}(j+1) - \frac{E_{i-1}(1)E_K(j+1)}{E_K(1)} \right], \quad E_K(1) \neq 0, \quad (65a)$$

such that (using (16) and (63b))

$$E_n(m) = (-1)^{n-m+1} \left( \frac{\alpha_m}{\beta_m} \dots \frac{\alpha_n}{\beta_n} \right)$$

$$\times \left( 1 + \sum_{q=1}^{\lfloor (n-m+1)/2 \rfloor} \sum_{(k_1, \dots, k_q) \in S_q(m+1, n)} (\sigma_{k_1} \dots \sigma_{k_q}) \right)$$

if  $m = 1, \dots, n-1, n = 2, \dots, K,$

$$\begin{aligned}
&= -\frac{\alpha_n}{\beta_n} \quad \text{if } m = n, \quad n = 1, \dots, K, \\
&= 1 \quad \text{if } m = n + 1, \quad n = 0, \dots, K, \\
&= 0 \quad \text{otherwise}
\end{aligned} \tag{65b}$$

and (using (11) and (63b))

$$\sigma_k = -\frac{\beta_{k-1}\gamma_k}{\alpha_{k-1}\alpha_k}. \tag{65c}$$

The set  $S_q(m+1, n)$  in (65b) is defined by (1a)–(1c).

It is known that for the matrix  $\Psi_K$  in (64), there exist four sequences  $\{u_i\}_{i=1}^K$ ,  $\{v_i\}_{i=1}^K$ ,  $\{r_i\}_{i=1}^K$ ,  $\{s_i\}_{i=1}^K$  [9,10,17] such that

$$\psi_{i,j} = \begin{cases} u_i v_j & \text{if } i > j, \\ r_i s_j & \text{if } i \leq j. \end{cases} \tag{66}$$

We can express  $\Psi_K$  as

$$\Psi_K = \left( (\Phi_K^T)^{-1} \right)^T, \tag{67}$$

where  $\Phi_K^T$  can be obtained from  $\Phi_K$  by interchanging the positions of  $\beta_j$  and  $\gamma_{j+1}$  in (62) for  $j = 1, \dots, K-1$ . Defining, without loss of generality,  $\gamma_{K+1}$  as

$$\gamma_{K+1} \triangleq 1, \tag{68}$$

and interchanging  $\beta_j$  and  $\gamma_{j+1}$  for  $j = 1, \dots, K$  in formula (65b) for  $E_n(m)$ , we obtain a quantity  $F_n(m)$  expressed as

$$\begin{aligned}
F_n(m) &= (-1)^{n-m+1} \left( \frac{\alpha_m}{\gamma_{m+1}} \dots \frac{\alpha_n}{\gamma_{n+1}} \right) \\
&\quad \times \left( 1 + \sum_{q=1}^{\lfloor (n-m+1)/2 \rfloor} \sum_{(k_1, \dots, k_q) \in S_q(m+1, n)} (\sigma_{k_1} \dots \sigma_{k_q}) \right) \\
&\quad \text{if } m = 1, \dots, n-1, \quad n = 2, \dots, K, \\
&= -\frac{\alpha_n}{\gamma_{n+1}} \quad \text{if } m = n, \quad n = 1, \dots, K, \\
&= 1 \quad \text{if } m = n+1, \quad n = 0, \dots, K, \\
&= 0 \quad \text{otherwise,}
\end{aligned} \tag{69}$$

where  $\sigma_k$ , which is invariant under the interchange of  $\beta_{k-1}$  and  $\gamma_k$ , is given by (65c). The elements of  $\Psi_K$  can alternatively be written in terms of  $F_n(m)$  as

$$\psi_{i,j} = \frac{1}{\gamma_{i+1}} \left[ F_{j-1}(i+1) - \frac{F_{j-1}(1)F_K(i+1)}{F_K(1)} \right], \quad F_K(1) \neq 0. \tag{70}$$

Observe from (65a) and (69) that, for  $m \leq n$ ,

$$F_n(m) = E_n(m) \frac{(\beta_m \cdots \beta_n)}{(\gamma_{m+1} \cdots \gamma_{n+1})}. \quad (71)$$

Therefore, combining (65a), (70) and (71), we obtain

$$\psi_{i,j} = \begin{cases} -\frac{F_{j-1}(1)F_K(i+1)}{\gamma_{i+1}F_K(1)} \\ = -\frac{E_{j-1}(1)E_K(i+1)}{\beta_i E_K(1)} \frac{(\gamma_{j+1} \cdots \gamma_i)}{(\beta_j \cdots \beta_{i-1})} & \text{if } i > j, \\ & F_K(1) \neq 0, \\ -\frac{E_{i-1}(1)E_K(j+1)}{\beta_j E_K(1)} & \text{if } i \leq j, E_K(1) \neq 0. \end{cases} \quad (72)$$

Thus, (72), along with (65b) and (69), gives an *explicit formula for the element in the  $i$ th row and  $j$ th column of the inverse of the tridiagonal matrix  $\Phi_K$  in (62).*

We see that (72) is consistent with the structure (66) of  $\Psi_K$ . Without loss of generality, substituting

$$\begin{aligned} r_i &= E_{i-1}(1), \\ s_j &= -\frac{E_K(j+1)}{\beta_j E_K(1)}, \\ u_i &= -\frac{F_K(i+1)}{\gamma_{i+1}F_K(1)} = s_i \frac{(\gamma_2 \cdots \gamma_i)}{(\beta_1 \cdots \beta_{i-1})}, \\ v_j &= F_{j-1}(1) = r_j \frac{(\beta_1 \cdots \beta_{j-1})}{(\gamma_2 \cdots \gamma_j)}, \end{aligned} \quad (73)$$

we obtain from Proposition 2 the recurrences

$$r_i = -\frac{1}{\beta_{i-1}}(\alpha_{i-1}r_{i-1} + \gamma_{i-1}r_{i-2}), \quad (74a)$$

$$v_j = -\frac{1}{\gamma_j}(\alpha_{j-1}v_{j-1} + \beta_{j-2}v_{j-2}) \quad (74b)$$

for sequences  $\{r_i\}$  and  $\{v_j\}$ , respectively, and from Proposition 3 the recurrences

$$s_j = -\frac{1}{\beta_j}(\alpha_{j+1}s_{j+1} + \gamma_{j+2}s_{j+2}), \quad (75a)$$

$$u_i = -\frac{1}{\gamma_{i+1}}(\alpha_{i+1}u_{i+1} + \beta_{i+2}u_{i+2}) \quad (75b)$$

for sequences  $\{s_j\}$  and  $\{u_i\}$ , respectively. Recurrences (74a), (74b), (75a) and (75b) agree with those in Theorem 2 of [9]. Note that (66) combined with (73) can also be used to obtain the tridiagonal matrix inverse.



The quantities  $\beta_K$  and  $\gamma_{K+1}$  which have been defined to be 1 cancel out due to multiplication in the expression for  $\psi_{i,j}$  in (72) as well as in the expressions for  $u_i, s_j$  in (73). Hence, we can also consider  $\beta_K$  and  $\gamma_{K+1}$  to be arbitrary nonzero complex quantities in the definition of  $E_n(m)$  in (65b) and  $F_n(m)$  in (69).

#### 4.2. An example

Consider the  $3 \times 3$  tridiagonal matrix

$$\Phi_3 = \begin{bmatrix} \alpha_1 & \beta_1 & 0 \\ \gamma_2 & \alpha_2 & \beta_2 \\ 0 & \gamma_3 & \alpha_3 \end{bmatrix}, \quad \beta_1, \beta_2 \neq 0, \gamma_2, \gamma_3 \neq 0,$$

which is the matrix in (62) with  $K = 3$ . To obtain the inverse of this matrix, we first obtain the sequences  $\{u_i\}_{i=1}^3, \{v_i\}_{i=1}^3, \{r_i\}_{i=1}^3, \{s_i\}_{i=1}^3$  using (73), and then substitute their expressions in (66).

We need the expressions for  $\sigma_2$  and  $\sigma_3$  which, from (65c), are given by

$$\sigma_2 = -\frac{\beta_1\gamma_2}{\alpha_1\alpha_2}, \quad \sigma_3 = -\frac{\beta_2\gamma_3}{\alpha_2\alpha_3}.$$

Now, from (73) and (65b), we have

$$r_1 = E_0(1) = 1,$$

$$r_2 = E_1(1) = -\frac{\alpha_1}{\beta_1},$$

$$r_3 = E_2(1) = \frac{\alpha_1\alpha_2}{\beta_1\beta_2}(1 + \sigma_2) = \frac{1}{\beta_1\beta_2}(\alpha_1\alpha_2 - \beta_1\gamma_2),$$

and

$$\begin{aligned} s_1 &= -\frac{E_3(2)}{\beta_1 E_3(1)} \\ &= -\frac{\frac{\alpha_2\alpha_3}{\beta_2\beta_3}(1 + \sigma_3)}{\beta_1 \left( -\frac{\alpha_1\alpha_2\alpha_3}{\beta_1\beta_2\beta_3} \right) (1 + \sigma_2 + \sigma_3)} \\ &= \frac{(\alpha_2\alpha_3 - \beta_2\gamma_3)}{(\alpha_1\alpha_2\alpha_3 - \beta_1\gamma_2\alpha_3 - \beta_2\gamma_3\alpha_1)}, \\ s_2 &= -\frac{E_3(3)}{\beta_2 E_3(1)} \end{aligned}$$

$$\begin{aligned}
&= -\frac{\left(-\frac{\alpha_3}{\beta_3}\right)}{\beta_2 \left(-\frac{\alpha_1 \alpha_2 \alpha_3}{\beta_1 \beta_2 \beta_3}\right) (1 + \sigma_2 + \sigma_3)} \\
&= -\frac{\beta_1 \alpha_3}{(\alpha_1 \alpha_2 \alpha_3 - \beta_1 \gamma_2 \alpha_3 - \beta_2 \gamma_3 \alpha_1)}, \\
s_3 &= -\frac{E_3(4)}{\beta_3 E_3(1)} \\
&= -\frac{1}{\beta_3 \left(-\frac{\alpha_1 \alpha_2 \alpha_3}{\beta_1 \beta_2 \beta_3}\right) (1 + \sigma_2 + \sigma_3)} \\
&= \frac{\beta_1 \beta_2}{(\alpha_1 \alpha_2 \alpha_3 - \beta_1 \gamma_2 \alpha_3 - \beta_2 \gamma_3 \alpha_1)}.
\end{aligned}$$

By interchanging the positions of  $\beta_{k-1}$  and  $\gamma_k$ ,  $k = 2, 3, 4$  in  $r_i$ , we obtain  $v_i$ , and by doing the same in  $s_i$ , we obtain  $u_i$ . Thus, from (73),

$$v_1 = F_0(1) = 1,$$

$$v_2 = F_1(1) = -\frac{\alpha_1}{\gamma_2},$$

$$v_3 = F_2(1) = \frac{\alpha_1 \alpha_2}{\gamma_2 \gamma_3} (1 + \sigma_2) = \frac{1}{\gamma_2 \gamma_3} (\alpha_1 \alpha_2 - \beta_1 \gamma_2),$$

and

$$\begin{aligned}
u_1 &= -\frac{F_3(2)}{\gamma_2 F_3(1)} \\
&= -\frac{\frac{\alpha_2 \alpha_3}{\gamma_3 \gamma_4} (1 + \sigma_3)}{\beta_1 \left(-\frac{\alpha_1 \alpha_2 \alpha_3}{\gamma_2 \gamma_3 \gamma_4}\right) (1 + \sigma_2 + \sigma_3)} \\
&= \frac{(\alpha_2 \alpha_3 - \beta_2 \gamma_3)}{(\alpha_1 \alpha_2 \alpha_3 - \beta_1 \gamma_2 \alpha_3 - \beta_2 \gamma_3 \alpha_1)}, \\
u_2 &= -\frac{F_3(3)}{\gamma_3 F_3(1)}
\end{aligned}$$

$$\begin{aligned}
&= -\frac{\left(-\frac{\alpha_3}{\gamma_4}\right)}{\gamma_3 \left(-\frac{\alpha_1 \alpha_2 \alpha_3}{\gamma_2 \gamma_3 \gamma_4}\right) (1 + \sigma_2 + \sigma_3)} \\
&= -\frac{\gamma_2 \alpha_3}{(\alpha_1 \alpha_2 \alpha_3 - \beta_1 \gamma_2 \alpha_3 - \beta_2 \gamma_3 \alpha_1)}, \\
u_3 &= -\frac{F_3(4)}{\gamma_4 F_3(1)} \\
&= -\frac{1}{\gamma_4 \left(-\frac{\alpha_1 \alpha_2 \alpha_3}{\gamma_2 \gamma_3 \gamma_4}\right) (1 + \sigma_2 + \sigma_3)} \\
&= \frac{\gamma_2 \gamma_3}{(\alpha_1 \alpha_2 \alpha_3 - \beta_1 \gamma_2 \alpha_3 - \beta_2 \gamma_3 \alpha_1)}.
\end{aligned}$$

Note that the expressions for  $r_1, r_2, r_3, s_1, s_2, s_3, u_1, u_2, u_3, v_1, v_2, v_3$  do not contain  $\beta_3$  or  $\gamma_4$ .

From (66), the diagonal elements of the inverse of  $\Phi_3$  are then given by

$$\begin{aligned}
\psi_{1,1} &= u_1 v_1 = \frac{(\alpha_2 \alpha_3 - \beta_2 \gamma_3)}{(\alpha_1 \alpha_2 \alpha_3 - \beta_1 \gamma_2 \alpha_3 - \beta_2 \gamma_3 \alpha_1)}, \\
\psi_{2,2} &= u_2 v_2 = \frac{\alpha_1 \alpha_3}{(\alpha_1 \alpha_2 \alpha_3 - \beta_1 \gamma_2 \alpha_3 - \beta_2 \gamma_3 \alpha_1)}, \\
\psi_{3,3} &= u_3 v_3 = \frac{(\alpha_1 \alpha_2 - \beta_1 \gamma_2)}{(\alpha_1 \alpha_2 \alpha_3 - \beta_1 \gamma_2 \alpha_3 - \beta_2 \gamma_3 \alpha_1)}.
\end{aligned}$$

The upper diagonal elements are given by

$$\begin{aligned}
\psi_{1,2} &= r_1 s_2 = -\frac{\beta_1 \alpha_3}{(\alpha_1 \alpha_2 \alpha_3 - \beta_1 \gamma_2 \alpha_3 - \beta_2 \gamma_3 \alpha_1)}, \\
\psi_{1,3} &= r_1 s_3 = \frac{\beta_1 \beta_2}{(\alpha_1 \alpha_2 \alpha_3 - \beta_1 \gamma_2 \alpha_3 - \beta_2 \gamma_3 \alpha_1)}, \\
\psi_{2,3} &= r_2 s_3 = -\frac{\alpha_1 \beta_2}{(\alpha_1 \alpha_2 \alpha_3 - \beta_1 \gamma_2 \alpha_3 - \beta_2 \gamma_3 \alpha_1)}.
\end{aligned}$$

Each of the lower diagonal elements  $\psi_{i,j}$  is obtained by interchanging  $\beta_{k-1}$  and  $\gamma_k$ ,  $k = 2, 3$  in the corresponding upper diagonal element  $\psi_{j,i}$ . Therefore,

$$\begin{aligned}
\psi_{2,1} &= u_2 v_1 = -\frac{\gamma_2 \alpha_3}{(\alpha_1 \alpha_2 \alpha_3 - \beta_1 \gamma_2 \alpha_3 - \beta_2 \gamma_3 \alpha_1)}, \\
\psi_{3,1} &= u_3 v_1 = \frac{\gamma_2 \gamma_3}{(\alpha_1 \alpha_2 \alpha_3 - \beta_1 \gamma_2 \alpha_3 - \beta_2 \gamma_3 \alpha_1)},
\end{aligned}$$

$$\psi_{3,2} = u_3 v_2 = -\frac{\alpha_1 \gamma_3}{(\alpha_1 \alpha_2 \alpha_3 - \beta_1 \gamma_2 \alpha_3 - \beta_2 \gamma_3 \alpha_1)}.$$

## 5. Characteristic polynomial

The tridiagonal matrix of (62) satisfies

$$\Phi_K = \text{diag}(\beta_1, \dots, \beta_K) \mathbf{T}_K, \quad (76)$$

where  $\mathbf{T}_K$  is given by (51) with

$$A_i = -\frac{\alpha_i}{\beta_i}, \quad 1 \leq i \leq K, \quad B_i = -\frac{\gamma_i}{\beta_i}, \quad 2 \leq i \leq K, \quad \beta_K = 1,$$

as in (63a), (63b). From (76) and Proposition 5, we get

$$\det(\Phi_K) = (-1)^K (\beta_1 \cdots \beta_K) C_K = (-1)^K (\beta_1 \cdots \beta_K) E_K(1). \quad (77)$$

Using (65b), this can be rewritten as

$$\det(\Phi_K) = (\alpha_1 \cdots \alpha_K) \left( 1 + \sum_{q=1}^{\lfloor K/2 \rfloor} \sum_{(k_1, \dots, k_q) \in S_q(2, K)} (\sigma_{k_1} \cdots \sigma_{k_q}) \right), \quad (78)$$

where  $\sigma_k = -((\beta_{k-1} \gamma_k)/(\alpha_{k-1} \alpha_k))$  as in (65c).

Now the characteristic polynomial of  $\Phi_K$  is given by

$$\det(\lambda \mathbf{I}_K - \Phi_K) = (-1)^K \det(\Phi_K - \lambda \mathbf{I}_K), \quad (79)$$

where  $\mathbf{I}_K$  denotes the  $K \times K$  identity matrix. By replacing  $\alpha_k$  by  $-(\lambda - \alpha_k)$  for  $k = 1, \dots, K$  in (79), we get

$$\begin{aligned} \det(\lambda \mathbf{I}_K - \Phi_K) &= (\lambda - \alpha_1) \cdots (\lambda - \alpha_K) \\ &\times \left[ 1 + \sum_{q=1}^{\lfloor K/2 \rfloor} \sum_{(k_1, \dots, k_q) \in S_q(2, K)} \eta_{k_1}(\lambda) \cdots \eta_{k_q}(\lambda) \right], \end{aligned} \quad (80a)$$

where

$$\eta_k(\lambda) = -\frac{\beta_{k-1} \gamma_k}{(\lambda - \alpha_{k-1})(\lambda - \alpha_k)}, \quad (80b)$$

and  $S_q(2, K)$  is given by (1a)–(1c). Thus, (80a) and (80b) give an explicit formula for the characteristic polynomial of a tridiagonal matrix.

For example, when  $K = 4$  we get

$$\begin{aligned} \det(\lambda \mathbf{I}_4 - \Phi_4) &= (\lambda - \alpha_1)(\lambda - \alpha_2)(\lambda - \alpha_3)(\lambda - \alpha_4) \\ &\times [1 + \eta_2(\lambda) + \eta_3(\lambda) + \eta_4(\lambda) + \eta_2(\lambda)\eta_4(\lambda)] \\ &= (\lambda - \alpha_1)(\lambda - \alpha_2)(\lambda - \alpha_3)(\lambda - \alpha_4) \\ &\quad - \beta_1 \gamma_2(\lambda - \alpha_3)(\lambda - \alpha_4) - \beta_2 \gamma_3(\lambda - \alpha_1)(\lambda - \alpha_4) \\ &\quad - \beta_3 \gamma_4(\lambda - \alpha_1)(\lambda - \alpha_2) + \beta_1 \gamma_2 \beta_3 \gamma_4. \end{aligned}$$

## 6. Orthogonal polynomials

Consider a set  $\{p_n(t)\}_{n \geq 0}$  of polynomials that are orthogonal on the interval  $[a, b]$  with respect to some nonnegative weight function  $w(t)$  [18]. The polynomial  $p_n(t)$  has degree  $n$ , and satisfies the second-order homogeneous difference equation [18,19]

$$\beta_n p_n(t) = (t - \alpha_n) p_{n-1}(t) - \gamma_n p_{n-2}(t), \quad n \geq 1, \quad (81)$$

where  $\beta_n, p_n \neq 0$ ,  $p_0(t)$  is some nonzero constant, and  $p_{-1}(t)$  is a function of  $t$ , which may be identically zero. The difference equation (81) can be rewritten as

$$p_n(t) = \frac{(t - \alpha_n)}{\beta_n} p_{n-1}(t) - \frac{\gamma_n}{\beta_n} p_{n-2}(t), \quad n \geq 1, \quad (82)$$

with initial values  $p_{-1}(t)$  and  $p_0(t)$ . Comparing (82) with the homogeneous difference equation (10), we can write

$$y_{n+2} = p_n(t), \quad n = 1, 2, \dots, \quad (83a)$$

$$A_n = \frac{(t - \alpha_n)}{\beta_n}, \quad B_n = -\frac{\gamma_n}{\beta_n}. \quad (83b)$$

From (12)–(14b), (83a) and (83b), the solution of (82) is given by

$$p_n(t) = \mathcal{C}_n(t) p_0(t) + \mathcal{D}_n(t) p_{-1}(t), \quad n \geq 0, \quad (84)$$

where

$$\mathcal{C}_0(t) = 1, \quad \mathcal{C}_1(t) = \frac{(t - \alpha_1)}{\beta_1}, \quad (85a)$$

$$\mathcal{D}_0(t) = 0, \quad \mathcal{D}_1(t) = -\frac{\gamma_1}{\beta_1}, \quad \mathcal{D}_2(t) = -\frac{\gamma_1(t - \alpha_2)}{\beta_1 \beta_2} \quad (85b)$$

and

$$\begin{aligned} \mathcal{C}_n(t) &= \frac{(t - \alpha_1) \cdots (t - \alpha_n)}{\beta_1 \cdots \beta_n} \\ &\times \left[ 1 + \sum_{q=1}^{\lfloor n/2 \rfloor} \sum_{(k_1, \dots, k_q) \in S_q(2, n)} \eta_{k_1}(t) \cdots \eta_{k_q}(t) \right] \quad \text{for } n \geq 2, \end{aligned} \quad (86a)$$

$$\begin{aligned} \mathcal{D}_n(t) &= -\frac{\gamma_1(t - \alpha_2) \cdots (t - \alpha_n)}{\beta_1 \cdots \beta_n} \\ &\times \left[ 1 + \sum_{q=1}^{\lfloor (n-1)/2 \rfloor} \sum_{(k_1, \dots, k_q) \in S_q(3, n)} \eta_{k_1}(t) \cdots \eta_{k_q}(t) \right] \quad \text{for } n \geq 3, \end{aligned} \quad (86b)$$

such that

$$\eta_k(t) = -\frac{\beta_{k-1}\gamma_k}{(t - \alpha_{k-1})(t - \alpha_k)}$$

as in (80b).

Note that  $\mathcal{C}_n(t)$  is a polynomial of degree  $n$  for  $n \geq 0$ , while  $\mathcal{D}_n(t)$  is a polynomial of degree  $n - 1$  for  $n \geq 1$ . Thus, (84) is an expression for  $p_n(t)$ ,  $n \geq 0$ , in terms of the recurrence coefficients  $\alpha_1, \dots, \alpha_n$ ,  $\beta_1, \dots, \beta_n$ ,  $\gamma_1, \dots, \gamma_n$  and initial values  $p_{-1}(t)$ ,  $p_0(t)$ .

Let

$$\mathbf{p}(t) = \begin{bmatrix} p_0(t) \\ \vdots \\ p_{K-1}(t) \end{bmatrix} \quad (87)$$

denote the  $K \times 1$  vector of orthogonal polynomials, and  $\Phi_K$  the tridiagonal matrix in (62) of coefficients  $\alpha_1, \dots, \alpha_K$ ,  $\beta_1, \dots, \beta_{K-1}$ ,  $\gamma_2, \dots, \gamma_K$  of the difference equation (81) for  $n = 1, \dots, K$ . We can rewrite (81) for  $n = 1, \dots, K$  as [19,20]

$$(t\mathbf{I}_K - \Phi_K)\mathbf{p}(t) = \beta_K p_K(t)\mathbf{e}_K + \gamma_1 p_{-1}(t)\mathbf{e}_1, \quad (88)$$

where  $\mathbf{e}_1$  and  $\mathbf{e}_K$  are the first and last columns, respectively, of  $\mathbf{I}_K$ . If  $t$  is not an eigenvalue of  $\Phi_K$ , we get from (88)

$$\mathbf{p}(t) = -\beta_K p_K(t)(\Phi_K - t\mathbf{I}_K)^{-1}\mathbf{e}_K - \gamma_1 p_{-1}(t)(\Phi_K - t\mathbf{I}_K)^{-1}\mathbf{e}_1. \quad (89)$$

Replacing each diagonal element  $\alpha_k$  by  $\alpha_k - t$  in  $\Phi_K$ , we obtain the matrix  $\Phi_K - t\mathbf{I}_K$ . Thus, each of the polynomials  $p_0(t), \dots, p_{K-1}(t)$  can be expressed in terms of the inverse of the tridiagonal matrix  $\Phi_K - t\mathbf{I}_K$  and the functions  $p_K(t)$  and  $p_{-1}(t)$ .

The vector  $(\Phi_K - t\mathbf{I}_K)^{-1}\mathbf{e}_K$  is the  $K$ th column of  $(\Phi_K - t\mathbf{I}_K)^{-1}$ . Replacing  $\alpha_k$  by  $-(t - \alpha_k)$ ,  $k = 1, \dots, K$  in (65b), we obtain from (72) the following expression for  $\psi_{n+1,K}(t)$ , the element in the  $(n + 1)$ th row and  $K$ th column of  $(\Phi_K - t\mathbf{I}_K)^{-1}$ :

$$\psi_{n+1,K}(t)$$

$$= -\frac{\frac{(t - \alpha_1) \cdots (t - \alpha_n)}{\beta_1 \cdots \beta_n} \left[ 1 + \sum_{q=1}^{\lfloor n/2 \rfloor} \sum_{(k_1, \dots, k_q) \in S_q(2, n)} \eta_{k_1}(t) \cdots \eta_{k_q}(t) \right]}{\beta_K \frac{(t - \alpha_1) \cdots (t - \alpha_K)}{\beta_1 \cdots \beta_K} \left[ 1 + \sum_{q=1}^{\lfloor K/2 \rfloor} \sum_{(k_1, \dots, k_q) \in S_q(2, K)} \eta_{k_1}(t) \cdots \eta_{k_q}(t) \right]}. \quad (90)$$

Similarly, by replacing  $\alpha_k$  by  $-(t - \alpha_k)$ ,  $k = 1, \dots, K$  in (69), we obtain from (72) the following expression for  $\psi_{n+1,1}(t)$ , the element in the  $(n + 1)$ th row and first column of  $(\Phi_K - t\mathbf{I}_K)^{-1}$ :

$$\psi_{n+1,1}(t) = - \frac{\frac{(t - \alpha_{n+2}) \cdots (t - \alpha_K)}{\gamma_{n+3} \cdots \gamma_{K+1}} \left[ 1 + \sum_{q=1}^{\lfloor (K-n-1)/2 \rfloor} \sum_{(k_1, \dots, k_q) \in S_q(n+3, K)} \eta_{k_1}(t) \cdots \eta_{k_q}(t) \right]}{\gamma_{n+2} \frac{(t - \alpha_1) \cdots (t - \alpha_K)}{\gamma_2 \cdots \gamma_{K+1}} \left[ 1 + \sum_{q=1}^{\lfloor K/2 \rfloor} \sum_{(k_1, \dots, k_q) \in S_q(2, K)} \eta_{k_1}(t) \cdots \eta_{k_q}(t) \right]}. \quad (91)$$

Now  $p_n(t)$  is the  $(n+1)$ th row of  $\mathbf{p}$  in (89). Therefore, combining (89), (90) and (91), we obtain, for  $n = 0, \dots, K-1$ ,

$$p_n(t) = \frac{(\beta_{n+1} \cdots \beta_K) \left[ 1 + \sum_{q=1}^{\lfloor n/2 \rfloor} \sum_{(k_1, \dots, k_q) \in S_q(2, n)} \eta_{k_1}(t) \cdots \eta_{k_q}(t) \right]}{(t - \alpha_{n+1}) \cdots (t - \alpha_K) \left[ 1 + \sum_{q=1}^{\lfloor K/2 \rfloor} \sum_{(k_1, \dots, k_q) \in S_q(2, K)} \eta_{k_1}(t) \cdots \eta_{k_q}(t) \right]} p_K(t) + \frac{(\gamma_1 \cdots \gamma_{n+1}) \left[ 1 + \sum_{q=1}^{\lfloor (K-n-1)/2 \rfloor} \sum_{(k_1, \dots, k_q) \in S_q(n+3, K)} \eta_{k_1}(t) \cdots \eta_{k_q}(t) \right]}{(t - \alpha_1) \cdots (t - \alpha_{n+1}) \left[ 1 + \sum_{q=1}^{\lfloor K/2 \rfloor} \sum_{(k_1, \dots, k_q) \in S_q(2, K)} \eta_{k_1}(t) \cdots \eta_{k_q}(t) \right]} p_{-1}(t), \quad (92)$$

where  $\eta_k(t)$  is given by (80b). Thus, (92) is an expression for the  $K$  intermediate polynomials  $p_n(t)$ ,  $0 \leq n \leq K-1$ , in terms of the recurrence coefficients  $\alpha_1, \dots, \alpha_K, \beta_1, \dots, \beta_K, \gamma_1, \dots, \gamma_K$  and boundary values  $p_{-1}(t), p_K(t)$ .

We now show how the general formula for  $p_n(t)$  in (84) can be used to obtain expressions for Legendre, Hermite and Chebyshev polynomials.

### 6.1. Legendre polynomials

Legendre polynomials can be generated by the difference equation

$$p_n(t) = \frac{(2n-1)}{n} t p_{n-1}(t) - \frac{(n-1)}{n} p_{n-2}(t), \quad n \geq 1, \quad (93)$$

with  $p_0(t) = 1$ . The interval of orthogonality is  $[-1, 1]$  and the weight function  $w(t) = 1$ . Comparing (93) with (82), we obtain

$$\alpha_n = 0, \quad \beta_n = \frac{n}{(2n-1)}, \quad \gamma_n = \frac{(n-1)}{(2n-1)}, \quad n \geq 1. \quad (94)$$

Since  $\gamma_1 = 0$ , we have  $\mathcal{D}_n(t) = 0$ . Therefore,  $p_{-1}(t)$  does not affect the solution of (93).

Now, from (80b),

$$\eta_k(t) = -\frac{(k-1)^2}{(2k-3)(2k-1)t^2}. \quad (95)$$

Using (86a), we can write the solution of (93) as

$$\begin{aligned} p_n(t) &= \mathcal{C}_n(t) \\ &= \frac{(2n)!}{2^n (n!)^2} \\ &\quad \times \left[ t^n + \sum_{q=1}^{\lfloor n/2 \rfloor} (-1)^q t^{n-2q} \sum_{(k_1, \dots, k_q) \in S_q(2, n)} \prod_{i=1}^q \frac{(k_i - 1)^2}{(2k_i - 3)(2k_i - 1)} \right]. \end{aligned} \quad (96)$$

Denote  $G_{n,q}$  as

$$G_{n,q} = \sum_{(k_1, \dots, k_q) \in S_q(2, n)} \prod_{i=1}^q \frac{(k_i - 1)^2}{(2k_i - 3)(2k_i - 1)}. \quad (97)$$

Applying property (3) on  $S_q(2, n)$  we get

$$\begin{aligned} S_q(2, n) &= S_q(2, n-1) \cup \{ (k_1, \dots, k_{q-1}, k_q) : k_q = n; \\ &\quad (k_1, \dots, k_{q-1}) \in S_{q-1}(2, n-2) \}, \end{aligned} \quad (98)$$

which implies that  $G_{n,q}$  in (97) follows the recurrence

$$G_{n,q} = G_{n-1,q} + \frac{(n-1)^2}{(2n-3)(2n-1)} G_{n-2,q-1}. \quad (99)$$

From (99), it can be shown by mathematical induction that

$$G_{n,q} = \frac{(n!)^2 (2n-2q)!}{(2n)! q! (n-q)! (n-2q)!}. \quad (100)$$

Combining (96), (97) and (100), we finally obtain

$$p_n(t) = \sum_{q=1}^{\lfloor n/2 \rfloor} \frac{(-1)^q (2n-2q)!}{2^n q! (n-q)! (n-2q)!} t^{n-2q}, \quad (101)$$

which is the expression for the Legendre polynomial of degree  $n$ . This can alternatively be written as

$$p_n(t) = \frac{1}{2^n n!} \frac{d^n}{dt^n} [(t^2 - 1)^n].$$



## 6.2. Hermite polynomials

Hermite polynomials can be generated by the recurrence

$$p_n(t) = tp_{n-1}(t) - (n-1)p_{n-2}(t), \quad n \geq 1, \quad (102)$$

with  $p_0(t) = 1$ . The interval of orthogonality is  $(-\infty, \infty)$  and the weight function  $w(t) = e^{-(t^2/2)}$ . Comparing (102) with (82), we get

$$\alpha_n = 0, \quad \beta_n = 1, \quad \gamma_n = (n-1), \quad n \geq 1. \quad (103)$$

Since  $\gamma_1 = 0$ , we have  $\mathcal{D}_n(t) = 0$ ; hence,  $p_{-1}(t)$  does not affect the solution of (102).

From (80b),

$$\eta_k(t) = -\frac{(k-1)}{t^2}. \quad (104)$$

We can then write the solution of (102) as

$$p_n(t) = \mathcal{C}_n(t) = t^n + \sum_{q=1}^{\lfloor n/2 \rfloor} (-1)^q t^{n-2q} \sum_{(k_1, \dots, k_q) \in S_q(2, n)} \prod_{i=1}^q (k_i - 1). \quad (105)$$

Denoting  $G_{n,q}$  as

$$G_{n,q} = \sum_{(k_1, \dots, k_q) \in S_q(2, n)} \prod_{i=1}^q (k_i - 1), \quad (106)$$

and applying property (3) on  $S_q(2, n)$  we obtain the recurrence

$$G_{n,q} = G_{n-1,q} + (n-1)G_{n-2,q-1}. \quad (107)$$

It can be shown by mathematical induction using this recurrence that

$$G_{n,q} = \frac{n!}{2^q q! (n-2q)!}. \quad (108)$$

Combining (105), (106) and (108), we get

$$p_n(t) = \sum_{q=0}^{\lfloor n/2 \rfloor} \frac{(-1)^q n!}{2^q q! (n-2q)!} t^{n-2q}, \quad (109)$$

which is the expression for the Hermite polynomial of degree  $n$ . This can alternatively be written as

$$p_n(t) = (-1)^n e^{-(t^2/2)} \frac{d^n}{dt^n} [e^{-(t^2/2)}].$$

## 6.3. Chebyshev polynomials

Chebyshev polynomials can be generated by the difference equation

$$p_n(t) = 2tp_{n-1}(t) - p_{n-2}(t), \quad n \geq 1, \quad (110)$$

whose coefficients do not vary with  $n$ . The interval of orthogonality is  $[-1, 1]$  and the weight function  $w(t) = 1/\sqrt{1-t^2}$ .

Here

$$\alpha_n = 0, \quad \beta_n = \frac{1}{2} = \gamma_n, \quad n \geq 1, \quad (111)$$

and

$$\eta_k(t) = -\frac{1}{4t^2}. \quad (112)$$

Since  $\eta_k(t)$  does not depend on  $k$ , we can apply property (2) on  $S_q(2, n)$  and  $S_q(3, n)$  in (86a) and (86b). This results in

$$|S_q(2, n)| = \binom{n-q}{q}, \quad |S_q(3, n)| = \binom{n-q-1}{q}, \quad (113)$$

followed by

$$\mathcal{C}_n(t) = (2t)^n + \sum_{q=1}^{\lfloor n/2 \rfloor} (-1)^q \binom{n-q}{q} (2t)^{n-2q}, \quad (114a)$$

$$\mathcal{D}_n(t) = -\mathcal{C}_{n-1}(t). \quad (114b)$$

For the Chebyshev polynomials of the *first kind*, we have

$$p_0(t) = 1, \quad p_{-1}(t) = t,$$

and therefore

$$\begin{aligned} p_n(t) &= \mathcal{C}_n(t) + t\mathcal{D}_n(t) \\ &= \mathcal{C}_n(t) - t\mathcal{C}_{n-1}(t) \\ &= \frac{1}{2} \left\{ (2t)^n + \sum_{q=1}^{\lfloor n/2 \rfloor} (-1)^q \left[ \binom{n-q}{q} + \binom{n-q-1}{q} \right] (2t)^{n-2q} \right\}. \end{aligned} \quad (115)$$

This can also be expressed as

$$p_n(t) = \cos(n \cos^{-1}(t)).$$

For the Chebyshev polynomials of the *second kind*, we have

$$p_0(t) = 1, \quad p_{-1}(t) = 0,$$

and therefore  $p_n(t) = \mathcal{C}_n(t)$ , where  $\mathcal{C}_n(t)$  is given by (114a). This can also be expressed as

$$p_n(t) = \frac{\sin((n+1) \cos^{-1}(t))}{\sin(\cos^{-1}(t))} = \frac{\sin((n+1) \cos^{-1}(t))}{\sqrt{1-t^2}}.$$

## 7. A cyclic tridiagonal system

Consider the second-order linear difference equation over the finite index interval  $[1, K + 2]$ ,  $K \geq 2$ , given by

$$\beta_n y_{n+2} = -\alpha_n y_{n+1} - \gamma_n y_n + \beta_n x_{n+2}, \quad 1 \leq n \leq K, \quad (116)$$

with variable complex coefficients  $\beta_n, \alpha_n, \gamma_n$ , where  $\beta_n, \gamma_n \neq 0$ , complex forcing term  $x_{n+2}$ , and complex boundary values  $y_1, y_{K+2}$ . The boundary values are periodic with period  $K$ , that is, we have the boundary conditions

$$y_1 = y_{K+1}, \quad y_{K+2} = y_2. \quad (117)$$

The  $K$  equations in (116) can be written in matrix form after applying the boundary conditions as

$$A_K \begin{bmatrix} y_2 \\ \vdots \\ y_{K+1} \end{bmatrix} = \text{diag}(\beta_1, \dots, \beta_K) \begin{bmatrix} x_3 \\ \vdots \\ x_{K+2} \end{bmatrix}, \quad (118a)$$

where

$$A_K = \begin{bmatrix} \alpha_1 & \beta_1 & & & \gamma_1 \\ \gamma_2 & \alpha_2 & \beta_2 & & \mathbf{0} \\ & \ddots & \ddots & \ddots & \\ \mathbf{0} & & \gamma_{K-1} & \alpha_{K-1} & \beta_{K-1} \\ \beta_K & & & \gamma_K & \alpha_K \end{bmatrix}, \quad (118b)$$

$$\beta_1, \dots, \beta_K \neq 0, \quad \gamma_1, \dots, \gamma_K \neq 0.$$

The system ((118a), (118b)) of equations is a *cyclic linear tridiagonal system* [21], and the matrix  $A_K$  in (118b) a *cyclic tridiagonal matrix*. The solution of system ((118a), (118b)), or alternatively, of the difference equation (116) with boundary conditions (117) for  $n = 1, \dots, K$  can be expressed as

$$\begin{bmatrix} y_2 \\ \vdots \\ y_{K+1} \end{bmatrix} = A_K^{-1} \begin{bmatrix} \beta_1 x_3 \\ \vdots \\ \beta_K x_{K+2} \end{bmatrix}. \quad (119)$$

Let

$$A_K^{-1} = [\omega_{i,j}]_{i,j=1}^K. \quad (120)$$

The difference equation (116) can be rewritten as

$$y_{n+2} = A_n y_{n+1} + B_n y_n + x_{n+2}, \quad 1 \leq n \leq K, \quad (121a)$$

as in (41) with

$$A_n = -\frac{\alpha_n}{\beta_n}, \quad B_n = -\frac{\gamma_n}{\beta_n}. \quad (121b)$$

From Proposition 4, the quantities  $y_3, \dots, y_{K+2}$  of recurrence (121a) can be expressed in terms of  $y_1, y_2$  as

$$y_{i+1} = C_{i-1}y_2 + D_{i-1}y_1 + \sum_{j=1}^{i-1} E_{i-1}(j+1)x_{j+2}, \quad 2 \leq i \leq K+1, \quad (122)$$

where  $E_n(m)$  is given by (65b), and from (13a), (13b), (14a), (14b) and (121b),

$$C_1 = -\frac{\alpha_1}{\beta_1}, \quad D_1 = -\frac{\gamma_1}{\beta_1}, \quad D_2 = \frac{\gamma_1\alpha_2}{\beta_1\beta_2}, \quad (123a)$$

$$C_n = (-1)^n \left( \frac{\alpha_1}{\beta_1} \cdots \frac{\alpha_n}{\beta_n} \right) \left( 1 + \sum_{q=1}^{\lfloor n/2 \rfloor} \sum_{(k_1, \dots, k_q) \in S_q(2, n)} (\sigma_{k_1} \cdots \sigma_{k_q}) \right) \\ \text{for } n \geq 2, \quad (123b)$$

$$D_n = (-1)^n \frac{\gamma_1}{\beta_1} \left( \frac{\alpha_2}{\beta_2} \cdots \frac{\alpha_n}{\beta_n} \right) \left( 1 + \sum_{q=1}^{\lfloor (n-1)/2 \rfloor} \sum_{(k_1, \dots, k_q) \in S_q(3, n)} (\sigma_{k_1} \cdots \sigma_{k_q}) \right) \\ \text{for } n \geq 3, \quad (123c)$$

such that  $S_q(L, U)$  is defined by (1a)–(1c), and  $\sigma_k = -((\beta_{k-1}\gamma_k)/(\alpha_{k-1}\alpha_k))$  as in (65c).

Putting  $i = K$  in (122) and applying the boundary condition  $y_{K+1} = y_1$  in the resulting equation, we get

$$(1 - D_{K-1})y_1 - C_{K-1}y_2 = \sum_{j=1}^{K-1} E_{K-1}(j+1)x_{j+2}.$$

Similarly, putting  $i = K+1$  and  $y_{K+2} = y_2$  in (122) results in

$$-D_K y_1 + (1 - C_K)y_2 = \sum_{j=1}^K E_K(j+1)x_{j+2}.$$

We can then express  $y_1$  and  $y_2$  in terms of  $x_3, \dots, x_{K+2}$  as

$$\begin{bmatrix} y_1 \\ y_2 \end{bmatrix} = \begin{bmatrix} 1 - D_{K-1} & -C_{K-1} \\ -D_K & 1 - C_K \end{bmatrix}^{-1} \begin{bmatrix} \sum_{j=1}^{K-1} E_{K-1}(j+1)x_{j+2} \\ \sum_{j=1}^K E_K(j+1)x_{j+2} \end{bmatrix}. \quad (124)$$

Substituting (124) in (122), we can express  $y_{i+1}$  as

$$y_{i+1} = \sum_{j=1}^K \omega_{i,j} \beta_j x_{j+2}, \quad (125)$$

which implies from (119) and (120) that  $\omega_{i,j}$  is the element in the  $i$ th row and  $j$ th column of  $A^{-1}$ . Thus, we can obtain an explicit formula for the inverse of a cyclic tridiagonal matrix by solving a boundary value problem with periodic boundary conditions.

## 8. Case of constant diagonals

When

$$\alpha_n = \alpha, \quad \beta_n = \beta, \quad \gamma_n = \gamma, \quad n = 1, \dots, K, \quad (126)$$

the tridiagonal matrix  $\Phi_K$  in (62) is *Toeplitz*, while the cyclic tridiagonal matrix  $A_K$  in (118b) is *circulant*, which also implies that it is Toeplitz. The quantities  $C_n$ ,  $D_n$  and  $E_n(m)$  in (65b) and (123a)–(123c) can now be expressed in terms of the Chebyshev polynomials of the second kind discussed in Section 6.3 as

$$E_n(m) = (-1)^{n-m+1} \left( \sqrt{\frac{\gamma}{\beta}} \right)^{n-m+1} p_{n-m+1} \left( \frac{\alpha}{2\sqrt{\gamma\beta}} \right),$$

$$C_n = E_n(1) = (-1)^n \left( \sqrt{\frac{\gamma}{\beta}} \right)^n p_n \left( \frac{\alpha}{2\sqrt{\gamma\beta}} \right),$$

$$D_n = (-1)^n \left( \sqrt{\frac{\gamma}{\beta}} \right)^{n+1} p_{n-1} \left( \frac{\alpha}{2\sqrt{\gamma\beta}} \right),$$

where, using (114a),

$$p_n \left( \frac{\alpha}{2\sqrt{\gamma\beta}} \right) = \left( \frac{\alpha}{\sqrt{\gamma\beta}} \right)^n \left[ 1 + \sum_{q=1}^{\lfloor n/2 \rfloor} (-1)^q \binom{n-q}{q} \left( \frac{\gamma\beta}{\alpha^2} \right)^q \right].$$

From (72), the elements of the inverse of the tridiagonal matrix  $\Phi_K$  in (62) under condition (126) can be expressed as

$$\psi_{i,j} = \begin{cases} \frac{(-1)^{i-j}}{\sqrt{\gamma\beta}} \left( \sqrt{\frac{\gamma}{\beta}} \right)^{i-j} \frac{p_{j-1} \left( \frac{\alpha}{2\sqrt{\gamma\beta}} \right) p_{K-i} \left( \frac{\alpha}{2\sqrt{\gamma\beta}} \right)}{p_K \left( \frac{\alpha}{2\sqrt{\gamma\beta}} \right)} & \text{if } i > j, \\ \frac{(-1)^{j-i}}{\sqrt{\gamma\beta}} \left( \sqrt{\frac{\beta}{\gamma}} \right)^{j-i} \frac{p_{i-1} \left( \frac{\alpha}{2\sqrt{\gamma\beta}} \right) p_{K-j} \left( \frac{\alpha}{2\sqrt{\gamma\beta}} \right)}{p_K \left( \frac{\alpha}{2\sqrt{\gamma\beta}} \right)} & \text{if } i \leq j, \\ E_K(1) \neq 0. \end{cases}$$

The inverse of the cyclic tridiagonal matrix  $A_K$  in (118b) under condition (126) can also be obtained from  $E_n(m)$ ,  $C_n$ ,  $D_n$  using the approach discussed in Section 7.

## 9. Conclusions

We have obtained explicit formulae for the elements of the inverse of a general tridiagonal matrix by deriving the explicit solution of a second-order linear non-homogeneous difference equation with variable coefficients, and then applying the solution to a boundary value problem with zero boundary values. Using the formula for the determinant, we have got an expression for the characteristic polynomial. A connection between the matrix inverse and orthogonal polynomials has also been established. It has also been shown that how an application of the solution of a second-order linear difference equation to a boundary value problem with periodic boundary conditions can yield the inverse of a cyclic tridiagonal matrix. In the simple case of a tridiagonal or a cyclic tridiagonal matrix with constant diagonals, the elements of the inverse can be expressed in terms of the Chebyshev polynomials of the second kind.

## Acknowledgments

The author thanks Dr. Y.V.S.S. Sanyasi Raju, Department of Mathematics, Indian Institute of Technology, Chennai, for his valuable suggestions. He also expresses his gratitude to the anonymous referee for his/her constructive comments.

## References

- [1] G. Meurant, A review on the inverse of symmetric tridiagonal and block tridiagonal matrices, *SIAM J. Matrix Anal. Appl.* 13 (3) (1992) 707–728.
- [2] S.-F. Xu, On the Jacobi matrix inverse eigenvalue problem with mixed given data, *SIAM J. Matrix Anal. Appl.* 17 (3) (1996) 632–639.
- [3] A. Bunse-Gerstner, R. Byers, V. Mehrmann, A chart of numerical methods for structured eigenvalue problems, *SIAM J. Matrix Anal. Appl.* 13 (2) (1992) 419–453.
- [4] C.F. Fischer, R.A. Usmani, Properties of some tridiagonal matrices and their application to boundary value problems, *SIAM J. Numer. Anal.* 6 (1) (1969) 127–142.
- [5] L. Lu, W. Sun, The minimal eigenvalues of a class of block-tridiagonal matrices, *IEEE Trans. Inform. Theory* 43 (2) (1997) 787–791.
- [6] G. D. Smith, *Numerical Solution of Partial Differential Equations: Finite Difference Methods*, second ed., Oxford University Press, New York, 1978.
- [7] R.M.M. Mattheij, M.D. Smooke, Estimates for the inverse of tridiagonal matrices arising in boundary-value problems, *Linear Algebra Appl.* 73 (1986) 33–57.
- [8] G.H. Golub, C.F. Van Loan, *Matrix Computations*, third ed., Johns Hopkins University Press, Baltimore, MD, 1996.
- [9] T. Yamamoto, Y. Ikebe, Inversion of band matrices, *Linear Algebra Appl.* 24 (1979) 105–111.
- [10] W.W. Barrett, A theorem on inverses of tridiagonal matrices, *Linear Algebra Appl.* 27 (1979) 211–217.
- [11] W.W. Barrett, P.J. Feinsilver, Inverses of banded matrices, *Linear Algebra Appl.* 41 (1981) 111–130.

- [12] F. Romani, On the additive structure of the inverses of banded matrices, *Linear Algebra Appl.* 80 (1986) 131–140.
- [13] R. Nabben, Two-sided bounds on the inverses of diagonally dominant tridiagonal matrices, *Linear Algebra Appl.* 287 (1999) 289–305.
- [14] W.G. Kelley, A.C. Peterson, *Difference Equations: An Introduction with Applications*, Academic Press, San Diego, CA, 1991.
- [15] S.N. Elaydi, *An Introduction to Difference Equations*, Springer, New York, 1996.
- [16] R.K. Mallik, On the solution of a second order linear homogeneous difference equation with variable coefficients, *J. Math. Anal. Appl.* 215 (1) (1997) 32–47.
- [17] Y. Ikebe, On inverses of Hessenberg matrices, *Linear Algebra Appl.* 24 (1979) 93–97.
- [18] G. Szego, *Orthogonal Polynomials*, fourth ed., AMS, Providence, RI, 1981.
- [19] S. Elhay, G.H. Golub, J. Kautsky, Jacobi matrices for sums of weight functions, *BIT* 32 (1992) 143–166.
- [20] S. Elhay, G.H. Golub, J. Kautsky, Updating and downdating of orthogonal polynomials with data fitting applications, *SIAM J. Matrix Anal. Appl.* 12 (2) (1991) 327–353.
- [21] C. Temperton, Algorithms for the solution of cyclic tridiagonal systems, *J. Comput. Physics* 19 (1975) 317–323.