

Recitation 8


Probability

Random Variables

From a probability space (S, p) , we can create a new probability space (S', p') , where S' is a partition of S , that is, $S' = \{P_1, P_2, \dots, P_n\}$, and where $p' = \Pr(P_i)$ for all i .

Visually, we can take a box partitioned into m outcomes, and then partition these m outcomes into n groups. The amount of area each group takes up is just the sum of the outcomes within the group, so this box with m groups is just like having a probability space with m outcomes.

This leads us to one of the biggest, most famous misnomers in all of mathematics: random variables. A **random variable** is a function from a set of outcomes S to $\mathbb{R}$ or $\mathbb{Z}$. We can think of this random variable as partitioning S , where each group is a set of outcomes that are all assigned the same value. We can also think of the random variable as assigning each group a different value. This diagram shows what we mean:



Here, $o_1, \dots, o_5$ are outcomes in S . Our random variable assigns two of them to 1730, one of them to 3.14, and two of them to 22. Thus, the random variable partitions the outcomes into 3 groups. Let's walk through some other more concrete examples of random variables.

Task 1

Let random variable X be on the coin flip sample space $\{H, T\}$, where $X(H) = 1$, and $X(T) = 0$. (This random variable is also known as the indicator random variable of event H .)

-
- a. If $\Pr(H) = 1/2$, and $\Pr(T) = 1/2$, then what is $\Pr(X = 1)$? How about $\Pr(X = 0)$?

- b. We could also have the random variable Y , where the domain of Y is S , all sequences of coin flips of length n , and where $Y(s) =$ the number of heads in s . If S is uniformly distributed, then what is $\Pr(Y = k)$, where $0 \leq k \leq n$?

Hint: You computed this value in last week's recitation.

- c. Let C_i be a random variable for the i th coin flip, s_i , in our sequence, s , of coin flips of length n , where $C_i(H) = 1$ and $C_i(T) = 0$. Let $C(s) = C_1(s_1) + C_2(s_2) + \dots C_n(s_n)$. Explain why $C(s) = Y(s)$.

- d. Explain why $\Pr(C = k) = \binom{n}{k} \Pr(C_i = 1)^k \Pr(C_i = 0)^{n-k}$.

- e. *Optional:* Given an expression for $\Pr(C \geq k)$.

- f. *Optional:* Compute $\Pr(C = k | C_1 = 1)$.

Checkpoint 1 — Call over a TA!

Expected Value

Intuitively, the expected value is the weighted average of values, kind of a mass center of the probability distribution.

More formally, the *expected value* of a random variable is denoted $\mathbb{E}[X]$ and is defined as

$$\mathbb{E}[X] = \sum_{s \in S} X(s) \Pr(s) = \sum_{r \in X(S)} r \Pr(X = r).$$

We define the *conditional expected values* as follows: Given that event E has occurred, the expectation of random variable X is

$$\mathbb{E}[X \mid E] = \sum_{r \in X(S)} r \Pr(X = r \mid E). \quad (1)$$

Moreover, the *linearity of expectation* can be very useful in calculating expected value: Given that Z , X , Y are three random variables defined on a sample space S and a and b are two real numbers such that $Z = aX + bY$, we know that $\mathbb{E}[Z] = a\mathbb{E}[X] + b\mathbb{E}[Y]$ must be true.

Let's practice this through a task:

Task 2

Tim and Joe are playing a game. They flip a fair coin 3 times. When the coin is heads, Joe pays \$1 to Tim; and when the coin is tails, Tim pays \$1 to Joe.

- a. Let G_i be a random variable representing what Tim gains on the i -th round. For instance, $G_3 = -1$ if the coin is tails.

What is the expected value of G_i ?

- b. Let G be a random variable that represents Tim's *total* gain in this game. What is the expected value of G ?

- c. What is the expected value of G if the coin is biased and the probability of heads is p ? in other words, generalize your solution from part b in terms of p .

-
-
- d. Tim and Joe are still using the biased coin from part c. Let H_1 be the event that the first coin is heads. What is $\mathbb{E}[G|H_1]$?

- e. Use your answers to calculate $\mathbb{E}[G]$ and $\mathbb{E}[G | H_1]$ when $p = 0.7$.

- f. *Optional:* Assume that $p = 0.7$ and let's say we want to change the game to make it "fair." If the flip is tails, then Tim pays a dollar to Joe—how much should Joe pay Tim on Heads so that for any number of flips we know $\mathbb{E}[G] = 0$?

Task 3

Tim and Joe are now playing a similar, but different, game. This time they flip a coin **2 times**. Let X be the random variable that is equal to the number of heads and Y the random variable that is equal to the number of tails. At the end of the game, Joe pays Tim X^2 dollars. Once again, let G be the random variable for Tim's gain.

- a. Calculate $\mathbb{E}[X]^2$ when the probability of heads is $p = 0.7$.

- b. Calculate $\mathbb{E}[G]$ when $p = 0.7$.

c. Does $\mathbb{E}[G] = \mathbb{E}[X]^2$?

d. Find an example that shows that $\mathbb{E}[XY] = \mathbb{E}[X]\mathbb{E}[Y]$ does not hold where X and Y are not independent variables.

Hint: Try X and Y as defined above.

e. *Optional:* Prove that if X and Y are two independent random variables, then $\mathbb{E}[XY] = \mathbb{E}[X]\mathbb{E}[Y]$.

Bayes Rule

The Bayes Rule can be summarized as

$$P(A|B) = \frac{P(B|A)P(A)}{P(B)}$$

where A and B are events and $P(B) \neq 0$.

Task 4

Assume Brown's CS department has an evaluation system for CS courses based on student evaluations. In any class, the students can fill the evaluation form and give a score of 0, 1, or 2 to the course. Let X be the random variable of this score. The students of CS0220 either like the course with probability $3/4$ (Event L) or they do not like the course with probability $1/4$ (Event NOT L).

Assume that the conditional probability distribution of X given L is

$$\Pr(X = 0 \mid L) = 1/8$$

$$\Pr(X = 1 \mid L) = 1/4$$

$$\Pr(X = 2 \mid L) = 5/8$$

and given that they do not like the course (NOT L) it is

$$\Pr(X = 0 \mid \text{NOT } L) = 9/10$$

$$\Pr(X = 1 \mid \text{NOT } L) = 1/10$$

$$\Pr(X = 2 \mid \text{NOT } L) = 0.$$

- a. If a student has given score of 0 to CS0220, what is the probability that they do not like the course?

- b. Use the definition of conditional expected value (Equation 1) and find $\mathbb{E}[X \mid \text{NOT } L]$

c. *Optional:* Find $\mathbb{E}[X]$.

Checkpoint 2 — Call over a TA!

Task 5

The Brown Review is a test-prep company that publishes books helping high school students prepare for the upcoming Accelerated Placement tests. Recently, they published a study with 20000 students nationwide showing that using their books to prepare for the AP Statistics exam resulted were 5% more likely to pass than those who studied using their rival, Karron's. However, The Brown Review did not publish all of their data, and you uncover the following data table:

	The Brown Review	Karron's	Total
Took a statistics class	$\frac{8550}{9000} = 95\%$	$\frac{4450}{4500} = 99\%$	$\frac{13000}{13500} = 96\%$
Did not take a statistics class	$\frac{750}{1000} = 75\%$	$\frac{4350}{5500} = 79\%$	$\frac{5100}{6500} = 78\%$
Total	$\frac{9300}{10000} = 93\%$	$\frac{8800}{10000} = 88\%$	$\frac{18100}{20000} = 90\%$

- a. Make an argument as to why The Brown Review appears to be a better choice for test prep.

- b. Make an argument as to why Karron's is the better choice.

- c. Given that it is very likely that a student who takes a class before the exam succeeds more often than students who did not take a class, how could the Brown Review manipulate sample sizes such that their final percentage looks higher?

This contradiction of conclusions is known as Simpson's Paradox, which you can read more about [here](#). This occurs when comparisons of two variables in separate groups yield one conclusion, but comparisons of the variables overall yield a different result due to a more significant trend working in the background, like how students taking or not taking the class determines outcomes.

- d. It is very common for news services to read the conclusion of a scientific study and report the final results directly. It is also extremely common for readers to look at a news headline and not read the article. How do these practices encourage misinformation? Why do details of a study matter?

-
-
- e. With the growing popularity of data analysis and machine learning, our society increasingly relies on statistical tools to explain the world we live in. Why is it extremely important to never take any conclusions we garner from these methods at face value? What are the dangers in becoming over-reliant on these methods to solve complex problems such as policing or policymaking?
-

Task 6

Joe goes to PVDonuts and gets four boxes of donuts for everyone. One box has two chocolate donuts, two boxes have one chocolate and one glazed, and one box has two glazed donuts.

- a. Joe picks a box at random, and gives a random donut to Tim. If that donut is chocolate, what is the probability the other donut in the box is glazed?

- b. Let's consider another scenario, where Tim picks a donut box at random. Before he opens it, Joe tells him one of the donuts in the box is chocolate.

What is the probability the other donut is glazed?

Hint: Your answer should be different from part a.

Optional: Task 7

Suppose that, in a certain family, the probability of each child being born with a cat allergy is $1/2$. You can assume one of the parents have it and it gets inherited with $1/2$ probability. Assume this family has two children, Jiahua and Ken.

- a. What is the probability that both Jiahua and Ken have a cat allergy?
- b. Consider a scenario where you go to the family's house and meet one of the children. They tell you they have a cat allergy, but not their name. What is the probability the other child is allergic?
- c. The child you meet says their name is Jiahua. Does the probability of the other child being allergic change?
- d. Given that at least one of the children is allergic and was born on a Tuesday, what is the probability the family has 2 allergic children? You may assume that the probability a child is born on a given day of the week is $1/7$.

Checkoff — Call over a TA!