

Backtracking, Modular Arithmetic, Multiplicative Inverse

Michael L. Littman

CS 0220 2021

June 18, 2021

Overview

The Greatest Common Divisor (8.2)

One Solution for All Water Jug Problems (8.2.3)

The Fundamental Theorem of Arithmetic (8.3)

Modular Arithmetic (8.5)

Turing's Code, Version 2 (8.5.1)

Arithmetic with a Prime Modulus (8.6)

Multiplicative Inverses (8.6.1)

Pulvarizing

Corollary: An integer is a linear combination of a and b iff it is a multiple of $\gcd(a, b)$.

Pulvarizing

Corollary: An integer is a linear combination of a and b iff it is a multiple of $\gcd(a, b)$.

Proof:

Let $g = \gcd(a, b)$. We showed $g = sa + tb$ for some s and t .

Pulvarizing

Corollary: An integer is a linear combination of a and b iff it is a multiple of $\gcd(a, b)$.

Proof:

Let $g = \gcd(a, b)$. We showed $g = sa + tb$ for some s and t . Any multiple of g is a linear combination of a and b :

Pulvarizing

Corollary: An integer is a linear combination of a and b iff it is a multiple of $\gcd(a, b)$.

Proof:

Let $g = \gcd(a, b)$. We showed $g = sa + tb$ for some s and t . Any multiple of g is a linear combination of a and b :

$$kg = k(sa + tb) = (ks)a + (kt)b.$$

Pulvarizing

Corollary: An integer is a linear combination of a and b iff it is a multiple of $\gcd(a, b)$.

Proof:

Let $g = \gcd(a, b)$. We showed $g = sa + tb$ for some s and t . Any multiple of g is a linear combination of a and b :

$$kg = k(sa + tb) = (ks)a + (kt)b.$$

We know $a = k_1g$ and $b = k_2g$ because

Pulvarizing

Corollary: An integer is a linear combination of a and b iff it is a multiple of $\gcd(a, b)$.

Proof:

Let $g = \gcd(a, b)$. We showed $g = sa + tb$ for some s and t . Any multiple of g is a linear combination of a and b :

$$kg = k(sa + tb) = (ks)a + (kt)b.$$

We know $a = k_1g$ and $b = k_2g$ because g is a common divisor of a and b .

Pulvarizing

Corollary: An integer is a linear combination of a and b iff it is a multiple of $\gcd(a, b)$.

Proof:

Let $g = \gcd(a, b)$. We showed $g = sa + tb$ for some s and t . Any multiple of g is a linear combination of a and b :

$$kg = k(sa + tb) = (ks)a + (kt)b.$$

We know $a = k_1g$ and $b = k_2g$ because g is a common divisor of a and b . Any linear combination of a and b is a multiple of g :

Pulvarizing

Corollary: An integer is a linear combination of a and b iff it is a multiple of $\gcd(a, b)$.

Proof:

Let $g = \gcd(a, b)$. We showed $g = sa + tb$ for some s and t . Any multiple of g is a linear combination of a and b :

$$kg = k(sa + tb) = (ks)a + (kt)b.$$

We know $a = k_1g$ and $b = k_2g$ because g is a common divisor of a and b . Any linear combination of a and b is a multiple of g :

$$s'a + t'b = s'(k_1g) + t'(k_2g) = (s'k_1 + t'k_2)g.$$

Pulvarizing

Corollary: An integer is a linear combination of a and b iff it is a multiple of $\gcd(a, b)$.

Proof:

Let $g = \gcd(a, b)$. We showed $g = sa + tb$ for some s and t . Any multiple of g is a linear combination of a and b :

$$kg = k(sa + tb) = (ks)a + (kt)b.$$

We know $a = k_1g$ and $b = k_2g$ because g is a common divisor of a and b . Any linear combination of a and b is a multiple of g :

$$s'a + t'b = s'(k_1g) + t'(k_2g) = (s'k_1 + t'k_2)g.$$

Mixing a and b in different combinations, we can only make multiples of g .

Pulvarizing

Corollary: An integer is a linear combination of a and b iff it is a multiple of $\gcd(a, b)$.

Proof:

Let $g = \gcd(a, b)$. We showed $g = sa + tb$ for some s and t . Any multiple of g is a linear combination of a and b :

$$kg = k(sa + tb) = (ks)a + (kt)b.$$

We know $a = k_1g$ and $b = k_2g$ because g is a common divisor of a and b . Any linear combination of a and b is a multiple of g :

$$s'a + t'b = s'(k_1g) + t'(k_2g) = (s'k_1 + t'k_2)g.$$

Mixing a and b in different combinations, we can only make multiples of g .

Note: The combinations are not unique: $sa + tb =$

Pulvarizing

Corollary: An integer is a linear combination of a and b iff it is a multiple of $\gcd(a, b)$.

Proof:

Let $g = \gcd(a, b)$. We showed $g = sa + tb$ for some s and t . Any multiple of g is a linear combination of a and b :

$$kg = k(sa + tb) = (ks)a + (kt)b.$$

We know $a = k_1g$ and $b = k_2g$ because g is a common divisor of a and b . Any linear combination of a and b is a multiple of g :

$$s'a + t'b = s'(k_1g) + t'(k_2g) = (s'k_1 + t'k_2)g.$$

Mixing a and b in different combinations, we can only make multiples of g .

Note: The combinations are not unique: $sa + tb = (s - b)a + (t + a)b$.

How to make four

$$(2, -1, 1) = \text{gcdcombo}(3, 5).$$

How to make four

$$(2, -1, 1) = \text{gcdcombo}(3, 5).$$

$$\text{So, } 2 \cdot 3 - 1 \cdot 5 = 1.$$

How to make four

$$(2, -1, 1) = \text{gcdcombo}(3, 5).$$

$$\text{So, } 2 \cdot 3 - 1 \cdot 5 = 1.$$

$$\text{To make 4: } 8 \cdot 3 - 4 \cdot 5 = 4.$$

How to make four

$$(2, -1, 1) = \text{gcdcombo}(3, 5).$$

$$\text{So, } 2 \cdot 3 - 1 \cdot 5 = 1.$$

$$\text{To make 4: } 8 \cdot 3 - 4 \cdot 5 = 4. = 3 \cdot 3 - 1 \cdot 5 = 4.$$

How to make four

$$(2, -1, 1) = \text{gcdcombo}(3, 5).$$

$$\text{So, } 2 \cdot 3 - 1 \cdot 5 = 1.$$

$$\text{To make 4: } 8 \cdot 3 - 4 \cdot 5 = 4. = 3 \cdot 3 - 1 \cdot 5 = 4. = -2 \cdot 3 + 2 \cdot 5 = 4.$$

How to make four

$$(2, -1, 1) = \text{gcdcombo}(3, 5).$$

$$\text{So, } 2 \cdot 3 - 1 \cdot 5 = 1.$$

$$\text{To make 4: } 8 \cdot 3 - 4 \cdot 5 = 4. = 3 \cdot 3 - 1 \cdot 5 = 4. = -2 \cdot 3 + 2 \cdot 5 = 4.$$

Fill 2 cups of size 5, then empty out 2 cups of size 3.

Simple recipe

Actually, even simpler. Repeat until target reached:

1. Fill the smaller jug.

Simple recipe

Actually, even simpler. Repeat until target reached:

1. Fill the smaller jug.
2. Pour all the water in the smaller jug into the larger jug.

Simple recipe

Actually, even simpler. Repeat until target reached:

1. Fill the smaller jug.
2. Pour all the water in the smaller jug into the larger jug. If, at any time, the larger jug becomes full, empty it out, and continue.

Simple recipe

Actually, even simpler. Repeat until target reached:

1. Fill the smaller jug.
2. Pour all the water in the smaller jug into the larger jug. If, at any time, the larger jug becomes full, empty it out, and continue.

Eventually reaches all possible values.

Simple recipe

Actually, even simpler. Repeat until target reached:

1. Fill the smaller jug.
2. Pour all the water in the smaller jug into the larger jug. If, at any time, the larger jug becomes full, empty it out, and continue.

Eventually reaches all possible values. Bomb defused!

Fundamental Theorem of Arithmetic

Theorem: Every integer greater than 1 is a product of a unique non-increasing sequence of primes.

Lemma: If p is a prime and $p|ab$, then $p|a$ or $p|b$.

Fundamental Theorem of Arithmetic

Theorem: Every integer greater than 1 is a product of a unique non-increasing sequence of primes.

Lemma: If p is a prime and $p|ab$, then $p|a$ or $p|b$.

Proof of Lemma: One case is if $\gcd(a, p) = p$. Then, the claim holds, because a is a multiple of p .

Fundamental Theorem of Arithmetic

Theorem: Every integer greater than 1 is a product of a unique non-increasing sequence of primes.

Lemma: If p is a prime and $p|ab$, then $p|a$ or $p|b$.

Proof of Lemma: One case is if $\gcd(a, p) = p$. Then, the claim holds, because a is a multiple of p .

Otherwise, $\gcd(a, p) \neq p$.

Fundamental Theorem of Arithmetic

Theorem: Every integer greater than 1 is a product of a unique non-increasing sequence of primes.

Lemma: If p is a prime and $p|ab$, then $p|a$ or $p|b$.

Proof of Lemma: One case is if $\gcd(a, p) = p$. Then, the claim holds, because a is a multiple of p .

Otherwise, $\gcd(a, p) \neq p$. In this case, $\gcd(a, p)$ must be 1, since 1 and p are the only positive divisors of p .

Fundamental Theorem of Arithmetic

Theorem: Every integer greater than 1 is a product of a unique non-increasing sequence of primes.

Lemma: If p is a prime and $p|ab$, then $p|a$ or $p|b$.

Proof of Lemma: One case is if $\gcd(a, p) = p$. Then, the claim holds, because a is a multiple of p .

Otherwise, $\gcd(a, p) \neq p$. In this case, $\gcd(a, p)$ must be 1, since 1 and p are the only positive divisors of p . Since $\gcd(a, p)$ is a linear combination of a and p , we have $1 = sa + tp$ for some s, t .

Fundamental Theorem of Arithmetic

Theorem: Every integer greater than 1 is a product of a unique non-increasing sequence of primes.

Lemma: If p is a prime and $p|ab$, then $p|a$ or $p|b$.

Proof of Lemma: One case is if $\gcd(a, p) = p$. Then, the claim holds, because a is a multiple of p .

Otherwise, $\gcd(a, p) \neq p$. In this case, $\gcd(a, p)$ must be 1, since 1 and p are the only positive divisors of p . Since $\gcd(a, p)$ is a linear combination of a and p , we have $1 = sa + tp$ for some s, t . Then, $b = s(ab) + (tb)p$; that is, b is a linear combination of ab and p .

Fundamental Theorem of Arithmetic

Theorem: Every integer greater than 1 is a product of a unique non-increasing sequence of primes.

Lemma: If p is a prime and $p|ab$, then $p|a$ or $p|b$.

Proof of Lemma: One case is if $\gcd(a, p) = p$. Then, the claim holds, because a is a multiple of p .

Otherwise, $\gcd(a, p) \neq p$. In this case, $\gcd(a, p)$ must be 1, since 1 and p are the only positive divisors of p . Since $\gcd(a, p)$ is a linear combination of a and p , we have $1 = sa + tp$ for some s, t . Then, $b = s(ab) + (tb)p$; that is, b is a linear combination of ab and p . Since p divides both ab and p , it also divides their linear combination, b .

Fundamental Theorem of Arithmetic

Theorem: Every integer greater than 1 is a product of a unique non-increasing sequence of primes.

Lemma: If p is a prime and $p|ab$, then $p|a$ or $p|b$.

Proof of Lemma: One case is if $\gcd(a, p) = p$. Then, the claim holds, because a is a multiple of p .

Otherwise, $\gcd(a, p) \neq p$. In this case, $\gcd(a, p)$ must be 1, since 1 and p are the only positive divisors of p . Since $\gcd(a, p)$ is a linear combination of a and p , we have $1 = sa + tp$ for some s, t . Then, $b = s(ab) + (tb)p$; that is, b is a linear combination of ab and p . Since p divides both ab and p , it also divides their linear combination, b . QED.

Proof of Fundamental Theorem of Arithmetic

Lemma: Let p be a prime. If $p \mid a_1 a_2 \cdots a_n$, then p divides some a_j .

Proof of Fundamental Theorem of Arithmetic

Lemma: Let p be a prime. If $p \mid a_1 a_2 \cdots a_n$, then p divides some a_i .

Proof: We've seen that every positive integer can be expressed as a product of primes.

Proof of Fundamental Theorem of Arithmetic

Lemma: Let p be a prime. If $p \mid a_1 a_2 \cdots a_n$, then p divides some a_i .

Proof: We've seen that every positive integer can be expressed as a product of primes. We need to show this expression is unique.

Proof of Fundamental Theorem of Arithmetic

Lemma: Let p be a prime. If $p|a_1a_2\cdots a_n$, then p divides some a_j .

Proof: We've seen that every positive integer can be expressed as a product of primes. We need to show this expression is unique. We proceed by contradiction:

Proof of Fundamental Theorem of Arithmetic

Lemma: Let p be a prime. If $p|a_1a_2\cdots a_n$, then p divides some a_j .

Proof: We've seen that every positive integer can be expressed as a product of primes. We need to show this expression is unique. We proceed by contradiction: Assume there exists positive integers that can be written as products of primes in more than one way.

Proof of Fundamental Theorem of Arithmetic

Lemma: Let p be a prime. If $p|a_1a_2\cdots a_n$, then p divides some a_j .

Proof: We've seen that every positive integer can be expressed as a product of primes. We need to show this expression is unique. We proceed by contradiction: Assume there exists positive integers that can be written as products of primes in more than one way. Take the smallest such integer n and let $n = p_1p_2\cdots p_j = q_1q_2\cdots q_k$ be the two decompositions.

Proof of Fundamental Theorem of Arithmetic

Lemma: Let p be a prime. If $p|a_1a_2\cdots a_n$, then p divides some a_i .

Proof: We've seen that every positive integer can be expressed as a product of primes. We need to show this expression is unique. We proceed by contradiction: Assume there exists positive integers that can be written as products of primes in more than one way. Take the smallest such integer n and let $n = p_1p_2\cdots p_j = q_1q_2\cdots q_k$ be the two decompositions. Arrange them in non-increasing order and assume without loss of generality that $p_1 \leq q_1$.

Proof of Fundamental Theorem of Arithmetic

Lemma: Let p be a prime. If $p|a_1a_2\cdots a_n$, then p divides some a_j .

Proof: We've seen that every positive integer can be expressed as a product of primes. We need to show this expression is unique. We proceed by contradiction: Assume there exists positive integers that can be written as products of primes in more than one way. Take the smallest such integer n and let $n = p_1p_2\cdots p_j = q_1q_2\cdots q_k$ be the two decompositions. Arrange them in non-increasing order and assume without loss of generality that $p_1 \leq q_1$. If $p_1 = q_1$, the remaining part of the product is smaller than n and different, which is a contradiction (n was the smallest).

Proof of Fundamental Theorem of Arithmetic

Lemma: Let p be a prime. If $p|a_1a_2\cdots a_n$, then p divides some a_i .

Proof: We've seen that every positive integer can be expressed as a product of primes. We need to show this expression is unique. We proceed by contradiction: Assume there exists positive integers that can be written as products of primes in more than one way.

Take the smallest such integer n and let

$n = p_1p_2\cdots p_j = q_1q_2\cdots q_k$ be the two decompositions. Arrange them in non-increasing order and assume without loss of generality that $p_1 \leq q_1$. If $p_1 = q_1$, the remaining part of the product is smaller than n and different, which is a contradiction (n was the smallest).

Note that all the p_i s are less than q_1 . But $q_1|n$ and $n = p_1p_2\cdots p_j$, so q_1 divides one of the p_i s, which contradicts the fact that q_1 is bigger than all them.

Proof of Fundamental Theorem of Arithmetic

Lemma: Let p be a prime. If $p|a_1a_2\cdots a_n$, then p divides some a_i .

Proof: We've seen that every positive integer can be expressed as a product of primes. We need to show this expression is unique. We proceed by contradiction: Assume there exists positive integers that can be written as products of primes in more than one way.

Take the smallest such integer n and let

$n = p_1p_2\cdots p_j = q_1q_2\cdots q_k$ be the two decompositions. Arrange them in non-increasing order and assume without loss of generality that $p_1 \leq q_1$. If $p_1 = q_1$, the remaining part of the product is smaller than n and different, which is a contradiction (n was the smallest).

Note that all the p_i s are less than q_1 . But $q_1|n$ and $n = p_1p_2\cdots p_j$, so q_1 divides one of the p_i s, which contradicts the fact that q_1 is bigger than all them. QED.

Congruence definition

Definition: a is *congruent to* b modulo n iff $\text{rem}(b, n) = \text{rem}(a, n)$.

Congruence definition

Definition: a is *congruent to* b modulo n iff $\text{rem}(b, n) = \text{rem}(a, n)$.
Equivalently, $n \mid (a - b)$.

Congruence definition

Definition: a is *congruent to* b modulo n iff $\text{rem}(b, n) = \text{rem}(a, n)$.

Equivalently, $n \mid (a - b)$.

We write $a \equiv b \pmod{n}$.

Congruence definition

Definition: a is *congruent to* b modulo n iff $\text{rem}(b, n) = \text{rem}(a, n)$.

Equivalently, $n \mid (a - b)$.

We write $a \equiv b \pmod{n}$.

$29 \equiv 15 \pmod{7}$ because

Congruence definition

Definition: a is *congruent to* b modulo n iff $\text{rem}(b, n) = \text{rem}(a, n)$.

Equivalently, $n \mid (a - b)$.

We write $a \equiv b \pmod{n}$.

$29 \equiv 15 \pmod{7}$ because $7 \mid (29 - 15)$, namely 14.

Congruence definition

Definition: a is *congruent to* b modulo n iff $\text{rem}(b, n) = \text{rem}(a, n)$.
Equivalently, $n \mid (a - b)$.

We write $a \equiv b \pmod{n}$.

$29 \equiv 15 \pmod{7}$ because $7 \mid (29 - 15)$, namely 14. Both have a remainder of 1 when divided by 7.

Congruence definition

Definition: a is *congruent to* b modulo n iff $\text{rem}(b, n) = \text{rem}(a, n)$.
Equivalently, $n \mid (a - b)$.

We write $a \equiv b \pmod{n}$.

$29 \equiv 15 \pmod{7}$ because $7 \mid (29 - 15)$, namely 14. Both have a remainder of 1 when divided by 7.

Equivalence relation—partitions the integers.

Congruence definition

Definition: a is *congruent to* b modulo n iff $\text{rem}(b, n) = \text{rem}(a, n)$.

Equivalently, $n \mid (a - b)$.

We write $a \equiv b \pmod{n}$.

$29 \equiv 15 \pmod{7}$ because $7 \mid (29 - 15)$, namely 14. Both have a remainder of 1 when divided by 7.

Equivalence relation—partitions the integers.

Transitivity, reflexivity, symmetry.

Basic modular algebra

In regular algebra,

$$a = b$$

Basic modular algebra

In regular algebra,

$$a = b$$

$$a + c = b + c.$$

Basic modular algebra

In regular algebra,

$$a = b$$

$$a + c = b + c.$$

Can we do the same in congruence-land?

$$a \equiv b \pmod{n}$$

Basic modular algebra

In regular algebra,

$$a = b$$

$$a + c = b + c.$$

Can we do the same in congruence-land?

$$a \equiv b \pmod{n}$$

$$a + c \equiv b + c \pmod{n}.$$

Basic modular algebra

In regular algebra,

$$a = b$$

$$a + c = b + c.$$

Can we do the same in congruence-land?

$$a \equiv b \pmod{n}$$

$$a + c \equiv b + c \pmod{n}.$$

Yes!

$$a \equiv b \pmod{n} \text{ iff}$$

Basic modular algebra

In regular algebra,

$$a = b$$

$$a + c = b + c.$$

Can we do the same in congruence-land?

$$a \equiv b \pmod{n}$$

$$a + c \equiv b + c \pmod{n}.$$

Yes!

$$a \equiv b \pmod{n} \text{ iff } n \mid (a - b) \text{ iff}$$

Basic modular algebra

In regular algebra,

$$a = b$$

$$a + c = b + c.$$

Can we do the same in congruence-land?

$$a \equiv b \pmod{n}$$

$$a + c \equiv b + c \pmod{n}.$$

Yes!

$$a \equiv b \pmod{n} \text{ iff } n \mid (a - b) \text{ iff } \exists k, kn = a - b \text{ iff}$$

Basic modular algebra

In regular algebra,

$$a = b$$

$$a + c = b + c.$$

Can we do the same in congruence-land?

$$a \equiv b \pmod{n}$$

$$a + c \equiv b + c \pmod{n}.$$

Yes!

$$a \equiv b \pmod{n} \text{ iff } n \mid (a - b) \text{ iff } \exists k, kn = a - b \text{ iff}$$

$$\exists k, kn = a - b + (c - c) \text{ iff}$$

Basic modular algebra

In regular algebra,

$$a = b$$

$$a + c = b + c.$$

Can we do the same in congruence-land?

$$a \equiv b \pmod{n}$$

$$a + c \equiv b + c \pmod{n}.$$

Yes!

$$a \equiv b \pmod{n} \text{ iff } n \mid (a - b) \text{ iff } \exists k, kn = a - b \text{ iff}$$

$$\exists k, kn = a - b + (c - c) \text{ iff } \exists k, kn = (a + c) - (b + c) \text{ iff}$$

Basic modular algebra

In regular algebra,

$$a = b$$

$$a + c = b + c.$$

Can we do the same in congruence-land?

$$a \equiv b \pmod{n}$$

$$a + c \equiv b + c \pmod{n}.$$

Yes!

$$a \equiv b \pmod{n} \text{ iff } n \mid (a - b) \text{ iff } \exists k, kn = a - b \text{ iff}$$

$$\exists k, kn = a - b + (c - c) \text{ iff } \exists k, kn = (a + c) - (b + c) \text{ iff}$$

$$n \mid ((a + c) - (b + c)) \text{ iff } a + c \equiv b + c \pmod{n}.$$

Multiplication is repeated addition, so we can also multiply both sides by a constant.

Basic modular algebra

In regular algebra,

$$a = b$$

$$a + c = b + c.$$

Can we do the same in congruence-land?

$$a \equiv b \pmod{n}$$

$$a + c \equiv b + c \pmod{n}.$$

Yes!

$$a \equiv b \pmod{n} \text{ iff } n \mid (a - b) \text{ iff } \exists k, kn = a - b \text{ iff}$$

$$\exists k, kn = a - b + (c - c) \text{ iff } \exists k, kn = (a + c) - (b + c) \text{ iff}$$

$$n \mid ((a + c) - (b + c)) \text{ iff } a + c \equiv b + c \pmod{n}.$$

Multiplication is repeated addition, so we can also multiply both sides by a constant. By transitivity, we can always add or multiply each side by values that are congruent!

Basic modular algebra

In regular algebra,

$$a = b$$

$$a + c = b + c.$$

Can we do the same in congruence-land?

$$a \equiv b \pmod{n}$$

$$a + c \equiv b + c \pmod{n}.$$

Yes!

$$a \equiv b \pmod{n} \text{ iff } n \mid (a - b) \text{ iff } \exists k, kn = a - b \text{ iff}$$

$$\exists k, kn = a - b + (c - c) \text{ iff } \exists k, kn = (a + c) - (b + c) \text{ iff}$$

$$n \mid ((a + c) - (b + c)) \text{ iff } a + c \equiv b + c \pmod{n}.$$

Multiplication is repeated addition, so we can also multiply both sides by a constant. By transitivity, we can always add or multiply each side by values that are congruent! “Clock arithmetic”.

Example

$$2x + 17 \equiv x + 31 \pmod{12}$$

Example

$$2x + 17 \equiv x + 31 \pmod{12}$$

$$2x \equiv x + 14 \pmod{12} \quad \text{add } -17 \text{ to both sides}$$

Example

$$2x + 17 \equiv x + 31 \pmod{12}$$

$$2x \equiv x + 14 \pmod{12} \quad \text{add } -17 \text{ to both sides}$$

$$2x \equiv x + 2 \pmod{12} \quad \text{add } 0 \text{ to left and } -12 \text{ to right}$$

Example

$$2x + 17 \equiv x + 31 \pmod{12}$$

$$2x \equiv x + 14 \pmod{12} \quad \text{add } -17 \text{ to both sides}$$

$$2x \equiv x + 2 \pmod{12} \quad \text{add } 0 \text{ to left and } -12 \text{ to right}$$

$$x \equiv 2 \pmod{12} \quad \text{add } -x \text{ to both sides}$$

Example

$$2x + 17 \equiv x + 31 \pmod{12}$$

$$2x \equiv x + 14 \pmod{12} \quad \text{add } -17 \text{ to both sides}$$

$$2x \equiv x + 2 \pmod{12} \quad \text{add } 0 \text{ to left and } -12 \text{ to right}$$

$$x \equiv 2 \pmod{12} \quad \text{add } -x \text{ to both sides}$$

Double check. $4 + 17 = 21$ vs. 33 . Difference is 12, check!

Example

$$2x + 17 \equiv x + 31 \pmod{12}$$

$$2x \equiv x + 14 \pmod{12} \quad \text{add } -17 \text{ to both sides}$$

$$2x \equiv x + 2 \pmod{12} \quad \text{add } 0 \text{ to left and } -12 \text{ to right}$$

$$x \equiv 2 \pmod{12} \quad \text{add } -x \text{ to both sides}$$

Double check. $4 + 17 = 21$ vs. 33 . Difference is 12, check!

$$3x + 4 \equiv 27 \pmod{11}$$

Example

$$2x + 17 \equiv x + 31 \pmod{12}$$

$$2x \equiv x + 14 \pmod{12} \quad \text{add } -17 \text{ to both sides}$$

$$2x \equiv x + 2 \pmod{12} \quad \text{add } 0 \text{ to left and } -12 \text{ to right}$$

$$x \equiv 2 \pmod{12} \quad \text{add } -x \text{ to both sides}$$

Double check. $4 + 17 = 21$ vs. 33 . Difference is 12, check!

$$3x + 4 \equiv 27 \pmod{11}$$

$$3x \equiv 23 \pmod{11} \quad \text{add } -4 \text{ to both sides}$$

Example

$$2x + 17 \equiv x + 31 \pmod{12}$$

$$2x \equiv x + 14 \pmod{12} \quad \text{add } -17 \text{ to both sides}$$

$$2x \equiv x + 2 \pmod{12} \quad \text{add } 0 \text{ to left and } -12 \text{ to right}$$

$$x \equiv 2 \pmod{12} \quad \text{add } -x \text{ to both sides}$$

Double check. $4 + 17 = 21$ vs. 33 . Difference is 12, check!

$$3x + 4 \equiv 27 \pmod{11}$$

$$3x \equiv 23 \pmod{11} \quad \text{add } -4 \text{ to both sides}$$

Kind of stuck because we don't (yet) have a "divide both sides by 3" rule.

So, what about division?

If $a \equiv b \pmod{n}$, can we divide both sides by c ?

So, what about division?

If $a \equiv b \pmod{n}$, can we divide both sides by c ?

$$7 \equiv 28 \pmod{3}$$

So, what about division?

If $a \equiv b \pmod{n}$, can we divide both sides by c ?

$$7 \equiv 28 \pmod{3}$$

$$1 \equiv 4 \pmod{3} \quad \text{divide by 7}$$

So, what about division?

If $a \equiv b \pmod{n}$, can we divide both sides by c ?

$$7 \equiv 28 \pmod{3}$$

$$1 \equiv 4 \pmod{3} \quad \text{divide by 7}$$

So, maybe? At least if the answers are integers?

So, what about division?

If $a \equiv b \pmod{n}$, can we divide both sides by c ?

$$7 \equiv 28 \pmod{3}$$

$$1 \equiv 4 \pmod{3} \quad \text{divide by 7}$$

So, maybe? At least if the answers are integers?

Is division even meaningful more generally?

Back to basics

Definition: The *multiplicative inverse* of a number x is a number x^{-1} such that: $x \cdot x^{-1} = 1$.

Back to basics

Definition: The *multiplicative inverse* of a number x is a number x^{-1} such that: $x \cdot x^{-1} = 1$.

Division by x is really multiplication by x^{-1} .

Back to basics

Definition: The *multiplicative inverse* of a number x is a number x^{-1} such that: $x \cdot x^{-1} = 1$.

Division by x is really multiplication by x^{-1} .

Over the reals, what values have inverses?

Back to basics

Definition: The *multiplicative inverse* of a number x is a number x^{-1} such that: $x \cdot x^{-1} = 1$.

Division by x is really multiplication by x^{-1} .

Over the reals, what values have inverses? Everybody but zero.

Back to basics

Definition: The *multiplicative inverse* of a number x is a number x^{-1} such that: $x \cdot x^{-1} = 1$.

Division by x is really multiplication by x^{-1} .

Over the reals, what values have inverses? Everybody but zero.

Over the integers, what values have inverses?

Back to basics

Definition: The *multiplicative inverse* of a number x is a number x^{-1} such that: $x \cdot x^{-1} = 1$.

Division by x is really multiplication by x^{-1} .

Over the reals, what values have inverses? Everybody but zero.

Over the integers, what values have inverses? Only 1 and -1 .

Back to basics

Definition: The *multiplicative inverse* of a number x is a number x^{-1} such that: $x \cdot x^{-1} = 1$.

Division by x is really multiplication by x^{-1} .

Over the reals, what values have inverses? Everybody but zero.

Over the integers, what values have inverses? Only 1 and -1 .

Over the integers mod n , what values have inverses?

Example, mod 10

	0	1	2	3	4	5	6	7	8	9
0	0	0	0	0	0	0	0	0	0	0
1	0	1	2	3	4	5	6	7	8	9
2	0	2	4	6	8	0	2	4	6	8
3	0	3	6	9	2	5	8	1	4	7
4	0	4	8	2	6	0	4	8	2	6
5	0	5	0	5	0	5	0	5	0	5
6	0	6	2	8	4	0	6	2	8	4
7	0	7	4	1	8	5	2	9	6	3
8	0	8	6	4	2	0	8	6	4	2
9	0	9	8	7	6	5	4	3	2	1

Example, mod 10

	0	1	2	3	4	5	6	7	8	9
0	0	0	0	0	0	0	0	0	0	0
1	0	1	2	3	4	5	6	7	8	9
2	0	2	4	6	8	0	2	4	6	8
3	0	3	6	9	2	5	8	1	4	7
4	0	4	8	2	6	0	4	8	2	6
5	0	5	0	5	0	5	0	5	0	5
6	0	6	2	8	4	0	6	2	8	4
7	0	7	4	1	8	5	2	9	6	3
8	0	8	6	4	2	0	8	6	4	2
9	0	9	8	7	6	5	4	3	2	1

What specific values have inverses?

Example, mod 10

	0	1	2	3	4	5	6	7	8	9
0	0	0	0	0	0	0	0	0	0	0
1	0	1	2	3	4	5	6	7	8	9
2	0	2	4	6	8	0	2	4	6	8
3	0	3	6	9	2	5	8	1	4	7
4	0	4	8	2	6	0	4	8	2	6
5	0	5	0	5	0	5	0	5	0	5
6	0	6	2	8	4	0	6	2	8	4
7	0	7	4	1	8	5	2	9	6	3
8	0	8	6	4	2	0	8	6	4	2
9	0	9	8	7	6	5	4	3	2	1

What specific values have inverses? 1, 3, 7, 9.

Example, mod 10

	0	1	2	3	4	5	6	7	8	9
0	0	0	0	0	0	0	0	0	0	0
1	0	1	2	3	4	5	6	7	8	9
2	0	2	4	6	8	0	2	4	6	8
3	0	3	6	9	2	5	8	1	4	7
4	0	4	8	2	6	0	4	8	2	6
5	0	5	0	5	0	5	0	5	0	5
6	0	6	2	8	4	0	6	2	8	4
7	0	7	4	1	8	5	2	9	6	3
8	0	8	6	4	2	0	8	6	4	2
9	0	9	8	7	6	5	4	3	2	1

What specific values have inverses? 1, 3, 7, 9.

What specific values do *not* have inverses?

Example, mod 10

	0	1	2	3	4	5	6	7	8	9
0	0	0	0	0	0	0	0	0	0	0
1	0	1	2	3	4	5	6	7	8	9
2	0	2	4	6	8	0	2	4	6	8
3	0	3	6	9	2	5	8	1	4	7
4	0	4	8	2	6	0	4	8	2	6
5	0	5	0	5	0	5	0	5	0	5
6	0	6	2	8	4	0	6	2	8	4
7	0	7	4	1	8	5	2	9	6	3
8	0	8	6	4	2	0	8	6	4	2
9	0	9	8	7	6	5	4	3	2	1

What specific values have inverses? 1, 3, 7, 9.

What specific values do *not* have inverses? 0, 2, 4, 5, 6, 8.

Example, mod 10

	0	1	2	3	4	5	6	7	8	9
0	0	0	0	0	0	0	0	0	0	0
1	0	1	2	3	4	5	6	7	8	9
2	0	2	4	6	8	0	2	4	6	8
3	0	3	6	9	2	5	8	1	4	7
4	0	4	8	2	6	0	4	8	2	6
5	0	5	0	5	0	5	0	5	0	5
6	0	6	2	8	4	0	6	2	8	4
7	0	7	4	1	8	5	2	9	6	3
8	0	8	6	4	2	0	8	6	4	2
9	0	9	8	7	6	5	4	3	2	1

What specific values have inverses? 1, 3, 7, 9.

What specific values do *not* have inverses? 0, 2, 4, 5, 6, 8.

General rule?

Example, mod 10

	0	1	2	3	4	5	6	7	8	9
0	0	0	0	0	0	0	0	0	0	0
1	0	1	2	3	4	5	6	7	8	9
2	0	2	4	6	8	0	2	4	6	8
3	0	3	6	9	2	5	8	1	4	7
4	0	4	8	2	6	0	4	8	2	6
5	0	5	0	5	0	5	0	5	0	5
6	0	6	2	8	4	0	6	2	8	4
7	0	7	4	1	8	5	2	9	6	3
8	0	8	6	4	2	0	8	6	4	2
9	0	9	8	7	6	5	4	3	2	1

What specific values have inverses? 1, 3, 7, 9.

What specific values do *not* have inverses? 0, 2, 4, 5, 6, 8.

General rule? a has an inverse (mod n) iff $\gcd(a, n) = 1$.

Back to solving

$$3x + 4 \equiv 27 \pmod{11}$$

Back to solving

$$3x + 4 \equiv 27 \pmod{11}$$

$$3x \equiv 23 \pmod{11} \quad \text{add } -4 \text{ to both sides}$$

Back to solving

$$3x + 4 \equiv 27 \pmod{11}$$

$$3x \equiv 23 \pmod{11} \quad \text{add } -4 \text{ to both sides}$$

Want to multiply both sides by $3^{-1} =$

Back to solving

$$3x + 4 \equiv 27 \pmod{11}$$

$$3x \equiv 23 \pmod{11} \quad \text{add } -4 \text{ to both sides}$$

Want to multiply both sides by $3^{-1} = 4$, since $3 \times 4 \equiv 1 \pmod{11}$.

Back to solving

$$3x + 4 \equiv 27 \pmod{11}$$

$$3x \equiv 23 \pmod{11} \quad \text{add } -4 \text{ to both sides}$$

Want to multiply both sides by $3^{-1} = 4$, since $3 \times 4 \equiv 1 \pmod{11}$.

$$3x \equiv 23 \pmod{11}$$

Back to solving

$$3x + 4 \equiv 27 \pmod{11}$$

$$3x \equiv 23 \pmod{11} \quad \text{add } -4 \text{ to both sides}$$

Want to multiply both sides by $3^{-1} = 4$, since $3 \times 4 \equiv 1 \pmod{11}$.

$$3x \equiv 23 \pmod{11}$$

$$4 \times 3x \equiv 4 \times 23 \pmod{11} \quad \text{multiply both sides by 4}$$

Back to solving

$$3x + 4 \equiv 27 \pmod{11}$$

$$3x \equiv 23 \pmod{11} \quad \text{add } -4 \text{ to both sides}$$

Want to multiply both sides by $3^{-1} = 4$, since $3 \times 4 \equiv 1 \pmod{11}$.

$$3x \equiv 23 \pmod{11}$$

$$4 \times 3x \equiv 4 \times 23 \pmod{11} \quad \text{multiply both sides by 4}$$

$$12x \equiv 92 \pmod{11} \quad \text{simplify}$$

Back to solving

$$3x + 4 \equiv 27 \pmod{11}$$

$$3x \equiv 23 \pmod{11} \quad \text{add } -4 \text{ to both sides}$$

Want to multiply both sides by $3^{-1} = 4$, since $3 \times 4 \equiv 1 \pmod{11}$.

$$3x \equiv 23 \pmod{11}$$

$$4 \times 3x \equiv 4 \times 23 \pmod{11} \quad \text{multiply both sides by 4}$$

$$12x \equiv 92 \pmod{11} \quad \text{simplify}$$

$$x \equiv 4 \pmod{11} \quad \text{congruence}$$

Back to solving

$$3x + 4 \equiv 27 \pmod{11}$$

$$3x \equiv 23 \pmod{11} \quad \text{add } -4 \text{ to both sides}$$

Want to multiply both sides by $3^{-1} = 4$, since $3 \times 4 \equiv 1 \pmod{11}$.

$$3x \equiv 23 \pmod{11}$$

$$4 \times 3x \equiv 4 \times 23 \pmod{11} \quad \text{multiply both sides by 4}$$

$$12x \equiv 92 \pmod{11} \quad \text{simplify}$$

$$x \equiv 4 \pmod{11} \quad \text{congruence}$$

Double check: 16 vs. 5, 11 divides the difference!