

CSCI 5451 Fall 2015

## Week 1 Notes

Professor Ellen Gethner

August 19, 2015

## Introduction

Overview

## Algorithm Definition?

## Tools

Goals

Particulars

## This Week's Topics

Outline of Topics

Principle of Mathematical Induction

# Graduate Algorithms: Preliminaries

- ▶ **Prerequisites courses.** While there is no official prerequisite course, you are expected to have command of Discrete Structures and Undergraduate Algorithms.

# Graduate Algorithms: Preliminaries

- ▶ **Prerequisites courses.** While there is no official prerequisite course, you are expected to have command of Discrete Structures and Undergraduate Algorithms.
- ▶ **Assignments.** There will be six homework assignments, none of which are handed in.

# Graduate Algorithms: Preliminaries

- ▶ **Prerequisites courses.** While there is no official prerequisite course, you are expected to have command of Discrete Structures and Undergraduate Algorithms.
- ▶ **Assignments.** There will be six homework assignments, none of which are handed in.
- ▶ **Quizzes.** Your mastery of each assignment will be assessed by way of a closed-book closed-notes quiz.

# Graduate Algorithms: Quizzes, continued

- ▶ See the online syllabus for the dates of each quiz.

# Graduate Algorithms: Quizzes, continued

- ▶ See the online syllabus for the dates of each quiz.
- ▶ There are no make-up quizzes, although your lowest score will be dropped.

# Graduate Algorithms: Quizzes, continued

- ▶ See the online syllabus for the dates of each quiz.
- ▶ There are no make-up quizzes, although your lowest score will be dropped.
- ▶ Quizzes are given only once in class on the date specified on the syllabus.



# Graduate Algorithms: Quizzes, continued

- ▶ See the online syllabus for the dates of each quiz.
- ▶ There are no make-up quizzes, although your lowest score will be dropped.
- ▶ Quizzes are given only once in class on the date specified on the syllabus.
- ▶ Quizzes count toward 30% of your final grade.

# Graduate Algorithms: Examinations

- ▶ Examination 1 is given once only (in class) on Tuesday, 13 October.

# Graduate Algorithms: Examinations

- ▶ Examination 1 is given once only (in class) on Tuesday, 13 October.
- ▶ Exam 1 counts toward 35% of your final grade.

# Graduate Algorithms: Examinations

- ▶ Examination 1 is given once only (in class) on Tuesday, 13 October.
- ▶ Exam 1 counts toward 35% of your final grade.
- ▶ Examination 2 is given once only (in class) on Tuesday, 1 December.

# Graduate Algorithms: Examinations

- ▶ Examination 1 is given once only (in class) on Tuesday, 13 October.
- ▶ Exam 1 counts toward 35% of your final grade.
- ▶ Examination 2 is given once only (in class) on Tuesday, 1 December.
- ▶ Exam 2 counts toward 35% of your final grade.

# Topics

- ▶ **Design and Analysis of Algorithms**, including but not limited to *Divide and Conquer*, *Greedy*, *Dynamic*, and many more.

# Topics

- ▶ **Design and Analysis of Algorithms**, including but not limited to *Divide and Conquer*, *Greedy*, *Dynamic*, and many more.
- ▶ **Basic Algorithms** including but not limited to *Sequences and sets*, *graph algorithms*, *algebraic algorithms* and many more.

# Topics

- ▶ **Design and Analysis of Algorithms**, including but not limited to *Divide and Conquer*, *Greedy*, *Dynamic*, and many more.
- ▶ **Basic Algorithms** including but not limited to *Sequences and sets*, *graph algorithms*, *algebraic algorithms* and many more.
- ▶ **Advanced Topics** including but not limited to *Approximation*, *NP-Completeness*, *Cryptography*, *Computational Geometry* and more.



# Some Synonyms for “Algorithm”

- ▶ Recipe, process, method, technique, procedure, routine, . . .

# Some Synonyms for “Algorithm”

- ▶ Recipe, process, method, technique, procedure, routine, . . .
- ▶ Which do you think best describes what an algorithm is supposed to be?

# From Donald Knuth's **The Art of Computer Programming**

Five items whose sum total describes all of the necessary attributes that an algorithm must have:

- ▶ **A1: Finiteness.** An algorithm must terminate after a finite number of steps.

# From Donald Knuth's **The Art of Computer Programming**

Five items whose sum total describes all of the necessary attributes that an algorithm must have:

- ▶ **A1: Finiteness.** An algorithm must terminate after a finite number of steps.
- ▶ **A2: Definiteness.** Each step must be precisely defined; the actions to be carried out must be rigorously and unambiguously defined.

# From Donald Knuth's **The Art of Computer Programming**

Five items whose sum total describes all of the necessary attributes that an algorithm must have:

- ▶ **A1: Finiteness.** An algorithm must terminate after a finite number of steps.
- ▶ **A2: Definiteness.** Each step must be precisely defined; the actions to be carried out must be rigorously and unambiguously defined.
- ▶ **A3: Input.** An algorithm has zero or more inputs that can be given before the algorithm runs and/or during its execution.

# From Donald Knuth's **The Art of Computer Programming**

Five items whose sum total describes all of the necessary attributes that an algorithm must have:

- ▶ **A1: Finiteness.** An algorithm must terminate after a finite number of steps.
- ▶ **A2: Definiteness.** Each step must be precisely defined; the actions to be carried out must be rigorously and unambiguously defined.
- ▶ **A3: Input.** An algorithm has zero or more inputs that can be given before the algorithm runs and/or during its execution.

# From Donald Knuth's **The Art of Computer Programming**

Five items whose sum total describes all of the necessary attributes that an algorithm should have:

- ▶ **A4: Output.** An algorithm has one or more outputs (that presumably have some relation to the input).

# From Donald Knuth's **The Art of Computer Programming**

Five items whose sum total describes all of the necessary attributes that an algorithm should have:

- ▶ **A4: Output.** An algorithm has one or more outputs (that presumably have some relation to the input).
- ▶ **A5: Effectiveness.** Loosely speaking, an algorithm is effective if its operations are all basic enough that they can be done in a finite amount of time by someone using pencil and paper.



# Goals

We will learn the following algorithm fundamentals:

- ▶ design

# Goals

We will learn the following algorithm fundamentals:

- ▶ design
- ▶ proof techniques (correctness)

# Goals

We will learn the following algorithm fundamentals:

- ▶ design
- ▶ proof techniques (correctness)
- ▶ efficiency:
  1. How fast does your algorithm run, say, with respect to the size of the input? Can you do better?
  2. Alternatively, can you prove that you **cannot** do better?

- ▶ **Design.** There are no rules: creativity is your best tool together with knowledge of basic algorithms upon which to build.

- ▶ **Design.** There are no rules: creativity is your best tool together with knowledge of basic algorithms upon which to build.
- ▶ **Proof Techniques.** Use of mathematics and common sense.

- ▶ **Design.** There are no rules: creativity is your best tool together with knowledge of basic algorithms upon which to build.
- ▶ **Proof Techniques.** Use of mathematics and common sense.
- ▶ **Analysis of Efficiency.** Use of mathematics and common sense.

# Goals for this week

- ▶ Mathematical Induction (a lengthy review from your Discrete Structures Class)

# Goals for this week

- ▶ Mathematical Induction (a lengthy review from your Discrete Structures Class)
- ▶ Fibonacci numbers



# Goals for this week

- ▶ Mathematical Induction (a lengthy review from your Discrete Structures Class)
- ▶ Fibonacci numbers
- ▶ The oldest algorithm on record.

# Goals for this week

- ▶ Mathematical Induction (a lengthy review from your Discrete Structures Class)
- ▶ Fibonacci numbers
- ▶ The oldest algorithm on record.
- ▶ Connections among the latter two items.

# Induction

- ▶  $P(n)$  is an assertion about the integer  $n$ .

# Induction

- ▶  $P(n)$  is an assertion about the integer  $n$ .
- ▶ For example “if  $n \geq 1$  then  $n^2 > n$ .”

# Induction

- ▶  $P(n)$  is an assertion about the integer  $n$ .
- ▶ For example “if  $n \geq 1$  then  $n^2 > n$ .”
- ▶ Generally speaking, our goal will be to prove that  $P(n)$  is true for all integers  $n \geq N$  for some fixed  $N \in \mathbb{Z}$ .

# Steps in an induction proof

1. **Base Cases.** Verify that  $P(N)$  is true for some fixed  $N \in \mathbb{Z}$ .

# Steps in an induction proof

1. **Base Cases.** Verify that  $P(N)$  is true for some fixed  $N \in \mathbb{Z}$ .
2. **Reality Check.** Check a few more values as well.

# Steps in an induction proof

1. **Base Cases.** Verify that  $P(N)$  is true for some fixed  $N \in \mathbb{Z}$ .
2. **Reality Check.** Check a few more values as well.
3. **Induction Hypothesis.** Assume that  $P(N)$ ,  $P(N+1)$ ,  $\dots$ ,  $P(k)$  are true for some fixed  $k \in \mathbb{Z}$  such that  $k \geq N$ .



# Steps in an induction proof

1. **Base Cases.** Verify that  $P(N)$  is true for some fixed  $N \in \mathbb{Z}$ .
2. **Reality Check.** Check a few more values as well.
3. **Induction Hypothesis.** Assume that  $P(N)$ ,  $P(N+1)$ ,  $\dots$ ,  $P(k)$  are true for some fixed  $k \in \mathbb{Z}$  such that  $k \geq N$ .
4. **Inductive Step.** Prove that  $P(k+1)$  is true.

# Steps in an induction proof

1. **Base Cases.** Verify that  $P(N)$  is true for some fixed  $N \in \mathbb{Z}$ .
2. **Reality Check.** Check a few more values as well.
3. **Induction Hypothesis.** Assume that  $P(N)$ ,  $P(N+1)$ ,  $\dots$ ,  $P(k)$  are true for some fixed  $k \in \mathbb{Z}$  such that  $k \geq N$ .
4. **Inductive Step.** Prove that  $P(k+1)$  is true.
5. **Conclusion.** Conclude that  $P(n)$  is true  $\forall n \geq N$ .

## Example One: sum of consecutive odd integers

- **Problem.** Find a closed-form formula for

$$f(n) = \sum_{i=1}^n (2i - 1)$$

## Example One: sum of consecutive odd integers

- **Problem.** Find a closed-form formula for

$$f(n) = \sum_{i=1}^n (2i - 1)$$

- and prove that your formula is correct.

## Example One: sum of consecutive odd integers

- **Problem.** Find a closed-form formula for

$$f(n) = \sum_{i=1}^n (2i - 1)$$

- and prove that your formula is correct.
- Note that  $f(n) = \sum_{i=1}^n (2i - 1) = 1 + 3 + 5 + \cdots + 2n - 1$ .

## Example One, continued

The first thing we should do is make a guess at a formula for  $f(n)$  so let's experiment.

- ▶  $f(1) = \sum_{i=1}^1 (2i - 1) = 2 \times 1 - 1 = 1.$

## Example One, continued

The first thing we should do is make a guess at a formula for  $f(n)$  so let's experiment.

- ▶  $f(1) = \sum_{i=1}^1 (2i - 1) = 2 \times 1 - 1 = 1.$

- ▶  $f(2) = \sum_{i=1}^2 (2i - 1) = (2 \times 1 - 1) + (2 \times 2 - 1) = 1 + 3 = 4.$

## Example One, continued

The first thing we should do is make a guess at a formula for  $f(n)$  so let's experiment.

- ▶  $f(1) = \sum_{i=1}^1 (2i - 1) = 2 \times 1 - 1 = 1.$

- ▶  $f(2) = \sum_{i=1}^2 (2i - 1) = (2 \times 1 - 1) + (2 \times 2 - 1) = 1 + 3 = 4.$

- ▶  $f(3) = \sum_{i=1}^3 (2i - 1) = 1 + 3 + 5 = 9.$



# Ready for a conjecture

**Conjecture.**  $f(n) = n^2 \quad \forall n \in \mathbb{Z}^+.$

## Proof by induction on $n$

The statement  $P(n)$  says that  $f(n) = n^2$  and we wish to prove that  $P(n)$  is true  $\forall n \in \mathbb{Z}^+$ .

- **Base Cases.** We verified that  $P(1)$ ,  $P(2)$ , and  $P(3)$  were all true on the previous slide.

# Proof by induction on $n$

The statement  $P(n)$  says that  $f(n) = n^2$  and we wish to prove that  $P(n)$  is true  $\forall n \in \mathbb{Z}^+$ .

- ▶ **Base Cases.** We verified that  $P(1)$ ,  $P(2)$ , and  $P(3)$  were all true on the previous slide.
- ▶ **Induction Hypothesis.** Assume that  $P(k)$  is true for some **fixed** integer  $k \geq 3$ .

## Induction proof, continued

- **Inductive Step.** We must prove that  $P(k + 1)$  is true. That is, we must prove that  $f(k + 1) = (k + 1)^2$

## Induction proof, continued

- ▶ **Inductive Step.** We must prove that  $P(k + 1)$  is true. That is, we must prove that  $f(k + 1) = (k + 1)^2$
- ▶ **Here goes.** By definition of the function  $f$ , we have

## Induction proof, continued

- ▶ **Inductive Step.** We must prove that  $P(k + 1)$  is true. That is, we must prove that  $f(k + 1) = (k + 1)^2$
- ▶ **Here goes.** By definition of the function  $f$ , we have
- ▶  $f(k + 1) = \sum_{i=1}^{k+1} (2i - 1)$

## Induction proof, continued

- ▶ **Inductive Step.** We must prove that  $P(k+1)$  is true. That is, we must prove that  $f(k+1) = (k+1)^2$
- ▶ **Here goes.** By definition of the function  $f$ , we have
- ▶  $f(k+1) = \sum_{i=1}^{k+1} (2i-1)$
- ▶  $= \sum_{i=1}^k (2i-1) + 2(k+1) - 1$

## Induction proof, continued

- ▶ **Inductive Step.** We must prove that  $P(k+1)$  is true. That is, we must prove that  $f(k+1) = (k+1)^2$
- ▶ **Here goes.** By definition of the function  $f$ , we have
- ▶  $f(k+1) = \sum_{i=1}^{k+1} (2i-1)$
- ▶  $= \sum_{i=1}^k (2i-1) + 2(k+1) - 1$
- ▶  $= k^2 + 2k + 2 - 1$  by the induction hypothesis and algebra



## Induction proof, continued

- ▶ **Inductive Step.** We must prove that  $P(k+1)$  is true. That is, we must prove that  $f(k+1) = (k+1)^2$
- ▶ **Here goes.** By definition of the function  $f$ , we have
- ▶  $f(k+1) = \sum_{i=1}^{k+1} (2i-1)$
- ▶  $= \sum_{i=1}^k (2i-1) + 2(k+1) - 1$
- ▶  $= k^2 + 2k + 2 - 1$  by the induction hypothesis and algebra
- ▶  $= k^2 + 2k + 1$

## Induction proof, continued

- ▶ **Inductive Step.** We must prove that  $P(k+1)$  is true. That is, we must prove that  $f(k+1) = (k+1)^2$
- ▶ **Here goes.** By definition of the function  $f$ , we have
- ▶  $f(k+1) = \sum_{i=1}^{k+1} (2i-1)$
- ▶  $= \sum_{i=1}^k (2i-1) + 2(k+1) - 1$
- ▶  $= k^2 + 2k + 2 - 1$  by the induction hypothesis and algebra
- ▶  $= k^2 + 2k + 1$
- ▶  $= (k+1)^2$ , as desired.

# Induction proof, completed

**Conclusion.**  $P(n)$  is true  $\forall n \in \mathbb{Z}^+$ . That is,  $f(n) = n^2 \forall n \in \mathbb{Z}^+$ .

QED

## Example Two: Counting Squares

**Problem.** How many squares are there in an  $n \times n$  grid-square?

**Notation.** Let  $f(n)$  denote the number of squares in an  $n \times n$  grid square.

## Accounting for small values of $n$ : $n = 1$




Figure: A  $1 \times 1$  grid square

- Looks like  $f(1) = 1$ .

## Accounting for small values of $n$ : $n = 1$




Figure: A  $1 \times 1$  grid square

- ▶ Looks like  $f(1) = 1$ .
- ▶ Not very, illuminating, though.

## Accounting for small values of $n$ : $n = 1$




Figure: A  $1 \times 1$  grid square

- ▶ Looks like  $f(1) = 1$ .
- ▶ Not very, illuminating, though.
- ▶ So keep experimenting!

## Accounting for small values of $n$ : $n = 2$




Figure: A  $2 \times 2$  grid square



## Accounting for small values of $n$ : $n = 2$




Figure: A  $2 \times 2$  grid square

## Accounting for small values of $n$ : $n = 2$



Figure: A  $2 \times 2$  grid square

## Accounting for small values of $n$ : $n = 2$




Figure: A  $2 \times 2$  grid square

## Accounting for small values of $n$ : $n = 2$




Figure: A  $2 \times 2$  grid square

## Accounting for small values of $n$ : $n = 2$




Figure: A  $2 \times 2$  grid square

## Accounting for small values of $n$ : $n = 2$




Figure: A  $2 \times 2$  grid square

- Looks like  $f(2) = 4 + 1$ .

## Accounting for small values of $n$ : $n = 2$




Figure: A  $2 \times 2$  grid square

- ▶ Looks like  $f(2) = 4 + 1$ .
- ▶ Curiouser and curiouser...

## Accounting for small values of $n$ : $n = 2$




Figure: A  $2 \times 2$  grid square

- ▶ Looks like  $f(2) = 4 + 1$ .
- ▶ Curiouser and curiouser...
- ▶ Keep experimenting!



## Accounting for small values of $n$ : $n = 3$




Figure: A  $3 \times 3$  grid square

## Accounting for small values of $n$ : $n = 3$




Figure: A  $3 \times 3$  grid square

## Accounting for small values of $n$ : $n = 3$




Figure: A  $3 \times 3$  grid square

## Accounting for small values of $n$ : $n = 3$




Figure: A  $3 \times 3$  grid square

## Accounting for small values of $n$ : $n = 3$




Figure: A  $3 \times 3$  grid square

## Accounting for small values of $n$ : $n = 3$




Figure: A  $3 \times 3$  grid square

## Accounting for small values of $n$ : $n = 3$




Figure: A  $3 \times 3$  grid square

## Accounting for small values of $n$ : $n = 3$




Figure: A  $3 \times 3$  grid square



## Accounting for small values of $n$ : $n = 3$




Figure: A  $3 \times 3$  grid square

## Accounting for small values of $n$ : $n = 3$




Figure: A  $3 \times 3$  grid square

## Accounting for small values of $n$ : $n = 3$




Figure: A  $3 \times 3$  grid square

## Accounting for small values of $n$ : $n = 3$




Figure: A  $3 \times 3$  grid square

## Accounting for small values of $n$ : $n = 3$




Figure: A  $3 \times 3$  grid square

## Accounting for small values of $n$ : $n = 3$




Figure: A  $3 \times 3$  grid square

## Accounting for small values of $n$ : $n = 3$




Figure: A  $3 \times 3$  grid square

## Accounting for small values of $n$ : $n = 3$




Figure: A  $3 \times 3$  grid square

- Looks like  $f(3) = 9 + 4 + 1$ .



## Accounting for small values of $n$ : $n = 3$




Figure: A  $3 \times 3$  grid square

- ▶ Looks like  $f(3) = 9 + 4 + 1$ .
- ▶ Curiouser and curiouser...

## Accounting for small values of $n$ : $n = 3$




Figure: A  $3 \times 3$  grid square

- ▶ Looks like  $f(3) = 9 + 4 + 1$ .
- ▶ Curiouser and curiouser...
- ▶ Time for a conjecture...

# Conjecture

The statement  $P(n)$ , namely our conjecture, is as follows:

**If  $f(n)$  is the number of squares in an  $n \times n$  grid square then**

$$f(n) = \sum_{i=1}^n i^2 = 1^1 + 2^2 + \cdots + n^2$$

# Induction Argument to verify that $P(n)$ is true $\forall n \in \mathbb{Z}^+$

- **Base Cases.** We have verified that  $P(1)$ ,  $P(2)$ , and  $P(3)$  are all true.

# Induction Argument to verify that $P(n)$ is true $\forall n \in \mathbb{Z}^+$

- ▶ **Base Cases.** We have verified that  $P(1)$ ,  $P(2)$ , and  $P(3)$  are all true.
- ▶ **Induction Hypothesis.** Assume that  $P(k)$  is true for some **fixed** integer  $k \geq 3$ .

# Induction Argument to verify that $P(n)$ is true $\forall n \in \mathbb{Z}^+$


- ▶ **Base Cases.** We have verified that  $P(1)$ ,  $P(2)$ , and  $P(3)$  are all true.
- ▶ **Induction Hypothesis.** Assume that  $P(k)$  is true for some **fixed** integer  $k \geq 3$ .
- ▶ **Inductive Step.** We must prove that  $P(k + 1)$  is true.

## Inductive Step

- ▶ By the induction hypothesis, we know that  $f(k) = 1 + 2^2 + 3^2 + \cdots + k^2$ .

# Inductive Step


- ▶ By the induction hypothesis, we know that  $f(k) = 1 + 2^2 + 3^2 + \cdots + k^2$ .
- ▶ Consider a  $(k + 1) \times (k + 1)$  grid square.






# Inductive Step

- ▶ Since the lower left  $k \times k$  (black) square has been accounted for by the induction hypothesis,




# Inductive Step

- ▶ Since the lower left  $k \times k$  (black) square has been accounted for by the induction hypothesis,




- ▶ it remains to count all possible new squares, namely only the ones that include the top row and right column of red squares.

# Counting the remaining squares




- ▶  $1 \times 1$ 's:  $(k + 1) + k = 2k + 1$

# Counting the remaining squares




- ▶  $1 \times 1$ 's:  $(k + 1) + k = 2k + 1$
- ▶  $2 \times 2$ 's:  $k + (k - 1) = 2k - 1$

# Counting the remaining squares




- ▶  $1 \times 1$ 's:  $(k+1) + k = 2k+1$
- ▶  $2 \times 2$ 's:  $k + (k-1) = 2k-1$
- ▶  $3 \times 3$ 's:  $(k-1) + (k-2) = 2k-3$

# Counting the remaining squares




- ▶  $1 \times 1$ 's:  $(k + 1) + k = 2k + 1$
- ▶  $2 \times 2$ 's:  $k + (k - 1) = 2k - 1$
- ▶  $3 \times 3$ 's:  $(k - 1) + (k - 2) = 2k - 3$
- ▶  $\vdots$

# Counting the remaining squares



# Counting the remaining squares




▶  
⋮

▶  $k \times k$ 's:  $(k - (k - 1)) + (k - (k - 2)) = 3$



# Counting the remaining squares




▶  $\vdots$

▶  $k \times k$ 's:  $(k - (k - 1)) + (k - (k - 2)) = 3$

▶  $k + 1 \times k + 1$ 's:  $(k - (k)) + (k - (k - 1)) = 1$

# Counting the remaining squares



►  $\vdots$

►  $k \times k$ 's:  $(k - (k - 1)) + (k - (k - 2)) = 3$

►  $k + 1 \times k + 1$ 's:  $(k - (k)) + (k - (k - 1)) = 1$

► Thus the total number of **new** squares is  
 $\sum_{i=1}^{k+1} (2i - 1) = (k + 1)^2$  by Example 1.

## Total number of squares in a $k + 1 \times k + 1$ grid square

- **Conclusion.** Thus, by induction, the total number of squares in a  $k + 1 \times k + 1$  grid square is

$$\left(\sum_{i=1}^k i^2\right) + (k + 1)^2 = \sum_{i=1}^{k+1} i^2$$

.

## Total number of squares in a $k + 1 \times k + 1$ grid square

- **Conclusion.** Thus, by induction, the total number of squares in a  $k + 1 \times k + 1$  grid square is

$$\left(\sum_{i=1}^k i^2\right) + (k + 1)^2 = \sum_{i=1}^{k+1} i^2$$

.

- That is,  $P(n)$  is true  $\forall n \in \mathbb{Z}^+$ .

## Total number of squares in a $k + 1 \times k + 1$ grid square

- ▶ **Conclusion.** Thus, by induction, the total number of squares in a  $k + 1 \times k + 1$  grid square is

$$\left(\sum_{i=1}^k i^2\right) + (k + 1)^2 = \sum_{i=1}^{k+1} i^2$$

.

- ▶ That is,  $P(n)$  is true  $\forall n \in \mathbb{Z}^+$ .

- ▶ **QED**

## New Topic: The Fibonacci Numbers

- ▶ For a brief history of Leonardo Pisano Fibonacci (approximately 1170-1230 A.D.), see <http://www-history.mcs.st-and.ac.uk/Biographies/Fibonacci.html>.

## New Topic: The Fibonacci Numbers

- ▶ For a brief history of Leonardo Pisano Fibonacci (approximately 1170-1230 A.D.), see <http://www-history.mcs.st-and.ac.uk/Biographies/Fibonacci.html>.
- ▶ The  $i$ th Fibonacci number is denoted by  $F_i$ , and

## New Topic: The Fibonacci Numbers

- ▶ For a brief history of Leonardo Pisano Fibonacci (approximately 1170-1230 A.D.), see <http://www-history.mcs.st-and.ac.uk/Biographies/Fibonacci.html>.
- ▶ The  $i$ th Fibonacci number is denoted by  $F_i$ , and
- ▶  $F_0 = 0$ ,  $F_1 = 1$ , and for all integers  $n > 1$  we define  $F_i = F_{i-1} + F_{i-2}$ .



## New Topic: The Fibonacci Numbers

- ▶ For a brief history of Leonardo Pisano Fibonacci (approximately 1170-1230 A.D.), see <http://www-history.mcs.st-and.ac.uk/Biographies/Fibonacci.html>.
- ▶ The  $i$ th Fibonacci number is denoted by  $F_i$ , and
- ▶  $F_0 = 0$ ,  $F_1 = 1$ , and for all integers  $n > 1$  we define  $F_i = F_{i-1} + F_{i-2}$ .
- ▶ For example, the first 12 numbers in the *Fibonacci sequence* are 1,1,2,3,5,8,13,21,34,55,89,144

# Fibonacci Numbers: some thoughts

- ▶ The fact that  $F_n$  is defined by way of  $F_{n-1}$  and  $F_{n-2}$  lends itself well to the use of induction for proving properties of the Fibonacci sequence.

# Fibonacci Numbers: some thoughts

- ▶ The fact that  $F_n$  is defined by way of  $F_{n-1}$  and  $F_{n-2}$  lends itself well to the use of induction for proving properties of the Fibonacci sequence.
- ▶ Let  $f(n) = F_n$ . How big is  $f(n)$ ?

# Fibonacci Numbers: some thoughts

- ▶ The fact that  $F_n$  is defined by way of  $F_{n-1}$  and  $F_{n-2}$  lends itself well to the use of induction for proving properties of the Fibonacci sequence.
- ▶ Let  $f(n) = F_n$ . How big is  $f(n)$ ?
- ▶ Generally speaking, we will study the growth of functions, namely *asymptotics*, with the intention of measuring the efficiency of our algorithms.

## Example 1

**Problem.** Show by induction that  $F_1 + F_3 + \cdots + F_{2n-1} = F_{2n}$  for all integers  $n \geq 1$ .

- ▶ What is the formulation of the statement  $P(n)$  for the above problem?

## Example 1

**Problem.** Show by induction that  $F_1 + F_3 + \cdots + F_{2n-1} = F_{2n}$  for all integers  $n \geq 1$ .

- ▶ What is the formulation of the statement  $P(n)$  for the above problem?
- ▶  $P(n)$  is the statement  $F_1 + F_3 + \cdots + F_{2n-1} = F_{2n}$ .

## Example 1

**Problem.** Show by induction that  $F_1 + F_3 + \cdots + F_{2n-1} = F_{2n}$  for all integers  $n \geq 1$ .

- ▶ What is the formulation of the statement  $P(n)$  for the above problem?
- ▶  $P(n)$  is the statement  $F_1 + F_3 + \cdots + F_{2n-1} = F_{2n}$ .
- ▶ **Base Cases.**

## Example 1

**Problem.** Show by induction that  $F_1 + F_3 + \cdots + F_{2n-1} = F_{2n}$  for all integers  $n \geq 1$ .

- ▶ What is the formulation of the statement  $P(n)$  for the above problem?
- ▶  $P(n)$  is the statement  $F_1 + F_3 + \cdots + F_{2n-1} = F_{2n}$ .
- ▶ **Base Cases.**
- ▶  $n = 1$  (Test  $P(1)$ ):  $F_1 = 1 = F_2 = F_{2 \times 1}$ . ✓



## Example 1

**Problem.** Show by induction that  $F_1 + F_3 + \cdots + F_{2n-1} = F_{2n}$  for all integers  $n \geq 1$ .

- ▶ What is the formulation of the statement  $P(n)$  for the above problem?
- ▶  $P(n)$  is the statement  $F_1 + F_3 + \cdots + F_{2n-1} = F_{2n}$ .
- ▶ **Base Cases.**
- ▶  $n = 1$  (Test  $P(1)$ ):  $F_1 = 1 = F_2 = F_{2 \times 1}$ . ✓
- ▶  $n = 2$  (Test  $P(2)$ ):  $F_1 + F_3 = 1 + 2 + 3 = F_4 = F_{2 \times 2}$ . ✓

## Example 1

**Problem.** Show by induction that  $F_1 + F_3 + \cdots + F_{2n-1} = F_{2n}$  for all integers  $n \geq 1$ .

- ▶ What is the formulation of the statement  $P(n)$  for the above problem?
- ▶  $P(n)$  is the statement  $F_1 + F_3 + \cdots + F_{2n-1} = F_{2n}$ .
- ▶ **Base Cases.**
  - ▶  $n = 1$  (Test  $P(1)$ ):  $F_1 = 1 = F_2 = F_{2 \times 1}$ . ✓
  - ▶  $n = 2$  (Test  $P(2)$ ):  $F_1 + F_3 = 1 + 2 + 3 = F_4 = F_{2 \times 2}$ . ✓
  - ▶  $n = 3$  (Test  $P(3)$ ):  
 $F_1 + F_3 + F_5 = 1 + 2 + 5 = 8 = F_6 = F_{2 \times 3}$ . ✓

# Induction Hypothesis and Inductive Step

- **Induction hypothesis.** Assume  $F_1 + F_3 + \cdots + F_{2k-1} = F_{2k}$  for some fixed integer  $k \geq 1$ .

# Induction Hypothesis and Inductive Step

- ▶ **Induction hypothesis.** Assume  $F_1 + F_3 + \cdots + F_{2k-1} = F_{2k}$  for some fixed integer  $k \geq 1$ .
- ▶ **Inductive Step.** We must prove that  $F_1 + F_3 + \cdots + F_{2k-1} + F_{2k+1} = F_{2(k+1)}$

# Induction Hypothesis and Inductive Step

- ▶ **Induction hypothesis.** Assume  $F_1 + F_3 + \cdots + F_{2k-1} = F_{2k}$  for some fixed integer  $k \geq 1$ .
- ▶ **Inductive Step.** We must prove that  $F_1 + F_3 + \cdots + F_{2k-1} + F_{2k+1} = F_{2(k+1)}$
- ▶ **Here goes.** The first  $k$  terms of  $F_1 + F_3 + \cdots + F_{2k-1} + F_{2k+1}$  can be simplified by the induction hypothesis.

# Induction Hypothesis and Inductive Step

- ▶ **Induction hypothesis.** Assume  $F_1 + F_3 + \cdots + F_{2k-1} = F_{2k}$  for some fixed integer  $k \geq 1$ .
- ▶ **Inductive Step.** We must prove that  $F_1 + F_3 + \cdots + F_{2k-1} + F_{2k+1} = F_{2(k+1)}$
- ▶ **Here goes.** The first  $k$  terms of  $F_1 + F_3 + \cdots + F_{2k-1} + F_{2k+1}$  can be simplified by the induction hypothesis.
- ▶ That is,  $F_1 + F_3 + \cdots + F_{2k-1} + F_{2k+1} = F_{2k} + F_{2k+1}$ .

# Induction Hypothesis and Inductive Step

- ▶ **Induction hypothesis.** Assume  $F_1 + F_3 + \cdots + F_{2k-1} = F_{2k}$  for some fixed integer  $k \geq 1$ .
- ▶ **Inductive Step.** We must prove that  $F_1 + F_3 + \cdots + F_{2k-1} + F_{2k+1} = F_{2(k+1)}$
- ▶ **Here goes.** The first  $k$  terms of  $F_1 + F_3 + \cdots + F_{2k-1} + F_{2k+1}$  can be simplified by the induction hypothesis.
- ▶ That is,  $F_1 + F_3 + \cdots + F_{2k-1} + F_{2k+1} = F_{2k} + F_{2k+1}$ .
- ▶ But by the definition of the Fibonacci numbers we have that  $F_{2k} + F_{2k+1} = F_{2k+2}$ , as desired.

## Make a conclusion

- **Conclusion.** By induction,  $P(n)$  is true  $\forall n \in \mathbb{Z}^+$ .



## Make a conclusion

- ▶ **Conclusion.** By induction,  $P(n)$  is true  $\forall n \in \mathbb{Z}^+$ .
- ▶ That is,  $F_1 + F_3 + \cdots + F_{2n-1} = F_{2n} \forall n \in \mathbb{Z}^+$

## Make a conclusion

- ▶ **Conclusion.** By induction,  $P(n)$  is true  $\forall n \in \mathbb{Z}^+$ .
- ▶ That is,  $F_1 + F_3 + \cdots + F_{2n-1} = F_{2n} \forall n \in \mathbb{Z}^+$
- ▶ **QED**

## Example 2

- **Problem.** For  $F_n$ , the  $n$ th Fibonacci number, Let  $P(n)$  be the statement:

$$F_{n+1}F_{n-1} - F_n^2 = (-1)^n$$

## Example 2

- ▶ **Problem.** For  $F_n$ , the  $n$ th Fibonacci number, Let  $P(n)$  be the statement:

$$F_{n+1}F_{n-1} - F_n^2 = (-1)^n$$

- ▶ We will verify that  $P(n)$  is true  $\forall n \in \mathbb{Z}^+$ .

## Example 2

- ▶ **Problem.** For  $F_n$ , the  $n$ th Fibonacci number, Let  $P(n)$  be the statement:

$$F_{n+1}F_{n-1} - F_n^2 = (-1)^n$$

- ▶ We will verify that  $P(n)$  is true  $\forall n \in \mathbb{Z}^+$ .
- ▶ We'll do a proof by induction.

$$P(n) : F_{n+1}F_{n-1} - F_n^2 = (-1)^n$$

► **Base Cases.**  $n = 1$  (test  $P(1)$ ):

$$F_2F_0 - F_1^2 = 1 \times 0 - 1^2 = -1 = (-1)^1. \checkmark$$

$$P(n) : F_{n+1}F_{n-1} - F_n^2 = (-1)^n$$

► **Base Cases.**  $n = 1$  (test  $P(1)$ ):

$$F_2F_0 - F_1^2 = 1 \times 0 - 1^2 = -1 = (-1)^1. \checkmark$$

►  $n = 2$  (test  $P(2)$ ):  $F_3F_1 - F_2^2 = 2 \times 1 - 1^2 = 1 = (-1)^2. \checkmark$

$$P(n) : F_{n+1}F_{n-1} - F_n^2 = (-1)^n$$

► **Base Cases.**  $n = 1$  (test  $P(1)$ ):

$$F_2F_0 - F_1^2 = 1 \times 0 - 1^2 = -1 = (-1)^1. \checkmark$$

►  $n = 2$  (test  $P(2)$ ):  $F_3F_1 - F_2^2 = 2 \times 1 - 1^2 = 1 = (-1)^2. \checkmark$

►  $n = 3$  (test  $P(3)$ ):  $F_4F_2 - F_3^2 = 3 \times 1 - 2^2 = -1 = (-1)^3. \checkmark$



$$P(n) : F_{n+1}F_{n-1} - F_n^2 = (-1)^n$$

- ▶ **Base Cases.**  $n = 1$  (test  $P(1)$ ):

$$F_2F_0 - F_1^2 = 1 \times 0 - 1^2 = -1 = (-1)^1. \checkmark$$

- ▶  $n = 2$  (test  $P(2)$ ):  $F_3F_1 - F_2^2 = 2 \times 1 - 1^2 = 1 = (-1)^2. \checkmark$

- ▶  $n = 3$  (test  $P(3)$ ):  $F_4F_2 - F_3^2 = 3 \times 1 - 2^2 = -1 = (-1)^3. \checkmark$

- ▶ **Induction Hypothesis.** Assume that  $P(k)$  is true for some fixed integer  $k \geq 1$ .

$$P(n) : F_{n+1}F_{n-1} - F_n^2 = (-1)^n$$

- ▶ **Base Cases.**  $n = 1$  (test  $P(1)$ ):  
 $F_2F_0 - F_1^2 = 1 \times 0 - 1^2 = -1 = (-1)^1$ . ✓
- ▶  $n = 2$  (test  $P(2)$ ):  $F_3F_1 - F_2^2 = 2 \times 1 - 1^2 = 1 = (-1)^2$ . ✓
- ▶  $n = 3$  (test  $P(3)$ ):  $F_4F_2 - F_3^2 = 3 \times 1 - 2^2 = -1 = (-1)^3$ . ✓
- ▶ **Induction Hypothesis.** Assume that  $P(k)$  is true for some fixed integer  $k \geq 1$ .
- ▶ That is, assume  $F_{k+1}F_{k-1} - F_k^2 = (-1)^k$  for some fixed integer  $k \geq 1$ .

$$P(n) : F_{n+1}F_{n-1} - F_n^2 = (-1)^n$$

- **Inductive Step.** We must prove that  $P(k+1)$  is true;

$$P(n) : F_{n+1}F_{n-1} - F_n^2 = (-1)^n$$

- ▶ **Inductive Step.** We must prove that  $P(k+1)$  is true;
- ▶ That is, we must prove that  $F_{k+2}F_k - F_{k+1}^2 = (-1)^{k+1}$

$$P(n) : F_{n+1}F_{n-1} - F_n^2 = (-1)^n$$

- ▶ **Inductive Step.** We must prove that  $P(k+1)$  is true;
- ▶ That is, we must prove that  $F_{k+2}F_k - F_{k+1}^2 = (-1)^{k+1}$
- ▶ **Proof.**

# Inductive Step, continued

- ▶ **Proof.**

# Inductive Step, continued

► **Proof.**

►  $F_{k+2}F_k - F_{k+1}^2$

# Inductive Step, continued

- ▶ **Proof.**

- ▶  $F_{k+2}F_k - F_{k+1}^2$

- ▶  $= (F_{k+1} + F_k)F_k - F_{k+1}^2$  (by the definition of the Fibonacci numbers)



# Inductive Step, continued

- ▶ **Proof.**

- ▶  $F_{k+2}F_k - F_{k+1}^2$

- ▶  $= (F_{k+1} + F_k)F_k - F_{k+1}^2$  (by the definition of the Fibonacci numbers)

- ▶  $= F_{k+1}F_k + F_k^2 - F_{k+1}^2$  (by algebra)

# Inductive Step, continued

- ▶ **Proof.**

- ▶  $F_{k+2}F_k - F_{k+1}^2$

- ▶  $= (F_{k+1} + F_k)F_k - F_{k+1}^2$  (by the definition of the Fibonacci numbers)

- ▶  $= F_{k+1}F_k + F_k^2 - F_{k+1}^2$  (by algebra)

- ▶  $= F_{k+1}(F_k - F_{k+1}) + F_k^2$

# Inductive Step, continued

► **Proof.**

►  $F_{k+2}F_k - F_{k+1}^2$

►  $= (F_{k+1} + F_k)F_k - F_{k+1}^2$  (by the definition of the Fibonacci numbers)

►  $= F_{k+1}F_k + F_k^2 - F_{k+1}^2$  (by algebra)

►  $= F_{k+1}(F_k - F_{k+1}) + F_k^2$

►  $= F_{k+1}(F_k - (F_k + F_{k-1})) + F_k^2$

# Inductive Step, continued

► **Proof.**

►  $F_{k+2}F_k - F_{k+1}^2$

►  $= (F_{k+1} + F_k)F_k - F_{k+1}^2$  (by the definition of the Fibonacci numbers)

►  $= F_{k+1}F_k + F_k^2 - F_{k+1}^2$  (by algebra)

►  $= F_{k+1}(F_k - F_{k+1}) + F_k^2$

►  $= F_{k+1}(F_k - (F_k + F_{k-1})) + F_k^2$

►  $= F_{k+1}(-F_{k-1}) + F_k^2$

## Inductive step, continued

$$\blacktriangleright = F_{k+1}(-F_{k-1}) + F_k^2$$

## Inductive step, continued

$$\blacktriangleright = F_{k+1}(-F_{k-1}) + F_k^2$$

$$\blacktriangleright = -(F_{k+1}F_{k-1} - F_k^2)$$

## Inductive step, continued

$$\blacktriangleright = F_{k+1}(-F_{k-1}) + F_k^2$$

$$\blacktriangleright = -(F_{k+1}F_{k-1} - F_k^2)$$

$$\blacktriangleright = -(-1)^k \text{ (by the induction hypothesis)}$$

## Inductive step, continued

$$\blacktriangleright = F_{k+1}(-F_{k-1}) + F_k^2$$

$$\blacktriangleright = -(F_{k+1}F_{k-1} - F_k^2)$$

$$\blacktriangleright = -(-1)^k \text{ (by the induction hypothesis)}$$

$$\blacktriangleright = (-1)^{k+1}, \text{ as desired.}$$



## Make a conclusion

**Conclusion.** By induction  $F_{n+1}F_{n-1} - F_n^2 = (-1)^n \quad \forall n \in \mathbb{Z}^+.$   
**QED**

# A generating function for the Fibonacci numbers

- ▶ For  $F_i$ , the  $i$ th Fibonacci number, define  $g(z) = \sum_{i=0}^{\infty} F_i z^i$

# A generating function for the Fibonacci numbers

- ▶ For  $F_i$ , the  $i$ th Fibonacci number, define  $g(z) = \sum_{i=0}^{\infty} F_i z^i$
- ▶  $= 0 + 1z + 1z^2 + 2z^3 + 3z^4 + 5z^5 + 8z^6 + \dots$

# A generating function for the Fibonacci numbers

- ▶ For  $F_i$ , the  $i$ th Fibonacci number, define  $g(z) = \sum_{i=0}^{\infty} F_i z^i$
- ▶  $= 0 + 1z + 1z^2 + 2z^3 + 3z^4 + 5z^5 + 8z^6 + \dots$
- ▶ The above is called a *formal power series*, which you've learned about in your calculus classes.

## Step 1: The Claim.

- ▶ **Claim 1.**  $g(z) = z + zg(z) + z^2g(z)$ .

## Step 1: The Claim.

- ▶ **Claim 1.**  $g(z) = z + zg(z) + z^2g(z)$ .
- ▶ To see why claim 1 is true, let's deconstruct each of  $zg(z)$  and  $z^2g(z)$  from the original definition of  $g(z)$  as a formal power series.

Regarding  $g(z) = \sum_{i=0}^{\infty} F_i z^i$

►  $zg(z) = z \sum_{i=0}^{\infty} F_i z^i$

Regarding  $g(z) = \sum_{i=0}^{\infty} F_i z^i$

►  $zg(z) = z \sum_{i=0}^{\infty} F_i z^i$

►  $= \sum_{i=0}^{\infty} F_i z^{i+1} \quad (1)$



Regarding  $g(z) = \sum_{i=0}^{\infty} F_i z^i$

- ▶  $zg(z) = z \sum_{i=0}^{\infty} F_i z^i$
- ▶  $= \sum_{i=0}^{\infty} F_i z^{i+1} \quad (1)$
- ▶ Now let's carry out the same procedure of  $z^2 g(z)$ .

Regarding  $g(z) = \sum_{i=0}^{\infty} F_i z^i$

- ▶  $zg(z) = z \sum_{i=0}^{\infty} F_i z^i$

- ▶  $= \sum_{i=0}^{\infty} F_i z^{i+1} \quad (1)$

- ▶ Now let's carry out the same procedure of  $z^2 g(z)$ .

- ▶  $z^2 g(z) = z^2 \sum_{i=0}^{\infty} F_i z^i = \sum_{i=0}^{\infty} F_i z^{i+2} \quad (2)$

Regarding  $g(z) = \sum_{i=0}^{\infty} F_i z^i$

- ▶  $zg(z) = z \sum_{i=0}^{\infty} F_i z^i$
- ▶  $= \sum_{i=0}^{\infty} F_i z^{i+1} \quad (1)$
- ▶ Now let's carry out the same procedure of  $z^2g(z)$ .
- ▶  $z^2g(z) = z^2 \sum_{i=0}^{\infty} F_i z^i = \sum_{i=0}^{\infty} F_i z^{i+2} \quad (2)$
- ▶ Next, we'll massage (1) and (2) to make the final algebra easier to digest when we finally compute  $z + zg(z) + z^2g(z)$ .

Index changes for  $= \sum_{i=0}^{\infty} F_i z^{i+1}$  and  $\sum_{i=0}^{\infty} F_i z^{i+2}$

- ▶ Note that  $= \sum_{i=0}^{\infty} F_i z^{i+1}$  is exactly the same as  $\sum_{i=1}^{\infty} F_{i-1} z^i$

Index changes for  $= \sum_{i=0}^{\infty} F_i z^{i+1}$  and  $\sum_{i=0}^{\infty} F_i z^{i+2}$

- ▶ Note that  $= \sum_{i=0}^{\infty} F_i z^{i+1}$  is exactly the same as  $\sum_{i=1}^{\infty} F_{i-1} z^i$
- ▶ And here's a little trick:  $\sum_{i=1}^{\infty} F_{i-1} z^i = \sum_{i=2}^{\infty} F_{i-1} z^i$ . Why??

Index changes for  $= \sum_{i=0}^{\infty} F_i z^{i+1}$  and  $\sum_{i=0}^{\infty} F_i z^{i+2}$

- ▶ Note that  $= \sum_{i=0}^{\infty} F_i z^{i+1}$  is exactly the same as  $\sum_{i=1}^{\infty} F_{i-1} z^i$
- ▶ And here's a little trick:  $\sum_{i=1}^{\infty} F_{i-1} z^i = \sum_{i=2}^{\infty} F_{i-1} z^i$ . Why??
- ▶ Similarly  $= \sum_{i=0}^{\infty} F_i z^{i+2}$  is exactly the same as  $\sum_{i=2}^{\infty} F_{i-2} z^i$

Index changes for  $= \sum_{i=0}^{\infty} F_i z^{i+1}$  and  $\sum_{i=0}^{\infty} F_i z^{i+2}$

- ▶ Note that  $= \sum_{i=0}^{\infty} F_i z^{i+1}$  is exactly the same as  $\sum_{i=1}^{\infty} F_{i-1} z^i$
- ▶ And here's a little trick:  $\sum_{i=1}^{\infty} F_{i-1} z^i = \sum_{i=2}^{\infty} F_{i-1} z^i$ . Why??
- ▶ Similarly  $= \sum_{i=0}^{\infty} F_i z^{i+2}$  is exactly the same as  $\sum_{i=2}^{\infty} F_{i-2} z^i$
- ▶ Thus  $z + zg(z) + z^2g(z) = z + \sum_{i=2}^{\infty} F_{i-1} z^i + \sum_{i=2}^{\infty} F_{i-2} z^i$

Index changes for  $= \sum_{i=0}^{\infty} F_i z^{i+1}$  and  $\sum_{i=0}^{\infty} F_i z^{i+2}$

- ▶ Note that  $= \sum_{i=0}^{\infty} F_i z^{i+1}$  is exactly the same as  $\sum_{i=1}^{\infty} F_{i-1} z^i$
- ▶ And here's a little trick:  $\sum_{i=1}^{\infty} F_{i-1} z^i = \sum_{i=2}^{\infty} F_{i-1} z^i$ . Why??
- ▶ Similarly  $= \sum_{i=0}^{\infty} F_i z^{i+2}$  is exactly the same as  $\sum_{i=2}^{\infty} F_{i-2} z^i$
- ▶ Thus  $z + zg(z) + z^2g(z) = z + \sum_{i=2}^{\infty} F_{i-1} z^i + \sum_{i=2}^{\infty} F_{i-2} z^i$
- ▶  $= z + \sum_{i=2}^{\infty} (F_{i-1} + F_{i-2}) z^i$



Index changes for  $= \sum_{i=0}^{\infty} F_i z^{i+1}$  and  $\sum_{i=0}^{\infty} F_i z^{i+2}$

- ▶ Note that  $= \sum_{i=0}^{\infty} F_i z^{i+1}$  is exactly the same as  $\sum_{i=1}^{\infty} F_{i-1} z^i$
- ▶ And here's a little trick:  $\sum_{i=1}^{\infty} F_{i-1} z^i = \sum_{i=2}^{\infty} F_{i-1} z^i$ . Why??
- ▶ Similarly  $= \sum_{i=0}^{\infty} F_i z^{i+2}$  is exactly the same as  $\sum_{i=2}^{\infty} F_{i-2} z^i$
- ▶ Thus  $z + zg(z) + z^2g(z) = z + \sum_{i=2}^{\infty} F_{i-1} z^i + \sum_{i=2}^{\infty} F_{i-2} z^i$
- ▶  $= z + \sum_{i=2}^{\infty} (F_{i-1} + F_{i-2}) z^i$
- ▶  $= F_0 z^0 + F_1 z + \sum_{i=2}^{\infty} F_i z^i = \sum_{i=0}^{\infty} F_i z^i$

Index changes for  $= \sum_{i=0}^{\infty} F_i z^{i+1}$  and  $\sum_{i=0}^{\infty} F_i z^{i+2}$

- ▶ Note that  $= \sum_{i=0}^{\infty} F_i z^{i+1}$  is exactly the same as  $\sum_{i=1}^{\infty} F_{i-1} z^i$
- ▶ And here's a little trick:  $\sum_{i=1}^{\infty} F_{i-1} z^i = \sum_{i=2}^{\infty} F_{i-1} z^i$ . Why??
- ▶ Similarly  $= \sum_{i=0}^{\infty} F_i z^{i+2}$  is exactly the same as  $\sum_{i=2}^{\infty} F_{i-2} z^i$
- ▶ Thus  $z + zg(z) + z^2g(z) = z + \sum_{i=2}^{\infty} F_{i-1} z^i + \sum_{i=2}^{\infty} F_{i-2} z^i$
- ▶  $= z + \sum_{i=2}^{\infty} (F_{i-1} + F_{i-2}) z^i$
- ▶  $= F_0 z^0 + F_1 z + \sum_{i=2}^{\infty} F_i z^i = \sum_{i=0}^{\infty} F_i z^i$
- ▶  $= g(z)$ , as claimed. This completes the proof of Claim 1.

## Step 2: some algebra

- By step 1, we have seen that  $g(z) = z + zg(z) + z^2g(z)$ , which implies that

## Step 2: some algebra

- ▶ By step 1, we have seen that  $g(z) = z + zg(z) + z^2g(z)$ , which implies that
- ▶  $g(z) - zg(z) - z^2g(z) = z$

## Step 2: some algebra

- ▶ By step 1, we have seen that  $g(z) = z + zg(z) + z^2g(z)$ , which implies that
- ▶  $g(z) - zg(z) - z^2g(z) = z$
- ▶  $\Rightarrow g(z)(1 - z - z^2) = z$

## Step 2: some algebra

- ▶ By step 1, we have seen that  $g(z) = z + zg(z) + z^2g(z)$ , which implies that
- ▶  $g(z) - zg(z) - z^2g(z) = z$
- ▶  $\Rightarrow g(z)(1 - z - z^2) = z$
- ▶  $\Rightarrow g(z) = \frac{z}{1-z-z^2}.$

### Step 3. Surprising connection between step 2 and the golden ratio

- ▶ Let  $\phi = \frac{1+\sqrt{5}}{2}$  and  $\phi' = \frac{1-\sqrt{5}}{2}$ .

### Step 3. Surprising connection between step 2 and the golden ratio

- ▶ Let  $\phi = \frac{1+\sqrt{5}}{2}$  and  $\phi' = \frac{1-\sqrt{5}}{2}$ .
- ▶  $\phi$  is called the golden mean or golden ratio (or divine proportion...).



### Step 3. Surprising connection between step 2 and the golden ratio

- ▶ Let  $\phi = \frac{1+\sqrt{5}}{2}$  and  $\phi' = \frac{1-\sqrt{5}}{2}$ .
- ▶  $\phi$  is called the golden mean or golden ratio (or divine proportion...).
- ▶ You verify that  $1 - z - z^2 = (1 - \phi z)(1 - \phi' z)$ .

## Step 4.

- ▶ Assuming that you have done the homework on the previous slide, we then have

## Step 4.

- ▶ Assuming that you have done the homework on the previous slide, we then have

- ▶  $\frac{z}{1-z-z^2} = \frac{1}{\sqrt{5}} \left( \frac{1}{1-\phi z} - \frac{1}{1-\phi' z} \right)$ . Verify this!

## Step 5. A reminder about geometric series.

- ▶ For any  $a, r \in \mathbb{R}$ , let  $S = a + ar + ar^2 + \cdots +$ , where

## Step 5. A reminder about geometric series.

- ▶ For any  $a, r \in \mathbb{R}$ , let  $S = a + ar + ar^2 + \cdots +$ , where
- ▶  $a$  is called the *first term* and  $r$  is called the *ratio*.

## Step 5. A reminder about geometric series.

- ▶ For any  $a, r \in \mathbb{R}$ , let  $S = a + ar + ar^2 + \cdots$ , where
- ▶  $a$  is called the *first term* and  $r$  is called the *ratio*.
- ▶ Then formally speaking (all issues of convergence aside), we have  $S = \frac{a}{1-r}$ .

Step 6. Recall that  $a + ar + ar^2 + \cdots = \frac{a}{1-r}$

- ▶ Working backwards in step 4, we see  $\frac{1}{1-\phi z} = 1 + \phi z + \phi^2 z^2 + \cdots$ , and similarly

Step 6. Recall that  $a + ar + ar^2 + \dots = \frac{a}{1-r}$

- ▶ Working backwards in step 4, we see  $\frac{1}{1-\phi z} = 1 + \phi z + \phi^2 z^2 + \dots$ , and similarly

- ▶  $\frac{1}{1-\phi' z} = 1 + \phi' z + \phi'^2 z^2 + \dots$



Step 6. Recall that  $a + ar + ar^2 + \dots = \frac{a}{1-r}$

- ▶ Working backwards in step 4, we see

$$\frac{1}{1-\phi z} = 1 + \phi z + \phi^2 z^2 + \dots, \text{ and similarly}$$

- ▶  $\frac{1}{1-\phi' z} = 1 + \phi' z + \phi'^2 z^2 + \dots$

- ▶  $\Rightarrow \frac{1}{1-\phi z} - \frac{1}{1-\phi' z} =$   
 $(\phi - \phi')z + (\phi^2 - \phi'^2)z^2 + \dots + (\phi^i - \phi'^i)z^i + \dots$

## Step 7.

- ▶ Mini-conclude that  $g(z) = \frac{1}{\sqrt{5}}\left(\frac{1}{1-\phi z} - \frac{1}{1-\phi' z}\right)$

## Step 7.

► Mini-conclude that  $g(z) = \frac{1}{\sqrt{5}} \left( \frac{1}{1-\phi z} - \frac{1}{1-\phi' z} \right)$

►  $\Rightarrow g(z) = \sum_{i=1}^{\infty} \frac{1}{\sqrt{5}} (\phi^i - (\phi')^i) z^i.$

## Step 8. Major Conclusion, at last!

- So far, from step 7, we have  $g(z) = \sum_{i=1}^{\infty} \frac{1}{\sqrt{5}}(\phi^i - (\phi')^i)z^i$ .

## Step 8. Major Conclusion, at last!

- ▶ So far, from step 7, we have  $g(z) = \sum_{i=1}^{\infty} \frac{1}{\sqrt{5}}(\phi^i - (\phi')^i)z^i$ .
- ▶ Our original definition of  $g(z)$  is  $g(z) = \sum_{i=0}^{\infty} F_i z^i$ .

## Step 8. Major Conclusion, at last!

- ▶ So far, from step 7, we have  $g(z) = \sum_{i=1}^{\infty} \frac{1}{\sqrt{5}}(\phi^i - (\phi')^i)z^i$ .
- ▶ Our original definition of  $g(z)$  is  $g(z) = \sum_{i=0}^{\infty} F_i z^i$ .
- ▶ Thus, since we have two descriptions of the same formal power series, the coefficients of each must be the same.

## Step 8. Major Conclusion, at last!

- ▶ So far, from step 7, we have  $g(z) = \sum_{i=1}^{\infty} \frac{1}{\sqrt{5}}(\phi^i - (\phi')^i)z^i$ .
- ▶ Our original definition of  $g(z)$  is  $g(z) = \sum_{i=0}^{\infty} F_i z^i$ .
- ▶ Thus, since we have two descriptions of the same formal power series, the coefficients of each must be the same.
- ▶ In particular,  $F_i = \frac{1}{\sqrt{5}}(\phi^i - (\phi')^i)$ , or equivalently,

## Step 8. Major Conclusion, at last!

- ▶ So far, from step 7, we have  $g(z) = \sum_{i=1}^{\infty} \frac{1}{\sqrt{5}}(\phi^i - (\phi')^i)z^i$ .
- ▶ Our original definition of  $g(z)$  is  $g(z) = \sum_{i=0}^{\infty} F_i z^i$ .
- ▶ Thus, since we have two descriptions of the same formal power series, the coefficients of each must be the same.
- ▶ In particular,  $F_i = \frac{1}{\sqrt{5}}(\phi^i - (\phi')^i)$ , or equivalently,
- ▶  $F_i = \frac{1}{\sqrt{5}}\left(\left(\frac{1}{1+\sqrt{5}}\right)^i - \left(\frac{1}{1-\sqrt{5}}\right)^i\right)$ .



$$F_i = \frac{1}{\sqrt{5}} \left( \left( \frac{1}{1+\sqrt{5}} \right)^i - \left( \frac{1}{1-\sqrt{5}} \right)^i \right).$$

- ▶ Since  $F_i$  is supposed to be the  $i$ th Fibonacci number, the above formula is supposed to produce an integer!!!

$$F_i = \frac{1}{\sqrt{5}} \left( \left( \frac{1}{1+\sqrt{5}} \right)^i - \left( \frac{1}{1-\sqrt{5}} \right)^i \right).$$

- ▶ Since  $F_i$  is supposed to be the  $i$ th Fibonacci number, the above formula is supposed to produce an integer!!!
- ▶ Could this be so?!

$$F_i = \frac{1}{\sqrt{5}} \left( \left( \frac{1}{1+\sqrt{5}} \right)^i - \left( \frac{1}{1-\sqrt{5}} \right)^i \right).$$

- ▶ Since  $F_i$  is supposed to be the  $i$ th Fibonacci number, the above formula is supposed to produce an integer!!!
- ▶ Could this be so?!
- ▶ Let's experiment with *Mathematica*.