

# CSCI 5451 Fall 2015

## Week 14 Notes

Professor Ellen Gethner

November 15, 2015

# Polygon Triangulation and Art Galleries


- ▶ **Problem.** Given a polygonal floor plan, what is the fewest number of guards<sup>1</sup> that must be posted

---

<sup>1</sup>guards have 360° vision

# Polygon Triangulation and Art Galleries

- ▶ **Problem.** Given a polygonal floor plan, what is the fewest number of guards<sup>1</sup> that must be posted
- ▶ in order that all points interior to the polygon are visible to at least one guard?



---


<sup>1</sup>guards have 360° vision

# Denver Art Museum (DAM)

- In case you were wondering about the existence of art galleries with odd floor plans, the Denver Art Museum downtown is the perfect example.



# DAM Floorplan



# Guarding Art Galleries

- ▶ The Problem as stated in the first slide is the famous **Art Gallery Problem**

# Guarding Art Galleries

- ▶ The Problem as stated in the first slide is the famous **Art Gallery Problem**
- ▶ and was proposed by Vic Klee in 1976<sup>2</sup>.

---

<sup>2</sup>in Mathematical Gems II, by Ross Honsberger

# Guarding Art Galleries

- ▶ The Problem as stated in the first slide is the famous **Art Gallery Problem**
- ▶ and was proposed by Vic Klee in 1976<sup>2</sup>.
- ▶ Since that time, there have been many variations suggested, and many applications as well.

---

<sup>2</sup>in Mathematical Gems II, by Ross Honsberger



# Guarding Art Galleries

- ▶ The Problem as stated in the first slide is the famous **Art Gallery Problem**
- ▶ and was proposed by Vic Klee in 1976<sup>2</sup>.
- ▶ Since that time, there have been many variations suggested, and many applications as well.
- ▶ See, for example, <https://imaginary.org/film/point-guards-and-point-clouds-solving-general-art-gallery-problems>

---

<sup>2</sup>in Mathematical Gems II, by Ross Honsberger

## Preliminary Definitions<sup>3</sup>, etc

- ▶ **Definition 1.** A **polygon** is a region in the plane bounded by a finite collection of line segments.

---

<sup>3</sup>Computational Geometry in C by J. O'Rourke

## Preliminary Definitions<sup>3</sup>, etc

- ▶ **Definition 1.** A **polygon** is a region in the plane bounded by a finite collection of line segments.
- ▶ More formally, let  $v_0, v_1, \dots, v_{n-1}$  be  $n$  points in the plane and

---

<sup>3</sup>Computational Geometry in C by J. O'Rourke

## Preliminary Definitions<sup>3</sup>, etc

- ▶ **Definition 1.** A **polygon** is a region in the plane bounded by a finite collection of line segments.
- ▶ More formally, let  $v_0, v_1, \dots, v_{n-1}$  be  $n$  points in the plane and
- ▶ let  $e_0 = v_0v_1, e_1 = v_1v_2, \dots, e_i = v_iv_{i+1}, \dots, e_{n-1} = v_{n-1}v_0$  be the  $n$  line segments that “connect the points.”

---

<sup>3</sup>Computational Geometry in C by J. O'Rourke

## Preliminary Definitions<sup>3</sup>, etc

- ▶ **Definition 1.** A **polygon** is a region in the plane bounded by a finite collection of line segments.
- ▶ More formally, let  $v_0, v_1, \dots, v_{n-1}$  be  $n$  points in the plane and
- ▶ let  $e_0 = v_0v_1, e_1 = v_1v_2, \dots, e_i = v_iv_{i+1}, \dots, e_{n-1} = v_{n-1}v_0$  be the  $n$  line segments that “connect the points.”
- ▶ Then these segments bound a **simple polygon** if and only if

---

<sup>3</sup>Computational Geometry in C by J. O'Rourke

## Preliminary Definitions<sup>3</sup>, etc

- ▶ **Definition 1.** A **polygon** is a region in the plane bounded by a finite collection of line segments.
- ▶ More formally, let  $v_0, v_1, \dots, v_{n-1}$  be  $n$  points in the plane and
- ▶ let  $e_0 = v_0v_1, e_1 = v_1v_2, \dots, e_i = v_iv_{i+1}, \dots, e_{n-1} = v_{n-1}v_0$  be the  $n$  line segments that “connect the points.”
- ▶ Then these segments bound a **simple polygon** if and only if
  1. line segments intersect only at endpoints, namely  $v_0, v_1, \dots, v_{n-1}$ , and

---

<sup>3</sup>Computational Geometry in C by J. O'Rourke

## Preliminary Definitions<sup>3</sup>, etc

- ▶ **Definition 1.** A **polygon** is a region in the plane bounded by a finite collection of line segments.
- ▶ More formally, let  $v_0, v_1, \dots, v_{n-1}$  be  $n$  points in the plane and
- ▶ let  $e_0 = v_0v_1, e_1 = v_1v_2, \dots, e_i = v_iv_{i+1}, \dots, e_{n-1} = v_{n-1}v_0$  be the  $n$  line segments that “connect the points.”
- ▶ Then these segments bound a **simple polygon** if and only if
  1. line segments intersect only at endpoints, namely  $v_0, v_1, \dots, v_{n-1}$ , and
  2.  $e_i \cap e_j = \emptyset$  whenever  $j \not\equiv i + 1 \pmod{n}$ .

---

<sup>3</sup>Computational Geometry in C by J. O'Rourke


## Preliminary Definitions<sup>3</sup>, etc

- ▶ **Definition 1.** A **polygon** is a region in the plane bounded by a finite collection of line segments.
- ▶ More formally, let  $v_0, v_1, \dots, v_{n-1}$  be  $n$  points in the plane and
- ▶ let  $e_0 = v_0v_1, e_1 = v_1v_2, \dots, e_i = v_iv_{i+1}, \dots, e_{n-1} = v_{n-1}v_0$  be the  $n$  line segments that “connect the points.”
- ▶ Then these segments bound a **simple polygon** if and only if
  1. line segments intersect only at endpoints, namely  $v_0, v_1, \dots, v_{n-1}$ , and
  2.  $e_i \cap e_j = \emptyset$  whenever  $j \not\equiv i + 1 \pmod{n}$ .
- ▶ That is, the union of the line segments form a closed non-self intersecting curve.




# Non-example

- ▶ A non-simple polygon is shown below.



# A Simple Polygon



## Visibility

- ▶ **Definition 2.** Let  $P$  be a simple polygon and suppose points  $x, y \in P$ .

# Visibility


- ▶ **Definition 2.** Let  $P$  be a simple polygon and suppose points  $x, y \in P$ .
- ▶ We say that **point**  $x$  **can see point**  $y$  (or that  $y$  is visible to  $x$ ) if and only if the closed line segment  $\overline{xy}$  is nowhere exterior to  $P$ .

# Visibility

- ▶ **Definition 2.** Let  $P$  be a simple polygon and suppose points  $x, y \in P$ .
- ▶ We say that **point**  $x$  **can see point**  $y$  (or that  $y$  is visible to  $x$ ) if and only if the closed line segment  $\overline{xy}$  is nowhere exterior to  $P$ .
- ▶ In other words,  $\overline{xy} \subset P$ . Remember that  $P$  is a region bounded by line segments.

# Visibility

- ▶ **Definition 2.** Let  $P$  be a simple polygon and suppose points  $x, y \in P$ .
- ▶ We say that **point  $x$  can see point  $y$**  (or that  $y$  is visible to  $x$ ) if and only if the closed line segment  $\overline{xy}$  is nowhere exterior to  $P$ .
- ▶ In other words,  $\overline{xy} \subset P$ . Remember that  $P$  is a region bounded by line segments.



## Visibility, continued

- ▶ And here is an example in which the line segment  $\overline{xy}$  grazes one of the vertices of  $P$ .

## Visibility, continued

- And here is an example in which the line segment  $\overline{xy}$  grazes one of the vertices of  $P$ .




Figure: point  $x$  is still visible to point  $y$



## Visibility, continued

- ▶ And finally, here is an example in which the line segment  $\overline{xy}$  has points that are exterior to  $P$  and hence  $x$  is not visible to  $y$ .

## Visibility, continued

- And finally, here is an example in which the line segment  $\overline{xy}$  has points that are exterior to  $P$  and hence  $x$  is not visible to  $y$ .




Figure: point  $x$  is *not* visible to point  $y$

## Points that cover a simple polygon

- **Definition 3.** Let the set of guards be called points; a set of points is said to **cover** a simple polygon if every point in the polygon is visible to at least one guard/point.

## Points that cover a simple polygon


- ▶ **Definition 3.** Let the set of guards be called points; a set of points is said to **cover** a simple polygon if every point in the polygon is visible to at least one guard/point.
- ▶ **Art Gallery Problem Restated:** Over all simple polygons on  $n$  vertices, what is the minimum number of guards needed to cover a polygon?

## Points that cover a simple polygon

- ▶ **Definition 3.** Let the set of guards be called points; a set of points is said to **cover** a simple polygon if every point in the polygon is visible to at least one guard/point.
- ▶ **Art Gallery Problem Restated:** Over all simple polygons on  $n$  vertices, what is the minimum number of guards needed to cover a polygon?
- ▶ **Remark.** The question is one of “worst case” since any convex polygon requires exactly one guard.

## Points that cover a simple polygon

- ▶ **Definition 3.** Let the set of guards be called points; a set of points is said to **cover** a simple polygon if every point in the polygon is visible to at least one guard/point.
- ▶ **Art Gallery Problem Restated:** Over all simple polygons on  $n$  vertices, what is the minimum number of guards needed to cover a polygon?
- ▶ **Remark.** The question is one of “worst case” since any convex polygon requires exactly one guard.



# Minimizing the number of guards

- **One Possible Approach.** Triangulate polygon  $P$  by connecting suitable pairs of vertices, and then place one guard in each triangle.




Figure: a simple polygon  $P$

# Minimizing the number of guards

- **One Possible Approach.** Triangulate polygon  $P$  by connecting suitable pairs of vertices, and then place one guard in each triangle.




Figure: a simple polygon  $P$  triangulated



# Minimizing the number of guards

- **One Possible Approach.** Triangulate polygon  $P$  by connecting suitable pairs of vertices, and then place one guard in each triangle.




Figure: a simple polygon  $P$  triangulated and guarded

## Motivation for a Theorem


- ▶ The only problem is that we have no guarantee that an arbitrary simple polygon actually **can** be triangulated.



## A Helpful Theorem

- ▶ **Theorem 3.1**<sup>4</sup>. Every simple polygon  $P$  with three or more vertices admits a triangulation. And the triangulation of a simple polygon with  $n$  vertices contains exactly  $n - 2$  triangles.


---

<sup>4</sup>from Computational Geometry, Algorithms and Applications by M. de Berg 

## A Helpful Theorem

- ▶ **Theorem 3.1**<sup>4</sup>. Every simple polygon  $P$  with three or more vertices admits a triangulation. And the triangulation of a simple polygon with  $n$  vertices contains exactly  $n - 2$  triangles.
- ▶ **Proof (by induction on  $n$ ).**


---

<sup>4</sup>from Computational Geometry, Algorithms and Applications by M. de Berg 

## A Helpful Theorem

- ▶ **Theorem 3.1**<sup>4</sup>. Every simple polygon  $P$  with three or more vertices admits a triangulation. And the triangulation of a simple polygon with  $n$  vertices contains exactly  $n - 2$  triangles.
- ▶ **Proof (by induction on  $n$ ).**
- ▶ **Base Case.** Let  $n = 3$ . Then a priori  $P$  is a triangle; thus the theorem is trivially true when  $n = 3$ .


---

<sup>4</sup>from Computational Geometry, Algorithms and Applications by M. de Berg 

## A Helpful Theorem

- ▶ **Theorem 3.1**<sup>4</sup>. Every simple polygon  $P$  with three or more vertices admits a triangulation. And the triangulation of a simple polygon with  $n$  vertices contains exactly  $n - 2$  triangles.
- ▶ **Proof (by induction on  $n$ ).**
- ▶ **Base Case.** Let  $n = 3$ . Then a priori  $P$  is a triangle; thus the theorem is trivially true when  $n = 3$ .
- ▶ **Induction Hypothesis.** Assume  $n > 3$  and suppose the theorem is true for all integers  $m$  such that  $3 \leq m < n$ .


---

<sup>4</sup>from Computational Geometry, Algorithms and Applications by M. de Berg 

## A Helpful Theorem

- ▶ **Theorem 3.1**<sup>4</sup>. Every simple polygon  $P$  with three or more vertices admits a triangulation. And the triangulation of a simple polygon with  $n$  vertices contains exactly  $n - 2$  triangles.
- ▶ **Proof (by induction on  $n$ ).**
- ▶ **Base Case.** Let  $n = 3$ . Then a priori  $P$  is a triangle; thus the theorem is trivially true when  $n = 3$ .
- ▶ **Induction Hypothesis.** Assume  $n > 3$  and suppose the theorem is true for all integers  $m$  such that  $3 \leq m < n$ .
- ▶ **Inductive Step.** And now let  $P$  be a simple polygon with  $n$  vertices.


---

<sup>4</sup>from Computational Geometry, Algorithms and Applications by M. de Berg 

## A Helpful Theorem

- ▶ **Theorem 3.1<sup>4</sup>.** Every simple polygon  $P$  with three or more vertices admits a triangulation. And the triangulation of a simple polygon with  $n$  vertices contains exactly  $n - 2$  triangles.
- ▶ **Proof (by induction on  $n$ ).**
- ▶ **Base Case.** Let  $n = 3$ . Then a priori  $P$  is a triangle; thus the theorem is trivially true when  $n = 3$ .
- ▶ **Induction Hypothesis.** Assume  $n > 3$  and suppose the theorem is true for all integers  $m$  such that  $3 \leq m < n$ .
- ▶ **Inductive Step.** And now let  $P$  be a simple polygon with  $n$  vertices.
- ▶ Our goal is to prove that  $P$  admits a triangulation.

---

<sup>4</sup>from Computational Geometry, Algorithms and Applications by M. de Berg 



## Proof of triangulation theorem, continued

- **Claim.** There exist two vertices  $v_i$  and  $v_j$  in  $P$  so that the line segment<sup>5</sup>  $\overline{v_i v_j} \subset P$ .

---

<sup>5</sup>Such a line segment is called a **diagonal** of polygon  $P$ . 

## Proof of triangulation theorem, continued

- ▶ **Claim.** There exist two vertices  $v_i$  and  $v_j$  in  $P$  so that the line segment<sup>5</sup>  $\overline{v_i v_j} \subset P$ .
- ▶ **Here's Why.** Let  $v$  be the vertex in  $P$  with the smallest  $x$ -coordinate (ie, leftmost vertex).

---

<sup>5</sup>Such a line segment is called a **diagonal** of polygon  $P$ . 

## Proof of triangulation theorem, continued

- ▶ **Claim.** There exist two vertices  $v_i$  and  $v_j$  in  $P$  so that the line segment<sup>5</sup>  $\overline{v_i v_j} \subset P$ .
- ▶ **Here's Why.** Let  $v$  be the vertex in  $P$  with the smallest  $x$ -coordinate (ie, leftmost vertex).
- ▶ If it so happens that there are two or more leftmost vertices in  $P$ , then choose the one with the smallest  $y$ -coordinate.

---

<sup>5</sup>Such a line segment is called a **diagonal** of polygon  $P$ . 

## Proof of triangulation theorem, continued


- ▶ **Claim.** There exist two vertices  $v_i$  and  $v_j$  in  $P$  so that the line segment<sup>5</sup>  $\overline{v_i v_j} \subset P$ .
- ▶ **Here's Why.** Let  $v$  be the vertex in  $P$  with the smallest  $x$ -coordinate (ie, leftmost vertex).
- ▶ If it so happens that there are two or more leftmost vertices in  $P$ , then choose the one with the smallest  $y$ -coordinate.
- ▶ Now suppose the two vertices that neighbor  $v$  are  $u$  and  $w$ .

---

<sup>5</sup>Such a line segment is called a **diagonal** of polygon  $P$ .

## Proof of triangulation theorem, continued

- ▶ **Claim.** There exist two vertices  $v_i$  and  $v_j$  in  $P$  so that the line segment<sup>5</sup>  $\overline{v_i v_j} \subset P$ .
- ▶ **Here's Why.** Let  $v$  be the vertex in  $P$  with the smallest  $x$ -coordinate (ie, leftmost vertex).
- ▶ If it so happens that there are two or more leftmost vertices in  $P$ , then choose the one with the smallest  $y$ -coordinate.
- ▶ Now suppose the two vertices that neighbor  $v$  are  $u$  and  $w$ .
- ▶ If  $\overline{uw} \subset P$ , then  $\overline{uw}$  is a diagonal of  $P$ , and we are done.




---

<sup>5</sup>Such a line segment is called a **diagonal** of polygon  $P$ .


## Proof of claim, continued

- **Else** there is at least one vertex of  $P$  contained in triangle  $uvw$ . Call it  $v'$ .



## Proof of claim, continued

- **Else** there is at least one vertex of  $P$  contained in triangle  $uvw$ . Call it  $v'$ .








## Proof of claim, continued


- In fact choose  $v'$  to be the vertex in triangle  $uvw$  that is farthest from the line through  $u$  and  $w$ .



- In that case,  $\overline{vv'} \subset P$ ; if not there is another vertex inside triangle  $uvw$  farther left than  $v'$ , a contradiction.

## Proof of claim, continued


- ▶ In fact choose  $v'$  to be the vertex in triangle  $uvw$  that is farthest from the line through  $u$  and  $w$ .



- ▶ In that case,  $\overline{vv'} \subset P$ ; if not there is another vertex inside triangle  $uvw$  farther left than  $v'$ , a contradiction.
- ▶ Thus  $\overline{vv'}$  is a diagonal of  $P$ , which completes the proof of the claim.

## Proof of claim, continued


- ▶ In fact choose  $v'$  to be the vertex in triangle  $uvw$  that is farthest from the line through  $u$  and  $w$ .



- ▶ In that case,  $\overline{vv'} \subset P$ ; if not there is another vertex inside triangle  $uvw$  farther left than  $v'$ , a contradiction.
- ▶ Thus  $\overline{vv'}$  is a diagonal of  $P$ , which completes the proof of the claim.
- ▶ **Back to the induction argument.**


## Induction, continued

- ▶ Any diagonal of a simple polygon  $P$  splits  $P$  into two simple polygons  $P_1$  and  $P_2$ .




## Induction, continued

- ▶ Any diagonal of a simple polygon  $P$  splits  $P$  into two simple polygons  $P_1$  and  $P_2$ .




## Induction, continued

- If  $P$  has  $n$  vertices, and  $P_1$  has  $m$  vertices, then  $P_2$  has  $n - m + 2$  vertices. Why?



## Induction, continued


- ▶ If  $P$  has  $n$  vertices, and  $P_1$  has  $m$  vertices, then  $P_2$  has  $n - m + 2$  vertices. Why?



- ▶ That is,  $|V(P_1)| + |V(P_1)| = n + 2$ .

## Induction, continued

- ▶ If  $P$  has  $n$  vertices, and  $P_1$  has  $m$  vertices, then  $P_2$  has  $n - m + 2$  vertices. Why?




- ▶ That is,  $|V(P_1)| + |V(P_2)| = n + 2$ .
- ▶ By the induction hypothesis, each of  $P_1$  and  $P_2$  admit triangulations  $\Rightarrow P$  admits a triangulation by glueing  $P_1$  and  $P_2$  back together.



## Induction, continued


- ▶ If  $P$  has  $n$  vertices, and  $P_1$  has  $m$  vertices, then  $P_2$  has  $n - m + 2$  vertices. Why?



- ▶ That is,  $|V(P_1)| + |V(P_2)| = n + 2$ .
- ▶ By the induction hypothesis, each of  $P_1$  and  $P_2$  admit triangulations  $\Rightarrow P$  admits a triangulation by glueing  $P_1$  and  $P_2$  back together.
- ▶ Moreover, by the induction hypothesis,  $P_1$  contains  $n - m$  triangles and  $P_2$  contains  $m - 2$  triangles,

## Induction, continued

- ▶ If  $P$  has  $n$  vertices, and  $P_1$  has  $m$  vertices, then  $P_2$  has  $n - m + 2$  vertices. Why?



- ▶ That is,  $|V(P_1)| + |V(P_2)| = n + 2$ .
- ▶ By the induction hypothesis, each of  $P_1$  and  $P_2$  admit triangulations  $\Rightarrow P$  admits a triangulation by glueing  $P_1$  and  $P_2$  back together.
- ▶ Moreover, by the induction hypothesis,  $P_1$  contains  $n - m$  triangles and  $P_2$  contains  $m - 2$  triangles,
- ▶ which means  $P$  contains  $n - m + m - 2 = n - 2$  triangles, as desired.

# Proof completed

- ▶ This completes the inductive step and thus the proof.

# Proof completed

- ▶ This completes the inductive step and thus the proof.
- ▶ **QED**

# Proof completed

- ▶ This completes the inductive step and thus the proof.
- ▶ **QED**
- ▶ As a consequence of Theorem 3.1, we have shown that an arbitrary simple polygon on  $n$  vertices can be guarded by  $n - 2$  guards.

## Proof completed

- ▶ This completes the inductive step and thus the proof.
- ▶ **QED**
- ▶ As a consequence of Theorem 3.1, we have shown that an arbitrary simple polygon on  $n$  vertices can be guarded by  $n - 2$  guards.

▶ **Can we do better??**

## Some Slight Tweaks for further Improvement

- ▶ Start with simple polygon  $P$  as before, and triangulate it, which we know can be done by Theorem 3.1.

## Some Slight Tweaks for further Improvement

- ▶ Start with simple polygon  $P$  as before, and triangulate it, which we know can be done by Theorem 3.1.
- ▶ **Question.** How many diagonals are there in any triangulation of  $P$ ?



## Some Slight Tweaks for further Improvement

- ▶ Start with simple polygon  $P$  as before, and triangulate it, which we know can be done by Theorem 3.1.
- ▶ **Question.** How many diagonals are there in any triangulation of  $P$ ?
- ▶ **Answer:**  $n - 3$ . Why?

## Some Slight Tweaks for further Improvement

- ▶ Start with simple polygon  $P$  as before, and triangulate it, which we know can be done by Theorem 3.1.
- ▶ **Question.** How many diagonals are there in any triangulation of  $P$ ?
- ▶ **Answer:**  $n - 3$ . Why?
- ▶ Use Euler's formula for embedded planar graphs: our triangulated  $P$  has  $n - 2 + 1$  faces,  $n + d$  edges (where  $d$  is the number of diagonals) and  $n$  vertices.

## Some Slight Tweaks for further Improvement

- ▶ Start with simple polygon  $P$  as before, and triangulate it, which we know can be done by Theorem 3.1.
- ▶ **Question.** How many diagonals are there in any triangulation of  $P$ ?
- ▶ **Answer:**  $n - 3$ . Why?
- ▶ Use Euler's formula for embedded planar graphs: our triangulated  $P$  has  $n - 2 + 1$  faces,  $n + d$  edges (where  $d$  is the number of diagonals) and  $n$  vertices.
- ▶ Thus by Euler's formula, we have
$$n - (n + d) + n - 1 = 2 \Rightarrow d = n - 3.$$

## Some Slight Tweaks for further Improvement

- ▶ Start with simple polygon  $P$  as before, and triangulate it, which we know can be done by Theorem 3.1.
- ▶ **Question.** How many diagonals are there in any triangulation of  $P$ ?
- ▶ **Answer:**  $n - 3$ . Why?
- ▶ Use Euler's formula for embedded planar graphs: our triangulated  $P$  has  $n - 2 + 1$  faces,  $n + d$  edges (where  $d$  is the number of diagonals) and  $n$  vertices.
- ▶ Thus by Euler's formula, we have
$$n - (n + d) + n - 1 = 2 \Rightarrow d = n - 3.$$
- ▶ Slightly Better Result: Put a camera on each diagonal, thus reducing the number of guards from  $n - 2$  to  $n - 3$ .

# Can we do better??

- ▶ **Observation.** Each diagonal of  $P$  “sees” two triangles, so the number of cameras can be reduced to about  $\frac{n}{2}$ .

# Can we do better??

- ▶ **Observation.** Each diagonal of  $P$  “sees” two triangles, so the number of cameras can be reduced to about  $\frac{n}{2}$ .
- ▶ **A new idea.** What about placing cameras on selected vertices instead of selected edges?

# Can we do better??

- ▶ **Observation.** Each diagonal of  $P$  “sees” two triangles, so the number of cameras can be reduced to about  $\frac{n}{2}$ .
- ▶ **A new idea.** What about placing cameras on selected vertices instead of selected edges?
- ▶ **Why should this help?** Each triangle sees *at least* two triangles, maybe more.

# Can we do better??

- ▶ **Observation.** Each diagonal of  $P$  “sees” two triangles, so the number of cameras can be reduced to about  $\frac{n}{2}$ .
- ▶ **A new idea.** What about placing cameras on selected vertices instead of selected edges?
- ▶ **Why should this help?** Each triangle sees *at least* two triangles, maybe more.
- ▶ The following lemma from graph theory is key to our next step towards success.



# Triangulated simple polygon graph

- ▶ **Lemma 1.** A triangulated simple polygon  $P$  has a vertex of degree 2.

## Triangulated simple polygon graph

- ▶ **Lemma 1.** A triangulated simple polygon  $P$  has a vertex of degree 2.
- ▶ **Proof.** Assume BWOC that there exists a triangulated simple polygon on  $n$  vertices that has no vertex of degree 2.

## Triangulated simple polygon graph

- ▶ **Lemma 1.** A triangulated simple polygon  $P$  has a vertex of degree 2.
- ▶ **Proof.** Assume BWOC that there exists a triangulated simple polygon on  $n$  vertices that has no vertex of degree 2.
- ▶ Then every vertex of  $P$  has degree three or more.

## Triangulated simple polygon graph

- ▶ **Lemma 1.** A triangulated simple polygon  $P$  has a vertex of degree 2.
- ▶ **Proof.** Assume BWOC that there exists a triangulated simple polygon on  $n$  vertices that has no vertex of degree 2.
- ▶ Then every vertex of  $P$  has degree three or more.
- ▶ By Theorem 3.1, once  $P$  has been triangulated, it contains  $E = n + n - 3 = 2n - 3$  edges.

# Triangulated simple polygon graph

- ▶ **Lemma 1.** A triangulated simple polygon  $P$  has a vertex of degree 2.
- ▶ **Proof.** Assume BWOC that there exists a triangulated simple polygon on  $n$  vertices that has no vertex of degree 2.
- ▶ Then every vertex of  $P$  has degree three or more.
- ▶ By Theorem 3.1, once  $P$  has been triangulated, it contains  $E = n + n - 3 = 2n - 3$  edges.
- ▶ Altogether we have  $3n \leq 2E \Rightarrow 3n \leq 2(2n - 3) \Rightarrow n \leq 6$ .

# Triangulated simple polygon graph

- ▶ **Lemma 1.** A triangulated simple polygon  $P$  has a vertex of degree 2.
- ▶ **Proof.** Assume BWOC that there exists a triangulated simple polygon on  $n$  vertices that has no vertex of degree 2.
- ▶ Then every vertex of  $P$  has degree three or more.
- ▶ By Theorem 3.1, once  $P$  has been triangulated, it contains  $E = n + n - 3 = 2n - 3$  edges.
- ▶ Altogether we have  $3n \leq 2E \Rightarrow 3n \leq 2(2n - 3) \Rightarrow n \leq 6$ .
- ▶ You can check the cases  $n = 3, 4, 5, 6$  by hand by producing all possible triangulations and show that in all cases, there exists a vertex of degree 2.

# Triangulated simple polygon graph

- ▶ **Lemma 1.** A triangulated simple polygon  $P$  has a vertex of degree 2.
- ▶ **Proof.** Assume BWOC that there exists a triangulated simple polygon on  $n$  vertices that has no vertex of degree 2.
- ▶ Then every vertex of  $P$  has degree three or more.
- ▶ By Theorem 3.1, once  $P$  has been triangulated, it contains  $E = n + n - 3 = 2n - 3$  edges.
- ▶ Altogether we have  $3n \leq 2E \Rightarrow 3n \leq 2(2n - 3) \Rightarrow n \leq 6$ .
- ▶ You can check the cases  $n = 3, 4, 5, 6$  by hand by producing all possible triangulations and show that in all cases, there exists a vertex of degree 2.
- ▶ **Conclusion.** Every simple triangulated polygon has a vertex of degree 2. **QED**

## Help from graph coloring

- **Theorem 5.** Let  $G$  be the graph representing a simple triangulated polygon. Then  $\chi(G) = 3$ .



## Help from graph coloring

- ▶ **Theorem 5.** Let  $G$  be the graph representing a simple triangulated polygon. Then  $\chi(G) = 3$ .
- ▶ **Proof by induction on the number of vertices.** The base case is that the smallest possible simple triangulated polygon is  $K_3$ ; clearly  $\chi(K_3) = 3$ .

## Help from graph coloring

- ▶ **Theorem 5.** Let  $G$  be the graph representing a simple triangulated polygon. Then  $\chi(G) = 3$ .
- ▶ **Proof by induction on the number of vertices.** The base case is that the smallest possible simple triangulated polygon is  $K_3$ ; clearly  $\chi(K_3) = 3$ .
- ▶ **Induction Hypothesis.** For some fixed integer  $k \geq 3$ , the graph  $G$  of any triangulated simple polygon on  $k$  vertices satisfies  $\chi(G) = 3$ .

## Help from graph coloring

- ▶ **Theorem 5.** Let  $G$  be the graph representing a simple triangulated polygon. Then  $\chi(G) = 3$ .
- ▶ **Proof by induction on the number of vertices.** The base case is that the smallest possible simple triangulated polygon is  $K_3$ ; clearly  $\chi(K_3) = 3$ .
- ▶ **Induction Hypothesis.** For some fixed integer  $k \geq 3$ , the graph  $G$  of any triangulated simple polygon on  $k$  vertices satisfies  $\chi(G) = 3$ .
- ▶ **Inductive Step.** We must now prove that the graph  $G$  representing any triangulated simple polygon on  $k + 1$  vertices satisfies  $\chi(G) = 3$ .

## Proof of Theorem 5, continued

- ▶ By Lemma 1,  $G$  contains a vertex of degree 2; call it  $v$ . Consider the graph  $G \setminus v$ , which is a simple triangulated polygon on  $k$  vertices, thus satisfying our induction hypothesis.

## Proof of Theorem 5, continued

- ▶ By Lemma 1,  $G$  contains a vertex of degree 2; call it  $v$ . Consider the graph  $G \setminus v$ , which is a simple triangulated polygon on  $k$  vertices, thus satisfying our induction hypothesis.
- ▶ Thus  $\chi(G \setminus v) = 3$ ; properly vertex color  $G \setminus v$  with three colors  $R$ ,  $G$ , and  $B$ .

## Proof of Theorem 5, continued

- ▶ By Lemma 1,  $G$  contains a vertex of degree 2; call it  $v$ . Consider the graph  $G \setminus v$ , which is a simple triangulated polygon on  $k$  vertices, thus satisfying our induction hypothesis.
- ▶ Thus  $\chi(G \setminus v) = 3$ ; properly vertex color  $G \setminus v$  with three colors  $R$ ,  $G$ , and  $B$ .
- ▶ WLOG assume the two neighbors of  $v$  are colored  $R$  and  $G$ . Then reinstate  $v$  and color it  $B$ .

## Proof of Theorem 5, continued


- ▶ By Lemma 1,  $G$  contains a vertex of degree 2; call it  $v$ . Consider the graph  $G \setminus v$ , which is a simple triangulated polygon on  $k$  vertices, thus satisfying our induction hypothesis.
- ▶ Thus  $\chi(G \setminus v) = 3$ ; properly vertex color  $G \setminus v$  with three colors  $R$ ,  $G$ , and  $B$ .
- ▶ WLOG assume the two neighbors of  $v$  are colored  $R$  and  $G$ . Then reinstate  $v$  and color it  $B$ .
- ▶ So far, we've shown that  $\chi(G) \leq 3$ , but since  $K_3$  is a subgraph of  $G$ , we also have  $\chi(G) \geq 3 \Rightarrow \chi(G) = 3$ .

## Proof of Theorem 5, continued

- ▶ By Lemma 1,  $G$  contains a vertex of degree 2; call it  $v$ . Consider the graph  $G \setminus v$ , which is a simple triangulated polygon on  $k$  vertices, thus satisfying our induction hypothesis.
- ▶ Thus  $\chi(G \setminus v) = 3$ ; properly vertex color  $G \setminus v$  with three colors  $R$ ,  $G$ , and  $B$ .
- ▶ WLOG assume the two neighbors of  $v$  are colored  $R$  and  $G$ . Then reinstate  $v$  and color it  $B$ .
- ▶ So far, we've shown that  $\chi(G) \leq 3$ , but since  $K_3$  is a subgraph of  $G$ , we also have  $\chi(G) \geq 3 \Rightarrow \chi(G) = 3$ .
- ▶ **Conclusion.** If  $G$  is the graph of a simple triangulated polygon, then  $\chi(G) = 3$ .



## 3-coloring a triangulated polygon



## Reducing the number of guards with Theorem 5

- **Theorem 6.** For any simple polygon  $P$  on  $n$  vertices, there exists a cover with  $\lfloor \frac{n}{3} \rfloor$  guards.

## Reducing the number of guards with Theorem 5

- ▶ **Theorem 6.** For any simple polygon  $P$  on  $n$  vertices, there exists a cover with  $\lfloor \frac{n}{3} \rfloor$  guards.
- ▶ **Remark.** Theorem 6 is way better than our previous best  $n - 3$  guards!!

# Reducing the number of guards with Theorem 5

- ▶ **Theorem 6.** For any simple polygon  $P$  on  $n$  vertices, there exists a cover with  $\lfloor \frac{n}{3} \rfloor$  guards.
- ▶ **Remark.** Theorem 6 is way better than our previous best  $n - 3$  guards!!
- ▶ **Proof.** By Theorem 5, we can 3-color the vertices of  $P$  after it has been triangulated. Do so.

# Reducing the number of guards with Theorem 5

- ▶ **Theorem 6.** For any simple polygon  $P$  on  $n$  vertices, there exists a cover with  $\lfloor \frac{n}{3} \rfloor$  guards.
- ▶ **Remark.** Theorem 6 is way better than our previous best  $n - 3$  guards!!
- ▶ **Proof.** By Theorem 5, we can 3-color the vertices of  $P$  after it has been triangulated. Do so.
- ▶ Let  $C_R$  be the set of vertices colored red,  $C_G$  be the set of vertices colored green, and  $C_B$  be the set of vertices colored blue.

# Reducing the number of guards with Theorem 5

- ▶ **Theorem 6.** For any simple polygon  $P$  on  $n$  vertices, there exists a cover with  $\lfloor \frac{n}{3} \rfloor$  guards.
- ▶ **Remark.** Theorem 6 is way better than our previous best  $n - 3$  guards!!
- ▶ **Proof.** By Theorem 5, we can 3-color the vertices of  $P$  after it has been triangulated. Do so.
- ▶ Let  $C_R$  be the set of vertices colored red,  $C_G$  be the set of vertices colored green, and  $C_B$  be the set of vertices colored blue.
- ▶ **Fact.**  $|C_R| + |C_G| + |C_B| = n$ . Why?

$$|C_R| + |C_G| + |C_B| = n$$

- By algebra, if  $n$  objects are distributed among three sets, at least one of those sets contains no more than  $\lfloor \frac{n}{3} \rfloor$  objects.

$$|C_R| + |C_G| + |C_B| = n$$

- ▶ By algebra, if  $n$  objects are distributed among three sets, at least one of those sets contains no more than  $\lfloor \frac{n}{3} \rfloor$  objects.
- ▶ Thus one of the three color classes of vertices, say WLOG  $C_R$ , contains at most  $\lfloor \frac{n}{3} \rfloor$  vertices.



$$|C_R| + |C_G| + |C_B| = n$$

- ▶ By algebra, if  $n$  objects are distributed among three sets, at least one of those sets contains no more than  $\lfloor \frac{n}{3} \rfloor$  objects.
- ▶ Thus one of the three color classes of vertices, say WLOG  $C_R$ , contains at most  $\lfloor \frac{n}{3} \rfloor$  vertices.
- ▶ Place a guard on each red vertex; since  $P$  is triangulated, every triangle is 3-colored, and thus  $P$  is covered.

$$|C_R| + |C_G| + |C_B| = n$$

- ▶ By algebra, if  $n$  objects are distributed among three sets, at least one of those sets contains no more than  $\lfloor \frac{n}{3} \rfloor$  objects.
- ▶ Thus one of the three color classes of vertices, say WLOG  $C_R$ , contains at most  $\lfloor \frac{n}{3} \rfloor$  vertices.
- ▶ Place a guard on each red vertex; since  $P$  is triangulated, every triangle is 3-colored, and thus  $P$  is covered.
- ▶ **QED**

# Can we do better??

NO

## Exercise

- ▶ Construct a polygon on  $n$  vertices for any  $n \geq 3$  that requires a cover of  $\lfloor \frac{n}{3} \rfloor$  vertices, thus proving that Theorem 6 is best possible.

## Exercise

- ▶ Construct a polygon on  $n$  vertices for any  $n \geq 3$  that requires a cover of  $\lfloor \frac{n}{3} \rfloor$  vertices, thus proving that Theorem 6 is best possible.
- ▶ A useful resource about Art Gallery Theorems:  
`http://cs.smith.edu/~orourke/books/ArtGalleryTheorems/art.html`

## Exercise

- ▶ Construct a polygon on  $n$  vertices for any  $n \geq 3$  that requires a cover of  $\lfloor \frac{n}{3} \rfloor$  vertices, thus proving that Theorem 6 is best possible.
- ▶ A useful resource about Art Gallery Theorems:  
`http://cs.smith.edu/~orourke/books/ArtGalleryTheorems/art.html`
- ▶ **Next.** Speaking of Art, we'll look at a project inspired by M.C. Escher's Artwork.