

Assignment Five

ECE 4200


- Provide credit to **any sources** other than the course staff that helped you solve the problems. This includes **all students** you talked to regarding the problems.
- You can look up definitions/basics online (e.g., wikipedia, stack-exchange, etc).
- **The due date is 3/8/2020, 23.59.59 ET.**
- Submission rules are the same as previous assignments.
- **Please write your net-id on top of every page. It helps with grading.**

Problem 1. (15 points). SVM's obtain *non-linear* decision boundaries by mapping the feature vectors $\vec{X} \in \mathbb{R}^d$ to a possibly high dimensional space via a function $\phi : \mathbb{R}^d \rightarrow \mathbb{R}^m$, and then finding a linear decision boundary in the new space.

We also saw that to implement SVM, it suffices to know the kernel function $K(\vec{X}_i, \vec{X}_j) = \phi(\vec{X}_i) \cdot \phi(\vec{X}_j)$, without even explicitly specifying the function ϕ .

Recall **Mercer's theorem**. K is a kernel function if and only if for any n vectors, $\vec{X}_1, \dots, \vec{X}_n \in \mathbb{R}^d$, and **any** real numbers c_1, \dots, c_n , $\sum_{i=1}^n \sum_{j=1}^n c_i c_j K(\vec{X}_i, \vec{X}_j) \geq 0$.

1. Prove the following half of Mercer's theorem (which we showed in class). If K is a kernel then $\sum_{i=1}^n \sum_{j=1}^n c_i c_j K(\vec{X}_i, \vec{X}_j) \geq 0$.
2. Let $d = 1$, and $x, y \in \mathbb{R}$. Is the function $K(x, y) = x + y$ a kernel?
3. Let $d = 1$, and $x, y \in \mathbb{R}$. Is $K(x, y) = xy + 1$ a kernel?
4. Suppose $d = 2$, namely the original features are of the form $\vec{X}_i = [\vec{X}_i^1, \vec{X}_i^2]$. Show that $K(\vec{X}, \vec{Y}) = (1 + \vec{X} \cdot \vec{Y})^2$ is a kernel function. This is called as **quadratic kernel**.
(**Hint:** Find a $\phi : \mathbb{R}^2 \rightarrow \mathbb{R}^m$ (for some m) such that $\phi(\vec{X}) \cdot \phi(\vec{Y}) = (1 + \vec{X} \cdot \vec{Y})^2$).
5. Consider the training examples $\langle [0, 1], 1 \rangle, \langle [1, 2], 1 \rangle, \langle [-1, 2], 1 \rangle, \langle [0, 11], 1 \rangle, \langle [3, 4], -1 \rangle, \langle [-3, 4], -1 \rangle, \langle [1, -1], -1 \rangle, \langle [-1, -1], -1 \rangle$. We have plotted the data points below.
 - Is the data **linearly classifiable** in the original 2-d space? If yes, please come up with *any* linear decision boundary that separates the data. If no, please explain why.
 - Is the data linearly classifiable in the feature space corresponding to the quadratic kernel. If yes, please come up with *any* linear decision boundary that separates the data. If no, please explain why.



Problem 2. (10 points). The Gaussian kernel (also called Radial Basis Function kernel (RBF)) is:

$$K(\vec{X}, \vec{Y}) = \exp\left(-\frac{\|\vec{X} - \vec{Y}\|_2^2}{2\sigma^2}\right),$$

where \vec{X}, \vec{Y} are feature vectors in d dimensions. Suppose $d = 1$, and $2\sigma^2 = 1$.

1. Design a function $\phi : \mathbb{R} \rightarrow \mathbb{R}^m$ that corresponds to Gaussian kernel for $d = 1$, and $2\sigma^2 = 1$.

Hint: Use Taylor series expansion for the exponential function.

2. What is the value of m you end up with?

Problem 3. (10 points). Let $f, h_i, 1 \leq i \leq n$ be real-valued functions and let $\alpha \in \mathbb{R}^n$. Let $L(z, \alpha) = f(z) + \sum_{i=1}^n \alpha_i h_i(z)$. In this problem, we will prove that the following two optimization problems are equivalent.

$$\min_z f(z) \quad (1)$$

$$\text{s.t. } h_i(z) \leq 0, i = 1, \dots, n.$$

$$\min_z \max_{\alpha \geq \mathbf{0}} L(z, \alpha) \quad (2)$$

Let (z^*, α^*) be the solution of (2) and let z_p^* be the solution of (1). Prove that:

$$L(z^*, \alpha^*) = f(z_p^*)$$

Hint: Use the fact that for any $z, \alpha \geq \mathbf{0}$, $L(z^*, \alpha^*) \geq L(z^*, \alpha)$ and $L(z^*, \alpha^*) \leq L(z, \alpha_z)$, where $\alpha_z = \arg \max_{\alpha \geq \mathbf{0}} L(z, \alpha)$.

You may follow the following steps but it is not required as long as your proof is correct.

1. Prove that $L(z^*, \alpha^*) \leq f(z_p^*)$

2. Prove that $L(z^*, \alpha^*) \geq f(z_p^*)$

Problem 4 (25 points) SVM Classification. Please refer to the Jupyter Notebook in the assignment, and complete the coding part in it! You can use sklearn SVM package: <https://scikit-learn.org/stable/modules/generated/sklearn.svm.SVC.html#sklearn.svm.SVC>