

# Simple R Functions

*Senhao Li*

*January 26, 2018*

1.

- (a) Write functions `tmpFn1` and `tmpFn2` such that if `xVec` is the vector  $(x_1, x_2, \dots, x_n)$ , then `tmpFn1(xVec)` returns vector  $(x_1, x_2^2, \dots, x_n^n)$  and `tmpFn2(xVec)` returns the vector  $(x_1, \frac{x_2^2}{2}, \dots, \frac{x_n^n}{n})$ .

---

Here is `tmpFn1`

```
tmpFn1 <- function(xVec){  
  return(xVec^(1:length(xVec)))  
}
```

## simple example

```
a <- c(2, 5, 3, 8, 2, 4)
```

```
b <- tmpFn1(a)
```

```
b
```

```
## [1]      2    25    27 4096    32 4096
```

and now `tmpFn2`

```
tmpFn2 <- function(xVec2){  
  
  n = length(xVec2)  
  
  return(xVec2^(1:n)/(1:n))  
}
```

```
c <- tmpFn2(a)
```

```
c
```

```
## [1]      2.0000    12.5000     9.0000 1024.0000     6.4000  682.6667
```

- (b) Now write a function `tmpFn3` which takes 2 arguments  $x$  and  $n$  where  $x$  is a single number and  $n$  is a strictly positive integer. The function should return the value of

$$1 + \frac{x}{1} + \frac{x^2}{2} + \frac{x^3}{3} + \dots + \frac{x^n}{n}$$

2. Write a function `tmpFn(xVec)` such that if `xVec` is the vector  $x = (x_1, \dots, x_n)$  then `tmpFn(xVec)` returns the vector of moving averages:

$$\frac{x_1 + x_2 + x_3}{3}, \frac{x_2 + x_3 + x_4}{3}, \dots, \frac{x_{n-2} + x_{n-1} + x_n}{3}$$

Try out your function. `tmpFn(c(1:5,6:1))`

```
tmpFn <- function(xVec){
  return((xVec[-c(length(xVec),length(xVec)-1)]+xVec[-c(1,length(xVec))]+xVec[-c(1,2)]))/3)
}
tmpFn(c(1:5,6:1))
```

```
## [1] 2.000000 3.000000 4.000000 5.000000 5.333333 5.000000 4.000000 3.000000
## [9] 2.000000
```

3. Consider the continuous function


$$f(x) = \begin{cases} x^2 + 2x + 3 & \text{if } x < 0 \\ x + 3 & \text{if } 0 \leq x < 2 \\ x^2 + 4x - 7 & \text{if } 2 \leq x \end{cases}$$

Write a function `tmpFn` which takes a single argument `xVec`. the function should return the vector the values of the function  $f(x)$  evaluated at the values in `xVec`.

Hence plot the function  $f(x)$  for  $-3 < x < 3$ .

```
tmpFn_3 <- function(x){
  ifelse(x < 0, x^2+2*x+3, ifelse(x < 2, x+3, x^2+4*x-7))
}

w <- seq(-3,3, len=1000)
plot(w, tmpFn_3(w), "l")
```



4. Write a function which takes a single argument which is a matrix. The function should return a matrix which is the same as the function argument but every odd number is doubled.

Hence the result of using the function on the matrix

$$\begin{bmatrix} 1 & 1 & 3 \\ 5 & 2 & 6 \\ -2 & -1 & -3 \end{bmatrix}$$

should be:

$$\begin{bmatrix} 2 & 2 & 6 \\ 10 & 2 & 6 \\ -2 & -2 & -6 \end{bmatrix}$$

```
tmpFn_4 <- function(mat){
  mat[mat%%2 == 1] <- 2 * mat[mat%%2 == 1]
  return(mat)
}
mat_4 <- matrix(c(1,1,3,5,2,6,-2,-1,-3),nr=3,byrow=T)
tmpFn_4(mat_4)
```

```
##      [,1] [,2] [,3]
## [1,]    2    2    6
## [2,]   10    2    6
## [3,]   -2   -2   -6
```

5. Write a function which takes 2 arguments  $n$  and  $k$  which are positive integers. It should return the  $n \times n$  matrix:

$$\begin{bmatrix} k & 1 & 0 & 0 & \cdots & 0 & 0 \\ 1 & k & 1 & 0 & \cdots & 0 & 0 \\ 0 & 1 & k & 1 & \cdots & 0 & 0 \\ 0 & 0 & 1 & k & \cdots & 0 & 0 \\ \vdots & \vdots & \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & 0 & 0 & \cdots & k & 1 \\ 0 & 0 & 0 & 0 & \cdots & 1 & k \end{bmatrix}$$

```
tmpFn_5 <- function(k,n){
  mat <- diag(k, nr=n)
  mat[abs(row(mat)-col(mat))==1] <- 1
  return(mat)
}
tmpFn_5(10,5)
```

```
##      [,1] [,2] [,3] [,4] [,5]
## [1,]   10    1    0    0    0
## [2,]    1   10    1    0    0
## [3,]    0    1   10    1    0
## [4,]    0    0    1   10    1
## [5,]    0    0    0    1   10
```

6. Suppose an angle  $\alpha$  is given as a positive real number of degrees.

If  $0 \leq \alpha < 90$  then it is quadrant 1. If  $90 \leq \alpha < 180$  then it is quadrant 2.

if  $180 \leq \alpha < 270$  then it is quadrant 3. if  $270 \leq \alpha < 360$  then it is quadrant 4.

if  $360 \leq \alpha < 450$  then it is quadrant 1.

And so on ...

Write a function `quadrant(alpha)` which returns the quadrant of the angle  $\alpha$ .

```
quadrant <- function(alpha){
  alpha%%360%%90+1
}
quadrant(390)
```

```
## [1] 1
```

7.

(a) Zeller's congruence is the formula:

$$f = ([2.6m - 0.2] + k + y + [y/4] + [c/4] - 2c) \bmod 7$$

where  $[x]$  denotes the integer part of  $x$ ; for example  $[7.5] = 7$ .

Zeller's congruence returns the day of the week  $f$  given:

$k$  = the day of the month

$y$  = the year in the century

$c$  = the first 2 digits of the year (the century number)

$m$  = the month number (where January is month 11 of the preceding year, February is month 12 of the preceding year, March is month 1, etc.)

For example, the date 21/07/1963 has  $m = 5, k = 21, c = 19, y = 63$ ;

the date 21/2/63 has  $m = 12, k = 21, c = 19, y = 62$ .

Write a function `weekday(day, month, year)` which returns the day of the week when given the numerical inputs of the day, month and year.

Note that the value of 1 for  $f$  denotes Sunday, 2 denotes Monday, etc.

```
weekday <- function(month, day, year) {
  month <- month - 2
  if(month <= 0) {
    month <- month + 12
    year <- year - 1
  }
  cen <- year %/% 100
  year <- year %/% 100
  tmp <- floor(2.6*month - 0.2) + day + year + year %/% 4 + cen %/% 4 - 2 * cen
  c("Sunday", "Monday", "Tuesday", "Wednesday", "Thursday", "Friday", "Saturday")[1+tmp%%7]
}
weekday(2, 12, 2018)
```

```
## [1] "Monday"
```

(b) Does your function work if the input parameters day, month, and year are vectors with the same length and valid entries?