

Assignment 1

Deriving the implicit formula for the cylinder

Ding Markus

Mamie Robin

Montial Charline

March 6, 2019

To get the implicit equation, the following one needs to be solved:

$$\|x - c'\| - r = 0$$

where x is the intersection of the ray and the cylinder, c' is the projection of this point onto the axis of the cylinder and r its radius.

To compute c' , let us define \vec{a} as its normalized axis, c as its center and \vec{b} as the vector pointing from c to x - i.e. $\vec{b} = x - c$. By computing the dot product between $\vec{b} \cdot \vec{a}$, we get the distance between c and c' since \vec{a} is normal - it's an orthogonal projection. We get c' by using the following:

$$c + (\vec{b} \cdot \vec{a}) \vec{a}$$

The fact that \vec{a} is already normalized is to be kept in mind. We finally get:

$$\|x - c - (\vec{b} \cdot \vec{a}) \vec{a}\| - r = 0$$

$$\begin{aligned} 0 &= \|x - c - (\vec{b} \cdot \vec{a}) \vec{a}\| - r \\ &= \left(x - c - (\vec{b} \cdot \vec{a}) \vec{a}\right)^2 - r^2 \end{aligned}$$

Now, let us replace x with the explicit ray equation $o + t\vec{d}$, where o is its origin point and \vec{d} its vector, and \vec{b} with its definition.

$$\begin{aligned} 0 &= \left(o + t\vec{d} - c - ((o + t\vec{d} - c) \cdot \vec{a}) \vec{a}\right)^2 - r^2 \\ &= \left(t\vec{d} + o - c - ((o + t\vec{d} - c) \cdot \vec{a}) \vec{a}\right)^2 - r^2 \\ &= \left(t\vec{d} + o - c - (t\vec{d} \cdot \vec{a}) \vec{a} - ((o - c) \cdot \vec{a}) \vec{a}\right)^2 - r^2 \\ &= \left(t(\vec{d} - (\vec{d} \cdot \vec{a}) \vec{a}) + o - c - ((o - c) \cdot \vec{a}) \vec{a}\right)^2 - r^2 \end{aligned}$$

Now, let's define $x = \vec{d} - (\vec{d} \cdot \vec{a}) \vec{a}$ and $y = o - c - ((o - c) \cdot \vec{a}) \vec{a}$. We thus get the following:

$$\begin{aligned} 0 &= (tx + y)^2 - r^2 \\ &= t^2 x^2 + t 2xy + y^2 - r^2 \end{aligned}$$

Let's remember that a vector squared is the result of applying the scalar product on itself. We finally get as values for our quadratic function:

$$\begin{aligned} A &= x^2 = \left(\vec{d} - (\vec{d} \cdot \vec{a}) \vec{a} \right) \cdot \left(\vec{d} - (\vec{d} \cdot \vec{a}) \vec{a} \right) \\ B &= 2xy = 2 \left(\vec{d} - (\vec{d} \cdot \vec{a}) \vec{a} \right) \cdot (o - c - ((o - c) \cdot \vec{a}) \vec{a}) \\ C &= y^2 - r^2 = (o - c - ((o - c) \cdot \vec{a}) \vec{a}) \cdot (o - c - ((o - c) \cdot \vec{a}) \vec{a}) - r^2. \end{aligned}$$