
Poisson Regression or Regression of Counts (& Rates)

Carolyn J. Anderson

Department of Educational Psychology

University of Illinois at Urbana-Champaign

Outline

Overview

● Outline

● GLMs for count data

● More Examples...

Poisson regression for counts

Poisson regression for rates

- GLMs for count data.
 - ◆ Poisson regression for counts.
 - ◆ Poisson regression for rates.
- Inference and model checking.
 - ◆ Wald, Likelihood ratio, & Score test.
 - ◆ Checking Poisson regression.
 - ◆ Residuals.
 - ◆ Confidence intervals for fitted values (means).
 - ◆ Overdispersion.
- Fitting GLMS (a little technical).
 - ◆ Newton-Raphson algorithm/Fisher scoring.
 - ◆ Statistic inference & the Likelihood function.
 - ◆ “Deviance”.
- Summary

GLMs for count data

Situation: response/outcome variable Y is a count.

Generalized linear models for counts have as it's random component **Poisson Distribution**.

Examples:

- Number of cargo ships damaged by waves (classic example given by McCullagh & Nelder, 1989).
- Number of deaths due to AIDs in Australia per quarter (3 month periods) from January 1983 – June 1986.
- Number of violent incidents exhibited over a 6 month period by patients who had been treated in the ER of a psychiatric hospital (Gardner, Mulvey, & Shaw, 1995).
- Daily homicide counts in California (Grogger, 1990).
- Foundings of day care centers in Toronto (Baum & Oliver, 1992).
- Political party switching among members of the US House of Representatives (King, 1988).

Overview

● Outline

● GLMs for count data

● More Examples...

Poisson regression for counts

Poisson regression for rates

More Examples...

Overview

● Outline

● GLMs for count data

● More Examples...

Poisson regression for counts

Poisson regression for rates

- Number of presidential appointments to the Supreme Court (King, 1987).
- Number of children in a classroom that a child lists as being their friend (unlimited nomination procedure, sociometric data).
- Number of hard disk failures at uiuc during a year.
- Number of deaths due to SARs (Yu, Chan & Fung, 2006).
- Number of arrests resulting from 911 calls.
- Number of orders of protection issued.

In some of these examples, we should consider “exposure” to the event. i.e., “ t ”.

e.g., hard disk failures: In this case, “exposure” could be the number of hours of operation. Rather than model the number of failures (i.e., counts), we would want to measure and model the failure “rate”

$$Y/t = \text{rate}$$

Poisson regression for counts

Overview

Poisson regression for counts

- Poisson regression for counts
- Components of GLM for Counts

- The Poisson loglinear model
- Interpretation of β
- Interpretation of β (continued)
- Example: Number of Deaths Due to AIDs
- Data: Number of Deaths Due to AIDs \times Month
- A Linear Model for AIDs Data
- A Look at the Bad Model

- (linear link)
- Back to Data but Plot $\log(y_i)$ by Month
- Poisson Log-Linear Model for Deaths
- Figure of Fitted log(count) from Log-linear
- Figure of Fitted log(count) from Log-linear
- Transform explanatory variable
- Poisson Regression with Transformed x
- How Large is Large in a Statistical Sense?
- Observed and Fitted Log(Counts)

- Observed and Fitted Counts
- Comparison of Fitted Counts
- Generalized Linear Models
- Observed and Fitted Counts

Response Variable is a count

Explanatory Variable(s):

- If they are categorical (i.e., you have a contingency table with counts in the cells), convention is to call them “**Loglinear models**”.
- If they are numerical/continuous, convention is to call them “**Poisson Regression**”

First, Y = count and then Y/t rate data.

Comonents of GLM for Counts

Overview

Poisson regression for counts

- Poisson regression for counts
- Comonents of GLM for Counts
- The Poisson loglinear model
- Interpretation of β
- Interpretation of β (continued)
- Example: Number of Deaths Due to AIDs
- Data: Number of Deaths Due to AIDs \times Month
- A Linear Model for AIDs Data
- A Look at the Bad Model (linear link)
- Back to Data but Plot $\log(y_i)$ by Month
- Poisson Log-Linear Model for Deaths
- Figure of Fitted $\log(\text{count})$ from Log-linear
- Figure of Fitted $\log(\text{count})$ from Log-linear
- Transform explanatory variable
- Poisson Regression with Transformed x
- How Large is Large in a Statistical Sense?
- Observed and Fitted Log(Counts)

- Observed and Fitted Counts
- Comparison of Fitted Counts
- Comparing Linear Models
- Observed and Fitted Counts

- **Random component:** Poisson distribution and model the expected value of Y , denoted by $E(Y) = \mu$.
- **Systematic component:** For now, just 1 explanatory variable x (later, we'll go over an example with more than 1).
- **Link:** We could use
 - ◆ Identity link, which gives us $\mu = \alpha + \beta x$
Problem: a linear model can yield $\mu < 0$, while the possible values for $\mu \geq 0$.
 - ◆ Log link (much more common) $\log(\mu)$.
which is the “natural parameter” of Poisson distribution, and the log link is the “canonical link” for GLMs with Poisson distribution.

The Poisson regression model for counts (with a log link) is

$$\log(\mu) = \alpha + \beta x$$

This is often referred to as “**Poisson loglinear model**”.

The Poisson loglinear model

Overview

Poisson regression for counts

- Poisson regression for counts
- Components of GLM for

Counts

- The Poisson loglinear model
- Interpretation of β
- Interpretation of β

(continued)

- Example: Number of Deaths Due to AIDs
- Data: Number of Deaths Due to AIDs \times Month

- A Linear Model for AIDs Data
- A Look at the Bad Model

(linear link)

- Back to Data but Plot

$\log(y_i)$ by Month

- Poisson Log-Linear Model for Deaths

- Figure of Fitted $\log(\text{count})$ from Log-linear

- Figure of Fitted $\log(\text{count})$ from Log-linear

- Transform explanatory variable

- Poisson Regression with Transformed x

- How Large is Large in a Statistical Sense?

- Observed and Fitted $\log(\text{Counts})$

- Observed and Fitted Counts

- Comparison of Fitted Counts

- Generalized Linear Models

- Observed and Fitted Counts

$$\log(\mu) = \alpha + \beta x$$

Since the log of the expected value of Y is a linear function of explanatory variable(s), and the expected value of Y is a multiplicative function of x :

$$\begin{aligned}\mu &= \exp(\alpha + \beta x) \\ &= e^\alpha e^{\beta x}\end{aligned}$$

What does this mean for μ ?

How do we interpret β ?

Interpretation of β

Overview

Poisson regression for counts

- Poisson regression for counts
- Components of GLM for Counts

- The Poisson loglinear model
- Interpretation of β
- Interpretation of β

(continued)

- Example: Number of Deaths Due to AIDs
- Data: Number of Deaths Due to AIDs \times Month

- A Linear Model for AIDs Data
- A Look at the Bad Model

(linear link)

- Back to Data but Plot

$\log(y_i)$ by Month

- Poisson Log-Linear Model for Deaths

- Figure of Fitted $\log(\text{count})$ from Log-linear

- Figure of Fitted $\log(\text{count})$ from Log-linear

- Transform explanatory variable

- Poisson Regression with Transformed x

- How Large is Large in a Statistical Sense?

- Observed and Fitted

Log(Counts)

- Observed and Fitted Counts
- Comparison of Fitted Counts

- Comparison of Fitted Counts
- Comparison of Fitted Counts

- Comparison of Fitted Counts

- Observed and Fitted Counts

$$\log(\mu) = \alpha + \beta x$$

Consider 2 values of x (x_1 & x_2) such that the difference between them equals 1. For example, $x_1 = 10$ and $x_2 = 11$:

$$x_2 = x_1 + 1$$

The expected value of μ when $x = 10$ is

$$\mu_1 = e^{\alpha} e^{\beta x_1} = e^{\alpha} e^{\beta(10)}$$

The expected value of μ when $x = x_2 = 11$ is

$$\begin{aligned} \mu_2 &= e^{\alpha} e^{\beta x_2} \\ &= e^{\alpha} e^{\beta(x_1+1)} \\ &= e^{\alpha} e^{\beta x_1} e^{\beta} \\ &= e^{\alpha} e^{\beta(10)} e^{\beta} \end{aligned}$$

A change in x has a multiplicative effect on the mean of Y .

Interpretation of β (continued)

When we look at a 1 unit increase in the explanatory variable (i.e., $x_2 - x_1 = 1$), we have

$$\mu_1 = e^\alpha e^{\beta x_1} \quad \text{and} \quad \mu_2 = e^\alpha e^{\beta x_1} e^\beta$$

- If $\beta = 0$, then $e^0 = 1$ and
 - ◆ $\mu_1 = e^\alpha$.
 - ◆ $\mu_2 = e^\alpha$.
 - ◆ $\mu = E(Y)$ is not related to x .
- If $\beta > 0$, then $e^\beta > 1$ and
 - ◆ $\mu_1 = e^\alpha e^{\beta x_1}$
 - ◆ $\mu_2 = e^\alpha e^{\beta x_2} = e^\alpha e^{\beta x_1} e^\beta = \mu_1 e^\beta$
 - ◆ μ_2 is e^β times larger than μ_1 .
- If $\beta < 0$, then $0 \leq e^\beta < 1$
 - ◆ $\mu_1 = e^\alpha e^{\beta x_1}$.
 - ◆ $\mu_2 = e^\alpha e^{\beta x_2} = e^\alpha e^{\beta x_1} e^\beta = \mu_1 e^\beta$.
 - ◆ μ_2 is e^β times smaller than μ_1 .

Overview

Poisson regression for counts

● Poisson regression for counts

● Components of GLM for

Counts

● The Poisson loglinear model

● Interpretation of β

● Interpretation of β

(continued)

● Example: Number of Deaths

Due to AIDs

● Data: Number of Deaths Due to AIDs \times Month

● A Linear Model for AIDs Data

● A Look at the Bad Model

(linear link)

● Back to Data but Plot

$\log(y_i)$ by Month

● Poisson Log-Linear Model for

Deaths

● Figure of Fitted $\log(\text{count})$

from Log-linear

● Figure of Fitted $\log(\text{count})$

from Log-linear

● Transform explanatory

variable

● Poisson Regression with

Transformed x

● How Large is Large in a

Statistical Sense?

● Observed and Fitted

$\log(\text{Counts})$

● Observed and Fitted Counts

● Comparison of Fitted Counts

● Comparing Linear Models

● Observed and Fitted Counts

Example: Number of Deaths Due to AIDs

Whyte, et al 1987 (Dobson, 1990) reported the number of deaths due to AIDS in Australia per 3 month period from January 1983 – June 1986.

y_i = number of deaths
 x_i = time point (quarter)

x_i	y_i	x_i	y_i
1	0	8	18
2	1	9	23
3	2	10	31
4	3	11	20
5	1	12	25
6	4	13	37
7	9	14	45

Data: Number of Deaths Due to AIDs × Month

Overview

Poisson regression for counts

● Poisson regression for counts

● Components of GLM for

Counts

● The Poisson loglinear model

● Interpretation of β

● Interpretation of β

(continued)

● Example: Number of Deaths

Due to AIDs

● Data: Number of Deaths Due to AIDs × Month

● A Linear Model for AIDs Data

● A Look at the Bad Model

(linear link)

● Back to Data but Plot

$\log(y_i)$ by Month

● Poisson Log-Linear Model for

Deaths

● Figure of Fitted $\log(\text{count})$

from Log-linear

● Figure of Fitted $\log(\text{count})$

from Log-linear

● Transform explanatory

variable

● Poisson Regression with

Transformed x

● How Large is Large in a

Statistical Sense?

● Observed and Fitted


Log(Counts)

● Observed and Fitted Counts

● Comparison of Fitted Counts

● Comparison of Fitted Counts

● Observed and Fitted Counts



A Linear Model for AIDs Data

Let's try a linear model:

$$\mu_i = \alpha + \beta x_i$$

The estimated parameters from GLM with a Poisson distribution and the identity link:

$$\hat{\mu}_i = -6.7355 + 2.4287x_i$$

In **SAS OUTPUT**, there's *strange* things such as

- Standard errors for estimated parameters equal to 0.
- Some 0's in the OBSTATS.

From **SAS LOG** file...

WARNING: The specified model did not converge.

ERROR: The mean parameter is either invalid or at a limit of its range for some observations.

What's wrong?...

Overview

Poisson regression for counts

● Poisson regression for counts

● Components of GLM for Counts

● The Poisson loglinear model

● Interpretation of β

● Interpretation of β

(continued)

● Example: Number of Deaths Due to AIDs

● Data: Number of Deaths Due to AIDs \times Month

● A Linear Model for AIDs Data

● A Look at the Bad Model

(linear link)

● Back to Data but Plot

$\log(y_i)$ by Month

● Poisson Log-Linear Model for Deaths

● Figure of Fitted $\log(\text{count})$ from Log-linear

● Figure of Fitted $\log(\text{count})$ from Log-linear

● Transform explanatory variable

● Poisson Regression with Transformed x

● How Large is Large in a Statistical Sense?

● Observed and Fitted Log(Counts)

● Observed and Fitted Counts

● Comparison of Fitted Counts

● Comparing Linear Models

● Observed and Fitted Counts

A Look at the Bad Model (linear link)

Overview

Poisson regression for counts

● Poisson regression for counts

● Components of GLM for Counts

● The Poisson loglinear model

● Interpretation of β

● Interpretation of β

(continued)

● Example: Number of Deaths Due to AIDs

● Data: Number of Deaths Due to AIDs \times Month

● A Linear Model for AIDs Data

● A Look at the Bad Model

(linear link)

● Back to Data but Plot

$\log(y_i)$ by Month

● Poisson Log-Linear Model for Deaths

● Figure of Fitted $\log(\text{count})$ from Log-linear

● Figure of Fitted $\log(\text{count})$ from Log-linear

● Transform explanatory variable

● Poisson Regression with Transformed x

● How Large is Large in a Statistical Sense?


● Observed and Fitted Log(Counts)

● Observed and Fitted Counts

● Comparison of Fitted Counts

● Comparison of Linear Models

● Observed and Fitted Counts



Back to Data but Plot $\log(y_i)$ by Month

Overview

Poisson regression for counts

● Poisson regression for counts

● Components of GLM for Counts

● The Poisson loglinear model

● Interpretation of β

● Interpretation of β

(continued)

● Example: Number of Deaths Due to AIDs

● Data: Number of Deaths Due to AIDs \times Month

● A Linear Model for AIDs Data

● A Look at the Bad Model

(linear link)

● Back to Data but Plot

$\log(y_i)$ by Month

● Poisson Log-Linear Model for Deaths

● Figure of Fitted $\log(\text{count})$ from Log-linear

● Figure of Fitted $\log(\text{count})$ from Log-linear

● Transform explanatory variable

● Poisson Regression with Transformed x

● How Large is Large in a Statistical Sense?


● Observed and Fitted Log(Counts)

● Observed and Fitted Counts

● Comparison of Fitted Counts

● Comparison of Linear Models

● Observed and Fitted Counts



(line is linear regression line)

Poisson Log-Linear Model for Deaths

Figure suggests a **log link** might work better:

$$\log(\hat{\mu}_i) = .3396 + .2565x_i$$

Overview

Poisson regression for counts

● Poisson regression for counts

● Components of GLM for Counts

- The Poisson loglinear model
- Interpretation of β
- Interpretation of β
- (continued)
- Example: Number of Deaths Due to AIDs
- Data: Number of Deaths Due to AIDs \times Month
- A Linear Model for AIDs Data
- A Look at the Bad Model (linear link)
- Back to Data but Plot $\log(y_i)$ by Month
- Poisson Log-Linear Model for Deaths
- Figure of Fitted log(count) from Log-linear
- Figure of Fitted log(count) from Log-linear
- Transform explanatory variable
- Poisson Regression with Transformed x
- How Large is Large in a Statistical Sense?
- Observed and Fitted Log(Counts)


- Observed and Fitted Counts
- Comparison of Fitted Counts
- Comparison of Fitted Counts
- Comparison of Fitted Counts
- Observed and Fitted Counts

$\hat{\mu}_i$ when Link is				$\hat{\mu}_i$ when Link is			
x_i	y_i	Log	Identity	x_i	y_i	Log	Identity
1	0	1.82	-4.21	8	18	10.93	12.69
2	1	2.35	-1.88	9	23	14.13	15.12
3	2	3.03	0.55	10	31	18.26	17.55
4	3	3.92	2.98	11	20	23.60	19.98
5	1	5.06	5.41	12	25	30.51	22.41
6	4	6.56	7.84	13	37	39.43	24.84
7	9	8.46	10.27	14	45	50.96	27.27

... and it looks like it fits much better.

Figure of Fitted log(count) from Log-linear

Observed & Fitted Values of Log(count)



Overview

Poisson regression for counts

● Poisson regression for counts

● Components of GLM for Counts

● The Poisson loglinear model

● Interpretation of β

● Interpretation of β

(continued)

● Example: Number of Deaths

Due to AIDs

● Data: Number of Deaths Due

to AIDs \times Month

● A Linear Model for AIDs Data

● A Look at the Bad Model

(linear link)

● Back to Data but Plot

$\log(y_i)$ by Month

● Poisson Log-Linear Model for

Deaths

● Figure of Fitted log(count)

from Log-linear

● Figure of Fitted log(count)

from Log-linear

● Transform explanatory

variable

● Poisson Regression with

Transformed x

● How Large is Large in a

Statistical Sense?

● Observed and Fitted

Log(Counts)

● Observed and Fitted Counts


● Comparison of Fitted Counts

● Comparison of Models

● Observed and Fitted Counts

Figure of Fitted log(count) from Log-linear

Observed & Fitted Counts



Pattern in residuals.

Overview

Poisson regression for counts

● Poisson regression for counts

● Components of GLM for Counts

● The Poisson loglinear model

● Interpretation of β

● Interpretation of β

(continued)

● Example: Number of Deaths Due to AIDs

● Data: Number of Deaths Due to AIDs \times Month

● A Linear Model for AIDs Data

● A Look at the Bad Model

(linear link)

● Back to Data but Plot

$\log(y_i)$ by Month

● Poisson Log-Linear Model for Deaths

● Figure of Fitted log(count) from Log-linear

● Figure of Fitted log(count) from Log-linear

● Transform explanatory variable

● Poisson Regression with Transformed x

● How Large is Large in a Statistical Sense?

● Observed and Fitted Log(Counts)

● Observed and Fitted Counts

● Comparison of Fitted Counts

● Comparison of Linear Models


● Observed and Fitted Counts

Transform explanatory variable

The number of deaths with low & high values of x_i are “over-fit” and number with middle x_i ’s are under-fit.

Transform $x_i \longrightarrow x_i^* = \log(x_i)$

Log(counts) x Log(month)



Poisson Regression with Transformed x

Overview

Poisson regression for counts

- Poisson regression for counts
- Components of GLM for Counts

● The Poisson loglinear model

- Interpretation of β
- Interpretation of β (continued)
- Example: Number of Deaths Due to AIDs
- Data: Number of Deaths Due to AIDs \times Month
- A Linear Model for AIDs Data
- A Look at the Bad Model (linear link)
- Back to Data but Plot $\log(y_i)$ by Month
- Poisson Log-Linear Model for Deaths
- Figure of Fitted log(count) from Log-linear
- Figure of Fitted log(count) from Log-linear
- Transform explanatory variable
- Poisson Regression with Transformed x
- How Large is Large in a Statistical Sense?
- Observed and Fitted Log(Counts)

- Observed and Fitted Counts
- Comparison of Fitted Counts
- Comparing Linear Models
- Observed and Fitted Counts

The estimated GLM with model

- **Random:** Y follows Poisson distribution.
- **Systematic:** $\alpha + \beta \log(x_i) = \alpha + \beta x_i^*$
- **Link:** $\text{Log} \longrightarrow \log(\mu)$.

As a log-linear model

$$\log(\hat{\mu}_i) = -1.9442 + 2.1748x_i^*$$

or equivalently, as a multiplicative model

$$\hat{\mu}_i = e^{-1.9442} e^{2.1748x_i^*}$$

Interpretation: For a 1 unit increase in $\log(\text{month})$, the estimated count increases by a factor of $e^{2.1748} = 8.80$

Is this “large”?

How Large is Large in a Statistical Sense?

Overview

Poisson regression for counts

- Poisson regression for counts
- Components of GLM for Counts

● The Poisson loglinear model

- Interpretation of β
- Interpretation of β (continued)
- Example: Number of Deaths Due to AIDs
- Data: Number of Deaths Due to AIDs \times Month
- A Linear Model for AIDs Data
- A Look at the Bad Model (linear link)
- Back to Data but Plot $\log(y_i)$ by Month
- Poisson Log-Linear Model for Deaths
- Figure of Fitted log(count) from Log-linear
- Figure of Fitted log(count) from Log-linear
- Transform explanatory variable
- Poisson Regression with Transformed x
- How Large is Large in a Statistical Sense?
- Observed and Fitted Log(Counts)

- Observed and Fitted Counts
- Comparison of Fitted Counts
- Comparing Linear Models
- Observed and Fitted Counts

SAS/GENMOD provides asymptotic standard errors (ASE, i.e., large sample) for the parameter estimates.


The ASE for $\hat{\beta}$ equals .2151, and

$$\hat{\beta} \pm 2(.2151) \longrightarrow (1.745, 2.605)$$

which suggests that this is large in a statistical sense.

Observed and Fitted Log(Counts)

Final Model for AIDs Data



Overview

Poisson regression for counts

- Poisson regression for counts

- Components of GLM for

Counts

- The Poisson loglinear model

- Interpretation of β

- Interpretation of β

(continued)

- Example: Number of Deaths

Due to AIDs

- Data: Number of Deaths Due to AIDs \times Month

- A Linear Model for AIDs Data

- A Look at the Bad Model

(linear link)

- Back to Data but Plot

$\log(y_i)$ by Month

- Poisson Log-Linear Model for

Deaths

- Figure of Fitted $\log(\text{count})$

from Log-linear

- Figure of Fitted $\log(\text{count})$

from Log-linear

- Transform explanatory

variable

- Poisson Regression with

Transformed x

- How Large is Large in a

Statistical Sense?

- Observed and Fitted

Log(Counts)

- Observed and Fitted Counts


- Comparison of Fitted Counts

- Comparison of Models

- Observed and Fitted Counts

Observed and Fitted Counts

Final Model for AIDs Data



Overview

Poisson regression for counts

- Poisson regression for counts
- Components of GLM for Counts
- The Poisson loglinear model

● Interpretation of β

- Interpretation of β

(continued)

- Example: Number of Deaths Due to AIDs
- Data: Number of Deaths Due to AIDs \times Month

- A Linear Model for AIDs Data
- A Look at the Bad Model

(linear link)

- Back to Data but Plot $\log(y_i)$ by Month

- Poisson Log-Linear Model for Deaths
- Figure of Fitted $\log(\text{count})$ from Log-linear

- Figure of Fitted $\log(\text{count})$ from Log-linear

- Transform explanatory variable
- Poisson Regression with Transformed x

- How Large is Large in a Statistical Sense?
- Observed and Fitted Log(Counts)

- Observed and Fitted Counts
- Comparison of Fitted Counts

- Comparison of Fitted Counts
- Comparison of Fitted Counts

- Observed and Fitted Counts
- Comparison of Fitted Counts

- Observed and Fitted Counts
- Comparison of Fitted Counts

- Comparison of Fitted Counts
- Comparison of Fitted Counts

- Comparison of Fitted Counts
- Comparison of Fitted Counts

- Observed and Fitted Counts
- Comparison of Fitted Counts

Comparison of Fitted Counts

Overview

Poisson regression for counts

● Poisson regression for counts

● Components of GLM for

Counts

● The Poisson loglinear model

● Interpretation of β

● Interpretation of β

(continued)

● Example: Number of Deaths

Due to AIDs

● Data: Number of Deaths Due to AIDs \times Month

● A Linear Model for AIDs Data

● A Look at the Bad Model

(linear link)

● Back to Data but Plot

$\log(y_i)$ by Month

● Poisson Log-Linear Model for Deaths

● Figure of Fitted $\log(\text{count})$ from Log-linear

● Figure of Fitted $\log(\text{count})$ from Log-linear

● Transform explanatory variable

● Poisson Regression with Transformed x

● How Large is Large in a Statistical Sense?

● Observed and Fitted Log(Counts)

● Observed and Fitted Counts

● Comparison of Fitted Counts

● Generalized Linear Models

● Observed and Fitted Counts

		$\log(x_i)$	x_i	x_i
x_i	y_i	Log	Log	Identity
1	0	.14	1.82	-4.21
2	1	.65	2.35	-1.88
3	2	1.56	3.03	0.55
4	3	2.92	3.92	2.98
5	1	4.74	5.06	5.41
6	4	7.05	6.56	7.84
7	9	9.86	8.46	10.27
8	18	13.17	10.93	12.69
9	23	17.02	14.13	15.12
10	31	21.40	18.26	17.55
11	20	26.33	23.60	19.98
12	25	31.82	30.51	22.41
13	37	37.87	39.43	24.84
14	45	44.49	50.96	27.27

Comparison in Log-Scale

Overview


Poisson regression for counts

- Poisson regression for counts
- Components of GLM for Counts
- The Poisson loglinear model

● Interpretation of β

- Interpretation of β (continued)
- Example: Number of Deaths Due to AIDs
- Data: Number of Deaths Due to AIDs \times Month
- A Linear Model for AIDs Data
- A Look at the Bad Model (linear link)
- Back to Data but Plot $\log(y_i)$ by Month
- Poisson Log-Linear Model for Deaths
- Figure of Fitted $\log(\text{count})$ from Log-linear
- Figure of Fitted $\log(\text{count})$ from Log-linear
- Transform explanatory variable
- Poisson Regression with Transformed x
- How Large is Large in a Statistical Sense?
- Observed and Fitted Log(Counts)

- Observed and Fitted Counts
- Comparison of Fitted Counts
- Comparison in Log-Scale
- Observed and Fitted Counts



Observed and Fitted Counts


Overview

Poisson regression for counts

- Poisson regression for counts
- Components of GLM for Counts
- The Poisson loglinear model
- Interpretation of β
- Interpretation of β (continued)

- Example: Number of Deaths Due to AIDs
- Data: Number of Deaths Due to AIDs \times Month
- A Linear Model for AIDs Data
- A Look at the Bad Model (linear link)
- Back to Data but Plot $\log(y_i)$ by Month
- Poisson Log-Linear Model for Deaths
- Figure of Fitted $\log(\text{count})$ from Log-linear
- Figure of Fitted $\log(\text{count})$ from Log-linear
- Transform explanatory variable
- Poisson Regression with Transformed x
- How Large is Large in a Statistical Sense?
- Observed and Fitted Log(Counts)

- Observed and Fitted Counts
- Comparison of Fitted Counts
- Comparison of Fitted Counts
- Observed and Fitted Counts



More Interpretation of Poisson Regression

Overview

Poisson regression for counts

- Poisson regression for counts
- Components of GLM for

Counts

- The Poisson loglinear model
- Interpretation of β

● Interpretation of β (continued)

- Example: Number of Deaths Due to AIDs
- Data: Number of Deaths Due to AIDs \times Month
- A Linear Model for AIDs Data
- A Look at the Bad Model (linear link)
- Back to Data but Plot $\log(y_i)$ by Month
- Poisson Log-Linear Model for Deaths
- Figure of Fitted log(count) from Log-linear
- Figure of Fitted log(count) from Log-linear
- Transform explanatory variable
- Poisson Regression with Transformed x
- How Large is Large in a Statistical Sense?
- Observed and Fitted Log(Counts)

- Observed and Fitted Counts
- Comparison of Fitted Counts
- Comparing Linear Models
- Observed and Fitted Counts

- The marginal effect of x_i (month period) on μ_i (expected number of deaths due to AIDS).

For a 1 unit increase in $\log(\text{month})$, the estimated count increases by a factor of $e^{2.1748} = 8.80$.

- Computed fitted values and compared them to the observed. (table and plots of this).
- Additional one: We can look at the predicted probability of number of deaths given value on x_i . (This is not too useful here, but would be of use in a predictive setting).

Counts follow a Poisson distribution, so

$$P(Y_i = y) = \frac{e^{-\mu_i} \mu_i^y}{y!}$$

According to our estimated model, probabilities that the number of deaths equals y_i for particular value(s) of x_i is

$$P(Y_i = y) = \frac{e^{-e^{(-1.9442 + 2.1748x_i^*)}} e^{(-1.9442 + 2.1748x_i^*)^y}}{y!}$$

Probabilities of Number of Deaths

$$P(Y_i = y) = \frac{e^{-e^{(-1.9442 + 2.1748x_i^*)}} e^{(-1.9442 + 2.1748x_i^*)^y}}{y!}$$

or since we already have $\hat{\mu}_i$ computed, we can use

$$P(Y_i = y) = \frac{e^{-\hat{\mu}_i} \hat{\mu}_i^y}{y!}$$

For example, consider quarter = 3 (and $\log(3) = 1.09861$), we have

$$\hat{\mu}(\text{quarter} = 3) = 1.5606$$

$$P(Y_3 = 0) = e^{-1.5606} (1.5606)^0 / 0! = .210$$

$$P(Y_3 = 1) = e^{-1.5606} (1.5606)^1 / 1! = .328$$

$$P(Y_3 = 2) = e^{-1.5606} (1.5606)^2 / 2! = .128$$

⋮

$$P(Y_3 = 10) = e^{-1.5606} (1.5606)^{10} / 10! = .000000253$$

Overivew

Poisson regression for counts

- Poisson regression for counts
- Comonents of GLM for Counts
- The Poisson loglinear model
- Interpretation of β

● Interpretation of β (continued)


- Example: Number of Deaths Due to AIDs
- Data: Number of Deaths Due to AIDs \times Month
- A Linear Model for AIDs Data
- A Look at the Bad Model (linear link)
- Back to Data but Plot $\log(y_i)$ by Month
- Poisson Log-Linear Model for Deaths
- Figure of Fitted $\log(\text{count})$ from Log-linear
- Figure of Fitted $\log(\text{count})$ from Log-linear
- Transform explanatory variable
- Poisson Regression with Transformed x
- How Large is Large in a Statistical Sense?
- Observed and Fitted Log(Counts)

- Observed and Fitted Counts
- Comparison of Fitted Counts
- Comparing Linear Models
- Observed and Fitted Counts

Example 2: Crab Data

Agresti (1996)'s horseshoe crab data.

- **Response** variable is the number of satellites a female horseshoe crab has (i.e., how many males are attached to her).
- **Explanatory** variable is the width of the female's back.



Overview

Poisson regression for counts

- Poisson regression for counts
- Components of GLM for Counts

- The Poisson loglinear model
- Interpretation of β

● Interpretation of β (continued)


- Example: Number of Deaths Due to AIDs
- Data: Number of Deaths Due to AIDs \times Month
- A Linear Model for AIDs Data
- A Look at the Bad Model (linear link)
- Back to Data but Plot $\log(y_i)$ by Month
- Poisson Log-Linear Model for Deaths
- Figure of Fitted $\log(\text{count})$ from Log-linear
- Figure of Fitted $\log(\text{count})$ from Log-linear
- Transform explanatory variable
- Poisson Regression with Transformed x
- How Large is Large in a Statistical Sense?
- Observed and Fitted $\log(\text{Counts})$

- Observed and Fitted Counts
- Comparison of Fitted Counts
- Comparing Linear Models
- Observed and Fitted Counts

A Smoother Look

The data were collapsed into 8 groups by their width (i.e., ≤ 23.25 , $23.25-24.25$, $24.25-25.25$, ..., > 29.25).

Mean count and width of Grouped Data



Overview

Poisson regression for counts

- Poisson regression for counts
- Components of GLM for Counts
- The Poisson loglinear model
- Interpretation of β
- Interpretation of β
- (continued)

● Example: Number of Deaths Due to AIDs

- Data: Number of Deaths Due to AIDs \times Month
- A Linear Model for AIDs Data
- A Look at the Bad Model (linear link)
- Back to Data but Plot $\log(y_i)$ by Month
- Poisson Log-Linear Model for Deaths
- Figure of Fitted $\log(\text{count})$ from Log-linear
- Figure of Fitted $\log(\text{count})$ from Log-linear
- Transform explanatory variable
- Poisson Regression with Transformed x
- How Large is Large in a Statistical Sense?
- Observed and Fitted $\log(\text{Counts})$

- Observed and Fitted Counts
- Comparison of Fitted Counts
- Comparison of Fitted Counts
- Observed and Fitted Counts

Estimated Poisson Regression for Crabs

Overview

Poisson regression for counts

- Poisson regression for counts
- Components of GLM for Counts

- The Poisson loglinear model
- Interpretation of β

- Interpretation of β (continued)

- Example: Number of Deaths Due to AIDs

- Data: Number of Deaths Due to AIDs \times Month

- A Linear Model for AIDs Data
- A Look at the Bad Model

(linear link)

- Back to Data but Plot

$\log(y_i)$ by Month

- Poisson Log-Linear Model for Deaths

- Figure of Fitted $\log(\text{count})$ from Log-linear

- Figure of Fitted $\log(\text{count})$ from Log-linear

- Transform explanatory variable

- Poisson Regression with Transformed x

- How Large is Large in a Statistical Sense?

- Observed and Fitted Log(Counts)

- Observed and Fitted Counts

- Comparison of Fitted Counts

- Comparing Linear Models

- Observed and Fitted Counts

$$\log(\hat{\mu}_i) = -3.3048 + .1640x_i$$

- The estimated ASE of $\hat{\beta} = .164$ equals .020, which is small relative to $\hat{\beta}$.
- Since $\hat{\beta} > 0$, the wider the female crab, the greater the expected number of satellites. Note: $\exp(.1640) = 1.18$.
- There is an outlier (with respect to the explanatory variable).
 - ◆ Question: how much does this outlier effect the fit of the model?
 - ◆ Answer: Remove it and re-estimate the model.

$$\log(\hat{\mu}_i) = -3.4610 + .1700x_i$$

and ASE of $\hat{\beta} = .1700$ equals .0216.

- ◆ So in this case, it doesn't have much effect. . . The same basic result holds (i.e., positive effect of width on number of satellites, $\hat{\beta}$ is "significant" and similar in value).

Poisson Regression with Identity Link

Overview

Poisson regression for counts

- Poisson regression for counts
- Components of GLM for Counts

- The Poisson loglinear model
- Interpretation of β
- Interpretation of β

(continued)

- Example: Number of Deaths Due to AIDs
- Data: Number of Deaths Due to AIDs \times Month

● A Linear Model for AIDs Data

- A Look at the Bad Model (linear link)
- Back to Data but Plot $\log(y_i)$ by Month
- Poisson Log-Linear Model for Deaths
- Figure of Fitted log(count) from Log-linear
- Figure of Fitted log(count) from Log-linear
- Transform explanatory variable
- Poisson Regression with Transformed x
- How Large is Large in a Statistical Sense?
- Observed and Fitted Log(Counts)

- Observed and Fitted Counts
- Comparison of Fitted Counts
- Comparing Linear Models
- Observed and Fitted Counts


- From the figure of collapsed data, it looks like either a linear or a log link might work.
- The estimated model with the linear link :

$$\hat{\mu}_i = -11.53 + .55x_i$$

- Since the effect on the number of expected satellites of female width (μ_i) is linear and $\hat{\beta} = .55 > 0$, as width increases by 1 cm, the expected count increases by .55.
- Question: Is the Poisson regression model with the linear or the logit link better for these data?
- Answer: Quick look but more formal later when we discuss model assessment (or read further in the text).

Log versus Identity Link for Crabs

••• data — log — identity



Overview

Poisson regression for counts

- Poisson regression for counts
- Components of GLM for Counts
- The Poisson loglinear model
- Interpretation of β
- Interpretation of β (continued)
- Example: Number of Deaths Due to AIDs
- Data: Number of Deaths Due to AIDs \times Month

● A Linear Model for AIDs Data

- A Look at the Bad Model (linear link)
- Back to Data but Plot $\log(y_i)$ by Month
- Poisson Log-Linear Model for Deaths
- Figure of Fitted $\log(\text{count})$ from Log-linear
- Figure of Fitted $\log(\text{count})$ from Log-linear
- Transform explanatory variable
- Poisson Regression with Transformed x
- How Large is Large in a Statistical Sense?
- Observed and Fitted Log(Counts)

- Observed and Fitted Counts
- Comparison of Fitted Counts
- Comparing Linear Models
- Observed and Fitted Counts

Poisson regression for rates

Overview

Poisson regression for counts

Poisson regression for rates

- Poisson regression for rates
- The Data
- Poisson Regression for Rates of Events
- As a Multiplicative Model
- Gardner, Mulvey, & Shaw (1995)
- Estimated Parameters
- SAS/PROC GENMOD
- Example 2: Lung Cancer
- Plot of the Rate by Age
- Plot of the log(Rate) by Age
- Model 1: Age and City both Nominal
- Parameter Estimates from Model 1
- Model 2: City Nominal & Age Numerical
- Model 2: Observed and Fitted Values
- Model 3: City Nominal & Age Quadratic
- Model 3: Observed and Fitted Values
- Model 4: Simpler city & Age Quadratic
- Model 4: Fitted and Observed
- Next Steps

Events occur over time (or space), and the length of time (or amount of space) can vary from observation to observation. Our model should take this into account.

Example: Gardner, Mulvey, & Shaw (1995), *Psychological Bulletin*, 118, 392–404.

Y = Number of violent incidents exhibited over a 6 month period by patients who had been treated in the ER of a psychiatric hospital.

During the 6 months period of the study, the individuals were primarily residing in the community. The number of violent acts depends on the opportunity to commit them; that is, the number of days out of the 6 month period in which a patient is in the community (as opposed to being locked up in a jail or hospital).

The Data

Overivew

Poisson regression for counts

Poisson regression for rates

- Poisson regression for rates
- The Data
- Poisson Regression for Rates of Events
- As a Multiplicative Model
- Gardner, Mulvey, & Shaw (1995)
- Estimated Parameters
- SAS/PROC GENMOD
- Example 2: Lung Cancer
- Plot of the Rate by Age
- Plot of the log(Rate) by Age
- Model 1: Age and City both Nominal
- Parameter Estimates from Model 1
- Model 2: City Nominal & Age Numerical
- Model 2: Observed and Fitted Values
- Model 3: City Nominal & Age Quadratic
- Model 3: Observed and Fitted Values
- Model 4: Simpler city & Age Quadratic
- Model 4: Fitted and Observed
- Next Steps

put histogram here.

Poisson Regression for Rates of Events

Y = count (e.g., number violent acts).

t = index of the time or space (e.g., days in the community).

The sample **rate** of occurrence is Y/t .

The expected value of the rate is

$$E(Y/t) = \frac{1}{t}E(Y) = \mu/t$$

The Poisson loglinear regression model for the expected rate of the occurrence of events is

$$\log(\mu/t) = \alpha + \beta x$$

$$\log(\mu) - \log(t) = \alpha + \beta x$$

$$\log(\mu) = \alpha + \beta x + \log(t)$$

The term “ $-\log(t)$ ” is an adjustment term and each individual may have a different value of t .

The term $-\log(t)$ is referred to as an “**offset**”.

Overview

Poisson regression for counts

Poisson regression for rates

- Poisson regression for rates
- The Data
- Poisson Regression for Rates of Events
- As a Multiplicative Model
- Gardner, Mulvey, & Shaw (1995)
- Estimated Parameters
- SAS/PROC GENMOD
- Example 2: Lung Cancer
- Plot of the Rate by Age
- Plot of the log(Rate) by Age
- Model 1: Age and City both Nominal
- Parameter Estimates from Model 1
- Model 2: City Nominal & Age Numerical
- Model 2: Observed and Fitted Values
- Model 3: City Nominal & Age Quadratic
- Model 3: Observed and Fitted Values
- Model 4: Simpler city & Age Quadratic
- Model 4: Fitted and Observed
- Next Steps

As a Multiplicative Model

Overview

Poisson regression for counts

Poisson regression for rates

- Poisson regression for rates
- The Data
- Poisson Regression for Rates of Events
- As a Multiplicative Model
- Gardner, Mulvey, & Shaw (1995)
- Estimated Parameters
- SAS/PROC GENMOD
- Example 2: Lung Cancer
- Plot of the Rate by Age
- Plot of the log(Rate) by Age
- Model 1: Age and City both Nominal
- Parameter Estimates from Model 1
- Model 2: City Nominal & Age Numerical
- Model 2: Observed and Fitted Values
- Model 3: City Nominal & Age Quadratic
- Model 3: Observed and Fitted Values
- Model 4: Simpler city & Age Quadratic
- Model 4: Fitted and Observed
- Next Steps

The Poisson loglinear regression model with a log link for rate data is

$$\log(\mu/t) = \alpha + \beta x$$

$$\mu/t = e^{\alpha} e^{\beta x}$$

$$\mu = t e^{\alpha} e^{\beta x}$$

The expected value of counts depends on both t and x , both of which are observations (i.e., neither is a parameter of the model).

Gardner, Mulvey, & Shaw (1995)

Overview

Poisson regression for counts

Poisson regression for rates

- Poisson regression for rates
- The Data
- Poisson Regression for Rates of Events
- As a Multiplicative Model
- Gardner, Mulvey, & Shaw (1995)
- Estimated Parameters
- SAS/PROC GENMOD
- Example 2: Lung Cancer
- Plot of the Rate by Age
- Plot of the log(Rate) by Age
- Model 1: Age and City both Nominal
- Parameter Estimates from Model 1
- Model 2: City Nominal & Age Numerical
- Model 2: Observed and Fitted Values
- Model 3: City Nominal & Age Quadratic
- Model 3: Observed and Fitted Values
- Model 4: Simpler city & Age Quadratic
- Model 4: Fitted and Observed
- Next Steps

- **Response variable** is rate of violent incidents, which equals the number of violent incident divided by the number of days an individual resided in the community. ($\bar{y} = 3.0$ with $s = 7.3$ and $\bar{t} = 154$ with $s = 42$ days).

■ Explanatory variables:

- ◆ Age ($\bar{x}_1 = 28.6$ years and $s_1 = 11.1$)
- ◆ Sum of 2 ER clinicians ratings of concern on a 0 – 5 scale, so x_2 ranges from 0 to 10. ($\bar{x}_2 = 2.9$ with $s_2 = 3.1$).
- ◆ History of previous violent acts, where

$x_3 = 0$ means no previous acts
 $= 1$ previous act either 3 days before *or* more than 3 days before
 $= 2$ previous acts both 3 days before *and* more than 3 days before

$$r(\text{concern, history}) = .55$$

$$r(\text{age, history}) = -.11$$

$$r(\text{age, concern}) = -.07$$

Estimated Parameters

Coefficient	Value	ASE	value/ASE
Intercept	-3.410	.0690	-49.29
Age	-.045	.0023	-19.69
Concern	.083	.0075	11.20
History	.420	.0380	11.26

put histogram of fitted and observed

Note: Poisson regression models for rate data are related to models for “survival times”.

Overview

Poisson regression for counts

Poisson regression for rates

- Poisson regression for rates
- The Data
- Poisson Regression for Rates of Events
- As a Multiplicative Model
- Gardner, Mulvey, & Shaw (1995)
- Estimated Parameters
- SAS/PROC GENMOD
- Example 2: Lung Cancer
- Plot of the Rate by Age
- Plot of the log(Rate) by Age
- Model 1: Age and City both Nominal
- Parameter Estimates from Model 1
- Model 2: City Nominal & Age Numerical
- Model 2: Observed and Fitted Values
- Model 3: City Nominal & Age Quadratic
- Model 3: Observed and Fitted Values
- Model 4: Simpler city & Age Quadratic
- Model 4: Fitted and Observed
- Next Steps

SAS/PROC GENMOD

DATA violent;

INPUT age history concern days count;

Ldays = LOG(days);

To fit model for rate data with “days” as t :

PROC GENMOD DATA=violent;

MODEL count = age history concern

/ LINK=log DIST=poisson OFFSET=Ldays;

To include interactions, change model statement

MODEL count = age history concern history*concern

/ LINK=log DIST=poisson OFFSET=Ldays;

If we want to treat “history” as a discrete (nominal) variable, you would include the statement

CLASS history;

If the model with “history” treated as a numerical variable almost as well as the model with “history” treated as a nominal discrete variable, which model would you prefer?

Overview

Poisson regression for counts

Poisson regression for rates

- Poisson regression for rates
- The Data
- Poisson Regression for Rates of Events
- As a Multiplicative Model
- Gardner, Mulvey, & Shaw (1995)
- Estimated Parameters
- SAS/PROC GENMOD
- Example 2: Lung Cancer
- Plot of the Rate by Age
- Plot of the log(Rate) by Age
- Model 1: Age and City both Nominal
- Parameter Estimates from Model 1
- Model 2: City Nominal & Age Numerical
- Model 2: Observed and Fitted Values
- Model 3: City Nominal & Age Quadratic
- Model 3: Observed and Fitted Values
- Model 4: Simpler city & Age Quadratic
- Model 4: Fitted and Observed
- Next Steps

Example 2: Lung Cancer

Overview

Poisson regression for counts


Poisson regression for rates

- Poisson regression for rates
- The Data
- Poisson Regression for Rates of Events
- As a Multiplicative Model
- Gardner, Mulvey, & Shaw (1995)
- Estimated Parameters
- SAS/PROC GENMOD
- Example 2: Lung Cancer
- Plot of the Rate by Age
- Plot of the log(Rate) by Age
- Model 1: Age and City both Nominal
- Parameter Estimates from Model 1
- Model 2: City Nominal & Age Numerical
- Model 2: Observed and Fitted Values
- Model 3: City Nominal & Age Quadratic
- Model 3: Observed and Fitted Values
- Model 4: Simpler city & Age Quadratic
- Model 4: Fitted and Observed
- Next Steps

Data are from Lindsey (1995) from Andersen (1977)

- **Response Variable:** Y = Number of cases of lung cancer and it follows a Poisson distribution.:
- **Explanatory Variables:**
 - ◆ City in Denmark (Fredericia, Horsens, Kolding, Vejle).
 - ◆ Age (40–54, 55–59, 60–64, 65–69, 70–74, >75).
- **Offset** = Population size of each age group of each city.
- We will model the rate of cases of lung cancer = Y/t .

Plot of the Rate by Age




Overview

Poisson regression for counts

Poisson regression for rates

- Poisson regression for rates
- The Data
- Poisson Regression for Rates of Events
- As a Multiplicative Model
- Gardner, Mulvey, & Shaw (1995)
- Estimated Parameters
- SAS/PROC GENMOD
- Example 2: Lung Cancer
- Plot of the Rate by Age
- Plot of the log(Rate) by Age
- Model 1: Age and City both Nominal
- Parameter Estimates from Model 1
- Model 2: City Nominal & Age Numerical
- Model 2: Observed and Fitted Values
- Model 3: City Nominal & Age Quadratic
- Model 3: Observed and Fitted Values
- Model 4: Simpler city & Age Quadratic
- Model 4: Fitted and Observed
- Next Steps

Plot of the log(Rate) by Age



Overview

Poisson regression for counts

Poisson regression for rates

- Poisson regression for rates
- The Data
- Poisson Regression for Rates of Events
- As a Multiplicative Model
- Gardner, Mulvey, & Shaw (1995)
- Estimated Parameters
- SAS/PROC GENMOD
- Example 2: Lung Cancer
- Plot of the Rate by Age
- Plot of the log(Rate) by Age
- Model 1: Age and City both Nominal
- Parameter Estimates from Model 1
- Model 2: City Nominal & Age Numerical
- Model 2: Observed and Fitted Values
- Model 3: City Nominal & Age Quadratic
- Model 3: Observed and Fitted Values
- Model 4: Simpler city & Age Quadratic
- Model 4: Fitted and Observed
- Next Steps

Model 1: Age and City both Nominal

Define

$$\text{Fredericia} = \begin{cases} 1 & \text{if city is Frederica} \\ 0 & \text{other city} \end{cases}$$

$$\text{Horsens} = \begin{cases} 1 & \text{if city is Horsens} \\ 0 & \text{other city} \end{cases}$$

$$\text{Kolding} = \begin{cases} 1 & \text{if city is Kolding} \\ 0 & \text{other city} \end{cases}$$

Define Dummy variables for the 6 age classes (groups).

Model 1:

$$\begin{aligned} \log(Y/\text{pop}) &= \alpha + \beta_1(\text{Fredericia}) + \beta_2(\text{Horsens}) + \beta_3(\text{Kolding}) \\ &= \beta_4(\text{Age1}) + \beta_5(\text{Age2}) + \beta_6(\text{Age3}) + \beta_7(\text{Age4}) \\ &\quad \beta_8(\text{Age5}) \end{aligned}$$

Overview

Poisson regression for counts

Poisson regression for rates

- Poisson regression for rates
- The Data
- Poisson Regression for Rates of Events
- As a Multiplicative Model
- Gardner, Mulvey, & Shaw (1995)
- Estimated Parameters
- SAS/PROC GENMOD
- Example 2: Lung Cancer
- Plot of the Rate by Age
- Plot of the log(Rate) by Age
- Model 1: Age and City both Nominal
- Parameter Estimates from Model 1
- Model 2: City Nominal & Age Numerical
- Model 2: Observed and Fitted Values
- Model 3: City Nominal & Age Quadratic
- Model 3: Observed and Fitted Values
- Model 4: Simpler city & Age Quadratic
- Model 4: Fitted and Observed
- Next Steps

Parameter Estimates from Model 1

Overview

Poisson regression for counts

Poisson regression for rates

- Poisson regression for rates
- The Data
- Poisson Regression for Rates of Events
- As a Multiplicative Model
- Gardner, Mulvey, & Shaw (1995)
- Estimated Parameters
- SAS/PROC GENMOD
- Example 2: Lung Cancer
- Plot of the Rate by Age
- Plot of the log(Rate) by Age
- Model 1: Age and City both Nominal
- Parameter Estimates from Model 1
- Model 2: City Nominal & Age Numerical
- Model 2: Observed and Fitted Values
- Model 3: City Nominal & Age Quadratic
- Model 3: Observed and Fitted Values
- Model 4: Simpler city & Age Quadratic
- Model 4: Fitted and Observed
- Next Steps

Parameter			Estimate	<i>df</i>	s.e.	X^2	<i>p</i>
Intercept		α	1	−4.48	0.21	423.33	< .01
city	Frederic	β_1	1	0.27	0.18	2.10	.15
city	Horsens	β_2	1	−0.05	0.19	0.09	.76
city	Kolding	β_3	1	−0.09	0.19	0.25	.62
city	Vejle		0	0.00	0.00	.	.
age	40-54	β_4	1	−1.41	0.25	32.18	< .01
age	55-59	β_5	1	−0.31	0.25	1.60	.21
age	60-64	β_6	1	0.09	0.23	0.18	.67
age	65-69	β_7	1	0.34	0.23	2.22	.14
age	70-74	β_8	1	0.43	0.23	3.34	.07
age	>75		0	0.00	0.00	.	.

Note: $G^2 = 23.45$, $df = 15$, $p = .08$

Model 2: City Nominal & Age Numerical

The mid-point of the age ranges were used (except for the last one, I used 75).

$$\begin{aligned}\log(Y/\text{pop}) &= \alpha + \beta_1(\text{Fredericia}) + \beta_2(\text{Horsens}) + \beta_3(\text{Kolding}) \\ &= \beta_4(\text{Age Mid-point})\end{aligned}$$

Parameter		Estimate	<i>df</i>	s.e.	X^2	<i>p</i>
Intercept	α	1	−8.22	0.44	349.18	< .01
city	Frederic β_1	1	0.24	0.18	1.72	0.19
city	Horsens β_2	1	−0.05	0.19	0.10	0.76
city	Kolding β_3	1	−0.10	0.19	0.28	0.60
city	Vejle	0	0.00	0.00	.	.
age-midpoint	β_4	1	0.05	0.00	75.62	< .01

Note: $G^2 = 46.45$, $df = 19$, $p < .01$

Overview

Poisson regression for counts

Poisson regression for rates

- Poisson regression for rates
- The Data
- Poisson Regression for Rates of Events
- As a Multiplicative Model
- Gardner, Mulvey, & Shaw (1995)
- Estimated Parameters
- SAS/PROC GENMOD
- Example 2: Lung Cancer
- Plot of the Rate by Age
- Plot of the log(Rate) by Age
- Model 1: Age and City both Nominal
- Parameter Estimates from Model 1
- Model 2: City Nominal & Age Numerical
- Model 2: Observed and Fitted Values
- Model 3: City Nominal & Age Quadratic
- Model 3: Observed and Fitted Values
- Model 4: Simpler city & Age Quadratic
- Model 4: Fitted and Observed
- Next Steps


Model 2: Observed and Fitted Values

Overview

Poisson regression for counts

Poisson regression for rates

- Poisson regression for rates
- The Data
- Poisson Regression for Rates of Events
- As a Multiplicative Model
- Gardner, Mulvey, & Shaw (1995)
- Estimated Parameters
- SAS/PROC GENMOD
- Example 2: Lung Cancer
- Plot of the Rate by Age
- Plot of the log(Rate) by Age
- Model 1: Age and City both Nominal
- Parameter Estimates from Model 1
- Model 2: City Nominal & Age Numerical
- Model 2: Observed and Fitted Values
- Model 3: City Nominal & Age Quadratic
- Model 3: Observed and Fitted Values
- Model 4: Simpler city & Age Quadratic
- Model 4: Fitted and Observed
- Next Steps



Model 3: City Nominal & Age Quadratic

$$\begin{aligned}\log(Y/\text{pop}) &= \alpha + \beta_1(\text{Fredericia}) + \beta_2(\text{Horsens}) + \beta_3(\text{Kolding}) \\ &= \beta_4(\text{Age Mid-point}) + \beta_5(\text{Age Mid-point})^2\end{aligned}$$

Overview

Poisson regression for counts

Poisson regression for rates

- Poisson regression for rates
- The Data
- Poisson Regression for Rates of Events
- As a Multiplicative Model
- Gardner, Mulvey, & Shaw (1995)
- Estimated Parameters
- SAS/PROC GENMOD
- Example 2: Lung Cancer
- Plot of the Rate by Age
- Plot of the log(Rate) by Age
- Model 1: Age and City both Nominal
- Parameter Estimates from Model 1
- Model 2: City Nominal & Age Numerical
- Model 2: Observed and Fitted Values
- Model 3: City Nominal & Age Quadratic
- Model 3: Observed and Fitted Values
- Model 4: Simpler city & Age Quadratic
- Model 4: Fitted and Observed
- Next Steps

Parameter			Estimate	<i>df</i>	s.e.	X^2	<i>p</i>
Intercept		α	1	−21.72	3.09	49.24	< .01
city	Frederic	β_1	1	0.27	0.18	2.13	0.14
city	Horsens	β_2	1	−0.05	0.19	0.09	0.76
city	Kolding	β_3	1	−0.10	0.19	0.26	0.61
city	Vejle		0	0.00	0.00	.	.
age-midpoint		β_4	1	0.50	0.10	24.91	< .01
age ²		β_5	1	−0.00	0.00	19.90	< .01

$$G^2 = 26.02, df = 18, p = .10.$$


Model 3: Observed and Fitted Values

Overview

Poisson regression for counts

Poisson regression for rates

- Poisson regression for rates
- The Data
- Poisson Regression for Rates of Events
- As a Multiplicative Model
- Gardner, Mulvey, & Shaw (1995)
- Estimated Parameters
- SAS/PROC GENMOD
- Example 2: Lung Cancer
- Plot of the Rate by Age
- Plot of the log(Rate) by Age
- Model 1: Age and City both Nominal
- Parameter Estimates from Model 1
- Model 2: City Nominal & Age Numerical
- Model 2: Observed and Fitted Values
- Model 3: City Nominal & Age Quadratic
- Model 3: Observed and Fitted Values
- Model 4: Simpler city & Age Quadratic
- Model 4: Fitted and Observed
- Next Steps



Model 4: Simpler city & Age Quadratic

Define Fredericia = $\begin{cases} 1 & \text{if city is Frederica} \\ 0 & \text{other city} \end{cases}$

$$\log(Y/\text{pop}) = \alpha + \beta_1(\text{Fredericia}) + \beta_2(\text{Age Mid-point}) + \beta_3(\text{Age Mid-point})^2$$

That is,

$$\log(Y/\text{pop}) = \begin{cases} \alpha + \beta_1 + \beta_2(\text{Age}) + \beta_3(\text{Age})^2 & \text{if Fredericia} \\ \alpha + \beta_2(\text{Age}) + \beta_3(\text{Age})^2 & \text{if other city} \end{cases}$$

Parameter			Estimate	df	s.e.	X ²	p
Intercept		α	1	−21.78	3.09	49.61	< .01
frederic	1	β_1	1	0.32	0.14	4.92	.03
frederic	0		0	0.00	0.00	.	.
age-midpoint		β_2	1	0.50	0.10	24.93	< .01
age ²		β_3	1	−0.00	0.00	19.91	< .01

Note: $G^2 = 26.2815$, $df = 20$, $p = .16$.

Overview

Poisson regression for counts

Poisson regression for rates

- Poisson regression for rates
- The Data
- Poisson Regression for Rates of Events
- As a Multiplicative Model
- Gardner, Mulvey, & Shaw (1995)
- Estimated Parameters
- SAS/PROC GENMOD
- Example 2: Lung Cancer
- Plot of the Rate by Age
- Plot of the log(Rate) by Age
- Model 1: Age and City both Nominal
- Parameter Estimates from Model 1
- Model 2: City Nominal & Age Numerical
- Model 2: Observed and Fitted Values
- Model 3: City Nominal & Age Quadratic
- Model 3: Observed and Fitted Values
- Model 4: Simpler city & Age Quadratic
- Model 4: Fitted and Observed
- Next Steps


Model 4: Fitted and Observed

Overview

Poisson regression for counts

Poisson regression for rates

- Poisson regression for rates
- The Data
- Poisson Regression for Rates of Events
- As a Multiplicative Model
- Gardner, Mulvey, & Shaw (1995)
- Estimated Parameters
- SAS/PROC GENMOD
- Example 2: Lung Cancer
- Plot of the Rate by Age
- Plot of the log(Rate) by Age
- Model 1: Age and City both Nominal
- Parameter Estimates from Model 1
- Model 2: City Nominal & Age Numerical
- Model 2: Observed and Fitted Values
- Model 3: City Nominal & Age Quadratic
- Model 3: Observed and Fitted Values
- Model 4: Simpler city & Age Quadratic
- Model 4: Fitted and Observed
- Next Steps



Next Steps

Statistical Inference for Poisson Regression...

Overivew

Poisson regression for counts

Poisson regression for rates

- Poisson regression for rates
- The Data
- Poisson Regression for Rates of Events
- As a Multiplicative Model
- Gardner, Mulvey, & Shaw (1995)
- Estimated Parameters
- SAS/PROC GENMOD
- Example 2: Lung Cancer
- Plot of the Rate by Age
- Plot of the $\log(\text{Rate})$ by Age
- Model 1: Age and City both Nominal
- Parameter Estimates from Model 1
- Model 2: City Nominal & Age Numerical
- Model 2: Observed and Fitted Values
- Model 3: City Nominal & Age Quadratic
- Model 3: Observed and Fitted Values
- Model 4: Simpler city & Age Quadratic
- Model 4: Fitted and Observed
- Next Steps