MONASH University

MONASH BUSINESS SCHOOL

ISSN 1440-771X

Department of Econometrics and Business Statistics

http://monash.edu/business/ebs/research/publications

Anomaly detection using surprisals

Rob J Hyndman, David T Frazier

October 2025

Working Paper no/yr

AACSB ACCREDITED EQUIS ACCREDITED ASSOCIATION OF MBAS ACCREDITED

Anomaly detection using surprisals

Rob J Hyndman
Department of Econometrics & Business Statistics
Monash University
Clayton 3800
Australia
Email: Rob.Hyndman@monash.edu
Corresponding author

David T Frazier
Department of Econometrics & Business Statistics
Monash University
Clayton 3800
Australia

2 October 2025

JEL classification: C10,C14,C46

Anomaly detection using surprisals

Abstract

TBC

Keywords: Extreme value theory; Generalized Pareto Distribution; Outlier detection; Tail bounds.

1 Introduction

Anomaly detection methods are often rather ad hoc and lack a solid theoretical foundation, or they make strong distributional assumptions that may not hold in practice. Here we define an anomaly as a point of low probability, and we propose a general framework for anomaly detection by estimating the probability of each observation, or one more anomalous. This approach allows for a unified treatment of anomaly detection that can be applied to any anomaly detection problem, provided a probability distribution can be defined on the space of observations. It allows for univariate and multivariate settings, and can be applied to both unconditional and conditional distributions.

The basic idea is to calculate the surprisal (equal to the negative log density) of each observation, and then estimate the tail probability from the surprisal distribution. This allows us to estimate the probability of observing a surprisal value as extreme or more extreme than a given observation, which can then be used to flag anomalies. The surprisal approach to anomaly detection provides a principled way to control the false positive rate when flagging anomalies, by setting a threshold on the estimated tail probabilities.

Let $\boldsymbol{y}_1, \ldots, \boldsymbol{y}_n$ be iid observations from a probability distribution F defined on an appropriate sample space. The *surprisal* of observation $\boldsymbol{y}_i$ is defined as (Stone 2022)

$$s_i = -\log f(\boldsymbol{y}_i),$$

where f is either the density or probability mass function of F. While these are better known as "log scores" in statistics (Good 1952), in this context we prefer the term "surprisal" (coined by Tribus 1961), as it captures the idea that observations with low density are more "surprising" (and therefore potentially anomalous) than those with high density. A large value of s_i indicates that $\boldsymbol{y}_i$ is not a likely value, and so is a potential anomaly. On the other hand, typical values will have low surprisals.

In information theory, the average surprisal is known as the entropy of a random variable (Cover & Thomas 2006; Stone 2022) and provides a measure of uncertainty.

Although the term "surprisal" may not be in common use amongst statisticians and other data scientists, the underlying idea is widely used. For example, if f is obtained from a model with parameters that need to be estimated from the data, then the sum of the surprisals is equal to minus the log likelihood of the data. Therefore, finding parameters that minimize the sum of the surprisals is equivalent to maximum likelihood estimation.

Let $G(s) = \Pr\{S \leq s\}$ be the distribution function of the surprisal values, where $S = -\log f(\boldsymbol{Y})$ and $\boldsymbol{Y} \sim F$. Then, the probability of observing a value with surprisal at least as large as s is given by $1 - G(s)$, which leads to a natural way to assign an anomaly score to each observation $\boldsymbol{y}_i$:

$$p_i = 1 - G(s_i) = 1 - G(-\log f(\boldsymbol{y}_i)).$$

We define anomalies as observations with $p_i < \alpha$, giving a false positive rate of α if all observations are from F. This is preferable to directly computing the probabilities from the tail of F, as it allows for anomalies to occur anywhere in the distribution of F, not just in its tails. For example, anomalies may lie between modes in a multimodal distribution, or in otherwise low-density regions of the distribution F. It is also preferable to a distance-based approach to anomaly detection (e.g., Wilkinson 2018), as it naturally handles skewed and heavy-tailed distributions.

The surprisal approach converts the problem of working with F, a potentially multivariate distribution (or even one on a manifold or some non-Euclidean space), to working with a univariate distribution G defined on $\mathbb{R}$. We can write

$$G(s) = \Pr\{-\log f(Y) \leq s\} = \Pr\{f(Y) \geq e^{-s}\}.$$

This makes clear that the surprisal probability is related to the highest density regions of the distribution (Hyndman 1996). In fact, every observation with $p_i < \alpha$ lies outside the $100(1-\alpha)\%$ HDR region of F.

We will usually not know F, and so we need to estimate it from data or from theory. Such estimates are particularly fragile in areas of low density, which is precisely where we are looking for anomalies. Further, we are seeking a method that does not make strong distributional assumptions.

We present three ways to compute the tail probabilities $\{p_1, \dots, p_n\}$ from the surprisal distribution:

(1) by using the same assumed or estimated distribution that was used to compute the surprisals;

(2) by using the empirical distribution function of the observed surprisals $\{s_1, \dots, s_n\}$; and

(3) by fitting a Generalized Pareto distribution to the most extreme surprisals.

Clearly, approach (1) makes strong distributional assumptions, as it requires that we have an accurate distribution of the original observations. The other two approaches allow for considerable mis-specification of the distribution used to compute the surprisals. Approach (2) requires only that the ordering of the surprisals is similar to what would have been obtained under the true distribution, while approach (3) requires that extreme value theory applies to the surprisal distribution.

We show that under weak conditions on the tail of the surprisal distribution, the maximum surprisal converges to one of the three types of extreme value distributions. This justifies fitting a Generalized Pareto Distribution (GPD) to the largest surprisal values, and using it to estimate tail probabilities.

An important precursor to this work is Kandanaarachchi & Hyndman (2022) who proposed estimating F by a leave-one-out kernel density estimate, and then fitting a Generalized Pareto Distribution (GPD) to the most extreme surprisal values. The underlying idea is that even if F is not particularly well-estimated, the tail of the surprisal distribution should still be well-approximated by a GPD. Hyndman, Kandanaarachchi & Turner (2025) provides some theoretical justification for this approach in the context of kernel density estimation.

Here we extend that idea to allow F to be any well-behaved distribution, including conditional distributions. Specifically, we fit a GPD to the largest β of $\{s_i\}_{i=1}^n$, obtaining estimates of the location, scale and shape parameters, (μ, σ, ξ). We calculate the probability of each observation or one more extreme, $p_i = \beta\left[1 - P(s_i \mid \hat{\mu}, \hat{\sigma}, \hat{\xi})\right]$, where P is the distribution function of the GPD. Observations not contained in the largest β of $\{s_i\}_{i=1}^n$ are assigned $p_i = \beta$. To justify this approach, we seek conditions on the surprisal distribution of $S = -\log f(\boldsymbol{Y})$, where $\boldsymbol{Y} \sim F$, similar to the Fisher-Tippett-Gnedenko (FTG) theorem (Coles 2001).

We illustrate all three approaches with several examples, including both simulated and real data.

> **Note**
>
> Outline of paper TBC

2 Empirical surprisal probabilities

Under approach (2) above, we estimate p_i as the proportion of surprisals larger than s_i. For large n, this provides a quick and convenient estimate of p_i that is accurate even when the assumed distribution F is not.

> **!** Important
>
> What can we say about the quality of the approximation?
>
> If the surprisal ranks under the true and estimated F distributions are equal, this is obviously ok.
>
> That should be true in the tails of any distribution — i.e., when y_i is greater than the largest mode of f, or less than the smallest mode of f.
>
> Can we say anything else?

3 Extreme value theory for the surprisal distribution

We consider three possible conditions on the surprisal distribution of $S = -\log f(\boldsymbol{Y})$ where f is the density or mass function of $\boldsymbol{Y}$. These conditions describe increasingly heavy tails:

1. where S is sub-Gaussian (e.g., when f is bounded above and below);
2. where S is sub-exponential (e.g., when f is Gaussian); and
3. where S has a polynomial moment (e.g., when f is Student-t).

We state these as three alternative assumptions.

Assumption 3.1 (Sub-Gaussian)**.** The random variable $S = -\log f(Y) \in \mathbb{R}$, and satisfies, for all $\lambda \in \mathbb{R}$, and some $\nu > 0$,

$$\mathbb{E}\exp\{\lambda(S - \mathbb{E}[S])\} \leq \exp\{\lambda^2\nu^2/2\}.$$

Assumption 3.2 (Sub-exponential)**.** The random variable S is sub-exponential with parameters ν and b, i.e.,

$$\mathbb{E}\exp\{\lambda(S - \mathbb{E}[S])\} \leq \exp\{\lambda^2\nu^2/2\} \quad \text{for all} \quad |\lambda| < 1/b.$$

Assumption 3.3 (Polynomial)**.** The random variable $|S|$ has polynomial moments of order $\alpha \geq 1$; i.e., $\mathbb{E}[|S|^p] \leq C^p$ for some $C > 0$ such that $C^p - 1 > 0$.

Under each of Assumptions 3.1–3.3, we seek a result on the limiting distribution of the maximum of the surprisals $M_n = \max\{s_1, \ldots, s_n\}$. If we can show that this distribution falls within the class of extreme value distributions, then we can fit a Generalized Pareto Distribution (GPD) to the largest surprisal values (Coles 2001).

3.1 S has sub-Gaussian tails

Let's first assume that the surprisals are well-behaved and have a sub-Gaussian tail. Assumption 3.1 will be satisfied in cases where the density f is bounded both above and below, such as with mass functions on a finite set. However, it may not be satisfied if the log density has no lower bound.

For example, consider the truncated normal distribution on $[a,b]$, where $Y \sim \mathcal{TN}(a,b,\mu,\sigma^2)$ and $-\infty < a < b < \infty$. Then

$$S = -\log(2\pi\sigma^2)/2 - \frac{1}{2\sigma^2}(Y-\mu)^2 - \log\{\Phi(b)-\Phi(a)\},$$

where Φ is the standard normal distribution function. Since $Y \in [a,b]$, S is bounded for all $\sigma > 0$ and hence S satisfies Assumption 3.1.

Under Assumption 3.1, we have the following result on the maximum of the surprisals.

Lemma 3.1. *Suppose $\boldsymbol{y}_1,\dots,\boldsymbol{y}_n$ are iid from a density f, and let $M_n = \max\{s_1,\dots,s_n\}$ where $s_i = -\log f(\boldsymbol{y}_i)$. If $S = -\log f(Y) \in \mathbb{R}$, and satisfies, for all $\lambda \in \mathbb{R}$, and some $\nu > 0$,*

$$\mathbb{E}\exp\{\lambda(S-\mathbb{E}[S])\} \le \exp\{\lambda^2\nu^2/2\},$$

, then, for $s \in \mathbb{R}_+$,

$$\Pr\{M_n - \mathbb{E}[S] \ge s\} \le 1 - \left(1 - e^{-\frac{1}{2}s^2/\nu^2}\right)^n.$$

Proof. For any $\lambda > 0$,

$$\begin{aligned}\Pr\{S \ge s\} = \Pr\left\{e^{\lambda(S-\mathbb{E}[S])} \ge e^{\lambda(s-\mathbb{E}[S])}\right\} &\le e^{-\lambda(s-\mu)}\mathbb{E}e^{\lambda(S-\mu)} \\ &\le e^{-\lambda(s-\mu)}e^{\lambda^2\nu^2/2}.\end{aligned}$$

Now, we can maximise the RHS of the above bound in λ, which yields the optimal value $\lambda = (s-\mu)/\{M^2\}$. Plugging this in we have the bound

$$\Pr\{S \ge s\} \le e^{-\frac{1}{2}(s-\mu)^2/\nu^2}.$$

The stated result now follows by noting that

$$\begin{aligned}\Pr(M_n - \mathbb{E}[S] \ge s) = 1 - \Pr\{M_n \le s\} &= 1 - \Pr\{S \le s\}^n \\ &= 1 - (1 - \Pr\{S \ge s\})^n \\ &\le 1 - \left(1 - e^{-\frac{1}{2}s^2/\nu^2}\right)^n.\end{aligned}$$

□

Corollary 3.1. *If Assumption 3.1 is satisfied, then, for all $s > 0$,*

$$\Pr\left(|M_n - \mathbb{E}[S]| \geq \sqrt{2\nu^2 s} + \sqrt{2\nu^2 \log(2n)}\right) \leq e^{-s}.$$

Proof. This follows directly from the union bound:

$$\begin{aligned}
\Pr\left(|M_n| \geq \sqrt{2\nu^2 s} + \sqrt{2\nu^2 \log(n)}\right) &\leq n \Pr\left\{|S| \geq \sqrt{2}\nu^2 s + \sqrt{2\nu^2 \log(n)}\right\} \\
&\leq 2ne^{-[\sqrt{2\nu^2 s} + \sqrt{2\nu^2 \log(n)}]^2/(2\nu^2)} \\
&= 2ne^{-\frac{1}{2\nu^2}(2s\nu^2) - \frac{2\nu^2 \log(2n)}{2\nu^2} - \frac{\sqrt{2\nu^2 s}\sqrt{2\nu^2 \log(2n)}}{2\nu^2}} \\
&\leq e^{-\sqrt{\log(2n)s} - s} \\
&\leq e^{-s}
\end{aligned}$$

The first result follows from the iid assumption on S and the union bound, while the second from the assumption that S is sub-Gaussian. The remaining inequalities follow from reorganising terms. □

Corollary 3.1 is useful as it clarifies that if the surprisal is sub-Gaussian, then the maximal surprisal will follow a reversed Weibull distribution, and thus is within the class of extreme value distributions.

3.2 S has an exponential moment

In cases where the underlying density has full support on $\mathbb{R}$ and the log density cannot be bounded from below, then the sub-Gaussianity condition in Assumption 3.1 may be violated, as the following result illustrates.

Lemma 3.2. *If $Y \sim \mathcal{N}(\mu, \sigma^2)$, then $S = -\log f(Y)$ is sub-exponential. In particular,*

$$\mathbb{E}e^{\lambda(S-1/2)+(\lambda/2)\log(2\pi\sigma^2)} \leq e^{\lambda^2/2+(\lambda/2)\log(2\pi\sigma^2)}, \quad \forall|\lambda| < 1/2.$$

Proof. Firstly, note that

$$S = -\log f(Y) = \frac{1}{2}\log(2\pi\sigma^2) + \frac{(Y-\mu)^2}{2\sigma^2}.$$

and that

$$\mu_S = \mathbb{E}[S] = -\int f(y)\log f(y)\mathrm{d}y = \frac{1}{2}\{1 + \log(2\pi\sigma^2)\}.$$

Define

$$X = \frac{1}{2}\frac{(Y-\mu)^2}{\sigma^2} - \frac{1}{2},$$

then $\mathbb{E}[X] = 0$. Now, note that

$$\mathbb{E}e^{\frac{\lambda}{2}(S-1)} = e^{\lambda/2\{\log(2\pi\sigma^2)\}} \frac{1}{\sqrt{2\pi\sigma^2}} \int e^{\frac{\lambda}{2}\left[\frac{(y-\mu)^2}{\sigma^2}-1\right]} e^{-\frac{(y-\mu)^2}{2\sigma^2}} \mathrm{d}y.$$

Consider the change of variables $z = (y-\mu)/\sigma$ so that we have

$$\mathbb{E}e^{\frac{\lambda}{2}(S-1)} = e^{\lambda/2\{\log(2\pi\sigma^2)\}} \frac{1}{\sqrt{2\pi}} \int e^{\frac{\lambda}{2}[z^2-1]} e^{-z^2/2} \mathrm{d}y = e^{\lambda/2\{\log(2\pi\sigma^2)\}} \frac{e^{-2(\lambda/2)}}{\sqrt{1-2(\lambda/2)}}$$

for all $\lambda < 1$. Further, it can be verified that

$$\frac{e^{-\lambda/2}}{\sqrt{1-2(\lambda/2)}} \leq e^{\lambda^2/2}, \quad \forall |\lambda| < 1/2.$$

Hence, when we have $Y \sim \mathcal{N}(\mu, \sigma^2)$, then $S = -\log f(Y)$ is sub-exponential with parameter $\nu = 1$ and $b = 2$. □

Therefore, to be able to apply extreme value theory to the surprisal values, we need a version of Lemma 3.1 that uses sub-exponentiality.

Lemma 3.3. *Suppose $\boldsymbol{y}_1, \ldots, \boldsymbol{y}_n$ are iid from a density f, and let $M_n = \max\{s_1, \ldots, s_n\}$ where $s_i = -\log f(\boldsymbol{y}_i)$. If Assumption 3.2 is satisfied, then, for $s \in \mathbb{R}_+$,*

$$\Pr(M_n - \mathbb{E}[S] \geq s) \leq 1 - \begin{cases} (1-e^{-\frac{1}{2}s^2/\nu^2})^n & \text{if } 0 \leq s \leq \nu^2/b \\ (1-e^{-s/2b})^n & \text{if } s > \nu^2/b \end{cases}.$$

Proof. The result follows a similar argument as for Lemma 3.1, up until we maximize the tail bound. In particular, for any $\lambda > 0$,

$$\begin{aligned} \Pr\{S - \mathbb{E}[S] \geq s\} = \Pr\left\{e^{\lambda(S-\mathbb{E}[S])} \geq e^{\lambda s}\right\} &\leq e^{-\lambda s}\mathbb{E}e^{\lambda(S-\mathbb{E}[S])} \\ &\leq e^{-\lambda s}e^{\lambda^2\nu^2/2}. \end{aligned}$$

Now, we must consider a constrained optimization to ensure the right hand side exists. In particular, we know from the proof of Lemma 3.1 that the unconstrained optimum is $\lambda = s/\nu^2$, which corresponds to the constrained optimum on the set $0 \leq t < \nu^2/b$. However, when $s \geq \nu^2/b$, the maximum is achieved at $\lambda = 1/b$, and putting this together yields

$$\Pr\{S - \mathbb{E}[S] \geq s\} \leq \begin{cases} e^{-\frac{1}{2}s^2/\nu^2} & \text{if } 0 \leq s \leq \nu^2/b; \\ e^{-s/2b} & \text{if } s > \nu^2/b. \end{cases}$$

The stated result again follows by noting that

$$\begin{aligned}\Pr\{M_n - \mathbb{E}[S] \geq s\} &= 1 - \Pr\{M_n \leq s\} \\ &= 1 - \Pr\{S \leq s\}^n \\ &\leq 1 - \begin{cases} (1 - e^{-\frac{1}{2}s^2/v^2})^n & \text{if } 0 \leq s \leq v^2/b; \\ (1 - e^{-s/2b})^n & \text{if } s > v^2/b. \end{cases}\end{aligned}$$

□

Results similar to Lemma 3.3 also hold for $\Pr\{M_n - \mathbb{E}[S] \leq s\}$, and for $\Pr\{|M_n - \mathbb{E}[S]| \geq s\}$ with an additional factor of 2, i.e.,

$$\Pr\{|M_n - \mathbb{E}[S]| \geq s\} \leq 1 - \begin{cases} (1 - 2e^{-\frac{1}{2}s^2/v^2})^n & \text{if } 0 \leq s \leq v^2/b \\ (1 - 2e^{-s/2b})^n & \text{if } s > v^2/b. \end{cases}$$

Corollary 3.2. *Under Assumption 3.2, for $s > v^2/b$, as $n \to \infty$*

$$\Pr(|M_n - \mathbb{E}[S]| \geq (2b)s + (2b)\log(2n)) \leq 1 - e^{-e^{-s}} + o(1/n).$$

Proof. Apply Assumption 3.2 to see that

$$\begin{aligned}\Pr(|M_n - \mathbb{E}[S]| \geq (2b)s + (2b)\log(2n)) &\leq 1 - \left\{1 - 2e^{-[2bs + 2b\log(2n)]/2b}\right\}^n \\ &= 1 - \left(1 - e^{-s}/n\right)^n \\ &= 1 - e^{e^{-s}} - \left[\left(1 - e^{-s}/n\right)^n - e^{e^{-s}}\right].\end{aligned}$$

The result now follows by noting that, for all $s > 0$,

$$\left[\left(1 - e^{-s}/n\right)^n - e^{e^{-s}}\right] = o(1) \leq 0.$$

□

Corollary 3.2 is useful as it shows that if the surprisal has a sub-exponential tail, then the resulting maximum surprisal tail probability converges to that of an extreme value distribution. In particular, we have that Corollary 3.2 implies

$$\sup_{s:s>v^2/b} \left|\Pr\left\{|M_n - \mu| \geq \left[2bs + 2b\log(2n)\right]\right\} - \left(1 - e^{e^{-s}}\right)\right| = o(1).$$

To see that this implies that $M_n - \mathbb{E}[S]$, appropriately standardized, converges to an extreme value distribution, we first recall that

$$\Pr\{X_n > x\} - \Pr\{X > x\} = 1 - \Pr\{X_n \le x\} - [1 - \Pr\{X \le x\}] = \Pr\{X \le x\} - \Pr\{X_n \le x\}.$$

Hence,

$$\sup_{x \in \mathcal{X}} |\Pr\{X_n \le x\} - \Pr\{X \le x\}| = \sup_{x \in \mathcal{X}} |\Pr\{X_n > x\} - \Pr\{X > x\}|,$$

so that when X_n and X are both continuous we see that Corollary 3.2 implies that the CDF of $(M_n - \mathbb{E}[S] - 2b\log(2n))/2b$ converges to that of the Type-1 extreme value distribution (i.e., Gumbel distribution); i.e., using Corollary 3.2 and the above we have that

$$\sup_{s : s > v^2/b} \left| \Pr\left(|M_n - \mu| \le [2bs + 2b\log(2n)]\right\} - e^{e^{-s}} \right| = o(1).$$

This is one notion of convergence in distribution.

3.3 S has a polynomial moment

Finally, we consider the case where the surprisal has a polynomial moment (e.g., Student-t random variables), and derive a corresponding result to Lemma 3.1.

Suppose $\boldsymbol{y}_1, \ldots, \boldsymbol{y}_n$ are iid from a density f, and let $M_n = \max\{s_1, \ldots, s_n\}$ where $s_i = -\log f(\boldsymbol{y}_i)$. If Assumption 3.3 is satisfied, then, for $s > 0$ and such that $s > C > 0$,

$$\Pr\left(|M_n - \mathbb{E}[S]| \ge s\right) \le 1 - (1 - C^p/s^p)^n.$$

Hence,

$$\sup_{s : s > C} \left| \Pr\left(|M_n - \mathbb{E}[S]| \le (s \cdot n^{1/p})\right) - e^{-s^{-p}} \right| = o(1).$$

Proof. By Markov's inequality,

$$\Pr\{|S - \mathbb{E}[S]| \ge s\} \le \frac{\mathbb{E}|S|^p}{s^p} \le \frac{C^p}{s^p}.$$

To derive the first stated result, again note that

$$\begin{aligned}
\Pr\{|M_n - \mathbb{E}[S]| \ge s\} &= 1 - \Pr\{|M_n - \mathbb{E}[S]| \le s\} \\
&= 1 - \Pr\{|S - \mathbb{E}[S]| \le s\}^n \\
&= 1 - (1 - \Pr\{|S - \mathbb{E}[S]| \ge s\})^n \\
&\le 1 - (1 - C^p/s^p)^n.
\end{aligned}$$

To derive the second stated result, apply the first result with $t = sn^{1/p}C > 1$, which clearly holds for some n large enough, to see that

$$\Pr\{|M_n - \mathbb{E}[S]| \geq t\} \leq 1 - \left(1 - \frac{C^p}{(Csn^{1/p})^p}\right)^n = 1 - \left(1 - s^{-p}/n\right)^n.$$

Now, recall that, for $x \geq 0$,

$$\lim_{n\to\infty} (1 - x/n)^n = e^{-x}.$$

Since $x \mapsto e^{-x}$ is bounded for all $x \geq 0$, it follows that

$$\sup_{s:s>C} \left|\Pr\{|M_n - \mathbb{E}[S]| \geq t\} - e^{-s^{-p}}\right| = o(1).$$

□

3.4 Three types theorem

Placing the three lemmas together gives the following "three types" Theorem.

Theorem 3.1. *Let $\boldsymbol{y}_1, \ldots, \boldsymbol{y}_n$ be an iid sequence from F, and let $S = -\log f(\boldsymbol{Y})$, where $Y \sim \boldsymbol{F}$. Further, let $M_n = \max\{s_1, \ldots, s_n\}$, where $s_i = -\log f(y_i)$.*

1. *If the surprisal satisfies Assumption 3.1, then*

$$\sup_{s:s>0} \left|\Pr\left\{\left|M_n - \mathbb{E}[S]\right| \geq \sqrt{2v^2 s} + \sqrt{2v^2 \log(2n)}\right\} - e^{-s}\right| = o(1).$$

2. *If the surprisal satisfies Assumption 3.2, then*

$$\sup_{s:s>1/b} \left|\Pr\left\{\left|M_n - \mathbb{E}[S]\right| \geq (2b)s + (2b)\log(2n)\right\} - e^{-e^{-s}}\right| = o(1).$$

3. *If the surprisal satisfies Assumption 3.3, then*

$$\sup_{s:s>C} \left|\Pr\left\{\left|M_n - \mathbb{E}[S]\right| \geq (Csn^{1/p})\right\} - e^{-s^{-p}}\right| = o(1).$$

Theorem 3.1 gives a corresponding 'three types' theorem for the behavior of surprisals and has a simple interpretation. If the surprisal has Gaussian like tails, then the maximum surprisal is a reversed Weibull; if it only has an exponential moment, then the maximum surprisal will be Gumbel; if the maximum surprisal only has a polynomial moment, then it is Fréchet. As each of these distributions is a member of the generalized extreme value family, then the upper tail of the distribution can be approximated by a Generalized Pareto Distribution (Coles 2001, Theorem 4.1).

4 Experiments

4.1 Standard normal values

The use of a GPD to estimate the tail probabilities of the surprisal distribution is designed to be robust to incorrect assumptions about the underlying distribution F. Suppose we generate 1000 observations from a N(0,1) distribution, but use a t_4 distribution to compute the surprisal values. The GPD approach should still give reasonable estimates of the tail probabilities, despite the incorrect distribution being used to compute the surprisals.

To study the effect of the assumed distribution, we compute the surprisal values using both the correct Normal density and an incorrect t_4 density. Then, for each set of surprisal values, we estimate the surprisal tail probabilities using the same distribution used to compute the surprisal values, and alternatively using a GPD fitted to the largest 100 surprisal values.




Figure 1: *Left: Observations are computed from a N(0,1) distribution. Right: Observations are computed from a t_4 distribution. In each cases, surprisals are computed using the correct distribution and the incorrect distribution, and tail probabilities are computed using the assumed distribution, the empirical distribution and a GPD applied to the surprisal values. This demonstrates that the surprisal probabilities computed using the empirical distribution or under the GPD are still relatively accurate, even when the wrong distribution is used to compute the surprisal values.*

The various estimates are displayed in the left panel of Figure 1 for observations greater than 2.5, showing how the GPD estimates are particularly accurate in the extreme tails, where we need them for anomaly detection. The true values are shown in orange ("Normal"). The two GPD estimates (blue and pink) are almost identical despite the incorrect distribution being used to compute the surprisals for one of them. When the incorrect distribution is also used to compute the probabilities (green), the results are very inaccurate.

In summary, even when we don't know the distribution of the data, we can use some incorrect assumed distribution to compute the surprisals, and then apply either a GPD or the empirical distribution to obtain good estimates of the surprisal probabilities.

The converse situation is shown in the right panel of Figure 1, where the data are generated from a t_4 distribution, and the surprisal values are computed using both the correct t_4 density and an incorrect N(0,1) density. Again, the GPD estimates of the surprisal probabilities are accurate in the extreme tails, even when the wrong distribution is used to compute the surprisal values.

4.2 Gamma values

- 990 from Gamma(2,2) and 10 from Gamma(2,r). Fit Gamma to all 1000 and compute surprisals. Fit GPD to largest 50 surprisals. Compare true and estimated tail probabilities.
- 990 from trivariate N(0,1) with 10 from ring at distance 2 from origin. Fit bivariate normal to all 1000 and compute surprisals. Fit GPD to largest 50 surprisals. Compare true and estimated tail probabilities.

5 Applications

- US COVID death rates: https://otexts.com/weird/09-timeseries.html#multivariate-time-series
- https://otexts.com/weird/06-density.html#gam-surprisals

6 Discussion

Software and reproducibility

All results presented here can be reproduced using the code available at https://github.com/robjhyndman/Surprisal_Theory. The analysis was conducted using R version 4.5.1 (R Core Team 2025), with the following R packages: weird (Hyndman 2024), tsibble (Wang et al. 2025), targets (Landau 2025, 2021), ggplot2 (Wickham et al. 2025; Wickham 2016), and other tidyverse (Wickham et al. 2019) packages.

Acknowledgements

Both authors are funded by the Australian Research Council through Discovery Project DP250100702. Rob Hyndman is a member of the Australian Research Council Industrial Transformation Training Centre in Optimisation Technologies, Integrated Methodologies, and Applications (OPTIMA), Project ID IC200100009.

References

Coles, S (2001). *An introduction to statistical modeling of extreme values*. Vol. 208. Springer Series in Statistics. London UK: Springer.

Cover, TM & JA Thomas (2006). *Elements of information theory*. 2nd. Hoboken, NJ: Wiley.

Good, IJ (1952). Rational decisions. *Journal of the Royal Statistical Society. Series B, Statistical methodology* **14** (1), 107–114.

Hyndman, RJ (1996). Computing and graphing highest density regions. *The American Statistician* **50**(2), 120–126.

Hyndman, RJ (2024). *weird: Functions and Data Sets for "That's Weird: Anomaly Detection Using R" by Rob J Hyndman*. Version 1.0.2. https://pkg.robjhyndman.com/weird-package/.

Hyndman, RJ, S Kandanaarachchi & K Turner (2025). *When lookout sees crackle: Anomaly detection via kernel density estimation*. Tech. rep. Monash University.

Kandanaarachchi, S & RJ Hyndman (2022). Leave-one-out kernel density estimates for outlier detection. *J Computational & Graphical Statistics* **31** (2), 586–599.

Landau, WM (2021). The targets R package: a dynamic Make-like function-oriented pipeline toolkit for reproducibility and high-performance computing. *Journal of Open Source Software* **6**(57), 2959.

Landau, WM (2025). *targets: Dynamic Function-Oriented 'Make'-Like Declarative Pipelines*. Version v1.11.3. https://cran.r-project.org/package=targets.

R Core Team (2025). *R: A Language and Environment for Statistical Computing*. R Foundation for Statistical Computing. Vienna, Austria. https://www.R-project.org/.

Stone, JV (2022). *Information theory: a tutorial introduction*. Sheffield, England: Sebtel Press.

Tribus, M (1961). *Thermodynamics and Thermostatics: An Introduction to Energy, Information and States of Matter, with Engineering Applications*. New York, USA: D. Van Nostrand.

Wang, E, D Cook, RJ Hyndman, M O'Hara-Wild, T Smith & W Davis (2025). *tsibble: Tidy Temporal Data Frames and Tools*. Version 1.1.6. https://tsibble.tidyverts.org.

Wickham, H (2016). *ggplot2: Elegant Graphics for Data Analysis*. New York, USA: Springer-Verlag. https://ggplot2-book.org/.

Wickham, H, M Averick, J Bryan, W Chang, LD McGowan, R François, G Grolemund, A Hayes, L Henry, J Hester, M Kuhn, TL Pedersen, E Miller, SM Bache, K Müller, J Ooms, D Robinson, DP Seidel, V Spinu, K Takahashi, D Vaughan, C Wilke, K Woo & H Yutani (2019). Welcome to the tidyverse. *Journal of Open Source Software* **4**(43), 1686.

Wickham, H, W Chang, L Henry, TL Pedersen, K Takahashi, C Wilke, K Woo, H Yutani, D Dunnington, T van den Brand & Posit, PBC (2025). *ggplot2: Create Elegant Data Visualisations Using the Grammar of Graphics*. Version v3.5.2. https://cran.r-project.org/package=ggplot2.

Wilkinson, L (2018). Visualizing Big Data Outliers Through Distributed Aggregation. *IEEE Transactions on Visualization and Computer Graphics* **24**(1), 256–266.