Anomaly detection using surprisals

Rob J Hyndman and David T Frazier

24 September 2025

Abstract

TBC

Keywords: Extreme value theory; Generalized Pareto Distribution; Outlier detection; Tail bounds.

1 Introduction

Let $y_1, \dots, y_n$ be observations from a probability distribution F. The *surprisal* of observation y_i is defined as (Stone 2022)

$$s_i = -\log f(y_i),$$

where f is either the density or probability mass function of F. While these are better known as "log scores" in statistics (Good 1952), in this context we prefer the term "surprisal" (coined by Tribus 1961), as it captures the idea that observations with low density are more "surprising" (and therefore potentially anomalous) than those with high density.

Let $G(s) = P(S \leq s)$ be the distribution function of the surprisal values, where $S = -\log f(Y)$ and $Y \sim F$. Then, the probability of observing a value with surprisal at least as large as s is given by $1 - G(s)$, which leads to a natural way to assign an anomaly score to each observation y_i:

$$p_i = 1 - G(s_i) = 1 - G(-\log f(y_i)).$$

Hence, one way to detect anomalies is to flag observations with $p_i < \alpha$, giving a false positive rate of α. This is preferable to directly computing the probabilities from the tail of F, as it allows for anomalies to occur anywhere in the distribution of F, not just in its tails. For example, anomalies may lie between modes in a multimodal distribution, or in otherwise low-density regions of the distribution F.

However, we will usually not know F, and so we need to estimate it from data or from theory. Such estimates are particularly fragile in areas of low density, which is precisely where we are looking for anomalies.

To address this, Kandanaarachchi & Hyndman (2022) proposed estimating F by a leave-one-out kernel density estimate, and then fitting a Generalized Pareto Distribution (GPD) to the most extreme surprisal values. Hyndman et al. (2025) provides some theoretical justification for this approach. The

underlying idea is that even if F is not particularly well-estimated, the tail of the surprisal distribution should still be well-approximated by a GPD.

Here we extend that idea to allow F to be any well-behaved distribution, including conditional distributions. Specifically, we fit a GPD to the largest β of $\{-\log f_i\}_{i=1}^n$, obtaining estimates of the location, scale and shape parameters, (μ, σ, ξ). We calculate the probability of each observation or one more extreme, $p_i = \beta\left[1 - P(-\log f_i \mid \hat{\mu}, \hat{\sigma}, \hat{\xi})\right]$, where P is the distribution function of the GPD. Observations not contained in the largest β of $\{-\log f_i\}_{i=1}^n$ are assigned $p_i = \beta$.

To justify this approach, we seek conditions on the surprisal distribution of $S = -\log f(Y)$, where $Y \sim F$, similar to the Fisher-Tippett-Gnedenko (FTG) theorem (Coles 2001).

2 Assumptions and preliminary results

Before establishing the main results, we need to place some assumptions on the data generating process.

Assumption 1. *The sequence y_i is iid from F.*

Under Assumption 1, we have that $s_i = -\log f(y_i)$ is also iid, and its distribution function is given by

$$G(s) = \Pr(S \leq s) = \Pr\{-\log f(Y) \leq s\} = \Pr\{f(Y) \geq e^{-s}\}.$$

This makes clear that the surprisal probability is related to the highest density regions of the distribution (Hyndman 1996). In fact, every observation with $p_i < \alpha$ lies outside the $100(1-\alpha)\%$ HDR region of F.

Next we need to place some conditions on the surprisal distribution. We consider three possibilities describing increasingly heavy tails: (1) where S is sub-Gaussian (which occurs when f is bounded above and below); (2) where S is sub-exponential; and (3) where S has a polynomial moment. We state these as three alternative assumptions.

Assumption 2 (Sub-Gaussian)**.** *The random variable $S = -\log f(Y) \in \mathbb{R}$, and satisfies, for all $\lambda \in \mathbb{R}$, and some $\nu > 0$,*

$$\mathbb{E}\exp\{\lambda(S - \mathbb{E}[S])\} \leq \exp\{\lambda^2\nu^2/2\}.$$

Assumption 3 (Sub-exponential)**.** *The random variable S is sub-exponential with parameters ν and b, i.e.,*

$$\mathbb{E}\exp\{\lambda(S - \mathbb{E}[S])\} \leq \exp\{\lambda^2\nu^2/2\} \quad \textit{for all} \quad |\lambda| < 1/b.$$

Assumption 4 (Polynomial)**.** *The random variable $|S|$ has polynomial moments of order $\alpha \geq 1$; i.e., $\mathbb{E}[|S|^p] \leq C^p$ for some $C > 0$ such that $C^p - 1 > 0$.*

Under Assumption 1, and one of Assumptions 2–4, we seek a result on the limiting distribution of the maximum of the surprisals $M_n = \max\{s_1, \ldots, s_n\}$. If we can show that this distribution falls within

the class of extreme value distributions, then we can fit a Generalized Pareto Distribution to the largest surprisal values (Coles 2001).

2.1 S has sub-Gaussian tails

Let's first assume that the surprisals are well-behaved and have a sub-Gaussian tail. Assumption 2 will be satisfied in cases where the density f is bounded both above and below, such as with mass functions on a finite set. However, it may not be satisfied if the density has no lower bound.

For example, consider the truncated normal distribution on $[a, b]$, where $Y \sim \mathcal{TN}(a, b, \mu, \sigma^2)$ and $-\infty < a < b < \infty$. Then

$$S = -\log(2\pi\sigma^2)/2 - \frac{1}{2\sigma^2}(Y-\mu)^2 - \log\{\Phi(b) - \Phi(a)\},$$

where Φ is the standard normal distribution function. Since $Y \in [a, b]$, S is bounded for all $\sigma > 0$ and hence S satisfies Assumption 2.

Under Assumptions 1 and 2, we have the following result on the maximum of the surprisals $M_n = \max\{s_1, \ldots, s_n\}$.

Lemma 1. *If Assumptions 1–2 are satisfied, then, for $s \in \mathbb{R}_+$,*

$$\Pr(M_n - \mathbb{E}[S] \geq s) \leq 1 - \left\{1 - e^{-\frac{1}{2}s^2/v^2}\right\}^n.$$

Proof. For any $\lambda > 0$,

$$\begin{aligned}\Pr\{S \geq s\} = \Pr\left\{e^{\lambda(S-\mathbb{E}[S])} \geq e^{\lambda(s-\mathbb{E}[S])}\right\} &\leq e^{-\lambda(s-\mu)}\mathbb{E}e^{\lambda(S-\mu)} \\ &\leq e^{-\lambda(s-\mu)}e^{\lambda^2 v^2/2}.\end{aligned}$$

Now, we can maximise the RHS of the above bound in λ, which yields the optimal value $\lambda = (s-\mu)/\{M^2\}$. Plugging this in we have the bound

$$\Pr\{S \geq s\} \leq e^{-\frac{1}{2}(s-\mu)^2/v^2}.$$

The stated result now follows by noting that

$$\begin{aligned}\Pr\left(M_n - \mathbb{E}[S] \geq s\right) = 1 - \Pr\{M_n \leq s\} &= 1 - \Pr\{S \leq s\}^n \\ &= 1 - (1 - \Pr\{S \geq s\})^n \\ &\leq 1 - (1 - e^{-\frac{1}{2}s^2/v^2})^n.\end{aligned}$$

□

Corollary 1. *If Assumptions 1–2 are satisfied, then, for all $s > 0$,*

$$\Pr\left(|M_n - \mathbb{E}[S]| \geq \sqrt{2v^2 s} + \sqrt{2v^2\log(2n)}\right) \leq e^{-s}.$$

Proof. This follows directly from the union bound:

$$\begin{aligned}
\Pr\Big(|M_n| \geq \sqrt{2\nu^2 s} + \sqrt{2\nu^2 \log(n)}\Big) &\leq n \Pr\left\{|S| \geq \sqrt{2}\nu^2 s + \sqrt{2\nu^2 \log(n)}\right\} \\
&\leq 2ne^{-[\sqrt{2\nu^2 s} + \sqrt{2\nu^2 \log(n)}]^2/(2\nu^2)} \\
&= 2ne^{-\frac{1}{2\nu^2}(2s\nu^2) - \frac{2\nu^2 \log(2n)}{2\nu^2} - \frac{\sqrt{2\nu^2 s}\sqrt{2\nu^2 \log(2n)}}{2\nu^2}} \\
&\leq e^{-\sqrt{\log(2n)s} - s} \\
&\leq e^{-s}
\end{aligned}$$

The first result follows from the iid assumption on S and the union bound, while the second from the assumption that S is sub-Gaussian. The remaining inequalities follow from reorganising terms. □

Corollary 1 is useful as it clarifies that if the surprisal is sub-Gaussian, then the maximal surprisal will follow a reversed Weibull distribution, and thus is within the class of extreme value distributions.

2.2 S has an exponential moment

In cases where the underlying density has full support on $\mathbb{R}$ and cannot be bounded from below, then the sub-Gaussianity condition in Assumption 2 may be violated, as the following result illustrates.

Lemma 2. *If $Y \sim \mathcal{N}(\mu, \sigma^2)$, then $S = -\log f(Y)$ is sub-exponential. In particular,*

$$\mathbb{E}e^{\lambda(S-1/2)+(\lambda/2)\log(2\pi\sigma^2)} \leq e^{\lambda^2/2+(\lambda/2)\log(2\pi\sigma^2)}, \quad \forall|\lambda| < 1/2.$$

Proof. Firstly, note that

$$S = -\log f(Y) = \frac{1}{2}\log(2\pi\sigma^2) + \frac{(Y-\mu)^2}{2\sigma^2}.$$

and that

$$\mu_S = \mathbb{E}[S] = -\int f(y)\log f(y)\mathrm{d}y = \frac{1}{2}\{1 + \log(2\pi\sigma^2)\}.$$

Define

$$X = \frac{1}{2}\frac{(Y-\mu)^2}{\sigma^2} - \frac{1}{2},$$

then $\mathbb{E}[X] = 0$. Now, note that

$$\mathbb{E}e^{\frac{\lambda}{2}(S-1)} = e^{\lambda/2\{\log(2\pi\sigma^2)\}}\frac{1}{\sqrt{2\pi\sigma^2}}\int e^{\frac{\lambda}{2}\left[\frac{(y-\mu)^2}{\sigma^2}-1\right]}e^{-\frac{(y-\mu)^2}{2\sigma^2}}\mathrm{d}y.$$

Consider the change of variables $z = (y-\mu)/\sigma$ so that we have

$$\mathbb{E}e^{\frac{\lambda}{2}(S-1)} = e^{\lambda/2\{\log(2\pi\sigma^2)\}}\frac{1}{\sqrt{2\pi}}\int e^{\frac{\lambda}{2}[z^2-1]}e^{-z^2/2}\mathrm{d}y = e^{\lambda/2\{\log(2\pi\sigma^2)\}}\frac{e^{-2(\lambda/2)}}{\sqrt{1-2(\lambda/2)}}$$

for all $\lambda < 1$. Further, it can be verified that

$$\frac{e^{-\lambda/2}}{\sqrt{1-2(\lambda/2)}} \leq e^{\lambda^2/2}, \quad \forall |\lambda| < 1/2.$$

Hence, when we have $Y \sim \mathcal{N}(\mu, \sigma^2)$, then $S = -\log f(Y)$ is sub-exponential with parameter $\nu = 1$ and $b = 2$. □

Therefore, to be able to apply extreme value theory to the surprisal values, we need a version of Lemma 1 that uses sub-exponentiality.

Lemma 3. *If Assumptions 1 and 3 are satisfied, then, for $s \in \mathbb{R}_+$,*

$$\Pr\left(M_n - \mathbb{E}[S] \geq s\right) \leq 1 - \begin{cases} (1 - e^{-\frac{1}{2}s^2/\nu^2})^n & \text{if } 0 \leq s \leq \nu^2/b \\ (1 - e^{-s/2b})^n & \text{if } s > \nu^2/b \end{cases}.$$

Proof. The result follows a similar argument as for Lemma 1, up until we maximize the tail bound. In particular, for any $\lambda > 0$,

$$\begin{aligned} \Pr\left\{S - \mathbb{E}[S] \geq s\right\} = \Pr\left\{e^{\lambda(S-\mathbb{E}[S])} \geq e^{\lambda s}\right\} &\leq e^{-\lambda s}\mathbb{E}e^{\lambda(S-\mathbb{E}[S])} \\ &\leq e^{-\lambda s}e^{\lambda^2\nu^2/2}. \end{aligned}$$

Now, we must consider a constrained optimization to ensure the RHS exists. In particular, we know that from the proof of Lemma 1 that the unconstrained optimum is $\lambda = s/\nu^2$, which corresponds to the constrained optimum on the set $0 \leq t < \nu^2/b$. However, when $s \geq \nu^2/b$, the maximum is achieved at $\lambda = 1/b$, and putting this together yields

$$\Pr\left\{S - \mathbb{E}[S] \geq s\right\} \leq \begin{cases} e^{-\frac{1}{2}s^2/\nu^2} & \text{if } 0 \leq s \leq \nu^2/b \\ e^{-s/2b} & \text{if } s > \nu^2/b \end{cases}.$$

The stated result again follows by noting that

$$\begin{aligned} \Pr\left\{M_n - \mathbb{E}[S] \geq s\right\} &= 1 - \Pr\left\{M_n \leq s\right\} \\ &= 1 - \Pr\left\{S \leq s\right\}^n \\ &\leq 1 - \begin{cases} (1 - e^{-\frac{1}{2}s^2/\nu^2})^n & \text{if } 0 \leq s \leq \nu^2/b \\ (1 - e^{-s/2b})^n & \text{if } s > \nu^2/b \end{cases}. \end{aligned}$$

□

Results similar to Lemma 3 also hold for $\Pr\left\{M_n - \mathbb{E}[S] \leq s\right\}$, and for $\Pr\left\{|M_n - \mathbb{E}[S]| \geq s\right\}$ with an

additional factor of 2, i.e.,

$$\Pr\{|M_n - \mathbb{E}[S]| \geq s\} \leq 1 - \begin{cases} (1 - 2e^{-\frac{1}{2}s^2/v^2})^n & \text{if } 0 \leq s \leq v^2/b \\ (1 - 2e^{-s/2b})^n & \text{if } s > v^2/b \end{cases}.$$

Corollary 2. *Under Assumptions 1 and 3, for $s > v^2/b$, as $n \to \infty$*

$$\Pr\left(|M_n - \mathbb{E}[S]| \geq (2b)s + (2b)\log(2n)\right) \leq 1 - e^{-e^{-s}} + o(1/n).$$

Proof. Apply Assumption 3 to see that

$$\begin{aligned} \Pr\left(|M_n - \mathbb{E}[S]| \geq (2b)s + (2b)\log(2n)\right) &\leq 1 - \left\{1 - 2e^{-[2bs + 2b\log(2n)]/2b}\right\}^n \\ &= 1 - \left(1 - \frac{e^{-s}}{n}\right)^n \\ &= 1 - e^{e^{-s}} - \left[\left(1 - \frac{e^{-s}}{n}\right)^n - e^{e^{-s}}\right]. \end{aligned}$$

The result now follows by noting that, for all $s > 0$,

$$\left[\left(1 - \frac{e^{-s}}{n}\right)^n - e^{e^{-s}}\right] = o(1) \leq 0.$$

□

Corollary 2 is useful as it shows that if the surprisal has a sub-exponential tail, then the resulting maximum surprisal tail probability converges to that of an extreme value distribution. In particular, we have that Corollary 2 implies

$$\sup_{s:s>v^2/b} \left|\Pr\left(|M_n - \mu| \geq [2bs + 2b\log(2n)]\right) - \left(1 - e^{e^{-s}}\right)\right| = o(1).$$

To see that this implies that $M_n - \mathbb{E}[S]$, appropriately standardized, converges to an extreme value distribution, we first recall that

$$\Pr\{X_n > x\} - \Pr\{X > x\} = 1 - \Pr\{X_n \leq x\} - [1 - \Pr\{X \leq x\}] = \Pr\{X \leq x\} - \Pr\{X_n \leq x\}.$$

Hence,

$$\sup_{x \in \mathcal{X}} |\Pr\{X_n \leq x\} - \Pr\{X \leq x\}| = \sup_{x \in \mathcal{X}} |\Pr\{X_n > x\} - \Pr\{X > x\}|,$$

so that when X_n and X are both continuous we see that Corollary 2 implies that the CDF of $(M_n - \mathbb{E}[S] - 2b\log(2n))/2b$ converges to that of the Type-1 extreme value distribution (i.e., Gumbel distribution); i.e., using Corollary 2 and the above we have that

$$\sup_{s:s>v^2/b} \left|\Pr\left(|M_n - \mu| \leq [2bs + 2b\log(2n)]\right\} - e^{e^{-s}}\right| = o(1).$$

This is one notion of convergence in distribution.

Recall that, for $Y \sim \mathcal{N}(\mu, \sigma^2)$, we have that $S = -\log f(Y)$ is sub-exponential with $\nu = 1$ and $b = 2$. Further,

$$\mu = \mathbb{E}[S] = -\int f(y) \log f(y) \mathrm{d}y = \frac{1}{2}\{1 + \log(2\pi\sigma^2)\}.$$

Hence, Lemma 3 immediately implies that, for $s > 1/2$,

$$\Pr\left\{ M_n \geq \frac{1}{2}\{1 + \log(2\pi\sigma^2)\} + 4s + 4\log(n) \right\} \leq 1 - (1 - e^{-s}/n)^n$$

or

$$\Pr\left\{ \left| M_n - \frac{1}{2}\{1 + \log(2\pi\sigma^2)\} \right| \geq 4s + 4\log(2n) \right\} \leq 1 - (1 - e^{-s}/n)^n$$

Consider that $\sigma^2 = 1$, then the mean surprisal is $\frac{1}{2}\{1 + \log(2\pi\sigma^2)\} = .919$. Given n observations, what is the probability that the absolute difference between the maximum surprisal and the mean is larger than four times the mean for various choices of n? Using the above we have that

$$\Pr\left\{ |M_n - \frac{1}{2}\{1 + \log(2\pi\sigma^2)\}| \geq 4 * \frac{1}{2}\{1 + \log(2\pi\sigma^2)\} + 4\log(2n) \right\} \leq 1 - \left(1 - \frac{e^{-n*.918}}{n}\right)^n$$

For $n = 10, 20, 50, 100$ the estimate of the probability is between .565 and .551. However, if we let $\sigma^2 = 10$, then the mean surprisal is about 3.74, and, for the same value of n, the probabilities range from about .225 to .223. This illustrates that the surprisal probability actually decreases as σ increases.

2.3 S has a polynomial moment

Finally, we consider the case where the surprisal only has a polynomial moment, and derive a corresponding result to Lemma 1.

Lemma 4. *If Assumptions 1 and 4 are satisfied, then, for $s > 0$ and such that $s > C > 0$,*

$$\Pr\left(|M_n - \mathbb{E}[S]| \geq s\right) \leq 1 - (1 - C^p/s^p)^n.$$

Hence,

$$\sup_{s:s>C} \left| \Pr\left(|M_n - \mathbb{E}[S]| \leq (s \cdot n^{1/p})\right) - e^{-s^{-p}} \right| = o(1).$$

Proof. By Markov's inequality,

$$\Pr\left\{|S - \mathbb{E}[S]| \geq s\right\} \leq \frac{\mathbb{E}|S|^p}{s^p} \leq \frac{C^p}{s^p}.$$

To derive the first stated result, again note that

$$\begin{aligned}\Pr\{|M_n - \mathbb{E}[S]| \geq s\} &= 1 - \Pr\{|M_n - \mathbb{E}[S]| \leq s\} \\ &= 1 - \Pr\{|S - \mathbb{E}[S]| \leq s\}^n \\ &= 1 - (1 - \Pr\{|S - \mathbb{E}[S]| \geq s\})^n \\ &\leq 1 - (1 - C^p/s^p)^n.\end{aligned}$$

To derive the second stated result, apply the first result with $t = sn^{1/p}C > 1$, which clearly holds for some n large enough, to see that

$$\Pr\{|M_n - \mathbb{E}[S]| \geq t\} \leq 1 - \left(1 - \frac{C^p}{(Csn^{1/p})^p}\right)^n = 1 - \left(1 - n^{-1}s^{-p}\right)^n.$$

Now, recall that, for $x \geq 0$,

$$\lim_{n\to\infty}\left(1 - \frac{x}{n}\right)^n = e^{-x}.$$

Since $x \mapsto e^{-x}$ is bounded for all $x \geq 0$, it follows that

$$\sup_{s:s>C}\left|\Pr\{|M_n - \mathbb{E}[S]| \geq t\} - e^{-s^{-p}}\right| = o(1).$$

□

Lemma 4 gives us a result on the behavior of surprisals that only possess a polynomial tail, e.g., student-t random variables.

2.4 Three types theorem

Placing the three lemmas together gives the following "three types" Theorem.

Theorem 1. *Let $y_1, \ldots, y_n$ be an iid sequence from F, and let $S = -\log f(Y)$, where $Y \sim F$. Further, let $M_n = \max\{s_1, \ldots, s_n\}$, where $s_i = -\log f(y_i)$.*

1. *If the surprisal satisfies Assumption 2, then*

$$\sup_{s:s>0}\left|\Pr\left\{|M_n - \mathbb{E}[S]| \geq \sqrt{2\nu^2 s} + \sqrt{2\nu^2\log(2n)}\right\} - e^{-s}\right| = o(1).$$

2. *If the surprisal satisfies Assumption 3, then*

$$\sup_{s:s>1/b}\left|\Pr\{|M_n - \mathbb{E}[S]| \geq (2b)s + (2b)\log(2n)\} - e^{-e^{-s}}\right| = o(1).$$

3. *If the surprisal satisfies Assumption 4, then*

$$\sup_{s:s>C}\left|\Pr\left\{|M_n - \mathbb{E}[S]| \geq (Csn^{1/p})\right\} - e^{-s^{-p}}\right| = o(1).$$

Theorem 1 gives a corresponding 'three types' theorem for the behavior of surprisals and has a simple interpretation. If the surprisal has Gaussian like tails, then the maximum surprisal is a reversed Weibull; if it only has an exponential moment, then the maximum surprisal will be Gumbel; if the maximum surprisal only has a polynomial moment, then it is Frechet.

3 Theorem 1 in Action

In this section, I have considered a few numerical experiments that illustrate the validity of Theorem 1. The essence of Theorem 1 is that, as $n \to \infty$, the distribution of M_n behaves like a generalized extreme value distribution,

$$\text{Gev}(x; \mu, \sigma, \xi) = \exp\left\{-\left(1 + \xi\frac{x-\mu}{\sigma}\right)^{-1/\xi}\right\},$$

defined when $1 + \xi(x-\mu)/\sigma > 0; \mu \in \Re, \sigma > 0$ and $\xi \in \mathbb{R}$. This distribution takes specific forms depending on the particular value of the shape parameter ξ and changes depending on this value.

1. The (reverse) Weibull distribution is obtained when $\xi < 0$.
2. The Gumble distirbution is obtained when $\xi = 0$.
3. The Frechet distribution is obtained when $\xi > 0$.

Thus, in order to validate the results of Theorem 1, all we need to show is that:

1. when S is sub-Gaussian, the distribution of the maximum surprisal M_n has shape parameter $\xi < 0$;
2. when S is sub-exponential, the distribution of the maximum surprisal M_n has shape parameter $\xi = 0$;
3. when S only has a polynomial moment, the distribution of the maximum surprisal M_n has shape parameter $\xi > 0$.

Example 1. *Let us return to the truncated normal distribution example. Consider that we observe data generated according to*

$$Y_i \overset{iid}{\sim} \mathcal{TN}(a = 1, b = 5, \mu = 2, \sigma = 1) \ (i = 1, \ldots, n).$$

In this example, since $Y \in [1, 5]$, it is simple to show that surprisal S_i is bounded. Hence, we would expect that $M_n = \max\{S_1, \ldots, S_n\}$ should eventually resemble a GEV distribution where the shape parameter $\xi < 0$. To verify this, we generate $R = 1000$ independent samples of size $n = 1000$ from the truncated normal distribution and for each $r = 1, \ldots, 1000$ calculate $M_n^{(r)} = \max\{S_1^{(r)}, \ldots, S_n^{(r)}\}$. We plot the histogram of $\{M_n^{(r)}\}_{r=1}^{1000}$ and compare this with the fitted GEV distribution in Fig. 1. The distribution seems very close to the (reversed) Weibull distribution, and indeed the estimated parameters from this experiment yield an estimated shape parameter $\hat{\xi}_R = -0.2002$, with a standard error of 0.0271, suggesting high confidence that the resulting estimated parameter is negative and that the distribution of M_n is indeed Weibull.




Figure 1: The estimated GEV parameters and standard errors are as follows: location: 5.6839 (0.0314); scale: 0.8676 (0.0229); shape: -0.20002 (0.0271).

Example 2. *Now, let us return to the normal example, which, from our theoretical results, we know should give rise to surprisals that exhibit Gumbel like behavior. To see that this is indeed the case, consider that we observe data generated according to*

$$Y_i \overset{iid}{\sim} \mathcal{N}(\mu = 1, \sigma = 1) \ (i = 1, \ldots, n).$$

In this example, we know that the surprisal S_i will have a sub-exponential tail and that $M_n = \max\{S_1, \ldots, S_n\}$ should eventually resemble a GEV distribution where the shape parameter $\xi = 0$. To verify this, we generate $R = 1000$ independent samples of size $n = 1000$ from the normal distribution with mean and variance both equal to unity, and for each $r = 1, \ldots, 1000$ we calculate $M_n^{(r)} = \max\{S_1^{(r)}, \ldots, S_n^{(r)}\}$. We plot the histogram of $\{M_n^{(r)}\}_{r=1}^{1000}$ and compare this with the fitted GEV distribution in Fig. 2. The estimated parameters from this experiment yield an estimated shape parameter $\hat{\xi}_R = 0.0012$, with a standard error of 0.0209, suggesting high confidence that the actual shape parameter is either zero or very close to it, and that the distribution of M_n is indeed Gumbel.




Figure 2: The estimated GEV parameters and standard errors are given by: location: 6.30016 (0.0332); scale: 0.945129961 (0.0239); shape: -0.0012 (0.0209).

Example 3. *For the last example, we consider the situation where the assumed model is Gaussian with mean and variance equal to unity, but where the observed data is actually generated from a student-t distribution with five degrees of freedom. In this case, the tails of the student-t ensure that the surprisal does not have an exponential moment, and, as discussed in Theorem 1, the distribution of the maximum surprisal, M_n, will be Frechet.*

To show that this is the case we generate $R = 1000$ independent samples of size $n = 1000$ from a student-t distribution with five degrees of freedom, but where the mean and variance both equal to unity, and for each $r = 1, \ldots, 1000$ we calculate $M_n^{(r)} = \max\{S_1^{(r)}, \ldots, S_n^{(r)}\}$. We plot the histogram of $\{M_n^{(r)}\}_{r=1}^{1000}$ and compare this with the fitted GEV distribution in Fig. 3. The estimated parameters from this experiment yield an estimated shape parameter $\hat{\xi}_R = 0.4557$, with a standard error of 0.0294, suggesting high confidence that the actual shape parameter is positive, and that the distribution of M_n is indeed Frechet.




Figure 3: The estimated GEV parameters and standard errors are given by: location: 24.0107 (0.3564); scale: 9.9537 (0.3329); shape: 0.4557 (0.0293).

The most interesting thing from these examples from a methodological perspective is that if we get the assumed distribution of the data wrong, then this will have meaningful consequence for the behavior of the surprisal we are calculating. In particular, as the student-t example shows, when we model the data using a distribution which is assumed to have thinner tails than those of the actual data, the surprisal can have a very long/thick tail. This immediately suggests that if our goal is to check for anomalies, we should be using distributions that are somewhat robust to outliers, e.g., student-t, etc. I've obtained similar results if you model the data with a Gaussian, but the actual data generation mechanism is exponential. In this case, the max surprisal is Frechet. However, if you had modeled the data with something like a heavy-tailed student-t, then the max surprisal is Gumble. This suggests that the model you specify is absolutely critical to understanding behavior of the max surprisal itself.

References

Coles, S. (2001), *An introduction to statistical modeling of extreme values*, Vol. 208 of *Springer Series in Statistics*, Springer, London UK.

Good, I. J. (1952), 'Rational decisions', *Journal of the Royal Statistical Society. Series B, Statistical methodology* **14**, 107–114.

Hyndman, R. J. (1996), 'Computing and graphing highest density regions', *The American Statistician* **50**(2), 120–126.

Hyndman, R. J., Kandanaarachchi, S. & Turner, K. (2025), When lookout sees crackle: Anomaly detection via kernel density estimation, Technical report, Monash University.

Kandanaarachchi, S. & Hyndman, R. J. (2022), 'Leave-one-out kernel density estimates for outlier detection', *J Computational & Graphical Statistics* **31**, 586–599.

Stone, J. V. (2022), *Information theory: a tutorial introduction*, Sebtel Press, Sheffield, England.

Tribus, M. (1961), *Thermodynamics and Thermostatics: An Introduction to Energy, Information and States of Matter, with Engineering Applications*, D. Van Nostrand, New York, USA.