

Forecast reconciliation

2. Perspectives on forecast reconciliation

Rob J Hyndman



MONASH University

Photo by Edvard Alexander Rølvaag on

robjhyndman.com/fr2023

Outline

- 1 Reconciliation via constraints
- 2 The geometry of forecast reconciliation
- 3 Game theory perspectives
- 4 Adding optimization constraints

Outline


- 1 Reconciliation via constraints
- 2 The geometry of forecast reconciliation
- 3 Game theory perspectives
- 4 Adding optimization constraints

Notation reminder

Every collection of time series with linear constraints can be written as

$$\mathbf{y}_t = \mathbf{S}\mathbf{b}_t$$

- \mathbf{y}_t = vector of all series at time t
- $y_{\text{Total},t}$ = aggregate of all series at time t .
- $y_{X,t}$ = value of series X at time t .
- \mathbf{b}_t = vector of most disaggregated series at time t
- \mathbf{S} = “summing matrix” containing the linear constraints.




$$\mathbf{y}_t = \begin{pmatrix} y_{\text{Total},t} \\ y_{A,t} \\ y_{B,t} \\ y_{C,t} \end{pmatrix} = \underbrace{\begin{pmatrix} 1 & 1 & 1 \\ 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}}_{\mathbf{S}} \underbrace{\begin{pmatrix} y_{A,t} \\ y_{B,t} \\ y_{C,t} \end{pmatrix}}_{\mathbf{b}_t}$$

Notation reminder

Every collection of time series with linear constraints can be written as

$$\mathbf{y}_t = \mathbf{S}\mathbf{b}_t$$

- \mathbf{y}_t = vector of all series at time t
- $y_{\text{Total},t}$ = aggregate of all series at time t .
- $y_{X,t}$ = value of series X at time t .
- \mathbf{b}_t = vector of most disaggregated series at time t
- \mathbf{S} = “summing matrix” containing the linear constraints.



- Base forecasts: $\hat{\mathbf{y}}_{T+h|T}$
- Reconciled forecasts: $\tilde{\mathbf{y}}_{T+h|T} = \mathbf{S}\mathbf{G}\hat{\mathbf{y}}_{T+h|T}$
- MinT:
 $\mathbf{G} = (\mathbf{S}'\mathbf{W}_h^{-1}\mathbf{S})^{-1}\mathbf{S}'\mathbf{W}_h^{-1}$
where \mathbf{W}_h is covariance matrix of base forecast errors.


Notation

Aggregation matrix

$$\mathbf{y}_t = \mathbf{S}\mathbf{b}_t$$

$$\begin{pmatrix} y_{\text{Total},t} \\ y_{A,t} \\ y_{B,t} \\ y_{C,t} \end{pmatrix} = \begin{pmatrix} 1 & 1 & 1 \\ 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} y_{A,t} \\ y_{B,t} \\ y_{C,t} \end{pmatrix}$$

$$\begin{pmatrix} \mathbf{a}_t \\ \mathbf{b}_t \end{pmatrix} = \begin{pmatrix} \mathbf{A} \\ \mathbf{I}_{n_b} \end{pmatrix} \mathbf{b}_t$$



Constraint matrix

$$\mathbf{C}\mathbf{y}_t = \mathbf{0}$$

$$\text{where } \mathbf{C} = \begin{bmatrix} 1 & -1 & -1 & -1 \end{bmatrix} \\ = \begin{bmatrix} \mathbf{I}_{n_a} & -\mathbf{A} \end{bmatrix}$$

Zero-constraint representation

Aggregation matrix A

$$\mathbf{y}_t = \begin{bmatrix} \mathbf{a}_t \\ \mathbf{b}_t \end{bmatrix} = \begin{bmatrix} \mathbf{A} \\ \mathbf{I}_{n_b} \end{bmatrix} \mathbf{b}_t = \mathbf{S} \mathbf{b}_t$$

Zero-constraint representation

Aggregation matrix A

$$\mathbf{y}_t = \begin{bmatrix} \mathbf{a}_t \\ \mathbf{b}_t \end{bmatrix} = \begin{bmatrix} \mathbf{A} \\ \mathbf{I}_{n_b} \end{bmatrix} \mathbf{b}_t = \mathbf{S} \mathbf{b}_t$$

Constraint matrix C

$$\mathbf{C} \mathbf{y}_t = \mathbf{0}$$

- Constraint matrix approach more general & more parsimonious.
- $\mathbf{C} = [\mathbf{I}_{n_a} \quad -\mathbf{A}]$.
- \mathbf{S} , \mathbf{A} and \mathbf{C} may contain any real values (not just 0s and 1s).

Zero-constraint representation

Assuming \mathbf{C} is full rank

$$\begin{aligned} \tilde{\mathbf{y}}_{T+h|T} &= \mathbf{M}\hat{\mathbf{y}}_{T+h|T} \\ \text{where } \mathbf{M} &= \mathbf{I} - \mathbf{W}_h \mathbf{C}' (\mathbf{C} \mathbf{W}_h \mathbf{C}')^{-1} \mathbf{C} \end{aligned}$$

- Originally proved by Byron (1978, 1979) for reconciling data.
- Re-discovered by Wickramasuriya, Athanasopoulos, and Hyndman (2019) for reconciling forecasts.
- $\mathbf{M} = \mathbf{S}\mathbf{G}$ (the MinT solution)
- Leads to more efficient reconciliation than using \mathbf{G} .

Outline

- 1 Reconciliation via constraints
- 2 The geometry of forecast reconciliation
- 3 Game theory perspectives
- 4 Adding optimization constraints

Projections in linear algebra

- A projection is a linear transformation \mathbf{M} such that $\mathbf{M}^2 = \mathbf{M}$.
- i.e., M is idempotent — it leaves its image unchanged.
- \mathbf{M} projects onto \mathfrak{s} if $\mathbf{M}\mathbf{y} = \mathbf{y}$ for all $\mathbf{y} \in \mathfrak{s}$.
- A projection is *orthogonal* if $\mathbf{M}' = \mathbf{M}$.
- If a projection is not orthogonal, it is called *oblique*.
- In regression, OLS is an orthogonal projection onto space spanned by predictors.

The coherent subspace

Coherent subspace


m -dimensional linear subspace $\mathfrak{s} \subset \mathbb{R}^n$ for which linear constraints hold for all $\mathbf{y} \in \mathfrak{s}$.

Hierarchical time series

An n -dimensional multivariate time series such that $\mathbf{y}_t \in \mathfrak{s} \quad \forall t$.

Coherent point forecasts

$\tilde{\mathbf{y}}_{t+h|t}$ is coherent if $\tilde{\mathbf{y}}_{t+h|t} \in \mathfrak{s}$.



$$\mathbf{y}_{Tot} = \mathbf{y}_A + \mathbf{y}_B$$

The coherent subspace

Coherent subspace

m -dimensional linear subspace $\mathfrak{s} \subset \mathbb{R}^n$ for which linear constraints hold for all $\mathbf{y} \in \mathfrak{s}$.

Hierarchical time series


An n -dimensional multivariate time series such that $\mathbf{y}_t \in \mathfrak{s} \quad \forall t$.

Coherent point forecasts

$\tilde{\mathbf{y}}_{t+h|t}$ is *coherent* if $\tilde{\mathbf{y}}_{t+h|t} \in \mathfrak{s}$.

Base forecasts

Let $\hat{\mathbf{y}}_{t+h|t}$ be vector of *incoherent* initial h -step forecasts.



$$\mathbf{y}_{Tot} = \mathbf{y}_A + \mathbf{y}_B$$

The coherent subspace

Coherent subspace

m -dimensional linear subspace $\mathfrak{s} \subset \mathbb{R}^n$ for which linear constraints hold for all $\mathbf{y} \in \mathfrak{s}$.

Hierarchical time series

An n -dimensional multivariate time series such that $\mathbf{y}_t \in \mathfrak{s} \quad \forall t$.

Coherent point forecasts


$\tilde{\mathbf{y}}_{t+h|t}$ is *coherent* if $\tilde{\mathbf{y}}_{t+h|t} \in \mathfrak{s}$.

Base forecasts

Let $\hat{\mathbf{y}}_{t+h|t}$ be vector of *incoherent* initial h -step forecasts.

Reconciled forecasts

Let ψ be a mapping, $\psi : \mathbb{R}^n \rightarrow \mathfrak{s}$.
 $\tilde{\mathbf{y}}_{t+h|t} = \psi(\hat{\mathbf{y}}_{t+h|t})$ “reconciles” $\hat{\mathbf{y}}_{t+h|t}$.



$$\mathbf{y}_{Tot} = \mathbf{y}_A + \mathbf{y}_B$$

Linear reconciliation

If $\psi(\mathbf{u}) = \mathbf{Mu}$ is a linear function, then

$$\tilde{\mathbf{y}}_{t+h|t} = \mathbf{M}\hat{\mathbf{y}}_{t+h|t}$$

OLS:
$$\mathbf{M} = \mathbf{S}(\mathbf{S}'\mathbf{S})^{-1}\mathbf{S}'$$
$$= \mathbf{I} - \mathbf{C}'(\mathbf{C}\mathbf{C}')^{-1}\mathbf{C}$$

MinT:
$$\mathbf{M} = \mathbf{S}(\mathbf{S}'\mathbf{W}_h^{-1}\mathbf{S})^{-1}\mathbf{S}'\mathbf{W}_h^{-1}$$
$$= \mathbf{I} - \mathbf{W}_h\mathbf{C}'(\mathbf{C}\mathbf{W}_h\mathbf{C}')^{-1}\mathbf{C}$$


Linear reconciliation

If $\psi(\mathbf{u}) = \mathbf{Mu}$ is a linear function, then

$$\tilde{\mathbf{y}}_{t+h|t} = \mathbf{M}\hat{\mathbf{y}}_{t+h|t}$$

OLS:
$$\mathbf{M} = \mathbf{S}(\mathbf{S}'\mathbf{S})^{-1}\mathbf{S}'$$
$$= \mathbf{I} - \mathbf{C}'(\mathbf{C}\mathbf{C}')^{-1}\mathbf{C}$$

MinT:
$$\mathbf{M} = \mathbf{S}(\mathbf{S}'\mathbf{W}_h^{-1}\mathbf{S})^{-1}\mathbf{S}'\mathbf{W}_h^{-1}$$
$$= \mathbf{I} - \mathbf{W}_h\mathbf{C}'(\mathbf{C}\mathbf{W}_h\mathbf{C}')^{-1}\mathbf{C}$$




Linear reconciliation

If $\psi(\mathbf{u}) = \mathbf{Mu}$ is a linear function, then

$$\tilde{\mathbf{y}}_{t+h|t} = \mathbf{M}\hat{\mathbf{y}}_{t+h|t}$$

OLS:
$$\mathbf{M} = \mathbf{S}(\mathbf{S}'\mathbf{S})^{-1}\mathbf{S}'$$
$$= \mathbf{I} - \mathbf{C}'(\mathbf{C}\mathbf{C}')^{-1}\mathbf{C}$$

MinT:
$$\mathbf{M} = \mathbf{S}(\mathbf{S}'\mathbf{W}_h^{-1}\mathbf{S})^{-1}\mathbf{S}'\mathbf{W}_h^{-1}$$
$$= \mathbf{I} - \mathbf{W}_h\mathbf{C}'(\mathbf{C}\mathbf{W}_h\mathbf{C}')^{-1}\mathbf{C}$$



Projections

Suppose \mathbf{M} is a projection onto \mathfrak{s} , then

- Coherent base forecasts are unchanged.
- Unbiased base forecasts remain unbiased.

- Orthogonal projections (i.e., OLS) lead to smallest possible adjustments of base forecasts.

Linear reconciliation

If $\psi(\mathbf{u}) = \mathbf{Mu}$ is a linear function, then

$$\tilde{\mathbf{y}}_{t+h|t} = \mathbf{M}\hat{\mathbf{y}}_{t+h|t}$$


$$\begin{aligned}\text{OLS:} \quad \mathbf{M} &= \mathbf{S}(\mathbf{S}'\mathbf{S})^{-1}\mathbf{S}' \\ &= \mathbf{I} - \mathbf{C}'(\mathbf{C}\mathbf{C}')^{-1}\mathbf{C}\end{aligned}$$

$$\begin{aligned}\text{MinT:} \quad \mathbf{M} &= \mathbf{S}(\mathbf{S}'\mathbf{W}_h^{-1}\mathbf{S})^{-1}\mathbf{S}'\mathbf{W}_h^{-1} \\ &= \mathbf{I} - \mathbf{W}_h\mathbf{C}'(\mathbf{C}\mathbf{W}_h\mathbf{C}')^{-1}\mathbf{C}\end{aligned}$$

Distance reducing property

If \mathbf{M} is an orthogonal projection onto \mathcal{S} :

$$\|\mathbf{y}_{t+h} - \tilde{\mathbf{y}}_{t+h|t}\| \leq \|\mathbf{y}_{t+h} - \hat{\mathbf{y}}_{t+h|t}\|$$



- Distance reduction holds for any realisation and any forecast.
- Other measures of forecast accuracy may be worse.
- Not necessarily the optimal₁₁ reconciliation.

Linear projections

$$\tilde{\mathbf{y}}_{T+h|T} = \mathbf{M}\hat{\mathbf{y}}_{T+h|T}$$

Variance

$$\mathbf{V}_h = \text{Var}[\mathbf{y}_{T+h} - \tilde{\mathbf{y}}_{T+h|T} \mid \mathbf{y}_1, \dots, \mathbf{y}_n] = \mathbf{M}\mathbf{W}_h\mathbf{M}'$$

where $\mathbf{W}_h = \text{Var}[\mathbf{y}_{T+h} - \hat{\mathbf{y}}_{T+h|T} \mid \mathbf{y}_1, \dots, \mathbf{y}_n]$.

Minimum trace (MinT) reconciliation

If \mathbf{M} is a projection, then the trace of \mathbf{V}_h is minimized when

$$\mathbf{M} = \mathbf{S}(\mathbf{S}'\mathbf{W}_h^{-1}\mathbf{S})^{-1}\mathbf{S}'\mathbf{W}_h^{-1} = \mathbf{I} - \mathbf{W}_h\mathbf{C}'(\mathbf{C}\mathbf{W}_h\mathbf{C}')^{-1}\mathbf{C}$$

Linear projections

$$\tilde{\mathbf{y}}_{T+h|T} = \mathbf{M}\hat{\mathbf{y}}_{T+h|T}$$

Variance


$\mathbf{V}_h = \text{Var}[\mathbf{y}_{T+h} - \tilde{\mathbf{y}}_{T+h|T} \mid \mathbf{y}_1, \dots, \mathbf{y}_n] = \mathbf{M}\mathbf{W}_h\mathbf{M}'$
where $\mathbf{W}_h = \text{Var}[\mathbf{y}_{T+h} - \hat{\mathbf{y}}_{T+h|T} \mid \mathbf{y}_1, \dots, \mathbf{y}_n]$.

Minimum trace (MinT) reconciliation

If \mathbf{M} is a projection, then the trace of \mathbf{V}_h is minimized when

$$\mathbf{M} = \mathbf{S}(\mathbf{S}'\mathbf{W}_h^{-1}\mathbf{S})^{-1}\mathbf{S}'\mathbf{W}_h^{-1} = \mathbf{I} - \mathbf{W}_h\mathbf{C}'(\mathbf{C}\mathbf{W}_h\mathbf{C}')^{-1}\mathbf{C}$$

- R is the most likely direction of deviations from \mathfrak{s} .
- Orange: in-sample errors



Linear projections

$$\tilde{\mathbf{y}}_{T+h|T} = \mathbf{M}\hat{\mathbf{y}}_{T+h|T}$$

Variance

$$\mathbf{V}_h = \text{Var}[\mathbf{y}_{T+h} - \tilde{\mathbf{y}}_{T+h|T} \mid \mathbf{y}_1, \dots, \mathbf{y}_n] = \mathbf{M}\mathbf{W}_h\mathbf{M}'$$


where $\mathbf{W}_h = \text{Var}[\mathbf{y}_{T+h} - \hat{\mathbf{y}}_{T+h|T} \mid \mathbf{y}_1, \dots, \mathbf{y}_n]$.

Minimum trace (MinT) reconciliation

If \mathbf{M} is a projection, then the trace of \mathbf{V}_h is minimized when

$$\mathbf{M} = \mathbf{S}(\mathbf{S}'\mathbf{W}_h^{-1}\mathbf{S})^{-1}\mathbf{S}'\mathbf{W}_h^{-1} = \mathbf{I} - \mathbf{W}_h\mathbf{C}'(\mathbf{C}\mathbf{W}_h\mathbf{C}')^{-1}\mathbf{C}$$

- R is the most likely direction of deviations from \mathfrak{s} .
- Grey: potential base forecasts



Linear projections

$$\tilde{\mathbf{y}}_{T+h|T} = \mathbf{M}\hat{\mathbf{y}}_{T+h|T}$$

Variance


$\mathbf{V}_h = \text{Var}[\mathbf{y}_{T+h} - \tilde{\mathbf{y}}_{T+h|T} \mid \mathbf{y}_1, \dots, \mathbf{y}_n] = \mathbf{M}\mathbf{W}_h\mathbf{M}'$
where $\mathbf{W}_h = \text{Var}[\mathbf{y}_{T+h} - \hat{\mathbf{y}}_{T+h|T} \mid \mathbf{y}_1, \dots, \mathbf{y}_n]$.

Minimum trace (MinT) reconciliation

If \mathbf{M} is a projection, then the trace of \mathbf{V}_h is minimized when

$$\mathbf{M} = \mathbf{S}(\mathbf{S}'\mathbf{W}_h^{-1}\mathbf{S})^{-1}\mathbf{S}'\mathbf{W}_h^{-1} = \mathbf{I} - \mathbf{W}_h\mathbf{C}'(\mathbf{C}\mathbf{W}_h\mathbf{C}')^{-1}\mathbf{C}$$

- R is the most likely direction of deviations from \mathfrak{s} .
- Grey: potential base forecasts
- Red: reconciled forecasts



Linear projections

$$\tilde{\mathbf{y}}_{T+h|T} = \mathbf{M}\hat{\mathbf{y}}_{T+h|T}$$

Variance


$\mathbf{V}_h = \text{Var}[\mathbf{y}_{T+h} - \tilde{\mathbf{y}}_{T+h|T} \mid \mathbf{y}_1, \dots, \mathbf{y}_n] = \mathbf{M}\mathbf{W}_h\mathbf{M}'$
where $\mathbf{W}_h = \text{Var}[\mathbf{y}_{T+h} - \hat{\mathbf{y}}_{T+h|T} \mid \mathbf{y}_1, \dots, \mathbf{y}_n]$.

Minimum trace (MinT) reconciliation

If \mathbf{M} is a projection, then the trace of \mathbf{V}_h is minimized when

$$\mathbf{M} = \mathbf{S}(\mathbf{S}'\mathbf{W}_h^{-1}\mathbf{S})^{-1}\mathbf{S}'\mathbf{W}_h^{-1} = \mathbf{I} - \mathbf{W}_h\mathbf{C}'(\mathbf{C}\mathbf{W}_h\mathbf{C}')^{-1}\mathbf{C}$$

- R is the most likely direction of deviations from \mathfrak{s} .
- Grey: potential base forecasts
- Red: reconciled forecasts



Outline

- 1 Reconciliation via constraints
- 2 The geometry of forecast reconciliation
- 3 Game theory perspectives
- 4 Adding optimization constraints

Game theory perspectives

Find the solution to the minimax problem

$$V = \min_{\tilde{\mathbf{y}} \in \mathfrak{s}} \max_{\mathbf{y} \in \mathfrak{s}} \{ \ell(\mathbf{y}, \tilde{\mathbf{y}}) - \ell(\mathbf{y}, \hat{\mathbf{y}}) \},$$

where ℓ is a loss function, and \mathfrak{s} is the coherent subspace.

- $V \leq 0$: reconciliation guaranteed to reduce loss.
- If $\ell(\mathbf{y}, \tilde{\mathbf{y}}) = (\mathbf{y} - \tilde{\mathbf{y}})' \Psi (\mathbf{y} - \tilde{\mathbf{y}})$, where Ψ is any symmetric pd matrix, then:
 - 1 $\tilde{\mathbf{y}} = \mathbf{S}(\mathbf{S}'\Psi\mathbf{S})^{-1}\mathbf{S}'\Psi\hat{\mathbf{y}}$ will always improve upon the base forecasts;
 - 2 The MinT solution $\tilde{\mathbf{y}} = \mathbf{S}(\mathbf{S}'\mathbf{W}_h^{-1}\mathbf{S})^{-1}\mathbf{S}'\mathbf{W}_h^{-1}\hat{\mathbf{y}}$ will optimise loss in expectation over any choice of Ψ .

Biased reconciliation

Regularized empirical risk minimization problem:

$$\min_{\mathbf{G}} \frac{1}{Nn} \|\mathbf{Y} - \hat{\mathbf{Y}}\mathbf{G}'\mathbf{S}'\|_F + \lambda \|\text{vec}\mathbf{G}\|_1,$$

- $N = T - T_1 - h + 1$, T_1 is minimum training sample size
- $\|\cdot\|_F$ is the Frobenius norm
- $\mathbf{Y} = [\mathbf{y}_{T_1+h}, \dots, \mathbf{y}_T]'$
- $\hat{\mathbf{Y}} = [\hat{\mathbf{y}}_{T_1+h|T_1}, \dots, \hat{\mathbf{y}}_{T|T-h}]'$
- λ is a regularization parameter.

When $\lambda = 0$, $\hat{\mathbf{G}} = \mathbf{B}'\hat{\mathbf{Y}}(\hat{\mathbf{Y}}'\hat{\mathbf{Y}})^{-1}$ where $\mathbf{B} = [\mathbf{b}_{T_1+h}, \dots, \mathbf{b}_T]'$.

Outline

- 1 Reconciliation via constraints
- 2 The geometry of forecast reconciliation
- 3 Game theory perspectives
- 4 Adding optimization constraints

Adding optimization constraints





Any approach to reconciliation based on optimisation uses a form of constrained optimisation since reconciled forecasts must lie on the coherent subspace. However, at times additional constraints may be implemented. The first is the case where reconciled forecasts must be non-negative. In general, even if base forecasts are constrained to be positive (which can be achieved by modelling on the log scale and back-transforming), there is no guarantee that the usual reconciliation approaches such as OLS and MinT will maintain the non-negativity of forecasts. To address this issue, the usual optimisation problem can be augmented with

ML and regularization




Bayesian versions

In-built coherence

References

-  Ben Taieb, S and B Koo (2019). Regularized regression for hierarchical forecasting without unbiasedness conditions. In: *Proceedings of the 25th ACM SIGKDD International Conference on Knowledge Discovery & Data Mining*. KDD '19. Anchorage, AK, USA: Association for Computing Machinery, pp.1337–1347.
-  Di Fonzo, T and D Girolimetto (2022). Forecast combination-based forecast reconciliation: Insights and extensions. *International Journal of Forecasting* **forthcoming**.
-  Eckert, F, RJ Hyndman, and A Panagiotelis (2021). Forecasting Swiss exports using Bayesian forecast reconciliation. *European J Operational Research* **291**(2), 693–710.
-  Panagiotelis, A, P Gamakumara, G Athanasopoulos, and RJ Hyndman (2021). Forecast reconciliation: A geometric view with new insights on bias correction. *International J Forecasting* **37**(1), 343–359.

References

-  van Erven, T and J Cugliari (2015). “Game-theoretically optimal reconciliation of contemporaneous hierarchical time series forecasts”. In: *Modeling and Stochastic Learning for Forecasting in High Dimension*. Ed. by A Antoniadis, JM Poggi, and X Brossat. Cham: Springer International Publishing, pp.297–317.
-  Wickramasuriya, SL, G Athanasopoulos, and RJ Hyndman (2019). Optimal forecast reconciliation for hierarchical and grouped time series through trace minimization. *J American Statistical Association* **114**(526), 804–819.
-  Wickramasuriya, SL, BA Turlach, and RJ Hyndman (2020). Optimal non-negative forecast reconciliation. *Statistics & Computing* **30**(5), 1167–1182.