

Two Displays, One Data Set

AP Statistics · Unit 1 · Topic 1.8 · B: 2026-09-22 · E: 2026-09-23 · TEACHER KEY

CED TETHER

- Enduring Understanding: UNC-1 Graphical representations and statistics allow us to identify and represent key features of data.
- Skills:
- **Skill 2.B** — Construct numerical or graphical representations of distributions.
- **Skill 2.A** — Describe data presented numerically or graphically.
- Learning Objectives and Essential Knowledge:
- UNC-1.L Represent summary statistics for quantitative data graphically. [Skill 2.B]
- **UNC-1.L.1** — Taken together, the minimum data value, the first quartile (Q1), the median, the third quartile (Q3), and the maximum data value make up the five-number summary.
- **UNC-1.L.2** — A boxplot is a graphical representation of the five-number summary (minimum, first quartile, median, third quartile, maximum). The box represents the middle 50% of data, with a line at the median and the ends of the box corresponding to the quartiles. Lines ("whiskers") extend from the quartiles to the most extreme point that is not an outlier, and outliers are indicated by their own symbol beyond this.
- UNC-1.M Describe summary statistics of quantitative data represented graphically. [Skill 2.A]
- **UNC-1.M.1** — Summary statistics of quantitative data, or of sets of quantitative data, can be used to justify claims about the data in context.
- **UNC-1.M.2** — If a distribution is relatively symmetric, then the mean and median are relatively close to one another. If a distribution is skewed right, then the mean is usually to the right of the median. If the distribution is skewed left, then the mean is usually to the left of the median.

Essential question: What information survives when individual observations are compressed into a box-plot?

DOK-3 skill: 4.B · **FRQ pattern:** what-a-boxplot-can-and-cannot-show-adjudicate

DOK rationale: Part (c) requires adjudicating two interpretations by coordinating the boxplot with the dotplot, distinguishing summaries from distribution features, and defending which display answers a decision question; no single calculation determines the response.

First take → rules + (a)–(c) → turn in

DOK: 1 → 3

Minutes: 45

Teacher does

- Board slide up at the bell. Read Omar's and Nia's claims aloud without confirming either.
- At minute 5, read the rules box; have students start (a).
- Board slide back up at minute 33. Circulate; ask students to point before they name a feature.
- Collect at the door. Score part (c) only, E/P/I.

Students do

- Commit to Omar or Nia from the two displays before any discussion.
- Use the rules box to read the quartiles and describe the dotplot in (a) and (b).
- Finish (a)–(c); complete the exit line; turn in.

Questions to ask

- Point to 32 minutes on the dotplot. How many dots are nearby?
- Could a different data set have this same five-number summary but no gap?
- Which features disappeared when the 24 dots became five numbers?
- For the coach's decision, do you need a center or the number of students above 30?

Adult role

Solo teacher. Circulate 5 pairs. Do not say 'bimodal' or confirm Nia; ask students to locate the median on both displays and name what disappears.

[WARN] LOOK FOR

- Treating the median as a mode or as a value an observation must equal.
- Claiming the boxplot displays the two clusters because its box is wide.
- Calling 55 an outlier without checking the 1.5 IQR fence.

SCORING PART (c) — E / P / I

Expected elements

- **nia-and-gap** (required) — Chooses Nia using the empty interval around 32
- **boxplot-limit** (required) — Explains that quartiles do not reveal the gap or concentration of individual observations
- **coach-decision** (required) — Chooses dotplot and uses 12 of 24 to reject more-than-half over 30

E	All three elements are correct and linked to this problem. Accept equivalent plain-language reasoning.
P	At least one element includes a correct evidence-to-reason link, but another is missing or incorrect.
I	No correct evidence-to-reason link; only a choice, copied numbers, or unsupported statements.

Common mistakes


- Treating the median as the most common value; no student took 32 minutes
- Calling the empty 27–37 interval a short bar even though the display is a dotplot
- Saying a boxplot shows every observation because it shows the minimum and maximum
- Calling 55 a formal 1.5 IQR outlier; the upper fence is 65.25
- Choosing the boxplot for the coach merely because it is more compact

[STAR] TODAY'S PROBLEM — annotated

The displays show practice-test times, in minutes, for 24 students. Two students interpret the boxplot.

Omar: “Most students took about 32 minutes—the middle of the box.”

Nia: “A boxplot cannot tell us whether most students took about 32 minutes.”



FIRST TAKE (5 min)

Who do you trust more, Omar or Nia? Point to one feature in either display.

Key: Not graded. Expect Omar; the empty interval around 32 is the reveal.

Rules callout “DOTPLOTS AND BOXPLOTS (keep on the page)” is printed on the student sheet.

DOK 1 (a) Read the median and the two quartiles from the boxplot.

Key: Median = 32, $Q_1 = 24$, $Q_3 = 40.5$ minutes.

DOK 2 (b) In two sentences, describe the dotplot’s groups and empty interval. Count how many students took more than 30 minutes.

Key: There are 12 times from 22–26 minutes and 11 from 38–44, plus one at 55. No values lie between 26 and 38; exactly 12 of 24 students took more than 30 minutes.

DOK 3 (c) Whose interpretation is better supported? Use a dotplot feature to explain what the boxplot hides. Choose a display for deciding whether more than half the students need over 30 minutes, and defend your choice.

Key: Nia: no observations are near 32, which lies between the two groups. The boxplot preserves quartiles but hides that gap; treating its median as where most observations gather is a misreading. Choose the dotplot for the coach: 12 of 24, exactly half, took over 30 minutes, so not more than half.

EXIT

One thing the rules box changed about my first take: _____