

NAME: _____

PERIOD: _____

DATE: _____

Two Production Lines, Two Sampling Plans

AP Statistics · Unit 4 · Topic 4.1 · B: 2027-01-25 · E: 2027-01-27

[STAR] TODAY'S PROBLEM

Suppose Lines A and B each contain 1,000 cycle times with population mean 40 seconds and standard deviation 12 seconds. Line A is approximately normal and an SRS of $n_A = 9$ is selected; Line B is strongly right-skewed and an SRS of $n_B = 36$ is selected. An auditor wants $P(\bar{X} > 46)$ for each.

Serena: “Only B can use a normal model because $36 \geq 30$; its probability is 0.00135. A is too small.”

Mateo: “Only A can use a normal model because its population is normal; its probability is 0.0668. B is too skewed.”

Sampling plans

Line	Shape	N	n	μ	σ
A	Normal	1000	9	40	12
B	Right-skewed	1000	36	40	12

FIRST TAKE — before we discuss (5 min)

Which plan seems more likely to give averages close to 40 seconds? Give one reason from the sample sizes or shapes.

[BOOK] MODEL SAMPLE MEANS, NOT INDIVIDUAL VALUES (keep on the page)

For an SRS of size n , $\mu_{\bar{X}} = \mu$ and $\sigma_{\bar{X}} = \sigma/\sqrt{n}$ when $n \leq 0.10N$ for sampling without replacement. The sampling distribution of $\bar{X}$ is approximately normal if the population is approximately normal (for any n) **or** if the population is not normal but $n \geq 30$ (Central Limit Theorem). A normal tail probability describes the mean of an entire random sample, not one individual value. For a normal probability, standardize with $z = (\text{value} - \text{mean})/\text{SD}$ and use the appropriate normal tail.

Work (a)–(c) from the rules box:

DOK 1 (a) State the mean of each sampling distribution and the 10% condition for using $\sigma_{\bar{X}} = \sigma/\sqrt{n}$.

DOK 2 (b) Verify the 10 percent condition, compute both sampling-distribution standard deviations, and calculate each normal tail probability when justified.

DOK 3 (c) In 3–4 sentences, explain what each student gets right and wrong. Use the two normality pathways to decide which probabilities are supported. Connect the different tail probabilities to the sampling SDs in (b).

Frame: For Line A, a normal model is _____ because _____; for Line B, it is _____ because _____.

Frame: The tail probabilities differ because 46 is _____ sampling SDs above 40 for Line A but _____ for Line B.

EXIT — turn this in

One thing the rules box changed about my first take: _____