

Two Analysts, One Planned Test

AP Statistics · Unit 3 · Topic 3.12 · B: 2026-12-15 · E: 2027-01-27 · TEACHER KEY

CED TETHER

- **Enduring Understanding VAR-6** — The normal distribution may be used to model variation.
- **Skill 1.F** — Identify null and alternative hypotheses.
- **Skill 1.E** — Identify an appropriate inference method for significance tests.
- **Skill 4.C** — Verify that inference procedures apply in a given situation.
- **LO VAR-6.H** — Identify the null and alternative hypotheses for a difference of two population proportions. [Skill 1.F]
- **EK VAR-6.H.1** — For a two-sample test for a difference of two proportions, the null hypothesis specifies a value of 0 for the difference in population proportions, indicating no difference or effect.
- **EK VAR-6.H.2** — The null hypothesis for a difference in proportions is: $H_0: p_1 = p_2$, or $H_0: p_1 - p_2 = 0$.
- **EK VAR-6.H.3** — A one-sided alternative hypothesis for a difference in proportions is $H_a: p_1 < p_2$ or $H_a: p_1 > p_2$. A two-sided alternative hypothesis is $H_a: p_1 \neq p_2$.
- **LO VAR-6.I** — Identify an appropriate testing method for the difference of two population proportions. [Skill 1.E]
- **EK VAR-6.I.1** — For a single categorical variable, the appropriate testing method for the difference of two population proportions is a two-sample z-test for a difference between two population proportions.
- **LO VAR-6.J** — Verify the conditions for making statistical inferences when testing a difference of two population proportions. [Skill 4.C]
- **EK VAR-6.J.1** — To make statistical inferences when testing a difference between population proportions, check for independence and approximately normal sampling distribution:
 - Independence: data collected using two independent, random samples or a randomized experiment; when sampling without replacement, check $n_1 \leq 10\%N_1$ and $n_2 \leq 10\%N_2$.
 - Shape (approximate normality): compute the combined (pooled) proportion $\hat{p}_c = (n_1\hat{p}_1 + n_2\hat{p}_2) / (n_1 + n_2)$; assuming H_0 is true, check that $n_1\hat{p}_c$, $n_1(1 - \hat{p}_c)$, $n_2\hat{p}_c$, and $n_2(1 - \hat{p}_c)$ are all ≥ 5 (or 10, per the predetermined value used).

Essential question: How do the original research question and study design determine a defensible two-proportion test setup?

DOK-3 skill: 4.C · **FRQ pattern:** predata-question-vs-postdata-tail-switch

DOK rationale: Part (c) uses the original research question and the timing of the tail choice to adjudicate competing test setups and explain how choosing a direction after observing results overstates evidence.

Do Now (first take) → Explore (video + follow-along) → Exit (finish + turn in) DOK: 1
→ 3 **Minutes: 45**

Teacher does

- Board slide up at the bell. Ask which setup matches the team's question without confirming the tail.
- Begin the 6.10 video follow-along at minute 5.
- Return to the board slide at minute 33. Circulate and make students separate the question chosen before data from the sign observed afterward.

- Collect at the door. Score part (c) only, E/P/I.

Students do

- Choose Nia's or Omar's setup and cite wording or design evidence.
- Complete the follow-along about hypotheses, procedure selection, pooling, and conditions.
- Finish (a)–(c), revise the first take in the exit line, and turn in the sheet.

Questions to ask

- What result would matter if the meter actually lowered completion?
- When was the direction of Omar's alternative chosen?
- Why are pooled expected counts appropriate under equality?
- What do random selection and random assignment each allow the team to claim?

Adult role

Solo teacher. Circulate 5 pairs. Require a setup, pooled-count check, and consequence of changing tails after seeing data.

[WARN] WATCH FOR

- Writing hypotheses with sample proportions instead of population parameters.
- Choosing a greater-than alternative only because the observed meter proportion is larger.
- Checking observed rather than pooled expected counts for the test.

SCORING PART (c) — E / P / I

Expected elements

- **judgment-1** (required) — Chooses Nia and states equality null and two-sided alternative with population parameters.
- **judgment-2** (required) — Ties either direction to the pre-data word change rather than the observed positive gap.
- **judgment-3** (required) — Explains that selecting a tail after observing the result changes the question and overstates evidence.

E	Meets all three checks with correct contextual reasoning; equivalent wording is acceptable.
P	Shows at least one substantive correct check, but has an omission or error that prevents E.
I	Shows none of the three checks with substantive correct reasoning; a choice alone is insufficient.

[STAR] TODAY'S PROBLEM — annotated

Suppose an online retailer has 50,000 active accounts. Before collecting data, its team asks, “Does a progress meter change the proportion of active accounts completing checkout?” It randomly selects 360 accounts and randomly assigns 180 to each screen. The table shows the outcomes. Let p_M and p_S be the corresponding population completion proportions.

Nia: “Use $H_0 : p_M - p_S = 0$ versus $H_a : p_M - p_S \neq 0$ because either direction is a change.”

Omar: “Because $\hat{p}_M$ is larger, use $H_a : p_M - p_S > 0$.”

Checkout outcomes

Screen	n	Complete	Other
Meter	180	126	54
Standard	180	108	72

FIRST TAKE (5 min)

When the team asks whether checkout changes, should a decrease count too? Use the original question to explain.

Key: Not graded. The positive observed difference is intentionally tempting; the research question must control the tail.

Rules callout “SET THE QUESTION, PARAMETERS, AND CONDITIONS (keep on the page)” is printed on the student sheet.

DOK 1 (a) Define p_M and p_S in context. Calculate $\hat{p}_M$, $\hat{p}_S$, and the observed difference $\hat{p}_M - \hat{p}_S$.

Key: Here p_M and p_S are the completion proportions for all active accounts under the meter and standard screens. Thus $\hat{p}_M = 126/180 = 0.70$, $\hat{p}_S = 108/180 = 0.60$, and the difference is 0.10.

DOK 2 (b) Name the test procedure. Calculate the pooled proportion and verify randomization, the 10 percent condition, and all four pooled expected counts.

Key: The appropriate procedure is a two-sample z -test for a difference of proportions. The 360 accounts were randomly selected and assigned; $360 \leq 0.10(50,000) = 5,000$. Under H_0 , $\hat{p}_c = 234/360 = 0.65$, so each group has pooled expected counts 117 and 63, all at least 10.

DOK 3 (c) In 3–4 sentences, choose and state the hypotheses. Use the original word change and when the direction was chosen. Explain how Omar’s choice after seeing the result changes the question and could overstate the evidence.

Key: Nia is defensible: $H_0 : p_M - p_S = 0$ versus $H_a : p_M - p_S \neq 0$. The question asked about change before data were collected, so either an increase or a decrease counts. The observed difference 0.10 cannot decide the research direction afterward. Omar’s right-tail switch omits decreases and makes the observed increase look stronger than it is for the original question.

EXIT

One thing the video changed about my first take: _____