

NAME: _____

PERIOD: _____

DATE: _____

Two Production Lines, Two Sampling Plans

AP Statistics · Topic 5.7 (CED 4.1) · B: 2027-01-11 · E: 2027-02-24

[STAR] TODAY'S PROBLEM

Suppose Lines A and B each contain 1,000 cycle times with population mean 40 seconds and standard deviation 12 seconds. Line A is approximately normal and is sampled with $n_A = 9$; Line B is strongly right-skewed and is sampled with $n_B = 36$. An auditor wants $P(\bar{X} > 46)$ for each.

Serena: “Only B can use a normal model because $36 \geq 30$; its probability is 0.00135. A is too small.”

Mateo: “Only A can use a normal model because its population is normal; its probability is 0.0668. B is too skewed.”

Sampling plans

Line	Shape	N	n	μ	σ
A	Normal	1000	9	40	12
B	Right-skewed	1000	36	40	12

FIRST TAKE — before the video (5 min)

Which parts of Serena's and Mateo's reasoning look defensible? Cite one shape or sample-size detail.

[BOOK] MODEL SAMPLE MEANS, NOT INDIVIDUAL VALUES (keep on the page)

For an SRS of size n , $\mu_{\bar{X}} = \mu$ and $\sigma_{\bar{X}} = \sigma/\sqrt{n}$ when $n \leq 0.10N$ for sampling without replacement. The sampling distribution of $\bar{X}$ is approximately normal if the population is approximately normal (for any n) **or** if the population is not normal but $n \geq 30$ (Central Limit Theorem). A normal tail probability describes the mean of an entire random sample, not one individual value.

Finish after the video:

DOK 1 (a) State the mean of each sampling distribution and the finite-population condition for using $\sigma_{\bar{X}} = \sigma/\sqrt{n}$.

DOK 2 (b) Verify the 10 percent condition, compute both sampling-distribution standard deviations, and calculate each normal tail probability when justified.

DOK 3 (c) Evaluate both claims using population shape, sample size, the two normality pathways, and sampling variability. Determine which probabilities are warranted, explain any difference between them, and trace what each incomplete rule would cause the auditor to discard.

Frame: For Line A, a normal model is _____ because _____; for Line B, it is _____ because _____.

Frame: The tail probabilities differ because 46 is _____ sampling SDs above 40 for Line A but _____ for Line B.

EXIT — turn this in

One thing the video changed about my first take: _____