


# 4 Continuous Random Variables

Together, discrete and continuous approaches form a powerful framework for modeling the world.

## 4.1 Probability density function

Continuous RVs!

An RV has a *continuous distribution* if its CDF is differentiable. Endpoints of CDF may be continuous but not differentiable. A continuous RV is a RV with a continuous distribution.



For a continuous RV  $X$  with CDF  $F$ , the PDF of  $X$  is derivative  $f$  of the CDF:  $f(x) = F'(x)$

The support of  $X$ : all  $x$  where  $f(x) > 0$ .

The PDF is kinda similar to PMF, but for PDF quantity of  $f(x)$  is **not a probability**. To obtain the probability, we need to **integrate** PDF.

We can be carefree about including or excluding endpoints as above for continuous RVs, but we must not be careless about this for discrete RVs.

Valid PDF of a continuous RV:

1. Nonnegative:  $f(x) \geq 0$
2. Integrates to 1:  $\int_{-\infty}^{\infty} f(x)dx = 1$

Example: logistic distribution.

$X \sim \text{Logistic}$ .

CDF:

$$F(x) = \frac{e^x}{1 + e^x}, x \in \mathbb{R}$$


PDF:

$$f(x) = \frac{e^x}{(1 + e^x)^2}, x \in \mathbb{R}$$

To find  $P(-2 < X < 2)$ , we need to integrate PDF from  $-2$  to  $2$ :

$$P(-2 < X < 2) = \int_{-2}^2 \frac{e^x}{(1 + e^x)^2} = F(2) - F(-2) \approx 0.76$$

Or  $P(-2 < X < 2)$  is indicated by the shaded area under the PDF and the height of the curly brace on the CDF.



## 4.2 Uniform distribution

A continuous RV  $U$  has the *Uniform distribution*  $X \sim \text{Unif}(a, b)$  on the interval  $(a, b)$  if its PDF is:

$$f(x) = \frac{1}{b-a} \quad \forall a < x < b,$$
$$f(x) = 0 \text{ otherwise}$$

The CDF is the accumulated area under the PDF:

$$F(x) = 0 \quad \forall x \leq a,$$
$$F(x) = \frac{x-a}{b-a} \quad a < x < b,$$
$$F(x) = 1 \quad \forall x \geq b.$$

$\text{Unif}(0, 1)$  is the standard Uniform.

For Uniform distributions, *probability is proportional to length*.

Location-scale transformation.

The RV  $Y$  has been obtained as a *location-scale transformation* of  $X$  if  $Y = \sigma X + \mu$ .  $\mu$  controls the location and  $\sigma$  controls the scale.

**Warning:** if  $Y$  is a linear function of  $X$ , the Uniformity is preserved, but if  $Y$  is defined as a *nonlinear* transformation of  $X$ ,  $Y$  will not be Uniform.

**Warning:** When using location-scale transformations, the shifting and scaling should be applied to the *random variables* themselves, not to their PDFs.

## 4.3 Universality of the Uniform distribution

Given a  $\text{Unif}(0, 1)$  RV, we can construct an RV with *any continuous distribution we want*.

Other names of the universality of Uniform:

- probability integral transform,

- inverse transform sampling,
- the quantile transformation,
- the fundamental theorem of simulation.

### Theorem:

$F$  is a CDF which is continuous function and strictly increasing on the support of distribution. This ensures that the inverse function  $F^{-1}$  exists as function  $(0, 1) \rightarrow \mathbb{R}$ . Results:

1. Let  $U \sim \text{Unif}(0, 1)$  and  $X = F^{-1}(U)$ . Then  $X$  is an RV with CDF  $F$ .
2. Let  $X$  be an RV with CDF  $F$ . Then  $F(X) \sim \text{Unif}(0, 1)$ .

What this theorem is saying about?

First part: Since  $F^{-1}$  is a function (**quantile function**),  $U$  is a RV, and a function of RV is RV,  $F^{-1}(U)$  is a RV; universality of the Uniform says its CDF is  $F$ .

Second part: reverse direction! Starting from RV  $X$  whose CDF is  $F$  and then creating RV  $\text{Unif}(0, 1)$ . Universality of the Uniform says that the distribution of  $F(X)$  is Uniform on  $(0, 1)$ .

Warning: potential notational collusion!

$F(x) = P(X \leq x)$  by definition, but  $F(X) = P(X \leq X) = 1$  is incorrect by definition. Rather, we should first find an expression for the CDF as a function of  $x$ , then replace  $x$  with  $X$  to obtain a random variable. For example, if the CDF of  $X$  is  $F(x) = 1 - e^{-x}$  for  $x > 0$ , then  $F(X) = 1 - e^{-X}$ .

## 4.4 Normal distribution