# AMORTIZED EXPLOITABILITY CERTIFICATES FOR LEARNED RESPONSE EQUILIBRIA

**Lekan Molu, Amazon IRG, Bellevue, WA 98004**
ogunmolu@amazon.com

## ABSTRACT

A fixed point of a learned response map shows that the model reproduces its input; it does not show that any player has low regret in the underlying game. We give a post-hoc exploitability certificate for arbitrary profiles in games with compact convex product action sets and differentiable costs that are convex in each player's own action. The certificate combines the learned map's fixed-point residual with regional upper bounds on the true gradients and on their discrepancy from the learned response operator, without requiring contraction, monotonicity, or equilibrium uniqueness. A finite cost-query construction is possible only when valid regularity bounds and an independent mean-zero sub-Gaussian noise model are supplied externally: finite cost values alone cannot establish those assumptions. Given them, finite differences on an $\eta$-cover produce a simultaneous $1-\alpha$ envelope for every profile in the region. The covering cost grows exponentially with the region's intrinsic dimension at fixed relative resolution, limiting uniform audits to structured regions with small covering numbers and substantial reuse. The certificate was valid in all 280 tested game–model–seed configurations, although its standard model-free entry was tighter in every pointwise configuration. Regional auditing instead trades tightness and upfront queries for later model-only evaluation. Contraction certification was neither necessary nor sufficient for low exploitability in the tested models, and a nonconvex counterexample shows that playerwise convexity is indispensable.

## 1 INTRODUCTION

Decision-making in multi-agent settings can be characterized as *non-cooperative games*, where heterogeneous and selfish entities, i.e. *agents or players*, act with independent decisions to optimize their own objective, irrespective of other players' goals. With applications that traverse socio-technical systems (Ekpo et al., 2026), regional science and transportation planning (Facchinei & Pang, 2003), multi-agent/robot path-finding (Zhang et al., 2024; Wang et al., 2024; Ren et al., 2021), and multi-agent reinforcement learning (Zhang et al., 2021), these games are often encumbered with incomplete or conflicting information and multiple conflicting goals. The (generalized) Nash equilibrium (Facchinei & Kanzow, 2010; Pavel & Hong, 2022; Li et al., 2022), is the solution concept used to characterize the strategic interactions and possible outcomes of such games. Modeling the players' interactions and analyzing the resulting equilibria beyond a small number of players proves intractable: the normal-form representation grows exponentially in the number of players, and even computing an approximate equilibrium is PPAD-hard (Rubinstein, 2015). Beyond tractability, for systems where cost functions are unavailable or players are boundedly rational, the Nash solution concept is practically ineffective for characterizing best responses.

This has stimulated the computation of equilibria from players' interaction data rather using structured learning principles (Scarabaggio et al., 2024). These data-driven formulations extend fixed-point theorems that analyze solution existence under convexity (Kakutani, 1941) to response analysis from data. Relaxations such as the first-order Nash equilibrium notion of Scarabaggio et al. (2026) retain the players' objectives while weakening the equilibrium concept, so that each player's action is a fixed point of its own projected gradient, with existence following from algebraic topology rather than from convex analysis. Scarabaggio et al. (2024) replace the objectives, characterizing best responses as a composition of differentiable neural network layers fitted to players' observed actions. Feasibility is

enforced by a projection of the network output onto each player's strategy set. Related formulations learn a parametric equilibrium map trained against the Nikaido–Isoda gap instead of regressing on observed play (Bemporad & Tatarenko, 2026a), or take the reacting object to be the data distribution rather than the response (Narang et al., 2023). This paper works in the second setting and certifies against the first: we measure the residual of the learned response map, which costs one forward pass, and bound through it the projected-gradient residual whose zeros define the first-order notion.

Suppose that $\mathcal{N} := \{1, \dots, N\}$ indexes the players and player $i$ chooses an action $x_i$ from a nonempty compact convex set $X_i \subseteq \mathbb{R}^{n_i}$. Learning-based Nash equilibrium (LBNE) (Scarabaggio et al., 2024) fits a map $f_i(x_{-i})$ predicting player $i$'s response to the other players' actions, $x_{-i}$. The *behavioral equilibrium* is a profile $x^\star$ for which $x_i^\star = f_i(x_{-i}^\star)$ for every $i \in \mathcal{N}$. Each $f_i$ composes a learned observer, a response head, and a Euclidean projection onto $X_i$, so that feasibility of the response is enforced by the projection. Given the continuity of each $f_i$ and the convexity and compactness of the product $X$, Scarabaggio et al. (2024) establish that the joint map $F = (f_1, \dots, f_N)$ admits a fixed point by Brouwer's theorem (Brouwer, 1911). However, convergence of the prescribed Picard–Banach iteration $x^{k+1} = F(x^k)$ is secured only under an aggregate contraction condition $\left(\sum_{i \in \mathcal{N}} \gamma_i^2\right)^{1/2} < 1$ on the playerwise Lipschitz constants $\gamma_i$, which excludes the nonexpansive case at which the iteration can fail to converge at all. Contraction hypotheses of this kind are replete across response-learning and decision-dependent formulations, where they buy uniqueness, a linear rate, and a perturbation bound $\varsigma/(1-q)$ under a uniform map error $\varsigma$ (Mignoni, 2025; Narang et al., 2023; Bauschke & Combettes, 2017).





Fig. 1: **A learned response map, and the game it is fitted to.** Both $G_i$ (6) and $\rho_i$ (9) are differences from the same $\hat{x}_i$; $\delta_i$ is the consistency error, the one discrepancy that cannot be formed from the model alone.

These guarantees constrain the learned map rather than the game. A contractive $F$ has a unique fixed point of the *learned* game, which need not be an equilibrium of the game that generated the data: the certificate bounds the sensitivity of a fitted network, whereas the quantity of interest is the regret a player would incur at the predicted profile. For arbitrary strategies $\hat{x} \in X$, a constant map $F \equiv \hat{x}$ is nonexpansive and has zero fixed-point residual at $\hat{x}$, while its exploitability is determined by the environment alone and is unconstrained by that residual. Enforcing contraction restricts the hypothesis class where the restriction is least affordable i.e. an interaction whose true best-response map is not contractive admits no contractive model since multi-agent retraining dynamics admit chaotic orbits (Piliouras & Yu, 2023).

While Scarabaggio et al. (2024) introduced the projected observer–head response map and showed that its fixed point solves a variational inequality of the learned game, Narang et al. (2023) obtained uniqueness and a linear rate for performatively stable equilibria under an analogous ratio condition. In a similar vein, Bemporad & Tatarenko (2026a) trained a network against the Nikaido–Isoda gap (Nikaidô & Isoda, 1955) of a surrogate game, while Bemporad & Tatarenko (2026b) instead enforce monotonicity of the learned game by construction. In contrast, Gürkan & Pang (2009) characterized when equilibria of approximating games converge to equilibria of the true game; however, the results are asymptotic with no explicit constant, and Bergerault et al. (2026) proved that the Nash gap, the exploitability of a profile, is *not* determined uniformly over games by a measure-matching error, establishing a divergence between prediction loss and any equilibrium guarantee. Neither approach yields a number that can be evaluated at a given profile *in-silico*: the first certifies the model under a hypothesis that may fail on the interaction of interest, and the second delimits what fitting can achieve without quantifying the shortfall anywhere. A certificate should therefore bound regret in the true game of the prediction model's profile, and should (ideally) be finite when the learned map does not contract.

Write $\rho_i$ for player $i$'s fixed-point residual, $\delta_i$ for the consistency error between the learned response operator and the true pseudo-gradient, and $\bar{\delta}_i$ for a bound on $\delta_i$ over the certified region (see Fig. 1). Our **contributions** are as follows: (1) We certify the exploitability of every profile $\hat{x} \in \Omega$ with a playerwise envelope, without requiring contraction, monotonicity, or equilibrium uniqueness. (2) We derive a quadratic exploitability bound in the distance to an interior best response, while the general envelope certificate remains linear in $\rho_i + \bar{\delta}_i$. (3) We show that the residual entry earns its cost only under amortization: $\bar{\delta}_i$ is enveloped once on a certified region using a finite covering, after which each residual costs one forward pass. (4) Across the two product-set games with analytic or numerically verified ground truth and seven response families, the bound holds in all 280 theorem-supported cells. The remaining 140 cells use a coupled feasibility constraint and are reported only as a stress diagnostic. Pooling the certifiable product-set cells with a coupling sweep, a small-gain contraction test admits a fraction 0.09 of the 1000 theorem-supported cells, against a fraction 0.524 whose exploitability is at or below 0.01, so that certification and equilibrium quality come apart.

The rest of this paper is organized as follows: Section 2 formulates the game and its learned response map, Section 3 states the certificate, Section 4 gives the envelope and its cost, and Section 5 reports the measurements. Proofs of our main results, discussions and limitations, as well as additional supporting empirical results are available in the appendices.

## 2 Problem Formulation

This section describes the problem formulation: we state assumptions, form this paper's notations and introduce conceptual machineries that enable our results.

### 2.1 Notations and Preliminaries

Let $\mathcal{N} = \{1, \dots, N\}$ denote players' indices. For each player $i$, let $x_i$ denote its action in a nonempty compact convex set $X_i \subseteq \mathbb{R}^{n_i}$. Player $i$ observes other players' actions $x_{-i} = (x_j)_{j \neq i}$ and incurs a local cost $J_i : X \to \mathbb{R}$ with $J = (J_1, \dots, J_N)$ as a game's cost profile. The joint action set is the product $X = \prod_{i \in \mathcal{N}} X_i \subseteq \mathbb{R}^d$, $d = \sum_{i \in \mathcal{N}} n_i$, and $X_{-i} = \prod_{j \neq i} X_j$. The action profile $x = (x_1, \dots, x_N) \in X$ is also written as $x = (x_i, x_{-i})$ when player $i$'s action is singled out. *We reserve $x^\star$ for an equilibrium and $\hat{x}$ for an arbitrary strategy profile at which a certificate is evaluated.* For vectors $v_1, \dots, v_N$ we write $\mathrm{col}(v_1, \dots, v_N)$ for the column vector obtained by stacking them. For a closed convex set $S$, the Euclidean projection $\Pi_S(x) := \arg\min_{y \in S} \|x - y\|$ is the unique element of $S$ nearest $x$; since $X$ is a product, $\Pi_X = (\Pi_{X_1}, \dots, \Pi_{X_N})$ acts blockwise. For differentiable $J_i$ we write $\nabla_{x_i} J_i(x)$ for its gradient in player $i$'s own action.

We now state the assumptions that underlie our theoretical machinery.

**Assumption 2.1** (Compact convex product actions)**.** Each action set $X_i$ is nonempty, compact, and convex, and the joint action set is the product $X = \prod_{i \in \mathcal{N}} X_i$.

**Assumption 2.2** (Playerwise convexity)**.** For each player $i$ and each $x_{-i} \in X_{-i}$, the cost $J_i(\cdot, x_{-i})$ is convex and continuously differentiable on $X_i$. The *pseudo-gradient* stacks the players' own-action gradients,

$$\Phi(x) \triangleq \mathrm{col}\left(\Phi_1(x), \dots, \Phi_N(x)\right) \in \mathbb{R}^d, \qquad \Phi_i(x) \triangleq \nabla_{x_i} J_i(x), \tag{1}$$

so that $\Phi_i(x) \in \mathbb{R}^{n_i}$ is player $i$'s block of the pseudo-gradient.

We assume no monotonicity of the pseudo-gradient (1) and no map contraction. The product structure in Assum. 2.1 makes $\Pi_X$ act blockwise and allows variation of a player's action under other players' fixed actions. Games whose feasibility sets couple across players fall outside the development.

All action coordinates and costs are nondimensionalized before applying the maps below: the unit of each action block is the chosen reference action scale, and each cost is divided by a fixed reference cost so that $x_i$ and $\Phi_i(x)$ have the same numerical units. Thus $x - \Phi(x)$ uses a unit proximal step. With physical units, every occurrence of $\Phi$ can equivalently be replaced by $\gamma\Phi$ for a stated step size $\gamma$; the paper fixes this normalization to $\gamma = 1$.

Players' regret measures a profile's quality. The *exploitability* of $x \in X$, equivalently the max-over-players variant of the Nikaido–Isoda gap (Nikaidô & Isoda, 1955), whose original form sums over

players rather than maximizing, is

$$\varepsilon(x) \triangleq \max_{i \in \mathcal{N}} \Big[ J_i(x) - \min_{y_i \in X_i} J_i(y_i, x_{-i}) \Big], \tag{2}$$

in which each bracket is nonnegative, since $x_i \in X_i$ is feasible for the inner minimization. Hence $x \in X$ is a Nash equilibrium precisely when $\varepsilon(x) = 0$, and $\varepsilon(x)$ is the smallest tolerance for which $x$ is an $\varepsilon$-Nash equilibrium (Radner, 1980).

**Definition 2.1** (Variational inequality)**.** Let $\mathcal{K} \subseteq \mathbb{R}^d$ be nonempty and let $T$ map $\mathbb{R}^d$ into itself. The *variational inequality problem*, denoted $\mathrm{VI}(\mathcal{K}, T)$, is to find a vector $x^\star \in \mathcal{K}$ such that

$$T(x^\star)^\top (y - x^\star) \geq 0 \quad \text{for all } y \in \mathcal{K}. \tag{3}$$

Geometrically, (3) asks that $T(x^\star)$ make an acute angle with every feasible direction emanating from $x^\star$ (Harker & Pang, 1990). Under Assumption 2.1 and 2.2 the equilibria coincide with the solutions of $\mathrm{VI}(X, \Phi)$ (Rosen, 1965; Facchinei & Pang, 2010). The *natural map* of $\mathrm{VI}(\mathcal{K}, T)$ is $x \mapsto x - \Pi_{\mathcal{K}}\big(x - T(x)\big)$; it vanishes exactly at the solutions of (3), and we call its norm the *natural residual* (Facchinei & Pang, 2003, §1.5).

**Definition 2.2** (Learned Response Map)**.** Given a dataset of players' observed actions, the response map of player $i$ composes an *observer* $h_i$, which summarizes the other players' actions in a feature vector $h_i(x_{-i})$; a *response head* $g_i(\cdot\,; \theta_i)$, a function approximator with parameters $\theta_i$; and a projection onto player $i$'s action set,

$$f_i(x_{-i}; \theta_i) = \Pi_{X_i}\big(g_i(h_i(x_{-i}); \theta_i)\big). \tag{4}$$

Writing $F(x) = \mathrm{col}\,\big(f_1(x_{-1}), \ldots, f_N(x_{-N})\big)$ for the joint response of all players, a *behavioral equilibrium* is a profile $x^\star \in X$ for which every player's prediction reproduces its own input,

$$x_i^\star = f_i(x_{-i}^\star; \theta_i) \quad \text{for all } i \in \mathcal{N}, \tag{5}$$

equivalently $x^\star = F(x^\star)$ (Scarabaggio et al., 2024).

Because $\Pi_{X_i}$ maps into $X_i$, we have $F(\mathbb{R}^d) \subseteq X$ and in particular $F : X \to X$, and because the projection onto a closed convex set is nonexpansive (Facchinei & Pang, 2003, §1.5), continuity of $f_i$ reduces to continuity of the learned part $g_i \circ h_i$. Under Assum. 2.1 the joint action set is nonempty, compact and convex, so that Brouwer's theorem (Brouwer, 1911) supplies a profile satisfying (5) whenever each $g_i \circ h_i$ is continuous.

The *induced operator* of the projected response map is

$$G(x) \triangleq x - F(x), \tag{6}$$

so $x - G(x) = F(x) \in X$ and therefore $F(x) = \Pi_X\big(x - G(x)\big)$. The fixed-point residual $x - F(x) = G(x)$ is thus the natural map of $\mathrm{VI}(X, G)$ (Facchinei & Pang, 2003, §1.5). The learned map is connected to the game via (i) the fixed-point residual of $F$, which one forward pass of the model computes; (ii) the operator-consistency error between (6) and (1), which cannot be formed without evaluating the true costs; and (iii) the natural residual of the variational inequality $\mathrm{VI}(X, \Phi)$. Only (i) can be computed without access to the game; any certificate that uses (ii) or (iii) must query the environment.

*Remark* 2.1 (A training loss does not bound $\delta$). Small prediction loss leaves the operator-consistency error uncontrolled: a response map fitted to observed actions constrains $G$ where data was observed, while $\Phi$ enters the exploitability (2) through its values on the region being certified. Bergerault et al. (2026) make the analogous point for multi-agent imitation learning: they exhibit games in which exact measure matching still leaves a large Nash gap, and prove that characterizing the Nash gap at a fixed measure-matching error is hard, recovering bounds only under additional structure on the expert equilibrium.

## 3 An Amortized Exploitability Certificate

Player $i$'s regret is controlled by its blockwise residual and its gradient magnitude, both defined in (7). Lemma 3.1 establishes the pointwise residuals with local constants for the true pseudo-gradient.

Lemma 3.2 produces a learned map, trading the variational residual for the fixed-point residual $\rho_i$ (see Fig. 1) and the consistency error $\delta_i$ of (9). Every result in this section is proved in Appendix C, and Appendix D collects what the constants and the rate do and do not give: what an average prediction error fails to control, whether the coefficients can be shrunk, and when the rate improves from linear to quadratic.

### 3.1 A BLOCKWISE GAP BOUND

For a profile $x \in X$ write the gradient magnitude, blockwise residual, and the diameter of each player's block as

$$M_i(x) \triangleq \|\Phi_i(x)\|, \;\; D_i \triangleq \operatorname{diam} X_i, \;\; r_i(x) \triangleq \big\|x_i - \Pi_{X_i}\big(x_i - \Phi_i(x)\big)\big\|. \tag{7}$$

Here $\operatorname{diam} X_i = \sup_{u,v \in X_i} \|u - v\|$, finite by Assum. 2.1. Each $r_i(x)$ is the norm of block $i$ of the right-hand side of the projected pseudo-gradient flow $\dot{x} = \Pi_X\big(x - \Phi(x)\big) - x$, so that every $r_i(x)$ vanishes exactly at the rest points of that flow: where the flow moves slowly, no player gains much by deviating.

**Lemma 3.1** (Blockwise gap bound)**.** *Suppose that Assumption 2.1 and 2.2 hold. Then for every $x \in X$,*

$$\varepsilon(x) \;\le\; \max_{i \in \mathcal{N}} \; \min\big\{\big(M_i(x) + D_i\big)\, r_i(x), \;\; M_i(x)\, D_i\big\}. \tag{8}$$

Lemma 3.1 follows from the projection inequality (Facchinei & Pang, 2003, §1.5). Its constants $M_i(x)$ and $D_i$ are local and blockwise: $M_i$ is evaluated at $x$ rather than maximized over $X$, and each refers only to player $i$'s own gradient and own action set. Error bounds of this kind ordinarily control the distance from $x$ to the solution set, for which the constant exists only under strong monotonicity (Yamashita & Fukushima, 1997). Equation (8) controls exploitability instead, for which compactness suffices. The second entry of the minimum, $M_i(x)D_i$, refers to no learned quantity and is available without a model, so that a certificate is informative only when it falls below $\max_i M_i(x)D_i$ — the comparison Corollary 3.1 makes precise.

### 3.2 TRANSFER TO A LEARNED MAP

Write $G_i(x) = x_i - f_i(x_{-i})$ for the $i$-th block of the induced operator (6), and define the blockwise residual $\rho_i$ and the consistency error $\delta_i$ by

$$\rho_i(x) \triangleq \big\|G_i(x)\big\|, \qquad \delta_i(x) \triangleq \big\|G_i(x) - \Phi_i(x)\big\|, \tag{9}$$

respectively.

**Lemma 3.2** (Blockwise residual transfer)**.** *Let Assum. 2.1 hold and let $\Phi_i$ be defined on $X$. Then $r_i(x) \le \rho_i(x) + \delta_i(x)$ for every $x \in X$ and every player $i$.*

**Proposition 3.1** (The learned map cannot improve a pointwise certificate)**.** *Let Assum. 2.1 hold and let the envelope of (12) be evaluated pointwise at $\hat{x}$, so that $\Omega = \{\hat{x}\}$, $\bar{M}_i = M_i(\hat{x})$, and $\bar{\delta}_i = \delta_i(\hat{x})$. If $M_i(\hat{x}) > 0$, then*

$$\big(M_i(\hat{x}) + D_i\big)\big(\rho_i(\hat{x}) + \delta_i(\hat{x})\big) \;\ge\; M_i(\hat{x})^2 + M_i(\hat{x})D_i \;>\; M_i(\hat{x})D_i, \tag{10}$$

*so player $i$'s minimum is attained by the model-free entry. If $M_i(\hat{x}) > 0$ for every player, the pointwise certificate reduces to $\max_i M_i(\hat{x})D_i$.*

Since forming $\delta_i(\hat{x})$ requires $\Phi_i(\hat{x})$, which already yields the model-free entry, the residual entry is worth its cost only when the consistency error is bounded once over a region rather than at the profile.

**Scope of the pointwise limitation.** The comparison is playerwise and strict when $M_i(\hat{x}) > 0$; if $M_i(\hat{x}) = 0$, both entries may vanish. It concerns a fresh pointwise audit, where evaluating $\delta_i(\hat{x})$ already requires the same true gradient that supplies the model-free entry. A regional envelope changes the comparison because it is fixed before the later profile is chosen and can be reused without another environment query.

### 3.3 AMORTIZING THE ENVELOPE OVER A REGION

The consistency error is a property of the model against the environment, so it can be bounded once on a region and reused. The residual is a property of the model alone, so it is free wherever it is evaluated. Separating the two is what makes the certificate worth computing.

**Definition 3.1** (Regional envelope)**.** Let $\Omega \subseteq X$ be closed. An *envelope* of a trained model on $\Omega$ is a pair of vectors $(\bar{M}, \bar{\delta})$ with, for every player $i$,

$$\bar{M}_i \geq \sup_{x \in \Omega} M_i(x), \qquad \bar{\delta}_i \geq \sup_{x \in \Omega} \delta_i(x). \tag{11}$$

**Theorem 3.1** (Amortized exploitability certificate)**.** *Let Assumption 2.1 and 2.2 hold, let $(\bar{M}, \bar{\delta})$ envelope a trained model on $\Omega \subseteq X$, and let $F$ be of the form (4). Then* every *profile $\hat{x} \in \Omega$ satisfies*

$$\varepsilon(\hat{x}) \;\leq\; \max_{i \in \mathcal{N}} \min\big\{\big(\bar{M}_i + D_i\big)\big(\rho_i(\hat{x}) + \bar{\delta}_i\big),\;\; \bar{M}_i D_i\big\}, \tag{12}$$

*and the residuals $\rho_i(\hat{x})$ require no access to the environment.*

When $(\bar{M}, \bar{\delta})$ is measured by probes rather than supplied by an oracle, (12) inherits the confidence of that joint envelope and holds with probability at least $1 - \alpha$ (Proposition 4.1); the same qualification attaches to Corollary 3.1.

**Corollary 3.1** (Non-vacuity threshold)**.** *Fix a player $i$ and suppose $\bar{M}_i + D_i > 0$, which holds whenever $X_i$ has more than one point. Then the residual entry of (12) improves on the model-free entry if and only if*

$$\rho_i(\hat{x}) + \bar{\delta}_i \;<\; \frac{\bar{M}_i D_i}{\bar{M}_i + D_i} \;\leq\; \min\{\bar{M}_i,\, D_i\}. \tag{13}$$

The threshold (13) is what player $i$'s residual entry must improve on over the model-free one; Theorem 3.1 holds either way. Because (12) reports a maximum over players, the certificate improves on $\max_i \bar{M}_i D_i$ only when (13) holds at every player attaining that maximum; clearing it for some other player buys nothing. It is built from $\bar{M}_i$ and $D_i$, which carry no learned quantity, hence training cannot relax it, and a lower training loss need not lower $\bar{\delta}_i$ either (Proposition D.1). Since $\rho_i \geq 0$, the envelope alone must clear the threshold, at an accuracy fixed by the game rather than by the model; through (17) that accuracy in turn fixes the covering radius $\eta$, and the covering size grows as $(\operatorname{diam}\Omega/\eta)^{\dim \Omega}$.

**Proposition 3.2** (A spurious stationary profile)**.** *Let $X_i = [-c, c]$ with $c > (1-\kappa)^{-1/2}$ and let*

$$J_i(x_i, x_{-i}) = \tfrac{1}{4}x_i^4 - \tfrac{1}{2}w_i(x_{-i})\,x_i^2, \qquad w_i(x_{-i}) = 1 + \tfrac{\kappa}{N-1}\sum_{j \neq i} x_j^2, \tag{14}$$

*with $\kappa \in [0, 1)$. Then $\Phi(0) = 0$, hence $r_i(0) = 0$ for every player, while $\varepsilon(0) = \frac{1}{4}$.*

The origin is a strict local maximum of each $J_i(\cdot, x_{-i})$ in its own argument, and it is the conditional mean of play whenever the data-generating process is symmetric across the two minimizers $\pm\sqrt{w_i(x_{-i})}$. A regressor fitted to conditional means therefore lands where the blockwise residual vanishes and exploitability is nonetheless $\frac{1}{4}$: without playerwise convexity, a vanishing residual certifies nothing. Under Assum. 2.2, multimodal play arises from multiple equilibria rather than from nonconvexity.

## 4 ENVELOPE COSTS

This section states the costs of the envelope in the sense of Definition 3.1 and the weaker guarantees available when those costs are omitted.

### 4.1 PROBING AT A PROFILE

Two access models bracket what a probe can learn. A *gradient oracle* returns $\Phi_i(x)$ directly; a *cost oracle* returns only the values $J_i(x)$, so that $\Phi_i$ must be estimated. We assume the cost oracle, the

access model of payoff-based equilibrium learning (Tatarenko & Kamgarpour, 2024). Differencing player $i$'s own coordinates at radius $\tau > 0$ gives

$$\big[\widehat{\Phi}_\tau(x)\big]_{i,k} = \frac{J_i(x + \tau e_{i,k}) - J_i(x - \tau e_{i,k})}{2\tau}, \tag{15}$$

in which $e_{i,k}$ is the unit vector along coordinate $k$ of player $i$'s block. Only own coordinates are differenced, since $\Phi_i = \nabla_{x_i} J_i$ involves no cross-partial, and the estimator costs $2d$ evaluations per profile, with $d = \sum_{i \in \mathcal{N}} n_i$. Costs returned by a simulator, or averaged over finitely many rollouts of a stochastic environment, are not exact, so we take a query at $x$ to return $J_i(x) + Z$ with $Z$ independent across queries, mean-zero and $\sigma$-sub-Gaussian. Equation (15) then carries both a smoothing bias and a noise term, and $\tau$ trades one against the other.

*Remark* 4.1 (Probes at the boundary). Since every $f_i$ ends in a projection onto $X_i$, certified profiles frequently lie on $\partial X$, and then $x \pm \tau e_{i,k}$ leaves $X$ where the costs are neither defined nor assumed smooth. Two remedies are available. Proposition 4.1 adopts the first, assuming each $J_i$ extends three times continuously differentiably to a $\tau$-neighborhood of $X$, letting its Taylor expansion run along the whole probe segment. When unavailable, we replace (15) by a one-sided difference on coordinates within $\tau$ of the boundary. This has bias $O(B_2\tau)$, with $B_2$ bounding the second derivative. Thus, the balanced radius on the coordinates becomes $\tau \asymp (\sigma/B_2)^{1/2}$ and their contribution to the penalty is $O\big(\sigma^{1/2} B_2^{1/2}\big)$, in both cases up to the $\sqrt{n_i}$ and logarithmic factors that (17) carries.

### 4.2 From probes to a regional envelope

Finite probes control a continuum only through supplied moduli of continuity. For every player, let $L_{\delta,i}$ and $L_{M,i}$ be valid Lipschitz upper bounds for $\delta_i$ and $M_i$ on $\Omega$. Because $G_i(x) = x_i - f_i(x_{-i})$, a valid bound for $L_{\delta,i}$ combines the identity block, a certified Lipschitz bound for the projected response $f_i$, and a valid bound for $\Phi_i$; learned-model weights alone do not bound the environment term. Proposition 4.1 takes these and the other regularity constants as inputs rather than estimating them from the covering data.

**Proposition 4.1** (Envelope from a finite covering). *Let $\Omega \subseteq X$ be nonempty and let $\mathcal{Z} \subseteq \Omega$ be a supplied finite Euclidean $\eta$-cover: every $x \in \Omega$ lies within distance $\eta$ of some $z \in \mathcal{Z}$. For every player $i$, suppose supplied constants $L_{\delta,i}$ and $L_{M,i}$ are valid Lipschitz bounds for $\delta_i$ and $M_i$ on $\Omega$. Let $\tau > 0$, suppose every required probe segment lies in a domain on which $J_i$ has a $C^3$ extension satisfying $|\partial^3 J_i / \partial x_{i,k}^3| \le B_3$, and let cost-query errors be independent, mean-zero, and $\sigma$-sub-Gaussian. Define*

$$E_i(\tau, \alpha) \triangleq \tfrac{1}{6} B_3 \tau^2 \sqrt{n_i} + \tfrac{\sigma}{\tau} \sqrt{n_i \log(2d|\mathcal{Z}|/\alpha)}. \tag{16}$$

*After the $2d|\mathcal{Z}|$ scalar cost evaluations in (15), the same event of probability at least $1 - \alpha$ simultaneously yields, for every player,*

$$\bar{M}_i = \max_{z \in \mathcal{Z}} \big\|\widehat{\Phi}_{\tau,i}(z)\big\| + L_{M,i}\eta + E_i(\tau, \alpha), \tag{17}$$

$$\bar{\delta}_i = \max_{z \in \mathcal{Z}} \big\|G_i(z) - \widehat{\Phi}_{\tau,i}(z)\big\| + L_{\delta,i}\eta + E_i(\tau, \alpha), \tag{18}$$

*and these quantities satisfy both inequalities in (11) for all players.*

Balancing the bias and noise terms in (17) at $\tau \asymp \big(\sigma\sqrt{\log(2d|\mathcal{Z}|/\alpha)}/B_3\big)^{1/3}$ leaves a penalty of order $\sqrt{n_i}\,\big(\log(2d|\mathcal{Z}|/\alpha)\big)^{1/3} \sigma^{2/3} B_3^{1/3}$, which vanishes with the oracle noise. The covering term $L_{\delta,i}\eta$ is a modeling cost, and at fixed relative resolution the covering number of a box-like $m$-dimensional region is $O((\operatorname{diam}\Omega/\eta)^m)$. A rigorous envelope is therefore affordable only for regions with small covering numbers, such as a line segment between two candidate profiles, the trajectory of a response iteration, or a neighborhood of a single deployment point.

*Remark* 4.2 (The finite-query construction is conditional on supplied regularity). Proposition 4.1 requires valid upper bounds $L_{\delta,i}$, $B_3$, and $L_{M,i}$, together with a justified independent mean-zero sub-Gaussian noise model. Finite cost values cannot establish these uniform regional quantities because they do not control behavior between queried points. A sampled violation can refute a proposed bound, but the absence of such a violation cannot validate it. A rigorous application must therefore obtain the constants from an analytic simulator, physical specifications, symbolic or interval bounds, a

certified environment class, a jointly certified environment–model class, or a separately justified noise model. Weight-derived bounds can control the learned response $f_i$ through the composite $g_i \circ h_i$ and the nonexpansive projection; the environment terms remain external assumptions. If these inputs are unavailable, the finite-cover construction is a diagnostic rather than a uniform certificate. Random sampling without an additional quantile or tolerance argument is also an empirical diagnostic, not a bound on the regional supremum.

## 5 EXPERIMENTS

We test soundness, the pointwise limitation, and regional amortization. Exploitability is computed from the analytic game, not the learned response. The seven responders are an affine baseline, two-layer 32–32 tanh/ReLU, spectral-normalized and Almost-Orthogonal-Layer (AOL) Lipschitz-certified networks (Prach & Lampert, 2022), and 64–64 conditional diffusion and flow models. Standard E1–E2 runs supply 240 noisy observational transitions; neural responders use Adam with batch size 128 for 200 epochs at learning rate $3 \times 10^{-3}$ (AOL: $5 \times 10^{-3}$), with least-squares, denoising, and flow-matching losses as appropriate. E8 uses its stated reduced training budget. A cell is one game, response family, and seed; comparisons are paired within cells, with 10,000-resample bootstrap intervals (Efron, 1979), Wilcoxon signed-rank tests (Wilcoxon, 1945) and Holm's step-down correction (Holm, 1979). The full protocol appears in Appendix F.

The learner and the auditor use different information. For each seed and player, training uses 240 pairs $(x_{-i}, x_i)$ obtained from sixty four-step trajectories of noisy simultaneous best-response play. Each trajectory starts from an independent uniform draw from the game's bounding box, projected onto the feasible set; these restarts broaden the input support relative to one long trajectory. The multiple-equilibrium experiment uses its game-specific multimodal response process. The learner sees actions but no costs or gradients. The theorem is otherwise training-agnostic. Training loss does not determine either regional envelope component: $\bar{M}_i$ concerns the true game, while $\bar{\delta}_i$ measures uniform agreement between that game and the trained response map. Cost-aware training is compatible with the certificate and may reduce $\bar{\delta}_i$; we study the complementary problem of independently auditing an already trained model.

**Bound and Tightness of Learned Map.** The tightness study fits seven response families over 20 seeds on three games: a coupled quadratic game whose coupling parameter fixes the monotonicity modulus exactly, a capacitated Cournot game whose capacities bind at equilibrium, and a Rosen game carrying one constraint shared across players. The quadratic and Cournot games have product action sets and test Theorem 3.1 directly: $2 \times 7 \times 20 = 280$ cells, all valid. The Rosen game violates the product requirement of Assum. 2.1; its 140 cells are reported only as a stress diagnostic and are excluded from theorem coverage. The shared game has an analytically specified best-response oracle, but its variational equilibrium is computed to the documented numerical tolerance rather than in closed form.

The standard model-free entry is active in every cell, with median tightness 108.7; the learned-residual entry has median 297.9(about 2.7 times larger). Here tightness is the certified bound divided by exact exploitability, so lower is better. Its paired difference from a fresh pointwise environment query is -0.001165in $\log_{10}$ tightness ($p_{\mathrm{Holm}} = 0.1519$); the test does not detect a difference. The model-free convexity/Cauchy–Schwarz entry is standard and is not our novelty. This experiment measures playerwise scaling, not a tightness improvement from learning; the learned response helps only after the consistency error is audited over a region and reused.

**Playerwise scaling.** The scaling study varies the player count $N$ while holding the observational-data budget fixed. At a fixed displacement from the reference equilibrium, the aggregate ratio of certificate to true exploitability increases from 38.15 at $N = 2$ to 1157 at $N = 256$. The corresponding blockwise ratio changes from 13.92 to 6.628. The aggregate ratio therefore grows by a factor of about thirty, whereas the blockwise ratio remains on the same order, which Fig. 2 shows at both action dimensions and Table 1 lists at scalar actions. Shading marks where the aggregated bound exceeds the model-free bound; the dash-dot rule marks where the responder fit becomes underdetermined.

The aggregate form uses one global constant and a joint residual, whereas the playerwise form uses (7); here the blockwise series equals the model-free entry. The fixed 240-transition budget makes the largest fits underdetermined, which can inflate learned-map residuals but not the model-free entry.





Fig. 2: **Joint constants degrade with the player count; playerwise ones do not.** Coupled quadratic game, (a) scalar and (b) two-dimensional actions, medians over 20 seeds, log–log, lower is tighter.

**Regional auditing creates the amortization regime.** At $\eta/\operatorname{diam}\Omega = 0.025$, the median cover contains 20 points on the one-dimensional segment and 8000 points on the three-dimensional box. The learned-residual entry binds on 0.5742 and 0.7266 of held-out profiles, with regional-to-pointwise ratios 1.788 and 1.725. Every tested profile satisfies the audited envelope; covering cost grows with region dimension. Fig. 3 shows the tradeoff.

E9 is a gradient-oracle audit, not an implementation of the noisy scalar-cost oracle in Section 4. The quadratic pseudo-gradient is affine, giving the exact modulus $L_{M,i} = \|A_{i,:}\|_2$. For $G_i = x_i - f_i(x_{-i})$, the audit uses the valid bound $L_{\delta,i} = \sqrt{1 + L_{f_i}^2} + \|A_{i,:}\|_2$, where $L_{f_i}$ is the responder's rigorous weight-derived Lipschitz bound. The midpoint grid uses $n_j = \lceil s_j\sqrt{m}/(2\eta)\rceil$ points on an edge of length $s_j$; its cell half-diagonal is at most $\eta$, so it is a Euclidean cover of the embedded $m$-dimensional region. No finite-difference $B_3$, $\sigma$, or $\tau$ enters E9. The conversion to $2d|\mathcal{Z}|$ scalar cost evaluations is an extrapolation under Proposition 4.1, not a measured cost-oracle experiment.

**Contraction and exploitability measure different properties.** Pooling the theorem-supported product-set cells from the tightness study with the coupling sweep gives 1000 model cells; the shared-constraint diagnostics are excluded. Small-gain certification accepts 0.09, while 0.524 have exploitability at most 0.01. The four contingency counts are 70, 20, 454, and 456; contraction is therefore neither necessary nor sufficient here. Generative families are excluded because they have no rigorous global Lipschitz bound.

**Multiplicity, solvers, and probe diagnostics.** On the playerwise-convex non-monotone game, generative responders are evaluated through conditional-mean maps, so the comparison is silent about generativity and capacity; their heads have roughly four times as many parameters yet recover fewer equilibria (Appendix E). The nonconvex witness remains a scope diagnostic: zero variational residual can coexist with nonzero exploitability. On the nonexpansive game, Picard reaches tolerance $10^{-8}$ in





Fig. 3: **Regional auditing trades upfront gradient queries for later model-only certificates.** Three-player quadratic game with two-dimensional actions and a tanh-MLP response; medians over 10 seeds and 256 profiles drawn uniformly from each region. The segment runs from the learned fixed point to a feasible profile obtained by displacing the reference equilibrium by $0.2$ block diameters and projecting. The coordinate boxes have intrinsic dimension one to three and are embedded in the six-dimensional joint action space; they vary their first $m$ coordinates over five percent of the corresponding coordinate ranges. (a) Median Euclidean $\eta$-cover size, equal to the oracle-call break-even against repeated pointwise gradient audits under the stated accounting. (b) Regional certificate divided by the fresh pointwise certificate; values above one show the tightness paid for reuse.

none of 20 runs, while Krasnosel'skiĭ–Mann and Halpern require medians of 126 and 5 iterations. Probe-radius sweeps reproduce the bias–noise tradeoff of Proposition 4.1; both are diagnostics, not replacements for its regularity assumptions.

## 6 Conclusion

A learned fixed point establishes model self-consistency, not low regret in the true game. Our certificate connects the two by combining a model-only residual with independently established regional bounds on true gradients and environment–model discrepancy. Any valid envelope yields profilewise exploitability guarantees without contraction, monotonicity, or equilibrium uniqueness. The finite cost-query construction is narrower: it requires externally certified regularity and noise assumptions and has a covering cost exponential in the audited region's intrinsic dimension. Its practical regime is therefore a structured region that is audited once and reused many times. In the pointwise experiments, the standard model-free entry bound all 280 configurations; the learned map's value appeared only in the regional regime, where it exchanged a looser bound and substantial upfront cost for later environment-query-free evaluation. Without the external assumptions, the finite-query procedure is diagnostic rather than a uniform certificate. Without playerwise convexity, a vanishing projected-gradient residual can coexist with exploitability $\frac{1}{4}$.

## REFERENCES


Yakov Babichenko. Query Complexity of Approximate Nash Equilibria. *Journal of the ACM*, 63(4): 1–24, 2016. doi: 10.1145/2908734.

Heinz H. Bauschke and Patrick L. Combettes. *Convex Analysis and Monotone Operator Theory in Hilbert Spaces*. CMS Books in Mathematics. Springer, Cham, 2nd edition, 2017. doi: 10.1007/978-3-319-48311-5.

Giuseppe Belgioioso, Peng Yi, Sergio Grammatico, and Lacra Pavel. Distributed Generalized Nash Equilibrium Seeking: An Operator-Theoretic Perspective. *IEEE Control Systems Magazine*, 42(4): 87–102, 2022. doi: 10.1109/MCS.2022.3171480.

Alberto Bemporad and Tatiana Tatarenko. Learning Approximate Solutions to Multiparametric Generalized Nash Equilibrium Problems, 2026a. arXiv preprint arXiv:2605.28757.

Alberto Bemporad and Tatiana Tatarenko. Learning Parametric Monotone Games, 2026b. arXiv preprint arXiv:2609.02494.

Antoine Bergerault, Volkan Cevher, and Negar Mehr. Matching Multiple Experts: On the Exploitability of Multi-Agent Imitation Learning, 2026. arXiv preprint arXiv:2602.21020.

Jérôme Bolte and Edouard Pauwels. Conservative set valued fields, automatic differentiation, stochastic gradient methods and deep learning. *Mathematical Programming*, 188(1):19–51, 2021. doi: 10.1007/s10107-020-01501-5.

L. E. J. Brouwer. Über Abbildung von Mannigfaltigkeiten. *Mathematische Annalen*, 71(1):97–115, 1911. doi: 10.1007/BF01456931.

Frank H. Clarke. *Optimization and Nonsmooth Analysis*. Canadian Mathematical Society Series of Monographs and Advanced Texts. John Wiley & Sons, New York, 1983. Reprinted as SIAM Classics in Applied Mathematics 5, 1990, doi:10.1137/1.9781611971309.

Stella Dafermos. Sensitivity analysis in variational inequalities. *Mathematics of Operations Research*, 13(3):421–434, 1988. doi: 10.1287/moor.13.3.421.

Alexander Davydov, Saber Jafarpour, and Francesco Bullo. Non-Euclidean Contraction Theory for Robust Nonlinear Stability. *IEEE Transactions on Automatic Control*, 67(12):6667–6681, 2022. doi: 10.1109/TAC.2022.3183966.

Roy Dong, Heling Zhang, and Lillian J. Ratliff. Approximate Regions of Attraction in Learning with Decision-Dependent Distributions. In *Proceedings of the 26th International Conference on Artificial Intelligence and Statistics (AISTATS)*, volume 206 of *Proceedings of Machine Learning Research*, pp. 11172–11184. PMLR, 2023.

Asen L. Dontchev and R. Tyrrell Rockafellar. Characterizations of strong regularity for variational inequalities over polyhedral convex sets. *SIAM Journal on Optimization*, 6(4):1087–1105, 1996. doi: 10.1137/S1052623495284029.

B. Efron. Bootstrap methods: Another look at the jackknife. *The Annals of Statistics*, 7(1):1–26, 1979. doi: 10.1214/aos/1176344552.

Promise Ekpo, Angelique Taylor, and Lekan Molu. A Generalized Nash Equilibrium-Seeking Scheme for Trauma Resuscitation. In *Proceedings of the IFAC World Congress*, Busan, Republic of Korea, 2026.

Francisco Facchinei and Christian Kanzow. Generalized Nash Equilibrium Problems. *Annals of Operations Research*, 175(1):177–211, 2010. doi: 10.1007/s10479-009-0653-x.

Francisco Facchinei and Jong-Shi Pang. *Finite-Dimensional Variational Inequalities and Complementarity Problems*. Springer Series in Operations Research and Financial Engineering. Springer, New York, 2003. doi: 10.1007/b97544. Two volumes, continuously paginated pp. 1–1234.

Francisco Facchinei and Jong-Shi Pang. Nash Equilibria: The Variational Approach. In Daniel P. Palomar and Yonina C. Eldar (eds.), *Convex Optimization in Signal Processing and Communications*, pp. 443–493. Cambridge University Press, Cambridge, 2010. doi: 10.1017/CBO9780511804458.013. Published online 3 December 2009.

Mahyar Fazlyab, Alexander Robey, Hamed Hassani, Manfred Morari, and George J. Pappas. Efficient and Accurate Estimation of Lipschitz Constants for Deep Neural Networks. In *Advances in Neural Information Processing Systems 32 (NeurIPS 2019)*, 2019.

Masao Fukushima. Equivalent differentiable optimization problems and descent methods for asymmetric variational inequality problems. *Mathematical Programming*, 53(1–3):99–110, 1992. doi: 10.1007/BF01585696.

Gül Gürkan and Jong-Shi Pang. Approximations of Nash Equilibria. *Mathematical Programming*, 117(1–2):223–253, 2009. doi: 10.1007/s10107-007-0156-y. Published online 21 August 2007.

Patrick T. Harker and Jong-Shi Pang. Finite-dimensional variational inequality and nonlinear complementarity problems: A survey of theory, algorithms and applications. *Mathematical Programming*, 48(1–3):161–220, 1990. doi: 10.1007/BF01582255.

John C. Harsanyi. Games with randomly disturbed payoffs: A new rationale for mixed-strategy equilibrium points. *International Journal of Game Theory*, 2(1):1–23, 1973. doi: 10.1007/BF01737554.

Sergiu Hart and Andreu Mas-Colell. Uncoupled Dynamics Do Not Lead to Nash Equilibrium. *American Economic Review*, 93(5):1830–1836, 2003. doi: 10.1257/000282803322655581.

Aaron J. Havens, Alexandre Araujo, Siddharth Garg, Farshad Khorrami, and Bin Hu. Exploiting Connections between Lipschitz Structures for Certifiably Robust Deep Equilibrium Models. In *Advances in Neural Information Processing Systems 36 (NeurIPS 2023)*, pp. 21658–21674, 2023.

Michael D. Hirsch, Christos H. Papadimitriou, and Stephen A. Vavasis. Exponential lower bounds for finding Brouwer fix points. *Journal of Complexity*, 5(4):379–416, 1989. doi: 10.1016/0885-064X(89)90017-4.

Josef Hofbauer and Ed Hopkins. Learning in perturbed asymmetric games. *Games and Economic Behavior*, 52(1):133–152, 2005. doi: 10.1016/j.geb.2004.06.006.

Sture Holm. A simple sequentially rejective multiple test procedure. *Scandinavian Journal of Statistics*, 6(2):65–70, 1979. doi: 10.2307/4615733.

Guoqiang Hu, Yipeng Pang, Chao Sun, and Yiguang Hong. Distributed Nash Equilibrium Seeking: Continuous-Time Control-Theoretic Approaches. *IEEE Control Systems Magazine*, 42(4):68–86, 2022. doi: 10.1109/MCS.2022.3171479.

Saber Jafarpour, Alexander Davydov, Anton V. Proskurnikov, and Francesco Bullo. Robust Implicit Networks via Non-Euclidean Contractions. In *Advances in Neural Information Processing Systems 34 (NeurIPS 2021)*, pp. 9857–9868, 2021.

Shizuo Kakutani. A generalization of Brouwer's fixed point theorem. *Duke Mathematical Journal*, 8 (3):457–459, 1941. doi: 10.1215/S0012-7094-41-00838-4.

Torbjörn Larsson and Michael Patriksson. A class of gap functions for variational inequalities. *Mathematical Programming*, 64(1–3):53–79, 1994. doi: 10.1007/BF01582565.

Tao Li, Guanze Peng, Quanyan Zhu, and Tamer Başar. The confluence of networks, games, and learning a game-theoretic framework for multiagent decision making over networks. *IEEE Control Systems Magazine*, 42(4):35–67, 2022. doi: 10.1109/MCS.2022.3171478.

Eric Mazumdar, Lillian J. Ratliff, and S. Shankar Sastry. On Gradient-Based Learning in Continuous Games. *SIAM Journal on Mathematics of Data Science*, 2(1):103–131, 2020. doi: 10.1137/18M1231298.

Richard D. McKelvey and Thomas R. Palfrey. Quantal Response Equilibria for Normal Form Games. *Games and Economic Behavior*, 10(1):6–38, 1995. doi: 10.1006/game.1995.1023.

Panayotis Mertikopoulos and Zhengyuan Zhou. Learning in games with continuous action sets and unknown payoff functions. *Mathematical Programming*, 173(1–2):465–507, 2018. doi: 10.1007/s10107-018-1254-8.

Nicola Mignoni. *Non-Cooperative Game Theoretical Control for Green and Efficient Energy Communities*. PhD thesis, Politecnico di Bari, Bari, Italy, 2025. URL https://hdl.handle.net/11589/284440. Chapter 6.

Adhyyan Narang, Evan Faulkner, Dmitriy Drusvyatskiy, Maryam Fazel, and Lillian J. Ratliff. Multiplayer Performative Prediction: Learning in Decision-Dependent Games. *Journal of Machine Learning Research*, 24(202):1–56, 2023. URL https://jmlr.org/papers/v24/22-0131.html.

Hukukane Nikaidô and Kazuo Isoda. Note on non-cooperative convex game. *Pacific Journal of Mathematics*, 5(5):807–815, 1955. doi: 10.2140/pjm.1955.5.807.

Jong-Shi Pang. Error bounds in mathematical programming. *Mathematical Programming*, 79(1–3): 299–332, 1997. doi: 10.1007/BF02614322.

Lacra Pavel and Yiguang Hong. Distributed games and nash equilibrium seeking in multiagent systems over networks: An introduction to the special issue. *IEEE Control Systems Magazine*, 42 (4):32–34, 2022. doi: 10.1109/MCS.2022.3171477.

Ji-Ming Peng. Equivalence of variational inequality problems to unconstrained minimization. *Mathematical Programming*, 78(3):347–355, 1997. doi: 10.1007/BF02614360.

Juan C. Perdomo, Tijana Zrnic, Celestine Mendler-Dünner, and Moritz Hardt. Performative Prediction. In *Proceedings of the 37th International Conference on Machine Learning (ICML)*, volume 119 of *Proceedings of Machine Learning Research*, pp. 7599–7609. PMLR, 2020. URL https://proceedings.mlr.press/v119/perdomo20a.html.

Georgios Piliouras and Fang-Yi Yu. Multi-Agent Performative Prediction: From Global Stability and Optimality to Chaos. In *Proceedings of the 24th ACM Conference on Economics and Computation (EC)*, pp. 1047–1074. ACM, 2023. doi: 10.1145/3580507.3597759.

Bernd Prach and Christoph H. Lampert. Almost-orthogonal layers for efficient general-purpose lipschitz networks. In *Computer Vision – ECCV 2022*, volume 13681 of *Lecture Notes in Computer Science*, pp. 350–365. Springer, 2022. doi: 10.1007/978-3-031-19803-8_21.

Roy Radner. Collusive behavior in noncooperative epsilon-equilibria of oligopolies with long but finite lives. *Journal of Economic Theory*, 22(2):136–154, 1980. doi: 10.1016/0022-0531(80)90037-X.

Zhongqiang Ren, Sivakumar Rathinam, and Howie Choset. Multi-objective conflict-based search for multi-agent path finding. In *IEEE International Conference on Robotics and Automation (ICRA)*, pp. 8786–8791, 2021. doi: 10.1109/ICRA48506.2021.9560985.

Stephen M. Robinson. Strongly regular generalized equations. *Mathematics of Operations Research*, 5(1):43–62, 1980. doi: 10.1287/moor.5.1.43.

J. B. Rosen. Existence and uniqueness of equilibrium points for concave $N$-person games. *Econometrica*, 33(3):520–534, 1965. doi: 10.2307/1911749.

Aviad Rubinstein. Inapproximability of Nash Equilibrium. In *Proceedings of the 47th Annual ACM Symposium on Theory of Computing (STOC)*, pp. 409–418, 2015. doi: 10.1145/2746539.2746578.

Paolo Scarabaggio, Nicola Mignoni, Raffaele Carli, and Mariagrazia Dotoli. Equilibrium Seeking in Learning-Based Noncooperative Nash Games. In *IEEE 63rd Conference on Decision and Control (CDC)*, pp. 210–215, Milan, Italy, December 2024. IEEE. doi: 10.1109/CDC56724.2024.10886614.

Paolo Scarabaggio, Nicola Mignoni, Raffaele Carli, and Mariagrazia Dotoli. Equilibria Existence in Games: A Perspective from the Lefschetz–Nielsen Theory. In *European Control Conference (ECC)*, pp. 2938–2943, Reykjavík, Iceland, July 2026. EUCA. ISBN 978-3-907144-13-8.

Tatiana Tatarenko and Maryam Kamgarpour. Payoff-based Learning of Nash Equilibria in Merely Monotone Games. *IEEE Transactions on Control of Network Systems*, 11(4):1790–1799, 2024. doi: 10.1109/TCNS.2024.3355035.

Aladin Virmaux and Kevin Scaman. Lipschitz Regularity of Deep Neural Networks: Analysis and Efficient Estimation. In *Advances in Neural Information Processing Systems 31 (NeurIPS 2018)*, 2018.

Anna von Heusinger and Christian Kanzow. Optimization reformulations of the generalized Nash equilibrium problem using Nikaido–Isoda-type functions. *Computational Optimization and Applications*, 43(3):353–377, 2009. doi: 10.1007/s10589-007-9145-6.

Fangji Wang, Han Zhang, Sven Koenig, and Jiaoyang Li. Efficient approximate search for multi-objective multi-agent path finding. In *International Conference on Automated Planning and Scheduling (ICAPS)*, volume 34, pp. 613–622, 2024. doi: 10.1609/icaps.v34i1.31524.

Ruigang Wang and Ian R. Manchester. Direct Parameterization of Lipschitz-Bounded Deep Networks. In *Proceedings of the 40th International Conference on Machine Learning (ICML)*, volume 202 of *Proceedings of Machine Learning Research*, pp. 36093–36110. PMLR, 2023.

Frank Wilcoxon. Individual comparisons by ranking methods. *Biometrics Bulletin*, 1(6):80–83, 1945. doi: 10.2307/3001968.

Ezra Winston and J. Zico Kolter. Monotone Operator Equilibrium Networks. In *Advances in Neural Information Processing Systems 33 (NeurIPS 2020)*, pp. 10718–10728, 2020. URL https://proceedings.neurips.cc/paper/2020/hash/798d1c2813cbdf8bcdb388db0e32d496-Abstract.html.

Nobuo Yamashita and Masao Fukushima. Equivalent unconstrained minimization and global error bounds for variational inequality problems. *SIAM Journal on Control and Optimization*, 35(1): 273–284, 1997. doi: 10.1137/S0363012994277645.

Han Zhang, Oren Salzman, Ariel Felner, Carlos Hernández Ulloa, and Sven Koenig. A-a*pex: Efficient anytime approximate multi-objective search. In *Symposium on Combinatorial Search (SoCS)*, volume 17, pp. 179–187, 2024. doi: 10.1609/socs.v17i1.31556.

Kaiqing Zhang, Zhuoran Yang, and Tamer Başar. Multi-agent reinforcement learning: A selective overview of theories and algorithms. In *Handbook of Reinforcement Learning and Control*, pp. 321–384. Springer, 2021. doi: 10.1007/978-3-030-60990-0_12.

# APPENDICES

## CONTENTS

**A Related Works and Distinctions** . . . . . **16**

**B Discussion and Limitations** . . . . . **19**

**C A Catalog of (Deferred) Proofs** . . . . . **22**

**D Constants and Rates of the Blockwise Bound** . . . . . **24**

D.1 Average prediction error does not control the consistency error . . . . . 24

D.2 Sharpness of the blockwise constants . . . . . 24

D.3 The quadratic regime . . . . . 25

**E Response Models and Recovering Multiple Equilibria** . . . . . **25**

**F Experimental Setup** . . . . . **25**

**G Robustness and Scope** . . . . . **28**

G.1 Invariance across response families and coupling . . . . . 28

G.2 Calibration of the central-difference estimator . . . . . 28

G.3 Scope: Convexity failure and an incomparable condition . . . . . 30

## A Related Works and Distinctions

**The Case for Certification** No uncoupled dynamic converges to Nash equilibrium across a family of games containing a neighborhood of a given game (Hart & Mas-Colell, 2003), and their witness lies inside the hypotheses adopted here: two players on unit disks, costs smooth and strongly convex in each player's own action, a unique strict equilibrium. As stated in Hart & Mas-Colell (2003), converging to equilibrium is a more stringent requirement than being in equilibrium. Certificates address being in equilibrium.

**Learned response maps.** Response-based models of interaction replace a player's optimization with a map fitted to its observed actions. Scarabaggio et al. (2024) introduced the projected observer–head architecture, established via Brouwer's theorem that a fixed point exists, and showed that such a fixed point solves a variational inequality; uniqueness and convergence of the response iteration follow from an aggregate contraction condition $\left(\sum_i \gamma_i^2\right)^{1/2} < 1$, relaxed to a Krasnosel'skiĭ iteration at the nonexpansive boundary. Their guarantees concern the learned game. The distinction drawn in this paper is that Theorem 3.1 bounds exploitability in the game that generated the data, at a profile that need not be an exact fixed point, and needs no condition on $\gamma_i$. A subsequent treatment from the same group combines projected maps, existence, and aggregate contraction in the same way (Mignoni, 2025, Chapter 6). Their variational inequality is stated in the learned map and is equivalent to the fixed-point condition itself, so that it constrains the game that generated the data nowhere. Discarding the learned component, (Scarabaggio et al., 2026) characterizes stationary profiles of the true costs as fixed points of the blockwise projected gradient and establish existence and homotopy persistence by Lefschetz–Nielsen theory. We measure the blockwise residual $r_i$ of the map.

**A different $\varepsilon$.** In the purification of Harsanyi (1973) the parameter scales a perturbation of the payoffs and the equilibrium is exact in the perturbed game; in quantal response (McKelvey & Palfrey, 1995) and in perturbed best-response learning (Hofbauer & Hopkins, 2005) it scales a smoothing and the equilibrium is an exact fixed point of a *modified* operator, whose exploitability in the original game is in general bounded away from zero. The certificate here bounds regret in the unmodified game, so convergence to a perturbed equilibrium implies nothing about (2), and conversely.

**Decision-dependent and performative games.** When the data distribution reacts to the deployed model, the fixed point of repeated retraining becomes the solution concept. Perdomo et al. (2020) introduced performative prediction and the repeated-risk-minimization fixed point, and Narang et al. (2023) extended it to several players, proving existence, uniqueness, and a linear rate for performatively stable equilibria when a sensitivity modulus divided by a strong-convexity modulus falls below one. That ratio plays the role $q$ plays here, and the resulting sensitivity bound carries the same $1/(1-\cdot)$ constant that Remark B.3 dispenses with. Outside the contractive regime the dynamics are not merely unanalyzed but genuinely irregular: Piliouras & Yu (2023) exhibit chaotic multi-agent retraining, and Dong et al. (2023) study regions of attraction where contraction holds only locally. These results motivate a certificate that does not degrade as the modulus approaches one.

**Certified operators and Lipschitz networks.** A parallel literature constrains what a network can do rather than enveloping what it did. Monotone operator equilibrium networks obtain well-posed implicit layers by construction (Winston & Kolter, 2020), and Jafarpour et al. (2021) certify implicit networks through a non-Euclidean, weighted one-sided Lipschitz condition that is strictly weaker than a Euclidean contraction; Davydov et al. (2022) develop the contraction theory behind it. Lipschitz bounds are enforced architecturally by direct parameterization (Wang & Manchester, 2023) and unified across several such constructions by Havens et al. (2023), or estimated post hoc by semidefinite relaxation (Fazlyab et al., 2019), exact computation being NP-hard (Virmaux & Scaman, 2018). Two cautions from this literature shape our treatment. Products of layerwise spectral norms bound the joint operator norm only loosely, and the Clarke chain rule for a composition holds as an inclusion rather than an equality (Clarke, 1983), so a layerwise regularity certificate is unsound. Automatic differentiation compounds the problem: it returns an element of a conservative field rather than of the Clarke Jacobian, and the two differ on activation and projection boundaries (Bolte & Pauwels, 2021). We therefore report sampled Jacobian quantities as lower-bound diagnostics and never as certificates.

**Error bounds and gap functions.** The inequality behind Lemma 3.1 is part of the error-bound theory for finite-dimensional variational inequalities (Facchinei & Pang, 2003, Ch. 6)(Pang, 1997). Gap and merit functions convert a variational inequality into an optimization problem and supply computable residuals: the regularized gap of Fukushima (1992), the gap-function class of Larsson & Patriksson

(1994), the D-gap of Peng (1997), the global error bounds of Yamashita & Fukushima (1997), and the Nikaido–Isoda reformulations for generalized games of von Heusinger & Kanzow (2009). These results run predominantly from a residual to *distance-from-solution* and require strong monotonicity for the constant to exist; our use runs instead to *exploitability*, which is why compactness suffices. Sensitivity of solutions under perturbation of the defining map is classical, through Robinson's strong regularity (Robinson, 1980), its characterization on polyhedral sets (Dontchev & Rockafellar, 1996), and the bound of Dafermos (1988) in which the contraction modulus is replaced by a strong-monotonicity modulus of the environment.

**Certifying a limit vs. a profile.** Mertikopoulos & Zhou (2018) work in our setting, with compact convex product action sets and payoffs only $C^1$ in each player's own action. However, what they certify is a limit point of dual averaging which is Nash only if the iterates converge and the game is pseudo-concave. Their global results require that the Nash set be variationally stable, the pseudo-gradient be monotone, or the game to admit a concave potential. Their one quantitative gap estimate is an $\mathcal{O}(1/\sqrt{n})$ rate for an ergodic average that they note may behave differently from the played sequence. Variational stability is adoptable and would furnish isolation and asymptotics. However, it cannot furnish a certificaten since it relates $\Phi(x)$ to an unknown equilibrium. Thus, with that equilibrium unknown it yields no computable number, and verifying it requires the game to be Hessian at the point one is trying to locate. For $C^2$ costs it is moreover equivalent to local asymptotic stability (Mazumdar et al., 2020), so that it excludes precisely the equilibria that gradient-based learners avoid from almost every initialization — the profiles most in need of an external certificate.

**Continuous-time equilibrium seeking.** A parallel literature drives dynamics to an equilibrium rather than certifying a profile. The survey of Hu et al. (2022) collects the consensus-based, passivity-based and gradient-free designs, and Belgioioso et al. (2022) give the operator-splitting counterpart for coupled constraints. Two features bear on this work. Their convergence certificates are Lyapunov functions of the form $\frac{1}{2}\|x - x^\star\|^2$, evaluable only once $x^\star$ is known, which is unavailable when the game is not; and their hypotheses begin at monotonicity of $\Phi$ and strengthen from there. Gap and merit functions for variational inequalities are cited in neither survey, and no $\varepsilon$-equilibrium notion appears in either. The single quantitative rate in Belgioioso et al. (2022) is an ergodic $\mathcal{O}(1/k)$ bound on a fixed-point residual, and their operator-to-game link holds at an exact zero of that residual, so that a positive residual is converted into a statement about the game nowhere in this line.

**Identifying a game from observed play.** Whether observed actions pin down a game is the question our consistency error confronts. Gürkan & Pang (2009) give conditions under which equilibria of approximating games converge to equilibria of the true game, via a multi-epiconvergence notion; the statements are asymptotic and contain no explicit constant, and the authors note the connection between their regularized Nikaido–Isoda function and variational gap functions without developing it quantitatively. Bemporad & Tatarenko (2026a) learn solution maps of parametric generalized games by training against the Nikaido–Isoda gap rather than regressing on precomputed equilibria, which requires only best-response data, and Bemporad & Tatarenko (2026b) instead secure monotonicity of the learned game by construction. Bergerault et al. (2026) prove the sharpest negative result in this direction: characterizing the Nash gap at a fixed measure-matching error is hard, and exact matching of observed play can still leave a large Nash gap. Their positive results impose structure on the expert, such as dominant-strategy behavior or continuity of best responses around the expert, to transfer distributional imitation error into a Nash gap. Our assumptions constrain different objects: playerwise convexity constrains the true one-shot game, and a regional envelope directly controls the learned operator's discrepancy from its pseudo-gradient. Neither assumption set implies the other. This distinction is why Remark 2.1 measures operator consistency rather than distributional fit, and why the environment access of Section 4 is stated as a cost rather than assumed away. Payoff-based equilibrium learning supplies the access model we adopt, in which players evaluate costs but do not differentiate them (Tatarenko & Kamgarpour, 2024).

**Generative responders.** Our set-valued responders draw on conditional generative models, which represent multimodal conditional distributions rather than conditional means. Proposition 3.2 shows why that distinction is not cosmetic for equilibrium computation where playerwise convexity fails. Where it holds, our measurements in Section 5 attribute recovery of multiple equilibria to model capacity rather than to a multimodal likelihood; we therefore do not argue that generativity is required. We are not aware of prior work analyzing diffusion policies, flow matching, or vision–language–action models as response correspondences in a game, though we have not surveyed those literatures systematically and state this as an impression rather than a priority claim. Two scope limits

apply when the certificate is read alongside them. Theorem 3.1 constrains the *environment* through Assum. 2.2, so that continuity of the learned map is not by itself sufficient; and under playerwise convexity each best-response set is convex, so the multimodality that motivates a generative responder must come from multiple equilibria rather than from a multimodal best response. Certifying a genuinely set-valued response, where the residual is defined for a sampler rather than for a map, remains open, as does certifying a generalist policy deployed simultaneously as several players, whose playerwise responses share parameters.

## B Discussion and Limitations

**The regional audit has two distinct burdens.** First, Proposition 4.1 requires externally justified regularity and noise bounds; finite cost queries do not infer them. Without those inputs, Theorem 3.1 remains valid for an envelope established by another method, but Proposition 4.1 does not construct one. Second, even with valid inputs, a uniform audit requires a finite covering. For an $m$-dimensional box, its size grows as $O((\operatorname{diam}\Omega/\eta)^m)$. The construction is therefore limited to regions with small covering numbers, such as a segment or a low-dimensional deployment neighborhood with controlled geometry.

In the amortization study, the tanh-MLP responder uses a three-player quadratic game with two-dimensional actions, ten seeds, and 256 held-out profiles in each audited region. The segment is one-dimensional. The three-dimensional box is embedded in the six-dimensional joint action space by varying only its first three coordinates, with distances and the $\eta$-cover computed in the induced Euclidean geometry. At $\eta/\operatorname{diam}\Omega = 0.025$, the median covering sizes are 20 on the segment and 8,000 on the embedded box. E9 actually uses an exact gradient oracle: one query returns one player's own-gradient block, so a pointwise audit costs $N$ block queries and a regional audit costs $N|\mathcal{Z}|$. Under the scalar cost-oracle model of Section 4, central differences would instead cost $2d$ evaluations per point and $2d|\mathcal{Z}|$ per cover. Either accounting gives an oracle-call break-even of $|\mathcal{Z}|$ later profiles when the same stencil is used. This comparison is conditional on already having the regularity constants and excludes cover construction, storage, training, model evaluation, and any rollout replication used to produce a cost value. The ratio of the regional bound to the fresh pointwise bound is computed first as a median over held-out profiles and then as a median over seeds; it is 1.788 on the segment and 1.725 on the embedded box. The construction buys reuse without new environment queries; it does not beat a fresh query in tightness. Replacing the covering by samples provides an empirical diagnostic unless a separate distributional guarantee is proved.

**Coupled feasibility sets are excluded.** Assum. 2.1 requires a product action set, and the requirement is load-bearing twice: it makes the projection act blockwise, and it permits one player's action to be varied while the others are held fixed. Generalized games with shared constraints therefore fall outside every result here. Our coupled-constraint experiment shows what the certificate does when its own hypotheses are violated, and is not a claim of coverage.

**Playerwise convexity is necessary, and it makes the set-valued question unanswerable here.** Proposition 3.2 shows that the blockwise residual certifies nothing once Assum. 2.2 fails. The assumption also removes the motivation for generative responders: under playerwise convexity each best-response set is convex, so that responses of the bimodal kind in Proposition 3.2 are excluded by hypothesis. The multimodality that survives arises from multiple *equilibria* of a playerwise-convex game, which non-monotone coupling produces readily. Our evaluation on those games (Section 5) reports what a *conditional-mean* responder recovers, since the residual $\rho_i$ of (9) requires a single-valued evaluation and the generative families expose one only through their mean map. It therefore speaks to neither generativity nor capacity: the comparison is between mean maps, and the generative heads carry roughly four times the parameters and still do worse. What it does show is that a linear regressor fitted to play pooled across basins lands between them on a fraction $0.724$ of instances, while nonlinear responders do not. Whether a certificate can be given for a genuinely set-valued response we leave open, and the open half is the metric one rather than the topological one. Existence for a response correspondence with nonempty closed convex values is classical (Kakutani, 1941; Facchinei & Kanzow, 2010); what is missing is a *residual*, since no notion of approximate fixed-pointness of a correspondence appears in the equilibrium-seeking literature we surveyed, and $\rho_i$ as defined here requires a single-valued evaluation.

**The quadratic rate is not a computable certificate.** Proposition D.2 governs the interior regime but requires distances to an $\arg\min$, hence $n$ constrained convex solves in the unknown costs. We report it as a rate and measure it only where ground truth is available. A computable surrogate would need interiority certified from probe data together with a curvature modulus, which we have not established.

**Sampled Jacobians are diagnostics, not certificates.** Where the experiments report operator norms of learned maps, rigorous bounds are computed from weights and sampled quantities are labeled as lower-bound diagnostics. The distinction is not pedantic: the Clarke chain rule for a composition holds only as an inclusion (Clarke, 1983), and automatic differentiation returns an element of a

conservative field rather than of the Clarke Jacobian, the two differing on activation and projection boundaries (Bolte & Pauwels, 2021). No certificate in this paper reads a sampled Jacobian.

*Remark* B.1 (Why certify rather than solve). The same oracle that supplies an envelope would also drive projected gradient play, so that the certificate must earn its place against simply solving the game. It does so on three counts. (i) An envelope is paid once and then certifies every profile in $\Omega$ for the price of a forward pass, whereas solving pays along every trajectory and yields one profile. (ii) Projected gradient play carries no convergence guarantee at our hypotheses, and the failure arrives one rung earlier than the absence of monotonicity: under monotone $\Phi$ the Lyapunov derivative is still nonpositive, but the set on which it vanishes gives only $\langle \Phi(x) - \Phi(x^\star), x - x^\star \rangle = 0$, which does not force $x = x^\star$, so the invariance argument does not close (Hu et al., 2022). Below monotonicity the dynamics are irregular rather than merely unanalyzed, and multi-player retraining outside the contraction regime admits chaotic orbits (Piliouras & Yu, 2023). (iii) The learned map is often the deliverable for reasons unrelated to equilibrium computation, such as simulating counterfactual play, so the question is whether the model one already holds is trustworthy where it is evaluated.

*Remark* B.2 (What the certificate does not claim). The bound (12) concerns profiles in $\Omega$. It does not assert that $\mathrm{Fix}(F)$ is a singleton, nor that a certified profile approximates any particular equilibrium. For a nonexpansive $F$ the fixed-point set is nonempty, closed, and convex, but need not be a singleton, as $F = \mathrm{Id}$ shows. We therefore write *a* behavioral equilibrium throughout, never *the* behavioral equilibrium.

*Remark* B.3 (Behavior at the contraction boundary). Classical fixed-point perturbation controls equilibrium displacement by $\varsigma/(1-q)$, for a uniform map error $\varsigma$ and a map of contraction modulus $q < 1$ (Bauschke & Combettes, 2017), and that constant diverges as $q \to 1$. The certificate (12) contains no $q$. Certifying a nonexpansive or expansive learned map is thus a regime the bound already covers, which is the operative difference from the aggregate-contraction condition of Scarabaggio et al. (2024) and from the sensitivity-to-curvature ratio governing performatively stable equilibria in Narang et al. (2023).

*Remark* B.4 (Regularity of the learned map certifies nothing). Let $F \equiv \hat{x}$ be constant for an arbitrary $\hat{x} \in X$. That map is nonexpansive and $\rho_i(\hat{x}) = 0$ for every player, while $\varepsilon(\hat{x})$ is as large as the game permits. Its induced operator satisfies $G_i(\hat{x}) = 0$, so $\delta_i(\hat{x}) = M_i(\hat{x})$ and the residual entry of (12) reads $(\bar{M}_i + D_i)\bar{M}_i$, which exceeds the model-free entry $\bar{M}_i D_i$. The constant map is therefore excluded by the model-free bound rather than by the envelope, and the certificate reports nothing better than the trivial bound for it. Regularity of a learned map carries no information about the game.

*Remark* B.5 (Remark on Proposition 4.1.). A strict contraction $F$ with modulus $q < 1$ gives geometric convergence of the Picard iteration $x^{k+1} = F(x^k) = \Pi_X\big(x^k - g(h(x^k))\big)$. To reduce the fixed-point residual below $\epsilon$, the number of value queries scales as

$$\mathcal{O}\left(\frac{\log(1/\epsilon)}{1-q}\right)$$

when $q$ is close to one. Equivalently, if $q = 1 - \varepsilon$ with $0 < \varepsilon \ll 1$, the cost is $\mathcal{O}(\varepsilon^{-1}\log(1/\epsilon))$. At the nonexpansive boundary $q = 1$, this contraction argument disappears. General value-oracle lower bounds for Brouwer fixed-point computation are exponential in the dimension and in the number of digits of accuracy, under the corresponding Lipschitz assumptions (Hirsch et al., 1989); related reductions yield query lower bounds for equilibrium computation (Babichenko, 2016). These results provide complexity context, rather than a bound on the exploitability certificate (12): our result certifies a profile without assuming contraction, whereas Picard iteration seeks a fixed point under contraction.

*Remark* B.6 (Sampling changes the guarantee). Replacing the covering by independent draws from a distribution $\nu$ on $\Omega$ removes the covering cost but also removes deterministic uniform coverage. Under a fixed deployment distribution and exchangeability, a separate order-statistic or tolerance theorem can provide a fresh-draw or quantile guarantee; such a result still does not bound the regional supremum. Because we prove no such distributional theorem here, we treat sample maxima only as empirical diagnostics.

**The boundary-stencil count is not a measurement.** The fraction of coordinates requiring the one-sided stencil of Remark 4.1 is 0 *by construction* rather than by measurement: the probe used in our runs is an unconstrained central difference that never switches stencil, so the number could not have come out otherwise. The companion quantity we can measure is the fraction of probe steps that leave the action set, which is 0 at every radius on the grid. A feasibility-respecting probe would

therefore not have switched either, but that is a property of these arenas and not evidence about the remedy.

## C A CATALOG OF (DEFERRED) PROOFS

This appendix catalogs the proofs of the results of Section 3 and Section 4, in order.

*Proof of Lemma 3.1.* Fix a player $i$ and let $y_i^\star \in \arg\min_{y_i \in X_i} J_i(y_i, x_{-i})$, which exists because $X_i$ is compact by Assum. 2.1 and $J_i(\cdot, x_{-i})$ is continuous by Assum. 2.2. Convexity of $J_i(\cdot, x_{-i})$ implies that $J_i(y_i^\star, x_{-i}) \geq J_i(x) + \langle \Phi_i(x), y_i^\star - x_i \rangle$. Hence, player $i$'s regret obeys

$$J_i(x) - J_i(y_i^\star, x_{-i}) \;\leq\; \big\langle \Phi_i(x),\, x_i - y_i^\star \big\rangle. \tag{19}$$

Set $p_i = \Pi_{X_i}\big(x_i - \Phi_i(x)\big)$ and $w_i = x_i - p_i$, so that $\|w_i\| = r_i(x)$ by (7). Projecting $x_i - \Phi_i(x)$ onto $X_i$ implies $\langle x_i - \Phi_i(x) - p_i,\, y_i - p_i \rangle \leq 0$ for every $y_i \in X_i$, the variational characterization of the projection (Facchinei & Pang, 2003, §1.5); equivalently, $p_i$ solves $\mathrm{VI}\big(X_i,\, u \mapsto u - (x_i - \Phi_i(x))\big)$ in the sense of Definition 2.1. It is easy to verify that $\langle \Phi_i(x),\, p_i - y_i \rangle \leq \langle w_i,\, p_i - y_i \rangle$; applying the Cauchy–Schwarz inequality with $\|p_i - y_i\| \leq D_i := \operatorname{diam} X_i$, since $p_i, y_i \in X_i$, gives

$$\langle \Phi_i(x),\, p_i - y_i \rangle \;\leq\; r_i(x)\, D_i \qquad \text{for every } y_i \in X_i. \tag{20}$$

Splitting (19) at $p_i$ as $\langle \Phi_i(x), x_i - y_i^\star \rangle = \langle \Phi_i(x), w_i \rangle + \langle \Phi_i(x), p_i - y_i^\star \rangle$, then bounding the first term by $M_i(x) r_i(x)$ with the Cauchy–Schwarz inequality and the second by (20), we find that the bound on the player $i$'s regret is $\big(M_i(x) + D_i\big) r_i(x)$, which is the first entry of (8). Separately, (19) and Cauchy–Schwarz give $J_i(x) - J_i(y_i^\star, x_{-i}) \leq M_i(x) \|x_i - y_i^\star\| \leq M_i(x) D_i$. Both bounds apply to the same player, hence player $i$'s regret is at most their minimum, and (8) follows by taking the maximum over players. Writing $a_i = \big(M_i(x) + D_i\big) r_i(x)$ and $b_i = M_i(x) D_i$, the inequality $\max_i \min\{a_i, b_i\} \leq \min\{\max_i a_i, \max_i b_i\}$ shows that taking the minimum after the maximum is valid but never tighter, and strict exactly when no single player has both $a_i \geq m$ and $b_i \geq m$ for $m = \min\{\max_j a_j, \max_j b_j\}$. □

*Proof of Lemma 3.2.* Since $x_i - G_i(x) = f_i(x_{-i}) \in X_i$, projection fixes this point. Inserting it into $r_i(x) = \|x_i - \Pi_{X_i}(x_i - \Phi_i(x))\|$ and using the triangle inequality gives $r_i(x) \leq \rho_i(x) + \|\Pi_{X_i}(x_i - G_i(x)) - \Pi_{X_i}(x_i - \Phi_i(x))\|$. Nonexpansiveness of $\Pi_{X_i}$ bounds the second term by $\|G_i(x) - \Phi_i(x)\| = \delta_i(x)$, proving the claim. □

*Proof of Proposition 3.1.* The reverse triangle inequality and $\rho_i(\hat{x}) = \|G_i(\hat{x})\|$ give $\rho_i(\hat{x}) + \delta_i(\hat{x}) \geq \|\Phi_i(\hat{x})\| = M_i(\hat{x})$. Multiplying by $M_i(\hat{x}) + D_i \geq 0$ yields the first inequality of (10), and the second is strict whenever $M_i(\hat{x}) > 0$, since then $M_i(\hat{x})^2 > 0$. □

*Proof of Theorem 3.1.* Fix $\hat{x} \in \Omega$. By (11), $M_i(\hat{x}) \leq \bar{M}_i$ and $\delta_i(\hat{x}) \leq \bar{\delta}_i$, and Lemma 3.2 gives $r_i(\hat{x}) \leq \rho_i(\hat{x}) + \bar{\delta}_i$. Both entries of (8) are nondecreasing in $M_i$ and in $r_i$ on the nonnegative orthant $\mathbb{R}_+$ since $D_i \geq 0$. Hence, substituting the enveloped upper bounds preserves each inequality. Applying Lemma 3.1 playerwise gives (12). The envelope is fixed before $\hat{x}$ is chosen, so the conclusion holds simultaneously for every $\hat{x} \in \Omega$. □

*Proof of Corollary 3.1.* Dividing $(\bar{M}_i + D_i)(\rho_i + \bar{\delta}_i) < \bar{M}_i D_i$ by it gives the first inequality of (13). For the second, $\bar{M}_i D_i / (\bar{M}_i + D_i) \leq \bar{M}_i$ because $D_i \leq \bar{M}_i + D_i$, and $\bar{M}_i D_i / (\bar{M}_i + D_i) \leq D_i$ because $\bar{M}_i \leq \bar{M}_i + D_i$; hence it is at most $\min\{\bar{M}_i, D_i\}$. □

*Proof of Proposition 3.2.* Differentiating (14) gives $\Phi_i = x_i\big(x_i^2 - w_i(x_{-i})\big)$, which vanishes at $x = 0$ for every $x_{-i}$. Since $0 \in X_i = [-c, c]$, the projection fixes it, so $r_i(0) = \|\Pi_{X_i}(0)\| = 0$ for every player. Each $J_i(\cdot, x_{-i})$ is even in $x_i$ with minimizers $\pm\sqrt{w_i(x_{-i})}$ and minimum value $-\frac{1}{4} w_i(x_{-i})^2$, while $J_i(0, x_{-i}) = 0$. At the origin $w_i(0) = 1$, hence every player's regret is $\frac{1}{4}$. The condition on $c$ places $\pm\sqrt{w_i}$ inside $X_i$ for every $x_{-i} \in X_{-i}$. □

*Proof of Proposition 4.1.* Fix a player $i$, a covering point $z \in \mathcal{Z}$ and a coordinate $k$ of player $i$'s block, and write $\varphi(t) = J_i(z + t e_{i,k})$, so that $\varphi'(0) = [\Phi_i(z)]_k$ is the quantity (15) estimates. The assumed extension makes $\varphi$ three times continuously differentiable on $[-\tau, \tau]$, hence Taylor's

theorem with Lagrange remainder gives $\varphi(\pm\tau) = \varphi(0) \pm \tau\varphi'(0) + \frac{1}{2}\tau^2\varphi''(0) \pm \frac{1}{6}\tau^3\varphi'''(\xi_\pm)$ for some $|\xi_\pm| \le \tau$. Subtracting the two expansions cancels the even orders, so that

$$\frac{\varphi(\tau) - \varphi(-\tau)}{2\tau} \; - \; \varphi'(0) \; = \; \frac{\tau^2}{12}\big(\varphi'''(\xi_+) + \varphi'''(\xi_-)\big). \tag{21}$$

Since $\varphi''' = \partial^3 J_i / \partial x_{i,k}^3$ is bounded by $B_3$ on the probe segment, the right side of (21) is at most $\frac{1}{6}B_3\tau^2$ in magnitude. Stacking the $n_i$ coordinates, player $i$'s bias vector has norm at most $\frac{1}{6}B_3\tau^2\sqrt{n_i}$.

For the noise, a difference of two independent $\sigma$-sub-Gaussian variables is $\sqrt{2}\sigma$-sub-Gaussian, so dividing by $2\tau$ makes each difference quotient $s$-sub-Gaussian about its mean with $s = \sigma/(\sqrt{2}\tau)$. There are $d|\mathcal{Z}|$ scalar estimates in all, one per coordinate and covering point, and each has two-sided tail $2\exp(-t^2/2s^2)$. Equating $2d|\mathcal{Z}|\exp(-t^2/2s^2)$ with $\alpha$ and solving for $t$, a union bound gives that every coordinatewise deviation is at most $s\sqrt{2\log(2d|\mathcal{Z}|/\alpha)} = \frac{\sigma}{\tau}\sqrt{\log(2d|\mathcal{Z}|/\alpha)}$ simultaneously, with probability at least $1-\alpha$; the $\sqrt{2}$ in $s$ cancels the one from the tail. On that event player $i$'s noise vector has norm at most $\frac{\sigma}{\tau}\sqrt{n_i\log(2d|\mathcal{Z}|/\alpha)}$.

Hence, on the same event and for every $z \in \mathcal{Z}$ at once, the triangle inequality bounds $\delta_i(z) = \|G_i(z) - \Phi_i(z)\|$ by $\|G_i(z) - \widehat{\Phi}_{\tau,i}(z)\| + E_i(\tau,\alpha)$ and bounds $M_i(z)$ by $\|\widehat{\Phi}_{\tau,i}(z)\| + E_i(\tau,\alpha)$. Finally, given $x \in \Omega$, choose $z \in \mathcal{Z}$ with $\|x - z\| \le \eta$. Lipschitzness gives $\delta_i(x) \le \delta_i(z) + L_{\delta,i}\eta$ and $M_i(x) \le M_i(z) + L_{M,i}\eta$. Taking the two maxima over the covering yields (17). The coordinatewise union bound already ranges over all $d$ coordinates and all covering points, so both envelope inequalities hold for every player on this one event. □

# D CONSTANTS AND RATES OF THE BLOCKWISE BOUND

Three analytic supplements to Lemma 3.1 and Theorem 3.1: what an average prediction error fails to control, whether the coefficients can be shrunk, and when the rate in $\rho_i + \delta_i$ improves from linear to quadratic.

## D.1 AVERAGE PREDICTION ERROR DOES NOT CONTROL THE CONSISTENCY ERROR

Remark 2.1 asserts that a small training loss leaves the operator-consistency error of Lemma 3.2 uncontrolled. The following one-player construction makes the failure explicit and elementary, and shows that the obstruction is a gap between an average and a supremum rather than anything about multi-player coupling.

**Proposition D.1** (Small mean-squared error, displaced fixed point)**.** *Let $X = [0, 1]$, let the reference response be $F_0 \equiv 0$, and for $0 < h < \frac{1}{4}$ let*

$$T_h(x) = \tfrac{1}{2} \max\big\{1 - |x - \tfrac{1}{2}|/h,\ 0\big\}. \tag{22}$$

*Then under observations uniform on $X$, $\mathbb{E}\big|T_h(x) - F_0(x)\big|^2 = h/6 \to 0$ as $h \to 0$, while $\sup_{x \in X} \big|T_h(x) - F_0(x)\big| = \frac{1}{2}$ for every $h$. The map $T_h$ retains the reference fixed point at $0$ and acquires a second fixed point at $\frac{1}{2}$, whose fixed-point residual is zero and whose distance from the reference fixed point is $\frac{1}{2}$.*

*Proof.* The triangle $T_h$ is supported on $[\frac{1}{2} - h, \frac{1}{2} + h]$ with peak $\frac{1}{2}$ at $x = \frac{1}{2}$, so its supremum is $\frac{1}{2}$ and $\int_0^1 T_h^2 = h/6$ by direct integration of the squared triangular profile. Evaluating (22) gives $T_h(0) = 0$ and $T_h(\frac{1}{2}) = \frac{1}{2}$, hence both points are fixed. The residual at $\frac{1}{2}$ vanishes because $\frac{1}{2}$ is a fixed point. □

To read this as a statement about the consistency error of Lemma 3.2, use a two-player decoupled game on $X_1 = X_2 = [0, 1]$ with $J_i(x) = \frac{1}{2}x_i^2$, hence $\Phi_i(x) = x_i$. Define the admissible response map $F(x_1, x_2) = (T_h(x_2), T_h(x_1))$. Then $G_i(x) = x_i - T_h(x_{-i})$ and $\delta_i(x) = T_h(x_{-i})$, so $\sup_X \delta_i = \frac{1}{2}$ while its squared expectation under the uniform distribution on $X$ is $h/6$. Both $(0, 0)$ and $(\frac{1}{2}, \frac{1}{2})$ are fixed points. On the full region $X$, $\bar{M}_i = D_i = 1$ and $\bar{\delta}_i = \frac{1}{2}$, so at the displaced fixed point the two certificate entries tie at 1. The bound itself still holds, since each player's regret there is $\frac{1}{8}$; what a small average error is compatible with is a displaced fixed point together with a regional learned-residual entry that improves on nothing.

Three readings follow. An average prediction error of $h/6$ is compatible with a uniform map error of $\frac{1}{2}$, so that no bound on the former controls the latter without a uniform regularity hypothesis; the spurious fixed point at $\frac{1}{2}$ has zero residual, which is Remark B.4 in miniature and for a different reason; and the construction does not contradict any contraction result, since $T_h$ has slope magnitude $1/(2h) > 1$ on its support. What is required to connect an average loss to the consistency error is a uniform statement, which is what the regional envelope of Proposition 4.1 supplies by probing a finite covering rather than by appealing to a training loss.

## D.2 SHARPNESS OF THE BLOCKWISE CONSTANTS

Neither coefficient in (8) can be rescaled downward. One limitation bounds what follows and is worth stating before it rather than after: the family below has $\varepsilon = MD$ at every parameter value, so the model-free entry binds throughout and the residual entry never does. What is established is therefore sharpness of the coefficients at profiles where a practitioner would report the model-free bound instead.

Take $n = 1$, $X = [0, 1]$, $J(x) = ax$ and evaluate at $x = 1$, so that $M = a$, $D = 1$ and $r = \min\{1, a\}$. The attainment ratio $\varepsilon/[(M + D)r]$ equals $1/(1 + a)$ for $a \le 1$, which tends to one as $a \to 0$ and so forbids replacing $D$ by $cD$ with $c < 1$; and it equals $a/(1 + a)$ for $a \ge 1$, which tends to one as $a \to \infty$ and so forbids replacing $M$ by $cM$. The argument excludes rescaling either coefficient and not a differently shaped constant.

### D.3 THE QUADRATIC REGIME

Whether the rate in (12) is linear or quadratic turns on whether best responses lie in the relative interior of the action sets. Write $\mathrm{BR}_i(x_{-i}) = \arg\min_{y_i \in X_i} J_i(y_i, x_{-i})$.

**Proposition D.2** (Quadratic certificate under interior best responses)**.** *Let Assum. 2.1 hold, let $J_i(\cdot, x_{-i})$ be differentiable with $L_i$-Lipschitz gradient $\Phi_i(\cdot, x_{-i})$ on $X_i$, and let some minimizer $b_i \in \mathrm{BR}_i(\hat{x}_{-i})$ lie in the relative interior of $X_i$ for every player $i$. With $\rho_i = \rho_i(\hat{x})$ and $\delta_i^{\mathrm{BR}} \triangleq \|g_i(h_i(\hat{x}_{-i})) - b_i\|$,*

$$\varepsilon(\hat{x}) \;\le\; \max_{i \in \mathcal{N}} \tfrac{L_i}{2} \left(\rho_i + \delta_i^{\mathrm{BR}}\right)^2. \tag{23}$$

*Proof.* Fix a player $i$. Since $b_i \in X_i$ we have $\Pi_{X_i} b_i = b_i$, hence nonexpansiveness of $\Pi_{X_i}$ gives $\|\hat{x}_i - b_i\| \le \rho_i + \delta_i^{\mathrm{BR}}$ as in Lemma 3.2. Minimality of $b_i$ gives $\langle \Phi_i(b_i, \hat{x}_{-i}), y_i - b_i \rangle \ge 0$ for every $y_i \in X_i$. Because $b_i$ lies in the relative interior of $X_i$, for each such $y_i$ the point $b_i - t(y_i - b_i)$ also lies in $X_i$ for all small $t > 0$, which supplies the reverse inequality; hence $\langle \Phi_i(b_i, \hat{x}_{-i}), y_i - b_i \rangle = 0$ for every $y_i \in X_i$, and in particular at $y_i = \hat{x}_i$. The descent lemma along the segment $[b_i, \hat{x}_i] \subseteq X_i$ then bounds player $i$'s regret by $\frac{L_i}{2}\|\hat{x}_i - b_i\|^2$. Taking the maximum over players gives (23). □

Two scope statements belong with Proposition D.2. Relative interiority, rather than topological interiority, is the sharp hypothesis: at a relative-interior point of a lower-dimensional $X_i$ the gradient need not vanish, but its inner product against every feasible direction does, which is all the proof uses. Playerwise convexity is not needed, hence (23) holds outside Assum. 2.2. What the refinement does require is a strictly stronger oracle than the envelope of Definition 3.1: $\delta_i^{\mathrm{BR}}$ is a distance to an $\arg\min$, hence obtaining it means solving $n$ constrained convex programs in the unknown costs. We therefore report (23) as the rate that governs the interior regime, not as a computable certificate, and Section 5 measures it only where ground truth is available.

## E RESPONSE MODELS AND RECOVERING MULTIPLE EQUILIBRIA

**Multiple equilibria and what recovers them.** Non-monotone coupling makes multiple equilibria the common case even under Assum. 2.2: of 120 playerwise-convex instances, 113 carry at least two distinct Nash equilibria, verified by active-set enumeration to a worst residual of $1.42 \times 10^{-14}$. A linear regressor fitted to play pooled across both basins lands between them rather than at either, recovering both equilibria on a fraction $0.431$ of instances. A unimodal nonlinear network recovers both on $0.862$, against $0.224$ for flow matching and $0.0517$ for diffusion.

These numbers do not separate capacity from generativity, and we do not read them as doing so. Every family is evaluated through a single-valued map, which for the two generative families is a conditional mean rather than a sample, so the comparison is between mean maps and the question of set-valuedness is not posed. The parameter counts run the other way besides: the generative heads carry roughly four times the parameters per player of the unimodal network and still recover fewer equilibria. What the measurement does establish is narrower and still useful: a linear responder fitted across basins lands between them, and sufficient capacity avoids that failure. We record it because it severs any empirical benefit claim from the observation that set-valued responses sit inside the hypotheses of Theorem 3.1; the latter is a scope statement, not a performance one.

**Set-valuedness where convexity fails.** On the nonconvex witness of Proposition 3.2 the conditional-mean profile has exploitability $0.25$ analytically while its variational residual is $0$, so the residual alone certifies nothing there. Mode-selected samples reduce exploitability from $0.256$ to $0.0286$ for diffusion ($p_{\mathrm{Holm}} = 1.91 \times 10^{-6}$) and from $0.252$ to $0.0175$ for flow matching ($p_{\mathrm{Holm}} = 1.91 \times 10^{-6}$), over 20 paired seeds. The certificate's hypotheses fail on this game and we never evaluate the bound as a claim about it.

## F EXPERIMENTAL SETUP

The studies named in Section 5 correspond to the modules of the released package as follows: the *tightness study* to `e1_tightness`, the *coupling sweep* to `e2_coupling`, the *dissociation study*

to `e3_incomparability`, the *set-valued witness* to `e4_set_valued`, the *multiplicity study* to `e5_multiplicity`, the *solver comparison* to `e6_solvers`, the *probe study* to `e7_probe`, the *scaling study* to `e8_scaling`, and the *amortization study* to `e9_amortization`.

Each game is chosen so that it isolates one hypothesis of Theorem 3.1 and supplies either analytic ground truth or a documented numerical oracle, so that a discrepancy is attributable to the bound rather than to an undocumented solver. We describe the five in turn, in each case stating what makes the instance hard and which claim it exercises.

**Coupled quadratic: monotonicity on a knob.** Set $Q_i = I$ and $A = I + \kappa(W + \frac{1}{2}Z)$ with $W$ symmetric and $Z$ antisymmetric, both with zero diagonal blocks, $W$ rescaled so that $\lambda_{\min}(W) = -1$ and $Z$ so that $\|Z\|_2 = 1$. Then

$$\tfrac{1}{2}\big(A + A^\top\big) = I + \kappa W, \qquad \text{hence} \qquad \lambda_{\min}\big(\tfrac{1}{2}(A + A^\top)\big) = 1 - \kappa, \tag{24}$$

so the monotonicity modulus is known analytically and the sweep crosses monotonicity exactly at $\kappa = 1$. The skew part keeps $A \neq A^\top$, so that the pseudo-gradient is not the gradient of any potential and the instance is not a potential game. This is the only family in which instances can be placed arbitrarily far on either side of the monotone boundary with the modulus known in closed form, which is what lets the coupling sweep attribute an outcome to non-monotonicity rather than to conditioning. It carries the claim that the certificate requires no monotonicity.

**Capacitated Cournot: an active projection.** Firms choose scalar quantities on $[0, \mathrm{cap}_i]$ with capacities alternating between $0.6$ and $5.0$, so that the tight capacities bind at equilibrium. The projection in $r_i$ is therefore active, which is exactly the regime Proposition 3.1 excludes by hypothesis. The game is monotone and unremarkable as economics; its role is to show that the interiority hypothesis in Proposition 3.1 is load-bearing rather than decorative, since on this game the learned map *can* sharpen a pointwise certificate.

**Rosen shared constraint: outside the product assumption.** Boxes intersected with a single coupled row $w^\top x \leq h$, so that player $i$ optimizes over $\mathcal{X}_i(x_{-i}) = \{x_i \in [\mathrm{lo}_i, \mathrm{hi}_i] : w_i^\top x_i \leq h - \sum_{j \neq i} w_j^\top x_j\}$. This violates the product requirement of Assum. 2.1, so every result here is formally silent about it. We include it to record what the certificate does when its own hypothesis fails, and we report those cells as a stress diagnostic and never as coverage. Two consequences are worth naming: the best response is a projection onto a box intersected with a halfspace, which is computed by exact bisection on the single multiplier and so needs no inner solver; and $D_i$ is conservative here, since it is the diameter of the box rather than of the coupled set.

**Bimodal coordination: convexity removed.** With $J_i$ and $w_i$ as in (14), $\partial^2 J_i / \partial x_i^2(0) = -w_i < 0$, so playerwise convexity fails. All $2^n$ sign patterns of $s = (1 - \kappa)^{-1/2}$ are equilibria in closed form, while the origin is a stationary point of $\Phi$ at which the variational residual vanishes and the exploitability is exactly $\frac{1}{4}$. This is the witness of Proposition 3.2. It tests the *necessity* of Assum. 2.2 and not the validity of the bound, and we never evaluate (12) as a claim about it.

**Nonexpansive boundary: monotone yet not contractive.** Let $J_i(x) = \frac{1}{2}(x_i - \sigma_i x_{i+1})^2$ cyclically with $\sigma_i = +1$ for $i < N - 1$ and $\sigma_{N-1} = -1$. Then $F(x) = Px$ with $P$ a signed cyclic permutation, so that $P^\top P = I$ and the best-response modulus is exactly one: nonexpansive, and not a contraction. The equilibrium is nevertheless unique and interior at the origin, because $P^N = (\prod_i \sigma_i) I = -I$ makes $I - P$ invertible. Picard iteration is $2N$-periodic and therefore never converges from any $x^0 \neq 0$, its residual staying exactly constant. This is the sharpest separation we can construct between monotonicity of the variational inequality and convergence of naive fixed-point iteration: the game is strongly monotone with a unique equilibrium, and the iteration still fails. It is the regime in which contraction-based guarantees say nothing while Theorem 3.1 still applies, and it is where the solver comparison is run.

**Ground truth and evaluation protocol.** Every game supplies a closed-form or verified-global best response, so that no inner minimization is approximated and the exploitability against which each bound is checked is exact rather than a solver output. Equilibria are verified to a residual below $10^{-10}$ when an equilibrium is required: the capacitated Cournot equilibrium by active-set enumeration, the

coupled-constraint variational equilibrium by projected extragradient to its documented tolerance, and the bimodal game's $2^n$ equilibria in closed form.

The nine experiments use the following fixed protocol. E1 tests bound validity and tightness on the three games above (420 cells, of which 280 satisfy the theorem assumptions); E2 sweeps eight coupling values on the product quadratic game (1,120 cells); E3 pools only the five families with rigorous Lipschitz bounds to compare joint and small-gain contraction certificates; E4 tests the nonconvex set-valued witness; E5 enumerates equilibria and compares responder families; E6 compares Picard, Krasnosel'ski–Mann, and Halpern on the analytic nonexpansive map; E7 varies finite-difference radius and oracle noise; E8 varies player count at a fixed observational-data budget; and E9 varies covering resolution and region dimension. Unless stated otherwise, E1–E2 use seeds 0–19, seven response families, 200 training epochs, and 240 observational transitions per seed. E3–E9 use the seed counts, tolerances, held-out profile counts, and correction procedures reported with their respective results files. All comparisons pair the same seed, game, family, and coupling wherever those factors are present; confidence intervals use $10{,}000$ paired bootstrap resamples, and Wilcoxon tests use Holm correction within the declared experiment family.

**Response families.** Seven response models receive the concatenated rival profile $x_{-i}$ and are fitted on observational play under a shared budget, so that no family is advantaged by a longer fit: a least-squares affine map, two-layer 32–32 tanh and ReLU networks, a spectral-normalized network, an Almost-Orthogonal-Layer (AOL) Lipschitz-certified network (Prach & Lampert, 2022), a conditional diffusion sampler, and conditional flow matching. The first five expose a rigorous Lipschitz bound computed from weights; the two samplers expose none, so that their certificate status is unknown and is recorded as not-certified rather than as passing. Capacity is not matched across the two groups. The deterministic heads use two hidden layers of width 32 and the generative ones two of width $64$ together with a time embedding, so that the generative group carries roughly four times the parameters per player — a mismatch that runs against, rather than for, any reading in which capacity explains the outcomes of Appendix E. All neural responders use Adam with batch size $128$ for $200$ epochs on the same $240$ noisy observational transitions per seed; the learning rate is $3 \times 10^{-3}$ except for AOL, which uses $5 \times 10^{-3}$. Affine and feedforward responders minimize mean-squared prediction error, diffusion uses denoising-score matching, and flow matching uses its conditional flow objective.

Table 1: **Certificate divided by true exploitability as players are added.** Coupled quadratic game, scalar actions, medians over 20 seeds; lower is tighter and one is exact. The aggregated column is the single-constant form $(M + D)(\rho + \delta)$, the blockwise column is Theorem 3.1. The aggregated form grows with $n$ because exploitability is a maximum over players whereas an aggregated residual is a joint norm. In this game the blockwise column coincides with the model-free bound $\max_i \bar{M}_i D_i$, hence it reports tightness of the model-free entry, not of the residual entry.

| players $n$ | 2 | 8 | 64 | 128 | 256 |
|---|---|---|---|---|---|
| aggregated | 38.15 | 84.85 | 431.1 | 696.1 | 1157 |
| blockwise | 13.92 | 10.78 | 7.862 | 7.176 | 6.628 |





Fig. 4: **Tightness by response family.** Certified bound over true exploitability, dots seeds and bars medians, log scale; panels are the quadratic, Cournot and shared-constraint games. Blue marks families with a rigorous Lipschitz bound, orange those without.

## G Robustness and Scope

Five figures, grouped evidence are reported here to support our main results. Fig. 4 and Fig. 5 test whether the pointwise result survives a change of architecture and game parameter. Fig. 6 calibrates the estimator supplying one term of the envelope. Fig. 7 and Fig. 8 bound the scope: the first exhibits a hypothesis failing, the second a familiar sufficient condition.

### G.1 Invariance across response families and coupling

Fig. 4 plots the certified bound of (12) over the true exploitability, one dot per seed and one bar per median, for seven response families on three games. The families sit close together: the widest within-game spread is a factor of 4.04. The reported certificate equals its model-free entry $\max_i \bar{M}_i D_i$ in all 280 product-set cells and in the 140 shared-constraint diagnostic cells; the latter are not theorem coverage. The measured fraction is 1. The plotted ratio therefore carries no learned-map contribution; families differ only through $\bar{M}_i$, which records where each fixed point lands.

Fig. 5 sweeps the coupling parameter, which fixes $\lambda_{\min}$ of the symmetrized pseudo-gradient Jacobian exactly, so that the grid crosses monotonicity at one. Validity is invariant: the bound holds in 1120 of 1120 cells across the grid, the non-monotone half included, which is the content of requiring no monotonicity. Tightness is not: its median falls monotonically from $3.01e+03$ to $9.86$. The sweep establishes robustness of validity and nothing about the quality of the bound. The bound holds at every coupling; the median tightness ratio, on the right axis, falls monotonically by a factor of about $300$ across the grid, so what the sweep shows to be stable is validity and not tightness. Certified fractions are shown for the joint-spectral and small-gain conditions.

### G.2 Calibration of the central-difference estimator

Fig. 6 reports the central-difference error in estimating $\Phi_i$ against the probe radius $\tau$ at five noise scales, with medians over twelve probe points at each radius and, at each nonzero scale, twenty-four noise replicates. Panel (a) has $B_3 > 0$ and exhibits the interior minimum that the bias–noise balance of (17) predicts; panel (b) sets $B_3 = 0$ exactly, where no interior minimum can arise and none does. Dotted verticals mark the balance radius that Proposition 4.1 predicts, one per noise scale. The observed minimizer over that radius is $2.14$, against a grid of multiplicative step $2.15$. The predicted





Fig. 5: **Coupling sweep across the monotone boundary.** Coupled quadratic game; the coupling parameter sets $\lambda_{\min}$ of the symmetrized pseudo-gradient Jacobian exactly, so the grid crosses monotonicity at one.





Fig. 6: **Probe error against probe radius.** Central-difference error in estimating $\Phi_i$ versus $\tau$, at several noise scales $\sigma$; medians over twelve probe points and twenty-four noise replicates each.

scaling in $\sigma$ is confirmed to within one grid step; the prefactor is not resolved, the grid being coarser than the discrepancy under test. Panel (a) is the multimodal arena, where $B_3 > 0$ and the predicted U-shape appears: bias of order $B_3\tau^2$ dominates at large $\tau$ and noise of order $\sigma/\tau$ at small, with the minimum near the balance point of Proposition 4.1. Panel (b) is a control with $B_3 = 0$ exactly, in which no interior minimum can occur and none does. The curves are diagnostics of the estimator, not certificates.





Fig. 7: **A vanishing residual is no evidence of a small gap.** Exploitability against variational residual on the bimodal witness of Proposition 3.2, log ordinate, 20 seeds per family. Filled markers are the conditional-mean fixed points, hollow markers the four mode-selected profiles per seed, and the star is the analytic witness, at residual exactly zero and exploitability $1/4$. Mode selection moves the cloud down and to the right of the conditional-mean profiles it improves on, so that a smaller gap is bought at a larger residual. Within the mode-selected cloud the two quantities are positively associated; it is the shift between the groups that the plane shows.

### G.3 Scope: Convexity failure and an incomparable condition

Fig. 7 compares the exploitability reached from a conditional-mean response with that reached from mode-selected samples, on the bimodal witness of Proposition 3.2, over 20 paired seeds.The structure of interest is at the origin, where the variational residual vanishes analytically while the exploitability is $0.25$. The learned conditional-mean fixed points land beside it, at $0.256$ and $0.252$, with measured residual $0.1834$ rather than the analytic zero, so that the degeneracy is approached and not attained. Mode selection lowers exploitability to $0.0286$ and $0.0175$ without repairing the hypothesis: recovery of the game's 4 equilibria is bimodal across runs, with more than half recovering none, so that no run is typical. Fig. 7 places all of this in the residual–exploitability plane, and the ordering there is the point. The witness sits at residual zero with exploitability $0.25$, while the mode-selected profiles reach those smaller gaps carrying a median residual of $0.3721$, larger than the $0.1834$ of the conditional-mean profiles they improve on. The comparison is paired and between groups rather than monotone: a fraction $0.825$ of mode-selected profiles carry both a smaller exploitability and a larger residual than the conditional-mean profile of the same seed, so that a smaller gap is bought at a larger residual. Within the mode-selected cloud the residual and the gap remain positively associated, so that the residual does not even order the profiles it fails to control.

Fig. 8 cross-classifies the joint-spectral against the small-gain condition over $1100$ certifiable cells pooled from the two sweeps: $106$ certified by both, $4$ by small gain alone, $0$ by the joint condition alone, and $990$ by neither. The empty cell is a fact about these fits rather than a theorem, since $\rho(\Gamma) \leq \|J_F\|_2$ held in all $1100$ of them with median slack $0.03969$, so that the direction separating the two conditions analytically is realized nowhere in this population. All $106$ doubly certified cells are affine. Reporting the joint-spectral condition alone would understate coverage on fits of this kind, which is not a dominance claim: both separating directions are realizable, and the two analytic witnesses accompanying the contingency distinguish an empty cell from an impossible one.





Fig. 8: **Joint-spectral against small-gain certification on trained models.** Contingency counts pooled over games and seeds, over 1100 certifiable trained models: 106 certified by both, 4 by small gain alone, 0 by the joint bound alone, and 990 by neither. The empty cell is a fact about these fits and not a dominance relation: $\rho(\Gamma) \leq \|J_F\|_2$ held in all 1100 of them, with a median gap of 0.040, so the skew-coupling direction that separates the two conditions analytically does not arise here at any threshold. This licenses no claim that either certificate dominates.