

Rigorous bounds on Lyapunov exponents of linked twist maps

Patrick Wright, Jitse Niesen and Rob Sturman

School of Mathematics, University of Leeds, Leeds, LS2 9JT, UK

E-mail: `r.sturman@leeds.ac.uk`

Abstract. Rigorous, elementary upper and lower bounds upon the Lyapunov exponents of a parametrised family of linked twist maps are given, and obtained explicitly for a specific range of parameter values. The method used to obtain the bounds utilises the existence of invariant cones for specific products of the underlying family of shear maps, and the return time partition of the overlap region of the two annuli. Improvements upon the accuracy of this method are then obtained by considering preceding sequences of matrices on the orbits.

1. Introduction

A fundamental concept in dynamical systems is stability, typically measured by the rate of growth of a quantity with time. Let $f : M \rightarrow M$ be a diffeomorphism of a compact manifold M , $x \in M$, $v \in T_x M$. Then the rate of expansion or contraction of infinitesimal perturbations in tangent space is given by

$$\lambda(x, v) = \lim_{n \rightarrow \infty} \frac{1}{n} \log \|D_x f^n v_0\| \quad (1)$$

whenever the limit exists, and where $D_x f^n$ is the Jacobian of f^n at x , given by the product

$$D_x f^n = D_{f^{n-1}(x)} f D_{f^{n-2}(x)} f \dots D_{f(x)} f D_x f.$$

It is well known that if f admits an invariant measure μ , the limit in (1) does indeed exist for μ -almost all x by the Oseledec multiplicative ergodic theorem [1]. Furthermore, if μ is ergodic, then $\lambda(x, v)$ takes on only a finite number of possible values, called Lyapunov exponents.

However, it is only in relatively rare instances that a Lyapunov exponent might be explicitly and analytically computed. When we assume μ to be ergodic, it appears that the Birkhoff ergodic theorem may be of use.

Theorem 1 (Birkhoff ergodic theorem [2]). *Let $f : M \rightarrow M$ be a μ -preserving ergodic dynamical system, and let $\phi \in \mathcal{L}^1$ be an observable function, $\phi : M \rightarrow \mathbb{R}$. Then*

$$\lim_{n \rightarrow \infty} \frac{1}{n} \sum_{i=0}^{n-1} \phi(f^i(x)) = \int_M \phi d\mu.$$

Thus if f is one-dimensional, theorem 1 can be applied directly, to write

$$\lambda(x, v) = \lim_{n \rightarrow \infty} \frac{1}{n} \log \|D_x f^n v\| = \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{i=0}^{n-1} \log |D_{f^i x} f| = \int_M \log |Df| d\mu(x).$$

In higher dimensions, a fundamental obstacle is that matrix norms are not multiplicative, and so Lyapunov exponents cannot be expressed using the Birkhoff ergodic theorem. Results such as the sub-additive or multiplicative ergodic theorems establish the existence of Lyapunov exponents, but give no practical method for efficiently calculating them. Only in particularly simple cases can Lyapunov exponents be calculated explicitly.

A typical example is given by the Arnold Cat Map [3], which can be considered the composition of horizontal and vertical shears. Let $\tilde{F}, \tilde{G}, \tilde{H} : \mathbb{T}^2 \rightarrow \mathbb{T}^2$ be such that $\tilde{F}(x, y) = (x + y, y)$, $\tilde{G}(x, y) = (x, y + x)$, $\tilde{H}(x, y) = \tilde{G} \circ \tilde{F} = (x + y, x + 2y)$. Lebesgue measure is an ergodic invariant measure for \tilde{H} , and since the Jacobian $D\tilde{H}$ is the matrix $\begin{pmatrix} 1 & 1 \\ 1 & 2 \end{pmatrix}$ for all $x \in \mathbb{T}^2$, the two Lyapunov exponents are λ and λ^{-1} , where $\lambda = \log((3 + \sqrt{5})/2)$ is trivially computed to be the logarithm of the larger eigenvalue of $D\tilde{H}$. In general, a uniformly hyperbolic map has the property that expansion at every iterate is governed by the Lyapunov exponent, expressed as

$$\|D_x f^n v_0\| \geq c e^{\lambda n} \|v_0\| \quad (2)$$

for all $n \geq 0$ and some $c > 0$, for each v_0 in the expanding subspace of T_M (with a corresponding expression for v_0 in the contracting subspace).

A much wider class of maps is created when the very strict requirement of (2) is relaxed to allow growth rates to vary along trajectories. Non-uniformly hyperbolic systems can be characterised as systems with non-zero Lyapunov exponents, according to Pesin theory [4]. A canonical example is a linked twist map [5, 6, 7, 8], and we give here a simple version defined on the torus \mathbb{T}^2 . Fix $p \in (0, 1)$ and define annuli P and Q by

$$P = \{(x, y) \in \mathbb{T}^2 : y \leq p\},$$

$$Q = \{(x, y) \in \mathbb{T}^2 : x \leq p\}.$$

Consider the maps $F : \mathbb{T}^2 \rightarrow \mathbb{T}^2$ and $G : \mathbb{T}^2 \rightarrow \mathbb{T}^2$ given by

$$F \begin{pmatrix} x \\ y \end{pmatrix} = \begin{cases} F_1 \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} 1 & \frac{1}{p} \\ 0 & 1 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} & \text{if } (x, y) \in P, \\ F_2 \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} & \text{otherwise.} \end{cases} \quad (3)$$

$$G \begin{pmatrix} x \\ y \end{pmatrix} = \begin{cases} G_1 \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} 1 & 0 \\ \frac{1}{p} & 1 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} & \text{if } (x, y) \in Q, \\ G_2 \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} & \text{otherwise.} \end{cases} \quad (4)$$

We define the map $H = G \circ F : P \cup Q \rightarrow P \cup Q$ as the linked twist map given by the composition of the restrictions of F and G to $P \cup Q$, and we define the region $S = P \cap Q = \{(x, y) \in \mathbb{T}^2 : x, y \leq p\}$. It is straightforward to see that the source of the hyperbolicity is the fact that when an iterate of $H = G_1 F_1$ (corresponding to an

orbit of H landing in S and remaining in S after F), the orbit feels a composition of horizontal and vertical shears, as in the Cat Map. It is also easy to understand the non-uniformity, since an orbit may fall arbitrarily close to a boundary of $P \setminus S$ or $Q \setminus S$, and thus experience arbitrarily long (non-hyperbolic) sequences for which $H^n = F_1^n$ or $H^n = G_1^n$.

Linked twist maps are ergodic [6] and Lebesgue measure-preserving, and so for μ -a.e. (x, y) there are a pair of Lyapunov exponents, λ and λ^{-1} . Other values of λ correspond to zero measure sets, such as fixed points and periodic orbits of H . For example, the point $(p/2, p/2)$ is a period-3 point of H , with $DH^3 = \begin{pmatrix} 1 & 2/p \\ 2/p & 1 + 4/p^2 \end{pmatrix}$, for which the Lyapunov exponent is given by $\frac{1}{3} \log(1 + 2/p^2 + \sqrt{4/p^2 + 4/p^4})$. The orbit for which $DH = DG_1DF_1 = \begin{pmatrix} 1 & 1/p \\ 1/p & 1 + 1/p^2 \end{pmatrix}$ at every iterate corresponds to a uniformly hyperbolic horseshoe of zero measure. This sequence maximises the entropy h_μ over all possible periodic sequences [9], and hence λ by the Pesin Entropy Formula [4], and so gives an immediate upper bound $\lambda \leq \lambda_{h_\mu} = \log(1 + 1/2p^2 + \sqrt{1/p^2 + 1/4p^4})$. When $p = 1$, the linked twist map clearly equates to the Arnold Cat Map \tilde{H} .

Computing λ is far more difficult, even for simple maps like \tilde{F} and \tilde{G} , when they are composed in any other way than periodically. For example, suppose map $\hat{H} : \mathbb{T}^2 \rightarrow \mathbb{T}^2$ is defined by choosing either \tilde{F} or \tilde{G} at random (say, each with probability $1/2$) at each iterate. Because \tilde{F} and \tilde{G} do not commute, computing

$$\lambda = \lim_{n \rightarrow \infty} \frac{1}{n} \mathbb{E} \log \|D_x \hat{H}^n v_0\|$$

is a famously challenging problem. The limit converges, by the Furstenberg-Kesten theorem [10], but [11] placed this problem in pride of place of subadditive ergodic theory. While numerical schemes to approximate λ exist [12], rigorous upper and lower bounds on Lyapunov exponents for \hat{H} were established by one of the present authors in [13]. The idea is that any orbit of \hat{H} is a sequence of randomly chosen maps, such as $GFFGGGFF \dots GGFFGGGFF$, and the corresponding Jacobian can be bracketed as

$$D_x \hat{H}^n = \prod_{i=0}^{k-1} D\tilde{G}^{b_i} D\tilde{F}^{a_i} = \prod_{i=0}^{k-1} \tilde{M}_{a_i, b_i},$$

and where the matrices \tilde{M}_{a_i, b_i} possess an invariant cone over which one may maximise and minimise norms. Then, knowing the distributions of the a_i s and b_i s, and hence the average length of a block, is sufficient to construct an explicit expression for upper and lower bounds.

An orbit of the linked twist map can be bracketed in the same way, but while the corresponding M_{a_i, b_i} still preserve an equivalent invariant cone, in this case we have, *a priori*, no knowledge of the distribution of the a_i s and b_i s, which depend deterministically on the initial condition. We will also need to know the relationship between the number n of iterates in the orbit of H , and the number k of the corresponding number of the M_{a_i, b_i} . The aim of the present paper is to establish these requirements.

The paper is organised as follows. In section 2 we state our main results, which are rigorous upper and lower bounds for the Lyapunov exponents of the linked twist map H . The required invariant cone is described in section 3, in which we also maximise

and minimise growth rates for vectors inside this cone. In section 4 we construct the return time partition necessary to obtain the distributions of a and b , which completes the proof. We adapt our techniques in section 5, using a more intricate geometrical construction, to improve both upper and lower bounds, and finally in section 6 we discuss the accuracy of the bounds, introduce other quantities for which this method can produce bounds, and comment on the rate of convergence of Lyapunov exponents for linked twist maps.

2. Statement of results

Since H^n is equivalent to a sequence of maps F_1 and G_1 , we can write

$$DH^n = \prod_{i=1}^k DG^{b_i} DF^{a_i} = \prod_{i=1}^k M_{a_i, b_i} \quad (5)$$

where

$$M_{a,b} = \begin{pmatrix} 1 & \frac{a}{p} \\ \frac{b}{p} & 1 + \frac{ab}{p^2} \end{pmatrix}. \quad (6)$$

For example, if (x, y) is such that

$$H^6(x, y) = G_1 F_1 G_1 F_2 G_1 F_1 G_1 F_1 G_1 F_2 G_1 F_1(x, y) = G_1 F_1 G_1^2 F_1 G_1 F_1 G_1^2 F_1(x, y), \quad (7)$$

then $DH^6 = M_{1,1} M_{1,2} M_{1,1} M_{1,2}$. Now, assuming $\|v_0\| = 1$, we have

$$\|DH^n v_0\| = \prod_{i=1}^k \frac{\|M_{a_i, b_i} v_{i-1}\|}{\|v_{i-1}\|}, \quad (8)$$

where $v_i = M_{a_i, b_i} v_{i-1}$. To obtain an upper (lower) bound on $\|DH^n v_0\|$, and consequently λ , we require an upper (lower) bound for $\|M_{a,b} v\|/\|v\|$ for each possible combination of a and b , and in section 3 we will establish such an upper bound $\phi(a, b)$ and a lower bound $\psi(a, b)$. We will also need to know the distribution of these sequences - the frequency with which they occur along an orbit - in order to calculate their overall contribution to the Lyapunov exponent. Note that the ergodicity of LTMs ([6], [8]) ensures that these frequencies are equal for μ a.e. orbit. We will discuss this distribution in detail in Section 4, but for now let $R(a, b)$ be the proportion of pairs (a_i, b_i) in (5) that equal (a, b) . Finally, let n_S be the average number of constituent matrices DF and DG in each M_{a_i, b_i} . Then we have

$$\lambda \leq \frac{1}{n_S} \sum_{a,b=1}^{\infty} R(a, b) \log \phi(a, b) = \Phi_H, \quad (9)$$

and

$$\lambda \geq \frac{1}{n_S} \sum_{a,b=1}^{\infty} R(a, b) \log \psi(a, b) = \Psi_H. \quad (10)$$

For example, if the pattern in (7) were to continue, with $M_{1,1}$ and $M_{1,2}$ alternating indefinitely, then the proportions are $R(1, 1) = R(1, 2) = \frac{1}{2}$ and the average number is $n_S = \frac{1}{2}(1 + 2) = \frac{3}{2}$, so the upper bound (9) reads $\lambda \leq \frac{1}{3}(\log \phi(1, 1) + \log \phi(1, 2))$.

In order for these bounds to converge, the frequency with which the $M_{a,b}$'s occur must decrease faster than the logarithm of the bounds $\phi(a,b)$ and $\psi(a,b)$ increase; in other words, longer sequences must be sparser within an orbit than shorter ones. We will find that this is indeed the case for the family of maps H . Furthermore, we will see that only a select few of the sequences $M_{a,b}$ are possible, and the frequency with which they occur within an orbit can be written inductively for 'large' a or b . The results are summarised in the following theorem.

Theorem 2. *Let H be the linked twist map defined above, and let p^* be the real root of $p^3 + p - 1 = 0$ (that is, $p^* \approx 0.682$). Define for $k \in \{1, 2, \infty\}$*

$$\Psi_{H,k} = \frac{1}{n_S} \sum_{a,b=1}^{\infty} R(a,b) \log \psi_k(a,b) \quad (11)$$

$$\Phi_{H,k} = \frac{1}{n_S} \sum_{a,b=1}^{\infty} R(a,b) \log \phi_k(a,b). \quad (12)$$

where the functions $\psi_k(a,b)$ and $\phi_k(a,b)$ are given in lemma 1, each $R(a,b)$ is a rational function in p given by lemmas 2 and 3, and $n_S = (2-p)/p$. Then for $p^* \leq p \leq 1$ the largest Lyapunov exponent λ of H satisfies

$$\Psi_{H,k} \leq |\lambda| \leq \Phi_{H,k}. \quad (13)$$

In [13] upper and lower bounds on Lyapunov exponents in the random case were improved by considering two iterates of the map at a time, which allowed a tighter cone to be used. In section 5 we provide the corresponding construction for this deterministic case, giving

Theorem 3. *Let H be the linked twist map defined above, and let \tilde{p} be the real root of $p^5 + 3p^3 - 3p^2 + p - 1 = 0$ (that is, $\tilde{p} \approx 0.8562$). Define for $k \in \{1, 2, \infty\}$*

$$\tilde{\Psi}_{H,k} = \frac{1}{n_S} \sum_{a_1, b_1, a_2, b_2=1}^{\infty} \tilde{R}(a_2, b_2, a_1, b_1) \log \tilde{\psi}_k(a_2, b_2, a_1, b_1) \quad (14)$$

$$\tilde{\Phi}_{H,k} = \frac{1}{n_S} \sum_{a_1, b_1, a_2, b_2=1}^{\infty} \tilde{R}(a_2, b_2, a_1, b_1) \log \tilde{\phi}_k(a_2, b_2, a_1, b_1). \quad (15)$$

where the functions $\tilde{\psi}_k(a_2, b_2, a_1, b_1)$ and $\tilde{\phi}_k(a_2, b_2, a_1, b_1)$ are given in lemma 4, each $\tilde{R}(a_2, b_2, a_1, b_1)$ is a rational function in p given by lemma 5, and $n_S = (2-p)/p$. Then for $\tilde{p} \leq p \leq 1$ the largest Lyapunov exponent λ of H satisfies

$$\tilde{\Psi}_{H,k} \leq \lambda \leq \tilde{\Phi}_{H,k}. \quad (16)$$

3. Invariant cones

In this section we discuss how to calculate the bounds $\phi(a,b)$ and $\psi(a,b)$, which requires finding bounds upon (8). To do this we make use of the existence of invariant cones ([14, 15, 16]) for the matrices $M_{a,b}$. Note that $M_{a,b}$ as given in (6) is diagonalizable with positive eigenvalues for any $a, b \geq 1$, and the eigenvectors of $M_{a,b}$ are given by

$$v_{\pm} = \begin{pmatrix} \frac{b}{p} \pm \sqrt{\frac{4b}{a} + \frac{b^2}{p^2}} \end{pmatrix}.$$

A cone is invariant for $M_{a,b}$ if and only if it contains v_+ , and its interior does not contain v_- [17]. A simple calculation shows that the cone $C(p)$ given by

$$C(p) = \{(x, y) \in T_x \mathbb{T}^2 : 0 \leq \frac{x}{y} \leq p\} \quad (17)$$

satisfies these conditions for any $a, b \geq 1$, and so C is an invariant cone for every $M_{a,b}$. Note that C is *not* an invariant cone for H , as its Jacobian matrix DH is not always hyperbolic.

In order to bound $\|H^N v\|$, we find upper and lower bounds for the quantity $\|M_{a,b} v\|/\|v\|$, for $v \in C$ and each possible a, b , using the ℓ_∞ , ℓ_1 and ℓ_2 norms.

Lemma 1. *The growth factors $\|M_{a,b} v\|_i/\|v\|_i$ for $i \in \{1, 2, \infty\}$ are bounded as*

$$\psi_i(a, b) \leq \frac{\|M_{a,b} v\|_i}{\|v\|_i} \leq \phi_i(a, b) \quad \text{for all } v \in C,$$

where the bounds are given by

$$\begin{aligned} \psi_1(a, b) &= 1 + \frac{ab + pa + p^2 b}{p^2(1 + p)} \\ \phi_1(a, b) &= 1 + \frac{a}{p} + \frac{ab}{p^2} \\ \psi_2(a, b) &= \min \left\{ \sqrt{\frac{(p + \frac{a}{p})^2 + (1 + b + \frac{ab}{p^2})^2}{1 + p^2}}, \sqrt{(\frac{a}{p})^2 + (1 + \frac{ab}{p^2})^2} \right\} \\ \phi_2(a, b) &= \sqrt{\lambda_{M_{a,b}^T M_{a,b}}^u} \\ \psi_\infty(a, b) &= 1 + \frac{ab}{p^2} \\ \phi_\infty(a, b) &= 1 + b + \frac{ab}{p^2}. \end{aligned}$$

Here, $\lambda_{M_{a,b}^T M_{a,b}}^u$ denotes the unstable eigenvalue of $M_{a,b}^T M_{a,b}$.

Proof. We first compute the vectors $v_{max}^i, v_{min}^i \in T_x \mathbb{T}^2$ that maximise and minimise the quantity $\|M_{a,b} v\|_i/\|v\|_i$ over the entire tangent space. By the definition of the spectral norm, we have that v_{max}^2 and v_{min}^2 are the unstable and stable eigenvectors of $M_{a,b}^T M_{a,b}$, respectively. Simple calculations show that

$$\begin{aligned} v_{max}^1 &= \begin{cases} (0, 1) & \text{if } \frac{a}{p} \geq \frac{b}{b+p}, \\ (1, 0) & \text{otherwise,} \end{cases} & v_{min}^1 &= \left(\frac{ab}{p^2} + 1, -\frac{b}{p} \right), \\ v_{max}^\infty &= (1, 1), & v_{min}^\infty &= \left(-\frac{ab}{p^2} - \frac{a}{p} - 1, \frac{b}{p} + 1 \right). \end{aligned}$$

Of all these vectors, v_{max}^2 is the only one inside the cone C , which yields the upper bound $\phi_2(a, b)$. In all other cases, we use that the norms vary monotonically between their minimum and maximum and thus the extrema are attained on the boundaries of C . Some further simple computations complete the proof. \square




Figure 1: The intersection $H^{-1}(S) \cap S = R_{1,1}$ is shown in dark grey. The pale grey quadrilaterals are the parts of $H^{-1}(S)$ which lie outside S , and which will return on a later iterate of H . Here we choose $p = 0.8$.

4. The return time distribution

A common technique in the study of hyperbolic dynamical systems is to inspect a return map to a hyperbolic set. That is, consider the map $H_S : S \rightarrow S$, defined by $H_S = H^k$, where k is such that, for $x \in S$, $H^k(x) \in S$ and $H^j(x) \notin S$ for $j = 1, \dots, k-1$. Such a return map induces a natural partition of S into regions of different return times k . This partition is central to the application of Young Tower techniques for computing mixing rates, and H_S was used in [18] to demonstrate that LTMs have polynomial decay of correlations. Here we observe that H_S partitions S into countably many open sets on which H_S is characterised by $DH_S = M_{a,b}$, where $a + b - 1 = k$. Typically in decay of correlation arguments only the partition sets for large k , and the rate at which they decrease in size, are of interest. Here we construct in detail the entire return time set, and use the notation

$$R_{a,b} = \{x | D_x H_S = M_{a,b}\},$$

that is, $R_{a,b}$ contains those points x for whom $H_S = G_1^b F_1^a$. We characterise the size of the sets $R_{a,b}$ using the conditional measure μ_S , and write

$$\mu_S(R_{a,b}) = \mu_L(R_{a,b})/p^2,$$

where μ_L is Lebesgue measure. Note that μ_S is an ergodic invariant measure for H_S [8].

In practice, to find the return time distribution, we calculate the pre-images of the set S under H , and record the sets of points for which these pre-images intersect S . For example, the set $R_{1,1}$ is the set $H^{-1}(S) \cap S$. Figure 1 shows the geometrical construction of this procedure, with all the coordinates of the intersections needed




Figure 2: The darker grey regions are the points which return to S after one ($R_{1,1}$) and two ($R_{1,2}$ and $R_{2,1}$) iterates of H . The lighter grey regions indicate the points yet to return, which are iterated backwards to obtain the later distribution elements. Of these, the quadrilateral will return on the next iterate to the central white quadrilateral ($R_{2,2}$), while the triangular regions form the sets $R_{n,1}$ and $R_{1,n}$ for $n \geq 3$. The circles mark points mentioned in the proof of lemma 2.

to calculate the area of this set. Figure 2 continues this procedure, to produce the regions which return to S under a second application of H^{-1} .

Lemma 2. *Let $p^3 + p - 1 \geq 0$. Then*


$$\begin{aligned}\mu_S(R_{1,1}) &= \frac{2p^3}{1+p^2}, \\ \mu_S(R_{1,2}) = \mu_S(R_{2,1}) &= \frac{-3p^4 + 2p^3 + 2p - 1}{2p(1+p^2)}, \\ \mu_S(R_{2,2}) &= \frac{(1-p)^2}{1+p^2}.\end{aligned}$$

Proof. These are all simple calculations based on the constructions shown in figures 1 and 2. We note that we give the condition $p^3 + p - 1 \geq 0$ (or $p \geq 0.682\dots$) since if this condition is not met, then the change in gradient which occurs at the points $(\frac{p^3+p-1}{p}, 1-p)$ and $(\frac{-p^3+p^2-2p+1}{p}, 2p-1)$ (marked with circles in figure 2) will instead be contained within the unreturned regions, and will continue into later iterates, making the return time distribution more complicated, but not notionally more difficult. \square

The set of unreturned points for $n \geq 2$ consists of $R_{2,2}$, plus four triangles, shown in blue in figure 2. Apart from $R_{2,2}$, the itineraries for such unreturned points are of the form $F^n G$ and FG^n , for $n \geq 3$. There are countably many such regions $R_{n,1}$ and $R_{1,n}$ but these are also simply calculated, and are shown in figure 3.

Lemma 3. *Let $n \geq 3$ and as before $p^3 + p - 1 \geq 0$. Then*

$$\mu_S(R_{n,1}) = \mu_S(R_{1,n}) = \frac{2(1-p)^2}{np(n-1)(n-2)}.$$


 Figure 3: The complete return time partition, shown here for $p = 0.8$.

Proof. The set $R_{n,1}$ consists of two triangles, one near $(1,0)$ and one near $(0,1)$. The first of these can be expressed as the difference between triangles T_n and T_{n-1} , where the corners of T_n have coordinates $(1,0)$, $(p, p(1-p)/n)$, $(p, p(1-p)/(n-1))$. The area of T_n is $p(1-p)^2/2n(n-1)$ and so $\mu_S(R_{n,1})$ is twice $p(1-p)^2/n(n-1)(n-2)$. The set $R_{1,n}$ is similar. \square

To complete the proof of theorem 2 we recognise that the required frequency with which $M_{a,b}$ appears along an orbit of H is exactly the relative size of the subset of S for which the return map H_S is equal to $G_1^b F_1^a$. That is, $R(a,b) = \mu_S(R_{a,b})$. This is an immediate consequence of theorem 1 with $\phi = \chi_{R_{a,b}}$. Finally, we require n_S . This can be computed directly from the expectation

$$\begin{aligned}
 n_S &= \langle a + b - 1 \rangle = \sum_{a,b} (a + b - 1) R(a,b) \\
 &= R(1,1) + 2(R(1,2) + R(2,1)) + 3R(2,2) + \sum_{n=3}^{\infty} n(R(1,n) + R(n,1)) \\
 &= \frac{-4p^4 + 7p^3 - 6p^2 + 7p - 2}{p(1+p^2)} + \frac{4(1-p)^2}{p} \sum_{n=3}^{\infty} \frac{1}{(n-1)(n-2)} \\
 &= \frac{-4p^4 + 7p^3 - 6p^2 + 7p - 2}{p(1+p^2)} + \frac{4(1-p)^2}{p} \sum_{m=1}^{\infty} \frac{1}{m(m+1)} \\
 &= \frac{-4p^4 + 7p^3 - 6p^2 + 7p - 2}{p(1+p^2)} + \frac{4(1-p)^2}{p} \\
 &= \frac{2}{p} - 1.
 \end{aligned}$$

Alternatively, we can apply Kac's lemma [19], which states that for an ergodic,




Figure 4: The partition depicting $\{x | H_S^2 = F_1^{a_2} G_1^{b_2} F_1^{a_1} G_1^{b_1}\}$ for $a_1, a_2, b_1, b_2 \in \{1, 2, 3, \dots\}$.

measure-preserving transformation, the expected first return time to a set S is $1/\mu(S)$, where μ is the normalized measure on the domain. Since $\mu(S) = p^2$ and $\mu(P \cap Q) = 1 - (1 - p)^2 = p(2 - p)$ we have $n_S = p(2 - p)/p^2 = 2/p - 1$, confirming the validity of the return time distribution in this section.

5. Improving the bounds

Here we discuss the circumstances under which we can consider narrower cones than C , in order to improve upon the bounds $\Psi_{H,k}$ and $\Phi_{H,k}$ in Theorem 2. In [13], the case of bounding random products of matrices, the corresponding modifications were relatively simple, as the frequency with which the matrix M_{a_2, b_2} preceded the matrix M_{a_1, b_1} in any orbit was independent of the choice of a_1 and b_1 . In the present deterministic case, however, this independence does not hold.

Instead, we consider which matrices M_{a_2, b_2} can precede each M_{a_1, b_1} . The benefit of doing so is to replace the cone C with its narrower image $M_{a_2, b_2}(C)$, improving the bounds of lemma 1.

Lemma 4. *We have*

$$\tilde{\psi}_k(a_2, b_2, a_1, b_1) \leq \frac{\|M_{a_1, b_1} v\|_k}{\|v\|_k} \leq \tilde{\phi}_k(a_2, b_2, a_1, b_1)$$

for $k \in \{1, 2, \infty\}$ and $v \in M_{a_2, b_2}(C)$, with

$$\begin{aligned}\tilde{\psi}_\infty &= 1 + \frac{a_1 b_1}{p^2} + \frac{a_2 b_1}{p^2 + a_2 b_2} \\ \tilde{\phi}_\infty &= 1 + \frac{a_1 b_1}{p^2} + \frac{b_1 p^2 + a_2 b_1}{p^2(1 + b_2) + a_2 b_2} \\ \tilde{\psi}_1 &= 1 + \frac{a_1 p^3 + (a_2 b_1 + a_1 b_1)p^2 + a_2 a_1 b_2 p + a_1 a_2 b_1 b_2}{p^2(a_2 b_2 + a_2 p + p^2)} \\ \tilde{\phi}_1 &= 1 + \frac{b_1 p^4 + (a_1 b_2 + a_1)p^3 + (b_2 a_1 b_1 + a_2 b_1 + a_1 b_1)p^2 + a_1 a_2 b_2 p + a_1 a_2 b_1 b_2}{p^2(p^3 + (b_2 + 1)p^2 + a_2 p + a_2 b_2)} \\ \tilde{\psi}_2 &= \min\{\xi_1, \xi_2\} \\ \tilde{\phi}_2 &= \begin{cases} \sqrt{\lambda_{M_{a_1, b_1}^T M_{a_1, b_1}}^u} & \text{if } v_{max}^2 \in M_{a_2, b_2}(C) \\ \max\{\xi_1, \xi_2\} & \text{otherwise,} \end{cases}\end{aligned}$$

where

$$\begin{aligned}\xi_1 &= \frac{\sqrt{\left(\frac{a_1(a_2 b_2 + p^2)}{p^3} + \frac{a_2}{p}\right)^2 + \left(\frac{a_2 b_1}{p^2} + \left(\frac{a_1 b_1}{p^2} + 1\right)\left(\frac{a_2 b_2}{p^2} + 1\right)\right)^2}}{\sqrt{\frac{a_2^2}{p^2} + \left(\frac{a_2 b_2}{p^2} + 1\right)^2}} \\ \xi_2 &= \frac{\sqrt{\left(\frac{b_1(a_2 + p^2)}{p^2} + \left(\frac{a_1 b_1}{p^2} + 1\right)\left(\frac{a_2 b_2}{p^2} + b_2 + 1\right)\right)^2 + \left(\frac{a_1(a_2 b_2 + p^2(b_2 + 1))}{p^3} + \frac{a_2}{p} + p\right)^2}}{\sqrt{\left(\frac{a_2}{p} + p\right)^2 + \left(\frac{a_2 b_2}{p^2} + b_2 + 1\right)^2}}\end{aligned}$$

Proof. Similarly to lemma 1, $\|M_{a_1, b_1} v\|_k / \|v\|_k$ changes monotonically in $M_{a_2, b_2}(C)$ in the l_1 and l_∞ cases. In the l_2 case, the maximising vector v_{max}^2 may or may not be contained in the cone $M_{a_2, b_2}(C)$. The calculations are again elementary. \square

To complete the improved bounds, we require the proportion of a typical orbit which sees M_{a_2, b_2} followed by M_{a_1, b_1} . To obtain this we calculate the measure of the set of points in R_{a_2, b_2} which map under the return map H_S into R_{a_1, b_1} , that is, $\mu(\{x | H_S^2 = F_1^{a_2} G_1^{b_2} F_1^{a_1} G_1^{b_1}\})$. Since H is μ -invariant we have

$$\mu(R_{a_2, b_2} \cap H_S^{-1}(R_{a_1, b_1})) = \mu(H_S(R_{a_2, b_2}) \cap R_{a_1, b_1})$$

and we use the notation $H_S(R_{a_2, b_2}) \cap R_{a_1, b_1} = \tilde{R}_{a_2, b_2, a_1, b_1}$. So we consider the partition in figure 3 and its forward image, which by the symmetry of the map is simply the original partition rotated through 90° . The partition and its image is shown in figure 4. The size of each region of this ‘improved partition’ can be calculated using simple geometry. The task is made simpler by the fact that the countably infinite regions of the original partition do not intersect with their own images, that is, for $n \geq 3$,

$$H_S(R_{1, n} \cup R_{n, 1}) \cap (R_{1, n} \cup R_{n, 1}) = \emptyset$$

(noted as Lemma 3.2 of [18]) and moreover the symmetry induces the relationship

$$\mu_S(\tilde{R}_{a_2, b_2, a_1, b_1}) = \mu_S(\tilde{R}_{b_1, a_1, b_2, a_2})$$

for all a_1, b_1, a_2, b_2 . The areas $\mu_S(\tilde{R}_{a_2, b_2, a_1, b_1})$ are given in:

Lemma 5.

$$\begin{aligned}
 \mu_S(\tilde{R}_{1,1,1,1}) &= \frac{p(12p^8 + 28p^6 - 12p^5 + 14p^4 - 16p^3 + 6p^2 - 4p + 2)}{2p^8 + 9p^6 + 12p^4 + 6p^2 + 1} \\
 \mu_S(\tilde{R}_{1,1,1,2}) &= \mu(\tilde{R}_{2,1,1,1}) \\
 &= -\frac{(10p^{10} + 4p^9 + 9p^8 - 10p^7 + p^6 - 24p^5 + 14p^4 - 12p^3 + 9p^2 - 2p + 1)}{4p^9 + 18p^7 + 24p^5 + 12p^3 + 2p} \\
 \mu_S(\tilde{R}_{1,1,2,1}) &= \mu(\tilde{R}_{1,2,1,1}) = -\frac{8p^6 - 4p^5 - 6p^3 + p^2 + 1}{4p^5 + 6p^3 + 2p} \\
 \mu_S(\tilde{R}_{1,1,2,2}) &= \mu(\tilde{R}_{2,2,1,1}) = \frac{(-p^3 + 4p^2 - 5p + 2)}{2(p^2 + 1)} \\
 \mu_S(\tilde{R}_{1,2,1,2}) &= \mu(\tilde{R}_{2,1,2,1}) = \frac{2p(p^2 - 2p + 1)}{2p^2 + 1} \\
 \mu_S(\tilde{R}_{1,2,2,1}) &= \frac{(-p^5 + 4p^4 - 9p^3 + 14p^2 - 12p + 4)}{p^4 + 5p^2 + 4} \\
 \mu_S(\tilde{R}_{1,2,2,2}) &= \mu(\tilde{R}_{2,2,2,1}) = \frac{p(p^4 - 4p^3 + 9p^2 - 10p + 4)}{2(p^4 + 5p^2 + 4)} \\
 \mu_S(\tilde{R}_{2,1,1,2}) &= \frac{4p^2(p^2 - 2p + 1)}{p^4 + 3p^2 + 1} \\
 \mu_S(\tilde{R}_{2,2,2,2}) &= \frac{p^2(p^2 - 2p + 1)}{p^4 + 5p^2 + 4} \\
 \mu_S(\tilde{R}_{n,1,1,1}) &= \mu_S(\tilde{R}_{1,n,1,1}) = \mu_S(\tilde{R}_{1,1,n,1}) = \mu_S(\tilde{R}_{1,1,1,n}) = \frac{2(1-p)^2}{np(n-1)(n-2)}.
 \end{aligned}$$

In each case, by theorem 1, we have $\tilde{R}(a_2, b_2, a_1, b_1) = \mu_S(\tilde{R}_{a_2, b_2, a_1, b_1})$.

Proof. The computation of these areas are all the result of simple geometry, illustrated in figure 4. \square


Replacing the functions ϕ_k , ψ_k and R of theorem 2 with $\tilde{\phi}_k$, $\tilde{\psi}_k$ and \tilde{R} completes the proof of theorem 3.

6. Discussion

6.1. Accuracy of the bounds


The bounds on λ given by theorems 2 and 3 can be computed easily and rapidly. Here we compare the numerical values of these bounds with values produced by a standard numerical algorithm [12] using Gram-Schmidt orthonormalisation to quantify the exponential growth of tangent vectors implicit in a positive Lyapunov exponent. The bounds $\Phi_{H,k}$ and $\Psi_{H,k}$ each involve an infinite sum, but truncating these at some large but finite limit give accurate evaluations. In particular, since every term in each sum is positive, any truncation of $\Psi_{H,k}$ is a strict lower bound, while each term decreases roughly as $n^{-3} \log n$.

Figure 5 shows the bounds $\Phi_{H,k}$ and $\Psi_{H,k}$ for each of the L_1 (red), L_2 (blue), L_∞ (green) norms. The numerically calculated value for λ , from 10^5 iterates of the



(a) All upper and lower bounds from theorem 2. (b) Sharpest bounds result from using l_2 for the upper bound, and l_1 for the lower.

Figure 5: (Colour online). Bounds of λ given by theorem 2. The black line shows the numerically computed value of λ . Upper bounds are shown as solid lines, lower bounds as dashed lines. As labelled, and in later figures, red lines originate from the l_1 norm, blue lines from the l_2 norm, and green lines from the l_∞ norm. The magenta line indicates the upper bound from maximising λ_{h_μ} .



(a) Distance between upper and lower bounds from theorem 2 shown dotted, and from theorem 3 as solid lines. (b) The sharpest bounds, using the l_2 norm, improve on the numerical calculation of 10^5 iterates.

Figure 6: (Colour online). Bounds of λ given by theorem 3. The black line shows the numerically computed value of λ .

standard orthonormalisation scheme, is shown as a solid black line. Upper bounds are shown as solid lines, and lower as dashed lines. Figure 5a shows that the tightest upper bound comes from the L_2 norm, while the tightest lower bound comes from the L_1 norm. The bound given by the topological entropy calculation of [9] is shown in magenta. Note that the upper bound $\Phi_{H,2}$ coincides with λ at $p = 1$. This is because the vector $v_{max}^2 \in C$ given in the proof of lemma 1 is equal to the expanding eigenvector of H when $p = 1$. The upper bounds for L_1 and L_∞ also coincide at $p = 1$, where the expressions in lemma 1 are equivalent. Figure 5b selects the best bounds, and shows that $\Phi_{H,2}$ is considerably tighter than $\Psi_{H,1}$, but still appears greater than the numerically calculated value for all $p < 1$.

Figure 6 shows the bounds $\tilde{\Phi}_{H,k}$ and $\tilde{\Psi}_{H,k}$. Here we plot $\Phi_{H,k} - \Psi_{H,k}$ and




Figure 7: Rigorous upper and lower bounds on the generalized Lyapunov exponent $\ell(q)$ for the linked twist map H , defined by equation (18), using the l_2 norm. The circle indicates $\ell(1)$, which corresponds to the topological entropy $\lambda_{h_\mu} = \log(1 + 1/2p^2 + \sqrt{1/p^2 + 1/4p^2})$.

$\tilde{\Phi}_{H,k} - \tilde{\Psi}_{H,k}$ (in figure 6a) to show the improvement in using the return partition and its image. Figure 6b shows the bounds using the L_2 norm, the tightest bounds, and illustrates that the envelope $\tilde{\Phi}_{H,2} - \tilde{\Psi}_{H,2}$ is often narrower than the uncertainty in the numerically calculated value of λ , even after 10^5 iterates.

6.2. Related quantities

The method described here could also be used to bound related quantities. For example, of interest in many applications is the *generalized Lyapunov exponent* $\ell(q)$, which gives the growth rate of the q -th moment of a matrix product norm [20]. The generalized Lyapunov exponent is used in large deviation theory, and is typically defined for products of random matrices. In this deterministic setting we have

$$\ell(q) = \lim_{n \rightarrow \infty} \frac{1}{n} \log \langle \|D_x f^n v_0\|^q \rangle \quad (18)$$

where the average is taken over the invariant measure μ . This is related to the standard Lyapunov exponent by $\lambda = \ell'(0)$. This quantity is notoriously difficult to compute numerically [21], because fluctuations along a finite orbit are magnified for large q . However, the expressions in theorems 2 and 3 can be easily adapted to give quick and accurate upper and lower bounds for $\ell(q)$, since the contributions from averaging over the invariant measure are given explicitly by $R(a, b)$. These bounds are shown, for the l_2 norm, in figure 7. Note that $\ell(q)$ is monotonic, since the existence of the invariant cone means that a vector is expanded by DH at every iterate of the map. Generalized Lyapunov exponents are closely related to the joint spectral radius, a quantity which can also be rigorously bounded, for each l_p norm, by this method.

6.3. Convergence of Lyapunov exponents

Finally, we comment on the difficulty of computing Lyapunov exponents numerically for linked twist maps. The standard algorithm [12] is not difficult to code, and as the system is low-dimensional and discrete time, is quick to run. However, as shown in figure 6b, a large number of iterates is required to improve on the rigorous bounds presented. Indeed the numerical calculation shows poor convergence to the expected values of λ , which should change smoothly with p . Typically, Lyapunov exponents converge according to the central limit theorem. Linked twist maps, however, have slow correlation decay (shown in [18] to be polynomial, with rate $1/n$). This results in a central limit theorem with a non-standard scaling factor of $\sqrt{n \log n}$, and so slower convergence than usual [22].

References

- [1] Oseledec V I 1968 *Transactions of the Moscow Mathematical Society* **19** 197–231
- [2] Birkhoff G D 1931 *Proceedings of the National Academy of Sciences* **17** 656–660
- [3] Arnold V and Avez A 1967 *Paris: Gauthier-Villars*
- [4] Pesin Y B 1977 *Russian Mathematical Surveys* **32** 55–114
- [5] Devaney R L 1978 *Proceedings of the American Mathematical Society* **71** 334–338
- [6] Burton R and Easton R W 1980 Ergodicity of linked twist maps *Global Theory of Dynamical Systems* (Springer) pp 35–49
- [7] Wojtkowski M 1980 *Annals of the New York Academy of sciences* **357** 65–76
- [8] Przytycki F 1983 *Annales scientifiques de l'Ecole normale supérieure* **16** 345–354
- [9] D'Alessandro D, Dahleh M and Mezić I 1999 *IEEE Transactions on Automatic Control* **44** 1852–1863
- [10] Furstenberg H and Kesten H 1960 *The Annals of Mathematical Statistics* **31** 457–469
- [11] Kingman J F C 1973 *Ann. Probab.* **1** 883–899
- [12] Parker T S and Chua L 2012 *Practical numerical algorithms for chaotic systems* (Springer Science & Business Media)
- [13] Sturman R and Thiffeault J L 2019 *Journal of Nonlinear Science* **29** 593–620
- [14] Alekseev V M 1968 *Mathematics of the USSR-Sbornik* **5** 73
- [15] Alekseev V M 1968 *Mathematics of the USSR-Sbornik* **6** 505
- [16] Alekseev V M 1969 *Mathematics of the USSR-Sbornik* **7** 1
- [17] Rodman L, Seyalioglu H and Spitkovsky I M 2010 *Linear Algebra and its Applications* **432** 911–926
- [18] Springham J and Sturman R 2014 *Ergodic Theory and Dynamical Systems* **34** 1724–1746
- [19] Kac M 1947 *Bulletin of the American Mathematical Society* **53** 1002–1010
- [20] Crisanti A, Paladin G and Vulpiani A 1988 *Journal of Statistical Physics* **53** 583–601
- [21] Vanneste J 2010 *Physical Review E* **81** 036701
- [22] Mohr L 2013 *Martingale central limit theorem and nonuniformly hyperbolic systems* Ph.D. thesis University of Massachusetts Amherst Open Access Dissertations. 810. <https://doi.org/10.7275/vd8y-jv94>https://scholarworks.umass.edu/open_access_dissertations/810