

# Co-occurrence of marine heat waves and cold spells in nearshore and offshore regions along South Africa

Robert W. Schlegel<sup>a,\*</sup>, Eric C. J. Oliver<sup>b,c</sup>, Thomas W. Wernberg<sup>d</sup>, Albertus J. Smit<sup>a</sup>

<sup>a</sup>*Department of Biodiversity and Conservation Biology, University of the Western Cape, Private Bag X17, Bellville 7535, South Africa*

<sup>b</sup>*ARC Centre of Excellence for Climate System Science, The University of New South Wales, Sydney, Australia*

<sup>c</sup>*Institute for Marine and Antarctic Studies, University of Tasmania, Hobart, Australia*

<sup>d</sup>*UWA Oceans Institute and School of Plant Biology, The University of Western Australia, Crawley, 6009 Western Australia, Australia*

---

## Abstract

The term marine heat wave (MHW) was first coined in 2013 with no central definition having been agreed upon before. This lack of a definition had led to an inability of different research groups to compare their findings on this phenomenon before 2013. In order to assuage this issue, a research team has recently created a definition for MHWs that will be valid anywhere in the world. We have taken this algorithm and applied it to the *in situ* time series available for the coast of South Africa that are longer than 10 years and with at least 90% complete daily records. It was also decided to apply the algorithm to cool temperatures and investigate the presence of marine cold spells (MCSs). We found that MHWs and MCSs can be found along the entire stretch of South Africa's coastline and with some temporal and spatial agreement between the largest events detected. MHWs occur more often, last longer than MCSs and have greater cumulative intensities. There was little variance in the cumulative intensity [ $^{\circ}\text{C} \times \text{days}$ ] around the mean for MHWs and MCSs however, several were much larger and there tended to be specific time series that displayed more dramatic results than others. The coastline was further divided into three sections (west, south, and east) to investigate the effect of geography on MHWs and MCSs and it was found that the south coast experiences more, longer and more intense MHWs and MCSs than the other two coastlines. The mechanism driving the higher intensity of events on the south coast, which is much greater than the other coasts, requires further study. The largest three MHWs of most time series along the coast of South Africa have occurred in the second half of the time series whereas the largest three MCSs have occurred in the first half. These same calculations were conducted for offshore temperatures from NOAA optimally interpolated sea surface temperature (OISST) data, too. It was found that the proportion of co-occurrence between *in situ* and OISST data ranged from 0.5–0.0 for each coastline with co-occurrence rates being the largest on the south coast. Few time series showed co-occurrence amongst the 50% largest events.

**Keywords:** marine heat waves, marine cold spells, OISST, *in situ* data, co-occurrence, climate change, extreme events, South Africa, coastal

---

\*Corresponding author

Email address: 3503570@myuwc.ac.za (Robert W. Schlegel)

## 1. Introduction

Over the past three decades, global-scale anthropogenically mediated warming has negatively affected marine and terrestrial realms with far reaching consequences for humanity and natural ecological functioning. Although climate change is generally understood as a gradual long-term rise in global mean surface temperature [39], which will continue for decades or centuries, it is generally the associated increase in frequency and severity of extreme events that affects humans and ecosystems alike in the short-term [12]. Impacts are often sudden with catastrophic consequences. Such extreme events include droughts, floods, wind storms, tropical cyclones, heat waves and cold spells. ‘Pulse’ events exceeding certain thresholds of frequency, intensity (extremeness), duration, timing and rate of onset (abruptness) can drive punctuated perturbations to species distributions, which eventually modify the structure and function of ecosystems [49, 1], and the recognition to focus more on events and less on trends has emerged as a recent direction of climate change research [26]. The focus of this paper is on marine thermal events that are extreme with respect to the seasonal climatology. They may be anomalously warm (marine heat waves, MHW; *sensu* Hobday et al. 2016), or anomalously cold (marine cold spells, MCS; introduced here). While MHWs are becoming reasonably well known by virtue of their increasing frequency and intensity, there is less information about the ecological effects of extreme cold events. There is also a paucity of information about their drivers.

The concept of heat waves is usually applied to atmospheric phenomena where vague definitions such as “a period of abnormally and uncomfortably hot and usually humid weather” are invoked [3], but there are also precise definitions based on statistical properties and other metrics of the temperature record that are relative to location and time of year (e.g. [35, 2, 18, 17]). Recent years have seen investigations of heat waves in the ocean due to them becoming more frequent over time (e.g. [32, 44, 47, 29, 10]). Well documented marine heat waves (MHW) have occurred in the Mediterranean in 2003 (e.g. [4, 37, 19]), off the coast of Western Australia in 2011 (e.g. [14, 40, 49]), in the north west Atlantic Ocean in 2012 (e.g. [36, 8, 7]) and now the “Blob” from 2014 to 2016 in the north east Pacific Ocean [5]. The extreme temperatures from these events, and others like them, may have wide ranging negative impacts upon the local ecology for the regions in which they occur. For example, the 2003 Mediterranean heat wave may have affected up to 80% of the gorgonian fan colonies in certain areas of this sea [19], whereas the 2011 event off the west coast of Australia has been recognized as being a driving factor in the regime shift there from temperate kelp forests to the beginnings of a coral reef system [49]. Because the inquiry into MHWs is a relatively new endeavour none of these studies provided adequate definitions for what constitutes a MHW, and to that end Hobday et al. (2016) [23] have defined it as “a prolonged discrete anomalously warm water event that can be described by its duration, intensity, rate of evolution, and spatial extent”. By applying the MHW definition to the aforementioned events, Hobday et al. (2016) [23] were able to derive statistical features of the MHWs, such as their frequency along a time series and maximum and cumulative intensity. Whereas extreme hot events may be demonstrably damaging to organisms and ecosystems, extreme cold events also have the potential to negatively impact organisms and ecosystems.

MCSs are analogous to MHWs, but they are of opposite sign. MCSs are projected to become less frequent under future climatic scenarios, but there are also examples of them becoming more frequent in some localities [20, 34]. They are frequently lethal [50] and are known to have caused mass fish [21, 22, 24] and invertebrate [22, 9] kills, the death of juvenile and sub-adult manatees [38, 33] as well as affecting organismal physiological tolerances, life history strategies, and habitat requirements [13]. Cold temperatures are therefore very important in setting species distribution limits, particularly limiting their range north- or southwards towards high latitudes [15], and the timing of the onset of the growing season [26]. At an ecosystem level there is still a paucity of information on effects of MCSs, but it is easy to postulate how population-level consequences might aggregate to drive whole ecosystem responses (e.g. [27, 1]). Indeed, the range contractions of ecosystem engineer species such as mussels have been shown to relate to MCSs (e.g. [15, 16]).

Some of the MCSs known to have had impacts on populations and ecosystems were caused by atmospheric cold spells that affect the intertidal biota (e.g. [21, 15]) and not by seawater. Here we focus on MCSs measured in seawater. This may imply local events (i.e. extreme atmospheric cold spells that perturb the seawater locally) or broad-scale drivers. The driver of the MCSs localised to the coast, as we have already suggested, is hypothesised to be coastal weather phenomena. But what mechanism might explain coastal MCSs originating from offshore or by oceanic processes? Large-scale atmospheric-oceanographic coupling is very likely being affected by global warming, which is projected to cause the intensification of upwelling favourable winds and consequently the intensification and increasing frequency of upwelling (see Garc  a-Reyes et al. 2015 for a review of this and alternative hypotheses). The question then is, could an intensification of upwelling be attributed to coastal MCSs, or are they linked to local coastal atmospheric forcing? Little research yet exists that investigates this question other than to link anoxia and other negative factors from problematic phytoplankton blooms caused by extreme upwelling events to create lethal conditions for species living within upwelling regions (e.g. [28]). Whereas anoxia is a problem attributable to phytoplankton blooms themselves [11] and not the extreme cold temperatures *per se*, if a relationship can be shown between MCSs and anoxia resulting from algal blooms it would provide extremely valuable insight into how coastal ecosystems respond to climatic change. Furthermore, since mass mortalities and ecosystem change may result directly from MCSs, a mechanistic understanding of their drivers will be invaluable. To this end it serves as a constructive first step to study the prevalence of MCSs with respect to different kinds of oceanic forcing mechanisms.

Hobday et al. (2016) [23] applied their MHW framework to  $\frac{1}{4}^\circ$  NOAA optimally interpolated sea surface temperature (hereafter referred to as OISST; [42]) data, but warned users to be cognisant that different data sets would provide different kinds of information pertaining to the heat waves. Our aims here are two-fold. Firstly, we apply the MHW (MCS) definition to datasets of *in situ* and gridded SST temperature time series collected at different scales along the South African coast for three different coastal sections, each variously forced by the Agulhas and Benguela Currents and regional aspects of the coastal bathymetry and geomorphology. These regional drivers of the thermal regime (east,


south and west coast) coupled with local modifications (coastal vs. offshore) can be expected to impart different thermal signatures on the temperature data sets and manifest in differences in the metrics of MHWs (MCSs). Secondly, we aim to discuss the significance of MHWs (MCSs) within the context of the data sets' inherent differences and the various dynamical properties that then emerge because of the regional oceanographic context, so as to provide a mechanistic understanding of the nature and origin of MHWs (MCSs) in three oceanographically distinct ocean/coastal regions.

To add a mechanistic understanding of the drivers of MHWs (MCSs) manifesting in the coastal environment, we hypothesised that coastal MHW (MCS) events could either be coupled with synoptic scale processes perturbing the offshore region at scales of 100s of kms, or originate solely at a local scale as isolated incidents. Investigating the former possibility required the assessment of concurrent gridded SSTs derived from daily OISST data product, extracted for the bounding boxes in Figure 1, averaged spatially, and lagged or led by a number of days relative to the onset of the events at the coast. This analysis centres around the top three MHWs (MCSs) ranked with respect to cumulative intensity for each of 21 coastal sites. The rates of co-occurrence of coastal with mesoscale MHWs (MCSs) are used in part to understand how many of the extreme events detected in all three coastal sections originate at the coast or are artefacts of warming (cooling) in the respective currents. We think that this approach will yield considerable insight into the nature and variability of the thermal regime of nearshore seawater.

## 2. Methods

### 2.1. Study region

The variety of oceanographic features around the *ca.* 2,700 km long South African coastline provides a natural testing bed for the potential effects of different forcing mechanisms on the occurrence of MHWs and MCSs. Annual mean (SD) coastal seawater temperatures range from 12.3 (1.2)°C at the western limit near the Namibian border (Site 1) to 24.4 (2.0)°C in the east near the Mozambican border (Site 21), and our study sites were selected to cover this full range (Figure 1). We classify the coastline into three regions based on their major oceanographic features, their temperature characteristics, and aspects of the underlying continental shelf. The first is the west coast region dominated by the Benguela Current, which forms an Eastern Boundary Upwelling System (EBUS) [25]. Seasonal upwelling is maintained by prevailing south-easterly trade winds. Evenly low temperatures are especially noticeable at upwelling cells over a relatively narrow continental shelf in the region northwards of the Cape Peninsula to Cape Columbine. The west coast defines a cool temperate regime, with the range of monthly mean temperatures at most sections intermediate between cold temperate and warm temperate [30]. The second region is the warm temperate (*sensu* [30]) east coast where the influence of the south-westerly flowing warm Agulhas Current flowing tightly along the narrow continental shelf (except for the Natal Bight) is strongly felt. This stretch of coastline is spatially homogeneous with respect to temperature and characterised by a moderate amount of seasonal variation. Although the Agulhas Current retroflects back into the southern Indian Ocean [25] just south of the much wider and



**Figure 1:** Map of South Africa showing the bathymetry (only the 250m isobath is indicated), the location of the *in situ* thermal time series shown with circles and approximations of the pixels used along the shore-normal transects from the daily  $\frac{1}{4}^\circ$  NOAA OISST [42] shown with black boxes. The SST field was derived from the JPL G1SST 1 km blended SST product and shows the state of the ocean on 2016-02-14. Sites 5, 6 and 7 are to the east of the Cape Peninsula and are situated along the shores of False Bay. The Agulhas Current along the east coast of the country is visualized here in a yellowish colour as a jet of relatively warmer water projecting in a south-westerly direction, and hugging the continental shelf. The blueish patches north of the Cape Peninsula represent upwelled water. Some upwelled water may also be present around Sites 14 (Tsitsikamma) and 15–16 (Port Elizabeth).

cooler Agulhas Bank [43], its influence extends as far west as False Bay (Sites 5–21; Figure 1). The  
 115 third coastal region is that overlying the Agulhas Bank. Although also warm temperate, it experiences  
 a much larger range in annual temperature and variability compared the west and east coast regions,  
 which is in part influenced by the retention and cooling of Agulhas Current water on the bank, the  
 presence of some current-driven upwelling cells along this coastline (Sites 15–17) [43], and because of  
 the effects of embayments and capes throughout the region.

## 120 2.2. Temperature data

We use two sources of seawater temperature data. The first dataset is comprised of 127 records of  
*in situ* temperature records of daily measurements for up to 40 years in duration with a mean duration  
 of *ca.* 19 years. Whereas these *in situ* time series are generally shorter than the recommended 30 year  
 minimum [23] and have some small amounts of missing data, it is our opinion that the benefit of using  
 125 *in situ* data over satellite data is that they give a better representation of the thermal characteristics  
 near the coastline, a region where satellite SST measurements have been shown to perform poorly (e.g.  
 [45, 6]). In a South African context, Smit et al. (2013) [46] have shown that satellite SST data display  
 a warm bias as large as 6°C over *in situ* temperatures in the nearshore environment. In an attempt to  
 compromise between the proscribed requirements in Hobday et al. (2016) [23] of a 30 year minimum  
 and no missing data, all time series under 10 years in length were eliminated. Next, our 127 time series  
 were screened and those missing more than 10% of their daily values were removed, leaving a total of  
 21 time series. Care was taken to select continuous series with as few as possible consecutive missing  
 values, since having regions in the data with more than two consecutive missing data points interferes  
 with the identification of the anomalous events (see below). These stations were classified into three  
 135 coastal sections defined by properties of their oceanography and biogeography [46]. The meta data for  
 these time series and the coastal sections they were aggregated into may be found in Table S1 and the  
 site localities are displayed spatially in Figure 1.

The second set of temperature data used in this study are the daily  $\frac{1}{4}^\circ$  NOAA optimally interpolated  
 sea surface temperature (OISST; [42]) derived from the Advanced Very High Resolution Radiometer  
 140 (AVHRR). To compare the OISST and *in situ* time series, shore-normal transects were drawn from  
 each of the 21 sites extending to the 200m isobath. The OISST data were then extracted at each of the  
 roughly  $25 \times 25$  km pixels along these transects, shown as black boxes in Figure 1. Where the shelf was  
 less than 25 km wide (Sites 17–21) the nearest ‘ocean pixel’ to the *in situ* time series coordinate was  
 used. The individual time series within each pixel were then averaged along each transect corresponding  
 145 to the 21 *in situ* sites. This produced 21 OISST time series that could then be analysed for MHWs  
 (MCSs) in the same way as the *in situ* data. Note that the OISST time series had valid data covering  
 1982–2014 which did not match exactly the coverage by individual *in situ* sites.

## 2.3. Defining and calculating MHWs and MCSs

MHWs are “discrete prolonged anomalously warm water events in a particular location.” Here we  
 150 introduce the opposite but analogous concept of a Marine Cold Spell (MCS), which is calculated in

The Supplemen-  
 tary material  
 must still be put  
 together.

**Table 1:** Metrics of MHWs and their descriptions as used by Hobday et al. (2016) [23]. In the case of MCSs, values are calculated with respect to the 10th percentile and absolute intensity values are reported.

Name [unit]	Definition
Count [no. events per year]	$n$ : number of MHWs per year
Duration [days]	$D$ : Consecutive period of time that temperature exceeds the threshold
Maximum intensity [°C]	$i_{max}$ : highest temperature anomaly value during the MHW
Mean intensity [°C]	$i_{mean}$ : mean temperature anomaly during the MHW
Cumulative intensity [°C x days]	$i_{cum}$ : sum of daily intensity anomalies

the same manner as a MHW, except that events are detected as deviations below a seasonally varying anomalously low threshold relative to the site’s climatology. Although MCS intensities are calculated as negative values (i.e. anomalies) they are reported here as absolute values.

A Python script (<https://github.com/ecjoliver/marineHeatWaves>; see Hobday et al. (2016) [23]) was used to calculate the MHWs and MCSs for both the *in situ* and OISST time series, producing the metrics in Table 1. The individual events detected and their attendant statistics were meaned into a series of annual values. These annual values were then meaned for each coastal section for later comparison.

To detect the individual events, a climatological mean and 90th and 10th percentiles were calculated for each day of the year by pooling all data within an 11-day window across all years. MHWs (MCSs) were detected as periods of time when temperatures exceeded the 90th (10th) percentile for at least five days. The implication is therefore that MHWs (MCSs) could develop in winter (summer) months. Since our *in situ* time series are of differing lengths we calculated the climatology over all available years; in the case of the OISST data, climatologies were calculated over a 30-year base period from 1982–2012. Furthermore, the algorithm found discrete events with well-defined start and end dates, but ‘breaks’ between events lasting  $\leq 2$  days followed by subsequent  $\geq 5$  day events were considered as continuous events. Once events were defined, a set of metrics were calculated including maximum and mean intensity (measured as anomalies relative to the climatological mean), duration (time between start and end dates), and cumulative intensity (the integrated intensity over the duration of the event, analogous to degree-heating-days).

Because MHWs (MCSs) are thus calculated by percentiles rather than maximum values centred around a window of time with respect to the Julian day, any time of year could be shown to be experiencing a MHW (MCS). This is an important consideration as unusually warm waters occurring during the winter months of a year, the time when many species need cold water for effective spawning spore release, can have a negative effect on the recruitment success of that population for the year [48].

It is important to understand that MHWs can result from a combination of atmospheric forcing and oceanic processes, but that the approach here aims only to shed light on the oceanic drivers by virtue of the inclusion of mesoscale OISST data linked with the coastal *in situ* data sets.

In order to better understand the potential impact mesoscale phenomena have on coastal events, the rates of co-occurrence between the MHWs (MCSs) found within each time series between the two datasets were compared. This was initially done by taking each event (warm and cold) within an *in situ* time series and looking for an event occurring within the OISST time series at the same site

within a certain period of time before the *in situ* date. These co-occurrence proportions were then used to describe how often the mesoscale oceanography off the coast pre-empted the extreme events occurring along the coastline. All events occurring on dates outside of the dates occurring within the matching time series were removed from this calculation. The sum of events found to occur within the same time frame were then divided by the total number of *in situ* events checked against the OISST data to produce a co-occurrence proportion. The proportions of co-occurrence were then recalculated controlling for the amount of lag used when comparing the two different datasets for concurrent events, as well as the directionality used for this comparison. In other words, a range of lag from 2–14 was used for each site to see how far apart events generally occurred and the lag period used was also applied only after the *in situ* date, as well as both before and after the date, effectively doubling the range of the lag. This allowed us to see how often the *in situ* event pre-empted the mesoscale event as well as seeing broadly the amounts of co-occurrence occurring between the two data sets.

Besides controlling for the length and direction of lag, the size of the events themselves (ranked by cumulative intensity) were compared. This was accomplished by controlling the pool of events with which to compare the datasets per site in steps of 10th percentiles. This progressively removed smaller events until only the larger events were being compared. This allowed us to track the co-occurrence of only the largest events, reducing the overall proportion of co-occurrence found within each site as caused by the smaller events occurring at similar times as large events.

The top three MHWs (MCSs) for each *in situ* and OISST time series as defined by cumulative intensity were also noted in order to visually compare the co-occurrence of events in detail, both within and between the different datasets.

Given that the anthropogenic forcing of climate change is predicted to increase the temperature of most of the ocean over time, it stands to reason that, as a function of the 90th and 10th percentiles, one would expect to see the larger MHWs near the end of the time series, and the larger MCSs near the beginning. This can be tracked visually by looking at the top three warm and cold events for each time series. Given that the OISST time series are greater than 30 years in length it is possible to discern the long term trends within the data apart from the noise of any inter-decadal patterns (Schlegel and Smit, in review). Using a simple linear model, the decadal trend in the annual occurrence of MHWs and MCSs was calculated for all of the OISST data as well as the *in situ* time series that were over 30 years long. The shorter time series simply had the proportion of MHWs or MCSs in the first half of the time series compared against those in the second half to show if there were more or less.

### 3. Results

#### 3.1. Events

One can see in Table 2 that the *in situ* time series showed that the typically cooler west coast experienced the most MHWs per year, and that these were longer and more intense on average than those along the other two coastal sections. Whereas the east coast experienced slightly more MCSs per year than the other two coastal sections, it is the volatile south coast that experienced the longest



**Table 2:** The mean(sd) annual values for event frequency, duration and intensity for MHWs and MCSs for each coastal section as calculated from the *in situ* time series. All individual events were first aggregated into annual means before being averaged into overall mean values for each coastal section.

coast	MHW [count]	duration [days]	intensity [°C]	MCS [count]	duration [days]	intensity [°C]
all	1.6(1.8)	9.3(5.1)	2.65(0.79)	1.5(1.7)	9.0(5.1)	2.79(1.09)
west	1.8(1.9)	9.1(3.9)	2.86(0.90)	1.5(1.9)	8.5(5.2)	2.32(0.58)
south	1.5(1.8)	9.8(6.1)	2.50(0.65)	1.5(1.6)	9.7(5.5)	3.08(1.22)
east	1.5(1.7)	7.7(2.2)	2.85(0.89)	1.6(1.6)	7.1(1.9)	2.37(0.67)

and most intense MCSs. There was no significant difference between the annual count of events for each coastal section ( $F=0.727$ ,  $df_1=2$ ,  $df_2=926$ ,  $p=0.48$ ) or between the count of MHWs and MCSs ( $F=0.732$ ,  $df_1=1$ ,  $df_2=926$ ,  $p=0.39$ ). There was a significant difference between the coastal sections for the duration of the events ( $F=8.907$ ,  $df_1=2$ ,  $df_2=594$ ,  $p<0.01$ ) but no significant difference between the event types (i.e. MHWs and MCSs) themselves ( $F=0.722$ ,  $df_1=1$ ,  $df_2=594$ ,  $p=0.39$ ). The intensity of events was significantly different between each coastal section ( $F=57.55$ ,  $df_1=2$ ,  $df_2=594$ ,  $p<0.01$ ) and the type of events ( $F=5372$ ,  $df_1=1$ ,  $df_2=594$ ,  $p<0.01$ ). Note that the degrees of freedom in the results for the count of events and duration/ intensity differ as there are many years from the data in which there were no events. These were given as "0" values for the count results as this value has meaning in this context however, any years in which no events occurred had their duration and intensity results listed as "NA" rather than "0" as this would be incorrect.

Results from the analysis of the OISST data (Table 3) show that there was a significant difference from the *in situ* data for count ( $F=50.27$ ,  $df_1=1$ ,  $df_2=2306$ ,  $p<0.01$ ) and duration ( $F=17.57$ ,  $df_1=1$ ,  $df_2=1612$ ,  $p<0.01$ ) of events but not mean intensity ( $F=0.117$ ,  $df_1=1$ ,  $df_2=1612$ ,  $p=0.73$ ). The mean annual count and duration of MHWs and MCSs were greater than their *in situ* counterparts for all coastal sections whereas the intensity of both event types on all coastal sections were less than the results of the *in situ* data. With this in mind we still see that the pattern of MHW and MCS event sizes along the coastline differed from the *in situ* data, too. The largest annual number of MHWs in the OISST data occurred on the warmer east coast whereas the longest MHWs occurred on the volatile south coast with the most intense events nearly split between the west and south coasts respectively. The cooler west coast saw the most frequent occurrence of MCSs with the OISST data. The longest MCSs occurred on the south coast, same as the *in situ* data however, the most intense MCSs were seen off the east coast. There was no significant difference between the annual count of events per coast ( $F=0.542$ ,  $df_1=2$ ,  $df_2=1380$ ,  $p=0.58$ ) or event type ( $F=0.155$ ,  $df_1=1$ ,  $df_2=1380$ ,  $p=0.69$ ). The duration of the events in the different coastal sections differed significantly from one another ( $F=9.534$ ,  $df_1=2$ ,  $df_2=1018$ ,  $p<0.01$ ) whereas the duration of the event types did not ( $F=0.055$ ,  $df_1=1$ ,  $df_2=1018$ ,  $p=0.81$ ). Lastly, as with the *in situ* data, the intensity of the events from the OISST dataset differed significantly between each coastal section ( $F=16.17$ ,  $df_1=2$ ,  $df_2=1018$ ,  $p<0.01$ ) and the type of event ( $F=17645$ ,  $df_1=1$ ,  $df_2=1018$ ,  $p<0.01$ ).

**Table 3:** The mean(sd) annual values for event frequency, duration and intensity for MHWs and MCSs for each coastal section as calculated from the OISST time series. All individual events were first aggregated into annual means before being averaged into overall mean values for each coastal section.

coast	MHW [count]	duration [days]	intensity [°C]	MCS [count]	duration [days]	intensity [°C]
all	2.2(2.1)	10.2(5.4)	1.72(0.33)	2.2(2.6)	10.2(5.1)	1.83(0.52)
west	2.1(1.8)	10.9(6.7)	1.75(0.41)	2.3(2.7)	9.8(6.6)	1.87(0.61)
south	2.2(2.1)	10.6(5.5)	1.74(0.29)	2.1(2.7)	10.7(5.0)	1.79(0.45)
east	2.5(2.3)	8.3(2.4)	1.64(0.33)	2.2(2.2)	9.4(3.4)	1.93(0.61)


**Table 4:** The three largest MHWs per coast from the *in situ* data. The coast column shows in which coastal section the event occurred. The site column gives the name of the site, as seen in Figure 2, which gives the index number necessary to find it’s location along the coastline in Figure 1. The start date column gives the day on which the event began and the duration (days) column shows how many days the event lasted for. The intMean column shows the mean intensity of the event and the intCum column shows the cumulative intensity, as calculated in Table 1.

coast	site	start date	duration [days]	intMean [°C]	intCum [°C x days]
west	Sea Point	1996-01-04	40	3.08	123.20
west	Sea Point	2005-05-21	39	2.56	99.66
west	Sea Point	1975-12-30	38	2.62	99.41
south	Muizenberg	1999-12-01	98	3.17	310.30
south	Mossel Bay	1993-06-25	97	1.77	171.30
south	Muizenberg	1999-10-20	35	4.47	156.40
east	Nahoon Beach	1995-10-14	18	5.18	93.31
east	Eastern Beach	1985-12-27	19	3.33	63.18
east	Orient Beach	1990-06-25	12	3.80	45.59

### 3.2. Top three events

The mean annual statistics shown in Table 2 and Table 3 give a broad overview of the events occurring along the coastline however, examining the largest MHWs and MCSs better aids in our understanding of which coastal sections show the most variability. The ranking of these events is based on the cumulative intensity (°C x days) statistic as explained in Table 1. The three largest MHWs that occurred within the *in situ* dataset were all on the south coast (Table 4). Two of these events occurred during 1999, making it a particularly hot year. The size of the south coast events are larger than those occurring along the west coast, with the largest three MHWs on the east coast being much smaller than those occurring on the south coast (Table 4). The cumulative intensity of the entire coastline and each section individually may be calculated for both datasets from Table 2 and Table 3 by multiplying the duration by the intensity. When calculating the mean cumulative intensity of MHWs from the *in situ* data for the entire coastline by the individual events and not the annual means we see that mean(sd) is 26.11(24.37)(°C x days).

As with the MHWs, the largest three MCSs from the *in situ* data were also found on the south coast (Table 5). Maintaining the pattern seen with the MHWs, the largest MCSs on the west coast were the next largest three events for the entire coastline with the three largest MCSs from the east coast being smaller than the other two coastal sections (Table 5). The mean(sd) cumulative intensity (°C x days) for MCSs over the entire coastline was 26.45(24.25)(°C x days). The cumulative intensity (°C x days) of events from each of the coastal sections in the *in situ* data were found to be significantly different



**Figure 2:** The daily temperature values for each *in situ* time series (grey) used in this study and the corresponding OISST time series (black) extracted for comparison as seen in Figure 1. The top three MHWs are indicated by circles (with the rank inside) for each site as judged by greatest cumulative intensity. The top three MCSs for each site are indicated by squares (with the rank inside). Sites 1 – 4 represent the west coast, sites 5 – 17 represent the south coast and sites 18 – 21 represent the east coast.

**Table 5:** The three largest MCSs per coast from the *in situ* data. Column descriptions may be found in the caption for Table 4.

coast	site	start date	duration [days]	intMean [°C]	intCum [°C x days]
west	Sea Point	1990-06-23	44	2.88	126.60
west	Sea Point	1983-06-10	39	2.84	110.90
west	Sea Point	2000-11-28	23	3.70	85.04
south	Muizenberg	1984-07-14	63	2.92	183.70
south	Muizenberg	1992-03-24	56	2.78	155.60
south	Ystervarkpunt	2000-05-11	51	2.94	150.10
east	Sodwana	2004-02-12	17	3.25	55.20
east	Orient Beach	1984-03-31	13	3.73	48.44
east	Orient Beach	1995-12-6	15	3.01	45.13

**Table 6:** The three largest MHWs per coast from the OISST data. Column descriptions may be found in the caption for Table 4.

coast	site	start date	duration [days]	intMean [°C]	intCum [°C x days]
west	Sea Point	1992-01-21	39	2.96	115.60
west	Hout Bay	1992-01-20	36	3.15	113.50
west	Kommetjie	2004-10-29	53	2.03	107.40
south	Knysna	1992-05-3	50	2.41	120.40
south	Fish Hoek	2004-10-30	53	1.92	101.60
south	Pollock Beach	1994-03-27	31	3.19	99.05
east	Nahoon Beach	2006-10-21	25	1.81	45.34
east	Eastern Beach	2000-06-24	26	1.58	41.12
east	Orient Beach	2000-06-24	26	1.58	41.12

( $F=7.377$ ,  $df_1=2$ ,  $df_2=1444$ ,  $p<0.01$ ) as well as the events themselves ( $F=1753$ ,  $df_1=1$ ,  $df_2=1444$ ,  $p<0.01$ ).

As can be seen in Figure 2, the three largest events occurring for each time series within the OISST dataset are largely different from the *in situ* dataset and show a greater amount of co-occurrence for neighbouring coastal stations than the corresponding *in situ* time series. This apparent difference in the cumulative intensity (°C x days) of events between the two datasets is significant ( $F=8.857$ ,  $df_1=1$ ,  $df_2=4493$ ,  $p<0.01$ ). The pattern seen in the *in situ* data of the largest MHWs and MCSs occurring on the south, west and east coasts respectively is not repeated with the OISST dataset. Whereas the largest MHW occurred on the south coast in the OISST data, the three largest MHWs from the west coast were larger than the second and third largest events from the south coast (Table 6). The three largest MHWs from the east coast again came in below the other coasts (Table 6). The mean(sd) cumulative intensity (°C x days) over the entire coastline for the MHWs from the OISST dataset was markedly lower than its *in situ* counterpart at 18.65(15.10)(°C x days).

The largest MCSs from the OISST data (Table 7) are much less clearly ranked than their *in situ* counterparts (Table 5). The three largest MCSs on the east coast, like with both datasets and both types of events, were smaller than the other two coasts (Table 7). All of the coastal sections show that at least two of their largest MCSs occurred at the same time at different sites. The coastal mean(sd) cumulative intensity (°C x days) of the MCSs from the OISST data were much closer to their *in situ* counterparts at 23.17(23.49)(°C x days). The difference in the cumulative intensity (°C x days) of events between the coastal sections in the OISST dataset were not found to be significant ( $F=0.503$ ,  $df_1=2$ ,  $df_2=3049$ ,  $p=0.60$ ) however, the difference between the MHWs and MCSs was ( $F=3472$ ,  $df_1=1$ ,  $df_2=3049$ ,  $p<0.01$ ).

**Table 7:** The three largest MCSs per coast from the OISST data. Column descriptions may be found in the caption for Table 4.

coast	site	start date	duration [days]	intMean [°C]	intCum [°C x days]
west	Kommetjie	2010-12-13	54	3.92	211.90
west	Hout Bay	2010-12-25	41	4.06	166.30
west	Sea Point	2010-12-25	41	3.78	154.90
south	Hamburg	1984-02-5	65	3.91	254.20
south	Storms River Mouth	1982-03-13	60	2.79	167.30
south	Tsitsikamma East	1982-03-13	60	2.79	167.30
east	Eastern Beach	2010-12-26	32	2.90	92.85
east	Orient Beach	2010-12-26	32	2.90	92.85
east	Eastern Beach	1984-02-24	22	3.97	87.26


The temperature values from the dates of the largest MHW and MCS for the west and south coasts from the *in situ* data may be seen concurrently with the temperature values from the matching OISST time series in Figure 3. One may see that when the largest events were occurring in the *in situ* data, nothing of note was occurring within the OISST data.

### 3.3. Co-occurrence rates

The proportion of co-occurrence found for MHWs and MCSs between the datasets for each site may be seen in Figure 4 and Figure 5 respectively. When using the lag windows before and after the *in situ* event and comparing all events (0<sup>th</sup> percentile) we see that as the width of lag increased from 2 to 14 days the mean(sd) proportion of co-occurrence for all sites increased linearly for MHWs (0.09(0.07) to 0.38(0.20)) and MCSs (0.10(0.05) to 0.30(0.13)). Using these same constraints we see that south coast sites had the largest mean(sd) increase in co-occurrence for MHWs (0.10(0.07) to 0.45(0.18)) and MCSs (0.11(0.06) to 0.34(0.14)) whereas the west coast sites showed the smallest increase for MHWs (0.07(0.03) to 0.28(0.06)) and MCSs (0.08(0.04) to 0.19(0.02)). With all variables controlled for in the same manner, the co-occurrence rates between the different coastal sections were not significantly different for MHWs at a 2 day lag ( $F=0.799$ ,  $df_1=2$ ,  $df_2=18$ ,  $p=0.46$ ) or a 14 day lag ( $F=2.374$ ,  $df_1=2$ ,  $df_2=18$ ,  $p=0.12$ ). There were no significant differences between the coastal sections for MCSs at a 2 day lag ( $F=0.671$ ,  $df_1=2$ ,  $df_2=18$ ,  $p=0.52$ ) or 14 day lag ( $F=2.357$ ,  $df_1=2$ ,  $df_2=18$ ,  $p=0.12$ ).

The directionality of the lag also affected the co-occurrence of events. Comparing all events (0<sup>th</sup> percentile) within a 14 day lag window before the *in situ* event gave higher mean(sd) rates of co-occurrence for MHWs (0.22(0.13)) than for the same lag window after the *in situ* event for MHWs (0.18(0.10)). This same comparison for MCSs showed that the lag window before the *in situ* event (0.16(0.09)) had slightly lower rates of co-occurrence than the lag window after the *in situ* event (0.17(0.08)). When the smaller events were screened from comparison and only the largest half of the events were used (50<sup>th</sup> percentile), the difference in mean(sd) co-occurrence proportions for MHWs event lessened at 0.16(0.11) before the *in situ* event and 0.15(0.12) after. The mean(sd) co-occurrence proportion of MCSs at this level was less when using a lag window before the *in situ* event at 0.05(0.08) than for a lag window after at 0.08(0.08).

There was no co-occurrence for the largest MCSs between the datasets, whereas several of the time series on the south coast showed co-occurrence for their most extreme MHWs. Interestingly, the rates



**Figure 3:** The temperature profiles of the largest MHW and MCS from the south and west coasts respectively. The left column shows the *in situ* event while the right column shows the OISST temperature values occurring on the same dates. The top row shows the largest MHW that occurred on the south coast while the second row shows the largest MHW that occurred on the west coast. The bottom two rows show the largest MCS from the *in situ* that occurred on the south and west coasts respectively.

of co-occurrence for these largest MHWs was greater when the *in situ* event preceded the OISST event.

### 3.4. Decadal trends in MHWs and MCSs

It was found that the mean(sd) decadal trend in occurrence of MHWs in the OISST dataset is 0.5(0.3) dec<sup>-1</sup> across all sites and -0.7(0.6) dec<sup>-1</sup> for MCSs. The decadal trends in MHW occurrence increase as one moves from the west to east coasts with the mean(sd) decadal MHW trend on the west coast being 0.3(0.3) dec<sup>-1</sup>, the south coast being 0.5(0.3) dec<sup>-1</sup> and the east coast coming in at 0.6(0.2) dec<sup>-1</sup>. Just as MHWs are occurring more frequently per decade on the east coast than the west, MCSs are decreasing more frequently on the east coast than the west. The decadal trend in MCSs on the west coast is -0.1(0.6) dec<sup>-1</sup>, -0.8(0.5) dec<sup>-1</sup> on the south coast and -0.9(0.2) dec<sup>-1</sup> on the east coast.

Perhaps the trends here would be better shown as a table?

Of the 21 *in situ* time series, only 4 of them were long enough to calculate decadal trends. Of these four time series the mean(sd) decadal trend for MHWs was found to be 0.3(0.5) dec<sup>-1</sup> and -0.1(0.5) dec<sup>-1</sup> for MCSs. There were two sites from the west coast and two from the south, excluding the east coast from a possible calculation of decadal change in MHWs and MCSs. The mean(sd) decadal trend for MHWs (MCSs) on the west coast was 0.1(0.5) dec<sup>-1</sup> (-0.2(0.7) dec<sup>-1</sup>) and 0.5(0.5) dec<sup>-1</sup> (0.1(0.4) dec<sup>-1</sup>) on the south.


The other 17 *in situ* time series had the sum of the annual count of events in the first half of each time series compared against the sum of the annual count of events in the second half. This produced a proportion value that could be used as an indicator as to how many more MHWs or MCSs were occurring in the second half of the time series. The mean(sd) proportion of MHWs occurring in the second half of the shorter time series was 1.7(1.3) whereas the proportion of MCSs was 0.8(0.6). The general pattern seen in the longer time series of increasing MHW occurrence and decreasing MCS occurrence was not seen in these shorter data. The mean(sd) proportion of MHWs (MCSs) in the second half of the time series on the west coast was 1.5(0.6) (1.8(0.6)), 2.1(1.4) (0.5(0.3)) on the south coast and 0.7(0.4) (1.0(0.8)) on the east coast.

## 4. Discussion

### 4.1. Events


Having never been calculated before, it was not yet known that every time series from each coastal section of South Africa experiences on average more than one MHW and one MCS per year. It was surprising to find that the mean intensity of MCSs on the south coast was significantly larger ( $p = XX$ ) than the west coast, which is in an EBUS. Even though the south coast is known to be the most volatile of the three coasts, it was not expected that the duration (days) of both the MHWs and MCSs occurring there would be significantly longer ( $p = XX$ ) than the other two coasts. From these results we may now hypothesise that there is an additional driver on the south coast affecting the extreme events there that is not present on the other two coastal sections. It was assumed that the east coast would experience the fewest and least intense events. Whereas the duration of its events

The significance values discussed here have been calculated via Post Hoc Tukey tests and only need to be inserted



**Figure 4:** Proportion of MHW co-occurrence between *in situ* and OISST datasets for each site where sites 1 – 4 represent the west coast, sites 5 – 17 the south coast and 18 – 21 the east coast. The left column denotes the proportion of co-occurrence when events in the OISST data occurring on or before the dates of the *in situ* events were used. The left column shows the proportion of co-occurrence when OISST events that occurred after the *in situ* event dates were compared. The central column shows the overall proportion of co-occurrence. The  $x$ -axis indicates the size of the events, based on percentiles, used for calculating the co-occurrence proportions. The days of lag used, from 2 – 14, are shown here in diminishing shades of red.





**Figure 5:** Proportion of MCS co-occurrence between *in situ* and OISST datasets for each site as seen for MHWs in Figure 4. The days of lag are shown here in diminishing shades of blue.

355 were significantly smaller than the other two coasts ( $p = XX$ ), the frequency and mean intensity of its MHWs and MCSs was not the smallest found. This means that every portion of the coastline has the potential to experience an event strong enough to affect its species assemblage and/or local ecology.

The difference between the *in situ* and OISST datasets was also striking. One may see in Table 2 and Table 3 that the patterns presented by the data are intrinsically different, it is not simply a matter  
360 of the statistical significance between the statistics. The OISST data show many weak events occurring often whereas the *in situ* dataset shows fewer, stronger events. This implies that events occurring in the different datasets are unrelated, and any co-occurrence is simply by chance. Besides this difference, both datasets tend to show that the south coast has the longer and more intense results than the other two coastal sections, but this too is not a consistent result. Before calculating the proportions of  
365 co-occurrence, it was already clear from these results that the events occurring within the OISST data would differ from the *in situ* data.

Another important difference found between the datasets was that MHWs are shown to be longer and more intense in the *in situ* dataset, whereas the OISST dataset shows MCSs being longer and more intense. This also supports the argument that the events detected by these two datasets are not  
370 the same, even when they are found to occur within similar time frames. It is also counter-intuitive to what we expected to find. One would assume that as the *in situ* data are measured at *ca.* 5m deep on average, which is below the bulk surface layer (*ca.* 0.5m) that the OISST data measure [41], they would be predisposed to picking up cold upwelling events and less exposed to thermal heating, which would appear as larger MCSs and smaller MHWs compared to the OISST data. The cause of this  
375 discrepancy warrants further research.

This apparent discrepancy also places doubt on the use of MCSs as a proxy for upwelling. If the *in situ* data had recorded longer and/or more intense MCSs than the OISST data it would have shown that the MCS algorithm was detecting more extreme cold events near the coastline, where upwelling is known to occur [25, 31]. Instead the results show that offshore MCS are, on average, longer and more  
380 intense. It is the suggestion of the authors that using the MCS algorithm to detect upwelling be done with extreme caution. The MCS algorithm detects cold events based on their intensity outside of a locally produced climatology, and because most upwelling occurs at seasonally predictable times, the cold events detected here are likely due to other factors.

#### 4.2. Top three events

385 It was hypothesised that the south coast would experience the most extreme events as measured by cumulative intensity, but it was unanticipated that these events would be so much larger than the other two coastal sections. On the opposite side of the coin, it was hypothesised that the east coast would experience the least extreme events however, besides the largest two MHWs recorded, very few of the events are greater than the coastal average.

390 The disagreement between the *in situ* and OISST datasets continued into the detection of the top three events along the coastline. The pattern of event sizes within the *in situ* data are very clear in that the south coast is much more volatile than the west and east coasts in that order. The OISST

data are less conclusive on whether the south or west coast experiences the most extreme events, but it is apparent from all of the analyses from both datasets that the east coast experiences very few extreme MHWs or MCSs. Indeed, these findings support the hypothesis that the east coast is the most stable of the three coasts in that in both datasets the most extreme events occurring here barely exceed the coastal mean for cumulative intensity.

The sites along the south coast could be further divided into those within False Bay (Sites 5–7) and those on the Agulhas Bank (Sites 8–17). False Bay, which is 50 km across, is situated within the transition zone between the Benguela and Agulhas Currents [46]. Many satellite temperatures products therefore inadequately resolve the SST within this body of water (cite.). This is problematic as it is important to precisely monitor the large ranges in temperature this area experiences (cite.) as it is important both ecologically (cite.) and to the many stakeholders that use this embayment. Two of the three largest MHWs and MCSs from the *in situ* dataset were recorded within False Bay, whereas only one large MHW and no MCSs were detected with the OISST dataset. This illustrates the problem of using satellite temperature data for coastal ecology.

The example of the discrepancies for the size of the events recorded in False Bay also serves to illustrate the usefulness of satellite SST data to detect events near the coastline. For example, Roberts (2004) [43] argues for a wind forced coastal upwelling cell near Tsitsikamma (Sites 12–14). That these three sites show greater cumulative intensities for MCSs than all but one time series for the OISST dataset supports the hypothesis of such a coastal upwelling cell. This is an intriguing use of the MCS algorithm to validate multiple competing hypotheses that as of yet may not have been able to be tested in any other way.

Need to relate MCSs not showing upwelling though...

#### 4.3. Co-occurrence rates

As one may see in Figure 4, when looking at the the lag window before the *in situ* events occurred, the rates of co-occurrence for MCSs are much lower than for the MHWs. This shows that more MHWs are likely being caused by meso-scale activity than MCSs, as was expected. This finding is supported further by comparing the rates of co-occurrence for MCS lagged before and after the *in situ* event occurred. More MCS from the OISST data are shown to occur after the *in situ* events for all coastal sections. The co-occurrence rates of MHWs before and after the *in situ* events are similar.

One may also infer from the result that the proportions of co-occurrence for time series on the south coast being much larger than the other two coasts is caused by the much higher level of influence from meso-scale phenomena occurring on the Agulhas Bank. We also see that there is a higher proportion of co-occurrence for the larger MHWs and MCSs on the south coast when a lag window after the *in situ* event is used (Figure 4). This supports the argument that events originating in the nearshore are then propagating out onto the Agulhas Bank and affecting the oceanography there more often than meso-scale events originating on the Agulhas Bank are affecting the nearshore environment. The overall low rates of co-occurrence for all three coastal sections reinforces the argument that it is not the meso-scale phenomena of the open ocean around the coastline that is causing extreme events in the nearshore.

The very low proportion of co-occurrence between the datasets, and the decline in the proportion as the smaller events are screened out is strong evidence against the hypothesis that meso-scale activity, both warm and cold, is causing nearshore extreme thermal events. The small increases in co-occurrence as outlined above do imply that there is some relationship between the inshore and offshore, but that some other variable(s) is having a greater effect on the inshore. This is likely atmospheric forcing (cite.).

#### 4.4. Climate change

As MHWs and MCSs are temperature related phenomena we would be remiss not to discuss the potential of our findings in relation to climate change. The count of the MHWs and MCSs occurring throughout the coastline is less telling in this regard than the trend in these events themselves. Although the *in situ* time series used in this investigation are too short to draw adequate conclusions on the trends seen in MHWs and MCSs, one can see in Figure 2 that most sites have their top three MCSs in the first half of the time series whereas the top three MHWs occur in the second half of the time series. As the algorithm used to calculate these events is based on percentiles, it stands to reason that as the mean temperature of South Africa's coastal waters increases by 0.1°C per decade on average (Schlegel and Smit, in press), that there will be an increase in MHWs and a decrease in MCSs. The gradual mean increase in temperature will cause the algorithm used here to be biased in its detection of MHWs as time progresses simply because temperatures are generally warmer in the later half of the time series therefore, the chances of the algorithm detecting a MHW increases because the base temperature from which the MHW will be fluctuating from will be greater than the beginning of the time series. Ultimately, for the species and ecosystems experiencing this increase in duress, the semantic argument of the viability of percentiles provides little solace.

must be updated to reflect the analyses done on decadal trends for MHWs and MCSs

But we can do this using the OISST data — and we should!

Not sure how to reference this...

Nice concluding statement here!

#### 4.5. Conclusion

Given that the MHW algorithm is based on the percentiles found within each time series and not on arbitrarily decided minimum or maximum thresholds, one will always find a certain number of MHWs and MCSs. This is evident in the results of our analysis (Table 2 and Table 3) in that every time series used from both datasets experiences on average at least one MHW and MCS per year. Within each dataset, but not between, the lengths of these events are similar throughout the coastline, regardless of the local oceanographic and geographic properties. It is the cumulative intensity of the events occurring on the different coastal sections that most clearly defines them. We expected to see the most intense MCSs on the west coast as this is part of an EBUS however, the south coast, a region dominated by the warmer Agulhas Current, but with some influence from the colder Benguela, had both the most intense and longest MHWs and MCS in the *in situ* data. Even though it had been hypothesized that the south coast would have intense events, the magnitude of intensity of the events that occurred here over the other coastlines is surprising.

This statement needs to be supported by the statistical results of an ANOVA.

Also perform ANOVA on the cumulative events to support this statement statistically.

We have also shown that MCSs are not a good indicator for upwelling. As upwelling tends to occur at seasonally predictable times, the MCS algorithm does not consider these events as anomalous.

Therefore the MCSs measured here are indicators of non-seasonal or atypical forcing, which is assumed to be largely atmospheric.

As the rates of co-occurrence between *in situ* and OISST data are generally low, this implies that some other force is contributing to extreme inshore events. This is likely due to atmospheric forcing and warrants further research to better understand what is driving the occurrence and intensity of these events.

## References

- [1] , ???? Knocking back invasions: variable resistance and resilience to multiple cold spells in native vs. nonnative fishes.
- [2] Alexander, L. V., Zhang, X., Peterson, T. C., Caesar, J., Gleason, B., Klein Tank, A. M. G., Haylock, M., Collins, D., Trewin, B., Rahimzadeh, F., Tagipour, A., Rupa Kumar, K., Revadekar, J., Griffiths, G., Vincent, L., Stephenson, D. B., Burn, J., Aguilar, E., Brunet, M., Taylor, M., New, M., Zhai, P., Rusticucci, M., Vazquez-Aguirre, J. L., 2006. Global observed changes in daily climate extremes of temperature and precipitation. *Journal of Geophysical Research Atmospheres* 111 (5).
- [3] American Meteorological Society, 2011. American Meteorological Society Glossary of Meteorology. URL <http://glossary.ametsoc.org/wiki/Flood>
- [4] Black, E., Blackburn, M., Harrison, R. G., Hoskins, B. J., Methven, J., 2004. Factors contributing to the summer 2003 European heatwave. *Weather* 59 (8), 217–223. URL <http://dx.doi.org/10.1256/wea.74.04>
- [5] Bond, N. A., Cronin, M. F., Freeland, H., Mantua, N., 2015. Causes and impacts of the 2014 warm anomaly in the NE Pacific.
- [6] Castillo, K. D., Lima, F. P., 2010. Comparison of in situ and satellite-derived (MODIS-Aqua/Terra) methods for assessing temperatures on coral reefs. *Limnology and Oceanography Methods* 8, 107–117. URL <papers3://publication/uuid/214CBD69-72B9-42FC-8EBF-00467926F564>
- [7] Chen, K., Gawarkiewicz, G., Kwon, Y.-O., Zhang, W. G., 2015. The role of atmospheric forcing versus ocean advection during the extreme warming of the Northeast U.S. continental shelf in 2012. *Journal of Geophysical Research: Oceans* 120, 1–16.
- [8] Chen, K., Gawarkiewicz, G. G., Lentz, S. J., Bane, J. M., 2014. Diagnosing the warming of the Northeastern U.S. Coastal Ocean in 2012: A linkage between the atmospheric jet stream variability and ocean response. *Journal of Geophysical Research: Oceans* 119 (1), 218–227.

- [9] Crisp, D. J., 1964. The effects of the severe winter of 1962-63 on marine life in Britain. *Journal of Animal Ecology* 33 (1), 165–210.  
URL <http://www.jstor.org/stable/10.2307/2355>
- [10] DeCastro, M., Gómez-Gesteira, M., Costoya, X., Santos, F., 2014. Upwelling influence on the number of extreme hot SST days along the Canary upwelling ecosystem. *Journal of Geophysical Research: Oceans* 119 (5), 3029–3040.
- [11] Diaz, R., Rosenberg, R., 2008. Spreading dead zones and consequences for marine ecosystems. *Science* 321 (5891), 926–929.  
URL <http://www.ncbi.nlm.nih.gov/pubmed/18703733>
- [12] Easterling, D. R., Meehl, G. A., Parmesan, C., Changnon, S. A., Karl, T. R., Mearns, L. O., 2000. Climate Extremes: observations, modeling, and impacts. *Science* 289 (5487), 2068–2074.  
URL <http://www.sciencemag.org/cgi/doi/10.1126/science.289.5487.2068>
- [13] Ellis, J. M., 2010. A Quantitative Assessment of the January 2010 Cold Spell Effect on Mangrove Utilizing Coral Reef Fishes from Biscayne National Park, Florida.  
URL [http://nsuworks.nova.edu/occ{}\\_stuetaid/377/](http://nsuworks.nova.edu/occ{}_stuetaid/377/)
- [14] Feng, M., McPhaden, M. J., Xie, S.-P., Hafner, J., 2013. La Niña forces unprecedented Leeuwin Current warming in 2011. *Scientific reports* 3, 1277.  
URL <http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=3572450&tool=pmcentrez&rendertype=abstract>
- [15] Firth, L. B., Knights, A. M., Bell, S. S., 2011. Air temperature and winter mortality: Implications for the persistence of the invasive mussel, *Perna viridis* in the intertidal zone of the south-eastern United States. *Journal of Experimental Marine Biology and Ecology* 400 (1-2), 250–256.
- [16] Firth, L. B., Mieszkowska, N., Grant, L. M., Bush, L. E., Davies, A. J., Frost, M. T., Moschella, P. S., Burrows, M. T., Cunningham, P. N., Dye, S. R., Hawkins, S. J., 2015. Historical comparisons reveal multiple drivers of decadal change of an ecosystem engineer at the range edge. *Ecology and Evolution* 5 (15), 3210–3222.
- [17] Fischer, E. M., Lawrence, D. M., Sanderson, B. M., 2011. Quantifying uncertainties in projections of extremes - a perturbed land surface parameter experiment. *Climate Dynamics* 37 (7-8), 1381–1398.
- [18] Fischer, E. M., Schär, C., 2010. Consistent geographical patterns of changes in high-impact European heatwaves. *Nature Geoscience* 3 (6), 398–403.  
URL <http://www.nature.com/doi/10.1038/ngeo866>
- [19] Garrabou, J., Coma, R., Bensoussan, N., Bally, M., Chevaldonn??, P., Cigliano, M., Diaz, D., Harmelin, J. G., Gambi, M. C., Kersting, D. K., Ledoux, J. B., Lejeusne, C., Linares, C., Marschal, C., P??rez, T., Ribes, M., Romano, J. C., Serrano, E., Teixido, N., Torrents, O., Zabala, M.,

- Zuberer, F., Cerrano, C., 2009. Mass mortality in Northwestern Mediterranean rocky benthic communities: Effects of the 2003 heat wave. *Global Change Biology* 15 (5), 1090–1103.
- [20] Gershunov, A., Douville, H., 2008. Extensive summer hot and cold extremes under current and possible future climatic conditions: Europe and North America. In: *Climate Extremes and Society*. pp. 74–98.
- [21] Gunter, G., 1941. Death of Fishes Due to Cold on the Texas Coast, January, 1940. *Ecology* 22 (2), 203–208.
- URL <http://www.esajournals.org/doi/abs/10.2307/1932218>
- [22] Gunter, G., 1951. Destruction of Fishes and Other Organisms on the South Texas Coast by the Cold Wave of January 28–February 3, 1951. *Ecology* 32 (4), 731–736.
- [23] Hobday, A. J., Alexander, L. V., Perkins, S. E., Smale, D. A., Straub, S. C., Oliver, E. C., Benthuyssen, J. A., Burrows, M. T., Donat, M. G., Feng, M., Holbrook, N. J., Moore, P. J., Scannell, H. A., Sen Gupta, A., Wernberg, T., feb 2016. A hierarchical approach to defining marine heatwaves. *Progress in Oceanography* 141, 227–238.
- URL <http://www.sciencedirect.com/science/article/pii/S0079661116000057>
- [24] Holt, S. A., Holt, G. J., 1983. Cold Death of Fishes at Port Aransas, Texas: January 1982. *The Southwestern Naturalist* 28 (4), 464–466 CR – Copyright © 1983 Southwestern A.
- URL <http://www.jstor.org/stable/3670832>
- [25] Hutchings, L., van der Lingen, C. D., Shannon, L. J., Crawford, R. J. M., Verheye, H. M. S., Bartholomae, C. H., van der Plas, a. K., Louw, D., Kreiner, a., Ostrowski, M., Fidel, Q., Barlow, R. G., Lamont, T., Coetzee, J., Shillington, F., Veitch, J., Currie, J. C., Monteiro, P. M. S., 2009. The Benguela Current: An ecosystem of four components. *Progress in Oceanography* 83 (1-4), 15–32.
- URL <http://dx.doi.org/10.1016/j.pocean.2009.07.046>
- [26] Jentsch, A., Kreyling J. and Beierkuhnlein, C., 2007. A New Generation of Climate-Change Experiments: Events, Not Trends 9295 (November 2015).
- [27] Kreyling, J., Beierkuhnlein, C., Ellis, L., Jentsch, A., 2008. Invasibility of grassland and heath communities exposed to extreme weather events - Additive effects of diversity resistance and fluctuating physical environment. *Oikos* 117 (10), 1542–1554.
- [28] Laboy-nieves, E. N., Klein, E., Conde, J. E., Losada, F., Cruz, J. J., Bone, D., 2001. Mass mortality of tropical marine communities in Morrocoy.pdf. *Bulletin of Marine Science* 68 (2), 163–179.
- [29] Lima, F. P., Wetthey, D. S., 2012. Three decades of high-resolution coastal sea surface temperatures reveal more than warming. *Nature Communications* 3, 704.
- URL <http://www.ncbi.nlm.nih.gov/pubmed/22426225> <http://dx.doi.org/10.1038/ncomms1713>

- [30] Lünning, K., 1990. Seaweds: their environment, biogeography and ecophysiology. Jhon Wiley and Sons. Wiley, New York (USA).  
 URL <http://scholar.google.co.za.ezproxy.ukzn.ac.za:2048/scholar?hl=en{&}q=Satellite+versus+in+situ+measurements+at+the+air?sea+interface{&}btnG={&}as{&}sdt=1,5{&}as{&}sdt=0{&}2C5{&}#}0>
- [31] Lutjeharms, J. R. E., Cooper, J., Roberts, M., 2000. Upwelling at the inshore edge of the Agulhas Current. *Continental Shelf Research* 20 (7), 737–761.  
 URL <http://www.sciencedirect.com/science/article/pii/S0278434399000928papers3://publication/uuid/8AEE7760-040F-4E0E-AE8A-FA07C224DFA4>
- [32] Mackenzie, B. R., Schiedek, D., 2007. Daily ocean monitoring since the 1860s shows record warming of northern European seas. *Global Change Biology* 13 (7), 1335–1347.
- [33] Marsh, H., O'Shea, T. J., Best, R. C., 1986. Research on Sirenians. *Ambio* 15 (3), 177–180.  
 URL <http://www.jstor.org/stable/4313244>
- [34] Matthes, H., Rinke, A., Dethloff, K., 2015. Recent changes in Arctic temperature extremes: warm and cold spells during winter and summer. *Environmental Research Letters* 10 (11), 114020.  
 URL <http://stacks.iop.org/1748-9326/10/i=11/a=114020?key=crossref.943405a086c6db32f307f68ffb159afd>
- [35] Meehl, G. a., 2004. More Intense, More Frequent, and Longer Lasting Heat Waves in the 21st Century. *Science* 305 (5686), 994–997.  
 URL <http://www.sciencemag.org/cgi/doi/10.1126/science.1098704>
- [36] Mills, K. E., Pershing, A. J., Brown, C. J., Chen, Y., Chiang, F.-S., Holland, D. S., Lehuta, S., Nye, J. a., Sun, J. C., Thomas, A. C., Wahle, R. a., 2012. Lessons From the 2012 Ocean Heat Wave in the Northwest Atlantic. *Oceanography* 26 (2), 60–64.
- [37] Olita, A., Sorgente, R., Natale, S., Gaberšek, S., Ribotti, A., Bonanno, A., Patti, B., 2007. Effects of the 2003 European heatwave on the Central Mediterranean Sea: surface fluxes and the dynamical response. *Ocean Science* 3 (2), 273–289.  
 URL <http://www.ocean-sci.net/3/273/2007/os-3-273-2007.html>
- [38] O'Shea, T. J., Beck, C. A., Bonde, R. K., Kochman, H. I., Odell, D. K., 1985. An analysis of manatee mortality patterns in Florida, 1976-81. *The Journal of Wildlife Management* 49 (1), 1–11.  
 URL <http://www.jstor.org/stable/3801830>
- [39] Pachauri, R. K., Meyer, L., Van Ypersele, J.-P., Brinkman, S., Van Kesteren, L., Leprince-Ringuet, N., Van Boxmeer, F., 2014. Climate Change 2014 Synthesis Report.
- [40] Pearce, A. F., Feng, M., 2013. The rise and fall of the "marine heat wave" off Western Australia during the summer of 2010/2011. *Journal of Marine Systems* 111-112, 139–156.



- [41] Reynolds, R. W., Rayner, N. A., Smith, T., 2002. An improved in situ and satellite SST analysis  
605 for climate. *Journal of Climate*.  
URL [http://ams.allenpress.com/perlserv/?request=get-abstract&doi=10.1175/1520-0442\(2002\)015{T1\textless}1609:AIISAS{T1\textgreater}2.0.CO;2papers3://publication/uuid/7532476F-59A5-492D-B57A-12B92961BA4C](http://ams.allenpress.com/perlserv/?request=get-abstract&doi=10.1175/1520-0442(2002)015{T1\textless}1609:AIISAS{T1\textgreater}2.0.CO;2papers3://publication/uuid/7532476F-59A5-492D-B57A-12B92961BA4C)
- [42] Reynolds, R. W., Smith, T. M., Liu, C., Chelton, D. B., Casey, K. S., Schlax, M. G., 2007.  
610 Daily high-resolution-blended analyses for sea surface temperature. *Journal of Climate* 20 (22), 5473–5496.  
URL <http://ams.allenpress.com/perlserv/?request=get-abstract&doi=10.1175/2007JCLI1824.1papers3://publication/doi/10.1175/2007JCLI1824.1>
- [43] Roberts, M. J., 2005. Chokka squid (*Loligo vulgaris reynaudii*) abundance linked to changes in  
615 South Africa’s Agulhas Bank ecosystem during spawning and the early life cycle. *ICES Journal of Marine Science* 62 (1), 33–55.
- [44] Selig, E., Casey, K., Bruno, J. F., 2010. New insights into global patterns of ocean temperature anomalies: implications for coral reef health and management. *Global Ecology and Biogeography* 9999 (9999).  
620 URL <http://dx.doi.org/10.1111/j.1466-8238.2009.00522.x>
- [45] Smale, D. A., Wernberg, T., 2009. Satellite-derived SST data as a proxy for water temperature in nearshore benthic ecology Peer reviewed article. *Marine Biology* 387, 27–37.  
URL <http://www.vliz.be/imis/imis.php?refid=144063&pp=printpapers3://publication/uuid/3F4EF52B-247E-4406-8AD9-419D5B46A4B5>
- 625 [46] Smit, A. J., Roberts, M., Anderson, R. J., Dufois, F., Dudley, S. F. J., Bornman, T. G., Olbers, J., Bolton, J. J., 2013. A coastal seawater temperature dataset for biogeographical studies: Large biases between in situ and remotely-sensed data sets around the coast of South Africa. *PLoS ONE* 8 (12).
- [47] Sura, P., 2011. A general perspective of extreme events in weather and climate.
- 630 [48] Wernberg, T., Russell, B. D., Moore, P. J., Ling, S. D., Smale, D. A., Campbell, A., Coleman, M. A., Steinberg, P. D., Kendrick, G. A., Connell, S. D., 2011. Impacts of climate change in a global hotspot for temperate marine biodiversity and ocean warming.  
URL <http://www.sciencedirect.com/science/article/pii/S0022098111000694>
- [49] Wernberg, T., Smale, D. A., Tuya, F., Thomsen, M. S., Langlois, T. J., de Bettignies, T., Bennett, S., Rousseaux, C. S., 2013. An extreme climatic event alters marine ecosystem structure in a  
635 global biodiversity hotspot. *Nature Climate Change* 3 (1), 78–82.  
URL <http://www.nature.com/articles/nclimate1627http://dx.doi.org/10.1038/nclimate1627>

[50] Woodward, F. I., 1987. Climate and Plant Distribution. Books.googlecom 154 (2), 174.

640 URL <http://www.cambridge.org/us/catalogue/catalogue.asp?isbn=9780521282147>