

Primary drivers of marine heatwaves in the Northwest Atlantic

Primary drivers

Robert W. Schlegel, Ke Chen, and Eric C. J. Oliver

May 28th, 2020

Marine heatwaves (MHWs)


Definition

“A prolonged discrete anomalously warm water event that can be described by its duration, intensity, rate of evolution, and spatial extent.”

(Hobday et al., 2016)

- The temperature at a location exceeds the 90% percentile of that calendar day
- This occurs at least 5 days in a row with no more than a 2 day gap thereafter

Schematic



(Eric Oliver, www.marineheatwaves.org)

How does heat enter/exit the ocean?

Surface heat flux (W/m^2):

- Q_s : The short wave energy radiated from the sun (shortwave radiation)
- Q_l : The net long-wave energy radiated back from the ocean (longwave radiation)
- Q_e : The heat loss by evaporation (latent heat flux)
- Q_h : The sensible heat loss by conduction (sensible heat flux)
- $Q_{\text{net}} = Q_s + Q_l + Q_e + Q_h$ (net positive downward heat flux)

Surface movement

- $\mathbf{u}_{\text{mix}} = (u_{\text{mix}}, v_{\text{mix}})$: Horizontal velocity averaged within the mixed layer
- $\mathbf{h} = (\mathbf{/x}, \mathbf{/y})$: Horizontal gradient operator
- $\mathbf{u}_{\text{mix}} \cdot \mathbf{h} T_{\text{mix}}$ = Temperature change due to horizontal advection

Mixed layer depth

- H : MLD (mixed-layer depth)
- T_{mix} : MLD temperature
- ρ_0 : Density of sea water
- c_p : Specific heat of sea water

Surface temperature

$$\delta T_{\text{mix}} / \delta t = ((Q_{\text{net}} - (Q_s(-H))) / \rho_0 c_p H) - \mathbf{u}_{\text{mix}} \cdot \nabla_h T_{\text{mix}} + (\text{res})$$

(Benthuysen et al., 2014; Chen et al., 2015; Oliver et al., 2017)

Known drivers of past MHWs

- Advection and heat flux are the primary drivers of SST
- For much of the coastal oceans the primary driver is abnormal movement (advection) of warm currents into the nearshore
- For large seas (i.e. the Mediterranean) warm air (heat flux) plays a more important role
- The primary drivers of MHWs are generally determined by analysing the contribution of heat budget terms to the occurrence of one large event at a time


Methods

What is problematic with the current methodology?

- Large MHWs are happening too rapidly to spend a year working on them one at a time
- Smaller events may also be important
- The use of self-organising maps (SOM) to identify MHW drivers has already been tested
- This technique allows a machine to clarify for us which drivers are most prominent during hundreds of MHWs simultaneously
- We are effectively asking these two questions:
 - 1 Are there recurrent environmental patterns during marine heatwaves?
 - 2 If so, can these be detected/quantified by the computer?

Study area

- The coastline of the Northwest Atlantic, divided into six regions based on their SST and SSS climatologies (Richaud et al., 2016)




Robert W. Schlegel, Ke Chen, and Eric C. J. Oliver

Primary drivers of marine heatwaves in the Northwest Atlantic

MHW detection

- All of the SST pixels within each region were averaged together into one time series
- MHWs were calculated from these six averaged time series
- A total of 289 MHWs were detected



Robert W. Schlegel, Ke Chen, and Eric C. J. Oliver


Primary drivers of marine heatwaves in the Northwest Atlantic

Data packets

- One data packet was created for each detected MHW
- These packets consisted of the mean anomalous air/sea state during that MHW
- The physical variables in these packets were: SST, U, V, t2m, U10, V10, MLD, Q_{net}
- All 289 data packets were then fed to the SOM to produce the 12 most common synoptic air/sea patterns

Results

Results



Robert W. Schlegel, Ke Chen, and Eric C. J. Oliver

Primary drivers of marine heatwaves in the Northwest Atlantic

Conclusions

Conclusions

- The nodes tell three main stories:
 - Warm GS + air pushing up from south along coast
 - Warm air sitting over entire coast
 - Warm air being pushed over the AO from the South/Southeast onto the coast
- A few nodes tell smaller stories:
 - Short lived storms that barely manage to perturb T_{mix} for long enough to qualify as a MHW
 - These are distinctly different from the patterns associated with intense MHWs
- Overall the most intense MHWs occur during Autumn/Winter when they match patterns that are normally seen in Summer
- It is likely that this SOM technique will not work on scales larger than one meso-scale feature at a time

Acknowledgements

Acknowledgements



References

References

- Benthuysen, J., Feng, M., and Zhong, L. (2014). Spatial patterns of warming off Western Australia during the 2011 Ningaloo Niño: quantifying impacts of remote and local forcing. *Continental Shelf Research* 91, 232–246. doi:10.1016/j.csr.2014.09.014.
- Chen, K., Gawarkiewicz, G., Kwon, Y.-O., and Zhang, W. G. (2015). The role of atmospheric forcing versus ocean advection during the extreme warming of the Northeast U.S. continental shelf in 2012. *Journal of Geophysical Research: Oceans* 120, 4324–4339. doi:10.1002/2014JC010547.
- Hobday, A. J., Alexander, L. V., Perkins, S. E., Smale, D. A., Straub, S. C., Oliver, E. C., et al. (2016). A hierarchical approach to defining marine heatwaves. *Progress in Oceanography* 141, 227–238.