

1 **Climate change in coastal waters: time series properties affecting trend**
2 **estimation**

3 Robert W. Schlegel* and Albertus J. Smit

4 *Department of Biodiversity and Conservation Biology, University of the Western Cape, Robert*
5 *Sobukwe Road, Bellville, 7535, Republic of South Africa*

6 * *Corresponding author address:* Robert W. Schlegel, Department of Biodiversity and Conservation
7 Biology, University of the Western Cape, Bellville, Republic of South Africa.
8 E-mail: 3503570@myuwc.ac.za

ABSTRACT

9 In South Africa 129 *in situ* temperature time series of up to 43 years are used
10 for investigations of the thermal characteristics of coastal seawater. They are
11 collected with handheld thermometers or underwater temperature recorders
12 (UTRs) and are recorded at precisions from 0.5 °C to 0.001 °C. Using the nat-
13 ural range of seasonal signals and variability for 84 of these time series, their
14 length, decadal trend and data precision were systematically varied before fit-
15 ting generalized least squares (GLS) models to study the effect these variables
16 have on trend detection. The variables that contributed most to accurate trend
17 detection in decreasing order were: time series length, decadal trend, vari-
18 ance, percentage of missing data (%NA) and measurement precision. Time
19 series > 30 years in length are preferred, and though larger decadal trends are
20 modeled more accurately, modeled significance (p -value) is largely affected
21 by the variance present. The risk of committing both type 1 and 2 errors
22 increases when $\geq 5\%$ NA is present. There is no appreciable effect on model
23 accuracy between measurement precision of 0.1 °C to 0.001 °C. Measurement
24 precisions of 0.5 °C require longer time series to give equally accurate model
25 results. The implication is that the thermometer time series in this dataset,
26 and others around the world, must be at least two years longer than their UTR
27 counterparts to be useful for decadal scale climate change studies. Further-
28 more, adding older lower precision UTR data to newer higher precision UTR
29 data within the same time series will increase their usefulness for this purpose.

30 1. Introduction

31 The roughly 3,000 km of South Africa's coastline is bordered by the Benguela and Agulhas
32 Currents (*e.g.* Roberts 2005; Hutchings et al. 2009), which, in combination with other nearshore
33 processes, affect the country's marine coastal ecosystems (Santos et al. 2012). A thorough un-
34 derstanding of these coastal processes is provided by several physical variables, with temperature
35 being one of the main determinants (*e.g.* Blanchette et al. 2008; Tittensor et al. 2010; Couce et al.
36 2012). The statistical properties of *in situ* seawater temperature time series representing the whole
37 coastline – such as the annual mean, minimum and maximum temperature, and the thermal range
38 and variance characteristics – vary greatly among coastal sections due to the varying influences
39 of the Benguela and Agulhas Currents. Based on these thermal properties, the coastline has been
40 classified into a cool temperate west coast, a warm temperate south coast and a sub-tropical east
41 coast (Smit et al. 2013). That the ocean temperature of these regions is changing has been re-
42 ported in recent years (Mead et al. 2013). For example, an increase of 0.55°C to $0.7^{\circ}\text{C dec}^{-1}$ has
43 been reported in the Agulhas Current (Rouault et al. 2009, 2010), while the southern Benguela has
44 decreased by $0.5^{\circ}\text{C dec}^{-1}$ during some parts of the year (Rouault et al. 2010).

45 The aforementioned climate change trends were derived from remotely-sensed gridded sea sur-
46 face temperature (SST) products. Whereas newer remotely-sensed gridded SST products are ap-
47 proaching high enough resolutions for use in coastal waters, older longer products that could be
48 used for the detection of long terms trends are not (*e.g.* Chao et al. 2009; Qiu et al. 2009; Vazquez-
49 Cuervo et al. 2013). A study by Smit et al. (2013) has also shown that remotely-sensed gridded
50 SST data have a warm bias as large as 6°C when compared to coastal *in situ* data. Nevertheless,
51 a widespread approach in coastal ecological research is to use satellite and/or model-generated
52 temperature data as a representation of SST along coastlines (*e.g.* Blanchette et al. 2008; Broitman

et al. 2008; Tyberghein et al. 2012). Either the dangers of applying gridded SSTs to the coast are not widely known or in many places in the world there simply are no suitable *in situ* coastal temperature time series available. It is for this reason that we strongly recommended the use of *in situ* data to support research conducted within 400 m of the shoreline.

Where records of *in situ* coastal seawater temperature do exist, the reliability of many of these datasets that could be used in place of the remotely-sensed SST data remains to be verified. Users of remotely-sensed SST data benefit from it being refined through a number of well documented validation and quality control processes (*e.g.* Reynolds and Smith 1994; Brown et al. 1999; Martin et al. 2012), whereas the standards and methods with which local *in situ* data from a single dataset are collected and refined may differ greatly. For example, there are currently seven organizations and/or governmental departments (hereafter referred to as bodies) contributing coastal seawater temperature data to the South African Coastal Temperature Network (SACTN). These bodies use different methods and instruments to collect their data as no national standard has been set. One consequence of this methodological disparity is that two thirds of the data were sampled with hand-held thermometers that are manually recorded at a data precision of 0.5 °C, as opposed to the current generation of underwater temperature recorders (UTRs), which have an instrument precision as fine as 0.001 °C. If these *in situ* temperature data are to be used together *in lieu* of remotely-sensed SST data, it is important that the characteristics of the contributing data sources are understood in terms of their ability to yield useful, reliable and accurate long-term measurements for use in climate change studies.

This prompted us to examine the 129 *in situ* time series that comprise the SACTN. The range of measurement precisions and statistical characteristics of this dataset were used to guide a series of enquiry-driven analyses into the suitability of the time series to yield statistically significant and accurate assessments of decadal temperature change. The length, decadal trend and data

77 precision of each time series were adjusted in a systematic manner, and this forms the core of our
78 analyses. Furthermore, the natural variability of each of the time series, which differ more-or-less
79 predictably between coastlines variously affected by the Benguela and Agulhas Currents, was also
80 entered into the analysis. Our aim was to assess the effect that each of these variables has on the
81 ability of a model to produce a robust estimate of time series decadal trend. The effect gaps in the
82 time series may have on the fitting of models was also investigated as many of the time series used
83 here have some missing data scattered throughout, which is unavoidable for a 20+ year time series
84 that is sampled by hand by a single technician at each site.

85 The study provides a better understanding of some of the characteristics of a time series that are
86 influential in the detection success of decadal trends in coastal ocean temperatures.

87 **2. Methods**

88 *a. Data Sources*

89 Our study lies within the political borders of South Africa's coastline and the location of each
90 point of collection may be seen in Figure 1. Of these 129 time series, 43 are recorded with UTRs
91 and the other 86 with hand-held mercury thermometers. The oldest currently running time series
92 began on January 1st, 1972; there are 11 total time series that started in the 70s, 53 more started in
93 the 80s, 34 began in the 90s, 18 in the 00s and 13 in the current decade.

94 The data are collected using two different methods and a variety of instruments. Hand-held
95 mercury thermometers (which are being phased out in favor of alcohol thermometers or electronic
96 instruments) are used in some instances at the shoreline, and represent seawater temperatures at
97 the surface. At other places, predominantly along the country's east coast, data are collected with
98 glass thermometers from small boats at the location of shark nets along the coast (Cliff et al. 1988).

99 Whereas both types of thermometers allow for a measurement precision of 0.1 °C, the recordings
100 are written down at a precision of 0.5 °C. Data at other localities are collected using delayed-mode
101 instruments that are permanently moored shallower than 10 m, but generally very close to the
102 surface below the low-water spring tide level.

103 Over the last 40+ years the electronic instruments used to measure coastal seawater temperatures
104 have changed and improved. The previous standard was the Onset Hobo UTR with a thermal
105 precision of 0.01 °C. The new standard currently being phased in is the Starmon Mini UTR. These
106 devices have a maximum thermal precision of $0.001\text{ °C} \pm 0.025\text{ °C}$ (<http://www.star-oddi.com/>).
107 Of the 43 UTR time series in this dataset, 30 were recorded at a precision of 0.001 °C for their
108 entirety, five UTR time series include older data that were recorded at a precision of 0.01 °C or
109 0.1 °C and so have been rounded down to match this level of precision. Eight additional UTR time
110 series have data recorded at a precision of only 0.1 °C.

111 The thermometer data are recorded manually and saved in an aggregated location at the head
112 offices of the collecting bodies. UTRs are installed and maintained by divers and data are retrieved
113 at least once annually. These data are digital and are downloaded to a hard drive at the respective
114 head offices of the collecting bodies.

115 *b. Data Management*

116 Each of the seven bodies contributing data to this study have their own method of data for-
117 matting. Steps are being taken towards a national standard as we move towards replacing all the
118 thermometer recordings with UTR devices; however, as of the writing of this article, one does not
119 yet exist. Data from each organization were formatted to a project-wide comma-separated values
120 (CSV) format with consistent column headers before any statistical analyses were performed. This
121 allowed for the same methodology to be used across the entire dataset, ensuring consistent analy-

sis. Before analyzing the data they were scanned for any values above 35 °C or below 0 °C. These data points were changed to NA, meaning ‘not available’, before including them in the SACTN dataset.

All analyses and data management performed in this paper were conducted with R version 3.3.1 (2016-06-21) (R Core Team 2013). The script and data used to conduct the analyses and create the figures seen in this paper may be found at https://github.com/schrob040/Trend_Analysis.

Any time series with a temporal precision finer than one day were averaged into daily values before being aggregated into the SACTN. A series of additional checks were then performed (*e.g.* removing long stretches in the time series without associated temperature recordings) and time series shorter than five calendar years, collected deeper than 10 m or missing more than 15% of their monthly values were removed. At the time of this analysis, this usable daily dataset consisted of 84 time series with a total of 819,499 days of data; monthly averages were then made from these daily data to create the 26,924 temperature values available for use in this study.

c. Systematic Analysis of Time Series

We used the 84 time series simply for their variance properties (composed of seasonal, inter-annual, decadal and noise components), which reflect that of the thermal environment naturally present along the roughly 3,000 km of South African coastline. Linear trends that may have been present in each time series were removed prior to the ensuing analysis by applying an ordinary least squares regression and keeping the detrended residuals as anomaly time series. In doing so we avoided the need to simulate a series of synthetic time series, whose variance components may not have been fully representative of that naturally present in coastal waters. These detrended anomaly time series (henceforth simply called ‘time series’) represent a range of time scales from 72 to 519 months in duration.

145 To each of the 84 time series we artificially added linear decadal trends of 0.00°C to
146 $0.20^{\circ}\text{C dec}^{-1}$. In other words, we now had time series that captured the natural thermal vari-
147 abilities around the coast, but with their decadal trends known *a priori*. The range of decadal
148 trends was selected based around the global average of 0.124°C from Kennedy et al. (2011) and
149 used in Stocker et al. (2013). Furthermore, in order to represent the instrumental precision of the
150 instruments used to collect these time series, we rounded each of these (84 time series \times 5 decadal
151 trends) to four levels of precision: 0.5°C , 0.1°C , 0.01°C and 0.001°C . Consequently, we had a
152 pool of 1,680 time series with which to work.

153 To gain further insight into the effect of time series length on trend detection, each time series
154 was first shortened to a minimum length of 5 years, starting in January so that the timing of the
155 seasonal signal for each time series would be equitable. After fitting the model (see *Time Series*
156 *Model*, below) to all 1,680 of the shortened time series, the next year of data for each time series
157 was added and the models fitted again. This process was iterated until the full length of each time
158 series was attained. For example, if a time series consisted of 12 full years of data, it would require
159 160 models (8 iterations of increasing length \times 5 decadal trends \times 4 levels of precision); similarly,
160 720 models would be applied to a 40 year time series. Considering the 84 time series available,
161 the total number of individual models required to capture each combination of variables quickly
162 increased to 36,220.

163 Our approach of fitting models to each of the semi-artificial time series that we generated allowed
164 us to study the effect that the relevant variables (time series length, natural variability, added
165 slope and level of measurement precision) has on the ability of the time series model to faithfully
166 detect the decadal thermal trend, which was known *a priori*. The primary results of interest in
167 these analyses were the significance (*p*-value) of the model fit, the accuracy of the decadal trend
168 determined by regression, as well as the error associated with the trend estimate.

169 *d. Time Series Model*

170 The selection of the appropriate model can greatly influence the ability to detect trends (Franzke
171 2012). Two broad approaches are widely used in climate change research (Stocker et al. 2013).
172 The first group of models estimates linear trends, and although linearity may not reflect reality (*i.e.*
173 trends are very frequently non-linear), these models do provide the convenience of producing an
174 easy to understand decadal trend (*e.g.* $0.106^{\circ}\text{C dec}^{-1}$; Wilks 2011; Stocker et al. 2013). The other
175 group accommodates non-linear trajectories of temperature through time by the use of higher-
176 degree polynomial terms or non-parametric smoothing splines, but the inconvenience comes from
177 not being able to easily compare models among sites (Wood 2006; Scinocca et al. 2010). Both
178 groups of models can accommodate serially correlated residuals, which is often the cause for much
179 criticism due to their effect on the uncertainty of the trend estimates (Von Storch 1999; Santer et al.
180 2008). For example, Generalized Least Squares (GLS; yielding estimates of linear trends) and
181 Generalized Additive Mixed Models (GAMM; non-linear fitting with no trend estimate provided)
182 can both capture various degrees of serial autocorrelation (Pinheiro and Bates 2006; Wood 2006).
183 Although our exploratory analysis assessed two parametrizations of each of the model groups, we
184 opted to proceed here with a GLS equipped with a second-order autoregressive AR(2) correlation
185 structure fitted using Restricted Maximum Likelihood (REML; Pinheiro and Bates 2006):

$$y_t = \beta_0 + \beta_1 x_t + \varepsilon_t$$

186 where the lag-2 autocorrelated residuals are given by

$$\varepsilon_t = \phi_1 \varepsilon_{t-1} + \phi_2 \varepsilon_{t-2} + w_t$$

187 and the white noise series is

$$w_t \sim \text{i.i.d. } N(0, \sigma^2)$$

This approach is similar to that of the IPCC, although the latter uses an AR(1) error term (Hartmann et al. 2013). Another difference from the IPCC approach is that we nested the autoregressive component within year. This modeling approach allowed us to assess how various properties of the detrended time series would affect the models' ability to detect trends by comparing the estimates of the trends against the known artificially added trends.

3. Results

The residuals for the base 84 detrended time series may be seen in Figure 2. From these detrended time series the length, decadal trend and precision variables were systematically manipulated as explained in the methods. It was found that the important variables affecting the accuracy of the slope detected by the GLS model, in decreasing order, were: i) time series length; ii) the size of the added decadal trend; iii) initial SD of the time series (after detrending but prior to adding artificial slopes); iv) the amount of NAs; and iv) measurement precision. These variables influence the model fits in a systematic manner.

As would be expected, the size of the decadal trend estimated by the GLS increases in direct proportion to the decadal trend which we added and therefore knew *a priori*. What is especially noteworthy in this analysis is that time series of longer duration more often result in trend estimates converging with the actual trend than those of shorter length (Figure 3). This effect is most evident from around 30 years. Furthermore, how well the estimated model trend converges with the actual trend is also very visible in the standard error (SE) of the trend estimate (Figure 4): models fitted to short time series always have modeled trends with larger SE compared to longer ones. The strength of this correlation is $r = 0.56$ ($p < 0.001$) and it remains virtually unchanged as the added decadal

209 trend increases. The p -values of the fitted models also vary in relation to time series duration
210 and to the steepness of the added decadal trend (Figure 5). It is usually the longer time series
211 equipped with steeper decadal trends that are able to produce model fits with estimated trends
212 that are statistically significant. Note, however, that this p -value tests the null hypothesis that the
213 estimated trend is no different from $0\text{ }^{\circ}\text{C dec}^{-1}$ at $p \leq 0.05$, and *not* that the slope is not different
214 from the added trend. Taken together, these outcomes show that although our GLS model can very
215 often result in trend estimates that *approach* the true trend, it is seldom that those estimates are
216 statistically significant in the sense that the estimated trends differ statistically from $0\text{ }^{\circ}\text{C dec}^{-1}$.

217 The variance of the detrended data is another variable that can affect model fitting, but its only
218 systematic influence concerns the SE of the trend estimate. Here, it acts in a manner that is
219 entirely consistent across all *a priori* trends (Figure 6). What we see is that as the variance of the
220 data increases (represented here as standard deviation, SD) the SE of the slope estimates increases
221 too. Moreover, it does so disproportionately more for time series of shorter duration. Again, as we
222 have seen with the estimated trend that converges to the true trend around 30 years, so too does
223 the initial SD of the data cease to be important in time series of around 3 decades in length.

224 The number of NAs permitted in any of our time series was limited to 15% per time series.
225 Twenty-five of the 84 time series have fewer than 1% NA. An additional 45 time series have up
226 to 5% NA, 10 have up to 10% NA and 4 have up to 15% NA. The mean number of NA for the data
227 is 2.65%. The relationship between %NA and the p -value of the models is shown in Figure 7.
228 At 2.5% or fewer NA their presence does not have any discernible effect on resultant p -values.
229 Progressively increasing the number of NAs above 5%, however, leads to a drastic improvement
230 of models fitted to series with no or gently increasing decadal trends (these generally have very
231 large p -values indicative of very poor fits, perhaps due to the presence of a very weak signal), and
232 a significant deterioration of models fitted to data with steep decadal trends (for these data, the

233 model generally fits better at low numbers of NAs, as suggested by the larger number of p -values
234 that approach 0.05). In other words, the more missing values (NA) there are in a time series with no
235 discernible decadal trend, the more likely a model is to erroneously detect one. On the other hand,
236 model results from time series that do have detectable decadal trends tend to produce fits that are
237 not significantly different from $0\text{ }^{\circ}\text{C dec}^{-1}$.

238 Regarding the effect that the level of measurement precision has on the GLS models, we see in
239 Figure 8 that decreasing the precision from $0.001\text{ }^{\circ}\text{C}$ to $0.01\text{ }^{\circ}\text{C}$ has an undetectable effect on any
240 differences in the modeled trends. The Root Mean Square Error (RMSE) between the slopes esti-
241 mated from $0.001\text{ }^{\circ}\text{C}$ and $0.01\text{ }^{\circ}\text{C}$ data is 0.001. The correspondence between the slopes estimated
242 for data reported at $0.5\text{ }^{\circ}\text{C}$ compared to that at $0.001\text{ }^{\circ}\text{C}$ decreases to a RMSE of 0.03.

243 The effect of decreasing data measurement precision from $0.001\text{ }^{\circ}\text{C}$ to $0.5\text{ }^{\circ}\text{C}$ has almost no
244 appreciable effect on any of the measures of variance presented in this study. The effect of mea-
245 surement precision on the accuracy of the modeled slope, however, becomes very pronounced
246 going from $0.1\text{ }^{\circ}\text{C}$ to $0.5\text{ }^{\circ}\text{C}$. This effect is larger on smaller decadal trends. For example, at a
247 trend of $0.05\text{ }^{\circ}\text{C dec}^{-1}$, the deviation from the true value of models fitted to data with a precision
248 of $0.1\text{ }^{\circ}\text{C}$ is negligible; however, the accuracy of the fitted model on data recorded at a precision
249 of $0.5\text{ }^{\circ}\text{C}$ with a real trend of $0.05\text{ }^{\circ}\text{C dec}^{-1}$ is 10.81% different on average (*i.e.* given a slope
250 of $0.05\text{ }^{\circ}\text{C dec}^{-1}$ the model detects slopes of $0.055\text{ }^{\circ}\text{C dec}^{-1}$). This accuracy of the models im-
251 proves to an average difference of 6.44% with a slope of $0.10\text{ }^{\circ}\text{C dec}^{-1}$, 2.24% at $0.15\text{ }^{\circ}\text{C dec}^{-1}$
252 and decreases slightly to 2.30% at $0.20\text{ }^{\circ}\text{C dec}^{-1}$. A precision of $0.5\text{ }^{\circ}\text{C}$ always provides clearly
253 less accurate modeled trends than at higher precisions; however, the current analysis did not high-
254 light one precision that consistently provides the most accurate estimate of the trends. This may
255 however become determinable in an analysis of synthetic data with variance structures that are
256 manipulated in a more consistent manner.

257 As the actual time series used to generate the data for this study are predominantly over 300
258 months in length and recorded at a data precision of 0.5 °C, we would be remiss not to investigate
259 the interaction between the increase in accuracy provided by a lengthy time series, against the
260 decrease caused by a data precision of 0.5 °C. In other words, at what point does a model fitted to a
261 longer time series, with less precise measurements (*e.g.* those taken by thermometers and reported
262 at a precision of 0.5 °C), become as accurate as a time series with more precise measurements (*e.g.*
263 UTRs)? Figure 8 shows how varied the modeled trends become when a precision of 0.5 °C is used,
264 and we see here that when these low resolution time series have a shallow slope of 0.05 °C dec⁻¹,
265 a fitted model requires 24 months of additional data on average to have a comparable level of
266 accuracy to a model fitted to data recorded at a precision of 0.1 °C. The difference in the required
267 time series length necessary for accurate detection decreases to 16 months when a larger slope
268 0.20 °C dec⁻¹ is present in the data.

269 An analysis with a large number of variables as shown here is bound to have a medley of complex
270 interactions between the various statistics being measured; however, much of the range seen in the
271 results of the GLS models can be well explained by the influence of one independent variable,
272 or two operating in concert, as we have shown above. The most important of these variables has
273 clearly been the length of the time series.

274 4. Discussion

275 The strongest finding of this analysis is that the accurate detection of long-term trends in time
276 series primarily concerns the length of a dataset. But there is also a host of nuances resulting from
277 time series length, the steepness of the decadal trend the model is asked to detect, the influence
278 of the SD of a time series, the amount of missing values and the precision at which the data have
279 been measured or recorded that interact with one-another and which must be considered.

Whereas time series with smaller variances (shown as SD in this study) generally produce model fits that are statistically significant (*i.e.* with decadal trends that are significantly different from $0^{\circ}\text{C dec}^{-1}$ at $p < 0.05$) and with smaller SE of the estimated trends after shorter lengths of time, we also see that increasing a time series' length beyond 25 years, but preferably beyond 30 years, will increase the likelihood of detecting a decadal temperature change even in very variable datasets. Detecting temperature change in highly variable coastal environments, such as those around the coast of South Africa and many temperate coastal environments globally, will therefore benefit from access to the longest possible time series available. This phenomenon is demonstrated in Figure 5, which uses symbols to show the time series binned by the three different coastal sections of South Africa (Smit et al. 2013). Of these three coastal sections the east coast is known to have the most stable thermal regime (*i.e.* with the smallest variance), with the south coast having the greatest variance. Long time series at sites of low variance result in great improvements in our ability to detect significant climate change trends, and this effect is most obvious in time series with steeper decadal trends. The selection of sites for long-term monitoring must therefore account for the location of study and necessitate adequate planning to collect a long enough time series.

The detection of long-term trends requires long-term data, a fact that is already firmly established in climate change research (Ohring et al. 2005; Stocker et al. 2013). The length of these time series is firmly under the control of the investigator with sufficient foresight and perseverance to plan the installation and management of new instrument networks that will yield usable results only after about three-quarters of a typical academic career has passed. Should such data already exist – and considering the scarcity of such long-term records that are already yielding benefits today – we must ensure that these sources of data are managed and curated with great care and diligence as they are practically irreplaceable. For this reason, it is essential that we understand the inherent strengths and weaknesses of such existing sources of data so that we may fully maximize their

utility and extract from them the model coefficients needed to detect decadal temperature trends, and know the accuracy of these estimates to the best of our ability. There are many time series < 20 years in length that should be avoided, where possible, for trend analysis. These will mature with time and their maintenance needs to be ensured going forward.

Aside from length, the most powerful time series have measurements that are taken regularly. The inclusion of too many missing values (NAs) in time series must be avoided. We have shown that permitting 5% NAs or more into our time series has a drastic and significant influence on the chance of committing a type 1 error (arriving at ‘false positive,’ *i.e.* detecting a trend when none exists) for time series with no or very gentle decadal trends. On the other hand, the inclusion of NAs in time series with a decadal trend present tends to cause an increase in the probability of committing a type 2 error (*i.e.* finding ‘false negatives’). Although modern UTR time series generally have fewer NAs than we should be concerned about – therefore with a low chance of committing type 1 or type 2 errors – the presence of NAs may seriously compromise some of the time series that are still being collected by hand using hand-held thermometers.

We have demonstrated clearly that as the steepness of an expected decadal trend increases, the ability for it to be modeled accurately increases, too. Our GLS model is generally not able to detect trends that are significantly different from $0^{\circ}\text{C dec}^{-1}$ unless a slope of $0.20^{\circ}\text{C dec}^{-1}$ exists. Very rarely were we able to produce significant model fits at shallower slopes. No trends with a slope < $0.05^{\circ}\text{C dec}^{-1}$ were found to be significant in this study. Based on the relationship between SD and the added decadal trend, we see that time series with an SD of 1.5°C (the bulk of the time series here) and a decadal trend of $0.10^{\circ}\text{C dec}^{-1}$ should consist of roughly 640 months of data before our GLS model would regularly be able to detect a significant trend ($p < 0.05$). This finding is somewhat discouraging as most global analyses of decadal SST change based on gridded SST products estimate a trend closer to $0.1^{\circ}\text{C dec}^{-1}$ (*e.g.* Stocker et al. 2013). This means that the

trends present in most time series representative of very variable coastal environments that exhibit the same variance structure as that of our data are probably unlikely to be detected as significant, even if they do indeed exist. In other words, the chance of committing a type 2 error is probably very real for such systems, unless time series > 50 years are available.

As 50 year coastal seawater temperature time series are probably very scarce, it is important to note that those measured at precisions of $0.1\text{ }^{\circ}\text{C}$ to $0.001\text{ }^{\circ}\text{C}$ require fewer months of data to detect long term trends. Based on the data presented here, we calculated that time series measured at a low precision ($0.5\text{ }^{\circ}\text{C}$) may require as much as an additional 24 months of data to accurately detect long-term trends. One of the motivators for this paper was to investigate the effect measurement precision has on a time series' ability to produce results useful for investigations of long-term climate change, and to validate the use of the low precision $0.5\text{ }^{\circ}\text{C}$ thermometer data. This is an important consideration as many studies investigating the effects of climate change (*e.g.* Grant et al. 2010; Scherrer and Körner 2010; Lathlean and Minchinton 2012) do use lower precision $0.1\text{ }^{\circ}\text{C}$ data. Whereas the precision of much of our data is below the current standard of $0.1\text{ }^{\circ}\text{C}$ required for climate change research (Ohring et al. 2005; Jarraud 2008), the length of the thermometer time series makes them a valuable asset. The average length of the thermometer time series in the SACTN, from which the 84 time series used in this study were drawn, is 349 months. The average length of the UTR time series is 167 months. Given this difference in the lengths of the time series, even after correcting for the negative effect of low measurement precision, the time series collected with thermometers are currently more useful for climate change research than the UTR time series within the SACTN. Because time series with data precisions of $0.1\text{ }^{\circ}\text{C}$ to $0.001\text{ }^{\circ}\text{C}$ produce comparable results, newer higher precision UTR data may be combined with older lower precision UTR data within the same time series without concern that the reduced overall data precision may have a negative impact on a model's ability to detect decadal trends. Extending

time series in this way will serve to make them more dependable as length is the primary criterion through which one should initially assess the potential to accurately detect a decadal trend before refining ones assumptions with any statistical analyses. A time series with data precision finer than 0.1 °C is therefore only necessary when an investigation requires that the decadal trend be known to an accuracy of 0.01 °C or finer (*e.g.* Karl et al. 2015).

It is important to take note of the accuracy of the models, not only to focus on the significance of their results. Indeed, the *p*-value given for the slope in a model does not show how well the model detects the true trend in the data (known *a-priori* in this study); rather, it tells us if the detected trend is significantly different from 0 °C dec⁻¹. This is not particularly useful for applying the results of climate change research more broadly to biotic interests. For example, of the 1,344 models (84 base time series × 4 decadal trends × 4 levels of precision) fitted to time series with decadal trends ≥ 0.05 °C dec⁻¹, 317 of these were accurate to within 10% of the decadal trend known *a priori*, but not significant ($p \geq 0.05$). That a long term trend does exist, may be accurately detected by a model and related to an observed change in the natural world – such as range expansion/contraction of coastal biota (Bolton et al. 2012; Straub et al. 2016; Wernberg et al. 2016) – is more important than whether or not the model can show if that trend is significantly different from 0 °C dec⁻¹ in a statistical sense.

We must mention also that much of the meta-data pertaining to the older temperature records used here have over time been lost. Unlike the bulk of the International Comprehensive Ocean-Atmosphere Data Set (ICOADS; Freeman et al. 2016), *in situ* coastal seawater temperature monitoring that started in the 1970s in South Africa was not developed with climate change research in mind, and comprehensive records that keep track of details of the instruments used, calibration, their turnover, change in monitoring methods and locations and so forth are not always available as per modern requirements (Aguilar et al. 2003). For studies of climate change *per se* this is a

serious limitation and it prevents us from knowing anything about the accuracy of the instruments or potential issues of drift (stability) that may have occurred. We do know however that all time series sampled with thermometers were sampled only with thermometers, and *vice versa* for the UTR time series, ensuring that the precisions of the measured data used in this study are correct. Moving forward with the further development of the SACTN and the establishment of a national standard of data collection and instrument maintenance, we are able to record and archive all these levels of pertinent meta-data, and allowing for the enforcement of SI traceability and the accurate measurement of instrument drift (Jarraud 2008). Nevertheless, the detrended anomaly time series used here were taken only for their variance properties, which we think accurately reflect those of the three different coastal sections. They provide a strong backbone for semi-artificial time series, and we have shown how important insights about model fitting could be derived from these data.

5. Conclusion

We draw several key conclusions:

1. There is a rapid increase in the accuracy and significance of modeled trends as time series lengths extend from 10 to 20 yr. This improvement slows from 20 to 30 yr, and as time series approach 40 yr in length the accuracy of models becomes nearly exact. Modeled trends from time series at or under 25 yr in length should be interpreted with extreme caution.
2. For our variable coastal seawater, a time series of 520 months in length is required to detect a decadal trend in line with the global average (*i.e.* near $0.1\text{ }^{\circ}\text{C dec}^{-1}$) with perfect accuracy; however, an additional 120 months of data is often required for the detected trend to be considered significant ($p \leq 0.05$).

- 397 3. The length of a time series required to detect a decadal trend at $0.10^{\circ}\text{C dec}^{-1}$ may rapidly
398 exceed 100 yr when a large amount of variance is present.
- 399 4. The larger the decadal trend within a time series, the more accurately it will be modeled
400 regardless of the amount of variance in the time series.
- 401 5. There is a complicated relationship between the accuracy of a trend fitted to a time series and
402 the %NA of that time series. As the %NA increases, so too does the chance of committing type
403 1 (with gentle trends) or type 2 errors (with steeper trends).
- 404 6. A measurement precision finer than 0.5°C is not required to confidently detect the long-term
405 trend in a time series; however, precisions at or finer than 0.1°C may reduce the length of
406 time required to accurately detect a long term trend, if one does exist, by as much as two yr.
- 407 7. Improving the precision of measurements finer than 0.1°C has almost no appreciable effect
408 on a models ability to detect a long-term trend, provided that the reported effect size matches
409 the level of precision by the instruments.

410 We understand that time series of >30 yr may be exceedingly rare. Therefore, as we move
411 forward as a scientific community investigating the issues of climate change, the continuity of
412 any current time series of sufficient length must be ensured as these commodities are practically
413 irreplaceable.

414 *Acknowledgments.* The authors would like to thank DAFF, DEA, EKZNW, KZNSB, SAWS and
415 SAEON for their contributions of the raw data used in this study. Without it, this article and the
416 SACTN would not be possible. This research was supported by NRF Grant (CPRR14072378735).
417 The authors report no financial conflicts of interests.

References

- Aguilar, E., I. Auer, M. Brunet, T. Peterson, and J. Wieringa, 2003: Guidelines on climate metadata and homogenization, wmo-td no. 1186 (wcdmp2). *World Meteorological Organization, Geneva*.
- Blanchette, C. A., C. Melissa Miner, P. T. Raimondi, D. Lohse, K. E. K. Heady, and B. R. Broitman, 2008: Biogeographical patterns of rocky intertidal communities along the Pacific coast of North America. *Journal of Biogeography*, **35** (9), 1593–1607.
- Bolton, J. J., R. J. Anderson, A. J. Smit, and M. D. Rothman, 2012: South African kelp moving eastwards: the discovery of *Ecklonia maxima* (Osbeck) Papenfuss at De Hoop Nature Reserve on the south coast of South Africa. *African Journal of Marine Science*, **34** (1), 147–151.
- Broitman, B. R., N. Mieszkowska, B. Helmuth, and C. A. Blanchette, 2008: Climate and recruitment of rocky shore intertidal invertebrates in the eastern North Atlantic. *Ecology*, **89** (11 Suppl), S81–90.
- Brown, O. B., P. J. Minnett, R. Evans, E. Kearns, K. Kilpatrick, A. Kumar, R. Sikorski, and A. Závody, 1999: MODIS Infrared Sea Surface Temperature Algorithm Algorithm Theoretical Basis Document Version 2.0. *University of Miami*, 31 098–33 149.
- Chao, Y., Z. Li, J. D. Farrara, and P. Hung, 2009: Blending sea surface temperatures from multiple satellites and *in situ* observations for coastal oceans. *Journal of Atmospheric and Oceanic Technology*, **26** (7), 1415–1426, doi:10.1175/2009JTECHO592.1.
- Cliff, G., S. F. J. Dudley, and B. Davis, 1988: Sharks caught in the protective gill nets off Natal, South Africa. 1. The sandbar shark *Carcharhinus plumbeus* (Nardo). *South African Journal of Marine Science*, **7** (1), 255–265.

- 439 Couce, E., A. Ridgwell, and E. J. Hendy, 2012: Environmental controls on the global distribution
440 of shallow-water coral reefs. *Journal of Biogeography*, **39** (8), 1508–1523.
- 441 Franzke, C., 2012: Nonlinear trends, long-range dependence, and climate noise properties of sur-
442 face temperature. *Journal of Climate*, **25** (12), 4172–4183, doi:10.1175/JCLI-D-11-00293.1.
- 443 Freeman, E., and Coauthors, 2016: ICOADS release 3.0: a major update to the historical marine
444 climate record. *International Journal of Climatology*, 1–22.
- 445 Grant, O. M., L. Tronina, J. C. Ramalho, C. Kurz Besson, R. Lobo-Do-Vale, J. Santos Pereira,
446 H. G. Jones, and M. M. Chaves, 2010: The impact of drought on leaf physiology of *Quercus*
447 *suber* L. trees: Comparison of an extreme drought event with chronic rainfall reduction. *Journal*
448 *of Experimental Botany*, **61** (15), 4361–4371, doi:10.1093/jxb/erq239.
- 449 Hartmann, D., and Coauthors, 2013: Observations: Atmosphere and surface supplementary mate-
450 rial. *in: Climate Change 2013: The Physical Science Basis*
- 451 Hutchings, L., and Coauthors, 2009: The Benguela Current: an ecosystem of four components.
452 *Progress in Oceanography*, **83** (1-4), 15–32, doi:10.1016/j.pocean.2009.07.046.
- 453 Jarraud, M., 2008: Guide to meteorological instruments and methods of observation (*wmo-no.* 8).
454 *World Meteorological Organisation, Geneva.*
- 455 Karl, T. R., and Coauthors, 2015: Possible artifacts of data biases in the recent global surface
456 warming hiatus. *Science*, **348** (6242), 1469–1472, doi:10.1126/science.aaa5632.
- 457 Kennedy, J. J., N. A. Rayner, R. O. Smith, M. Saunby, and D. E. Parker, 2011: Reassessing biases
458 and other uncertainties in sea-surface temperature observations measured *in situ* since 1850, Part
459 1: measurement and sampling uncertainties. *Journal of Geophysical Research Atmospheres*,
460 **116**.

- 461 Lathlean, J. A., and T. E. Minchinton, 2012: Manipulating thermal stress on rocky shores to pre-
462 dict patterns of recruitment of marine invertebrates under a changing climate. *Marine Ecology*
463 *Progress Series*, **467**, 121–136, doi:10.3354/meps09996.
- 464 Martin, M., and Coauthors, 2012: Group for High Resolution Sea Surface temperature (GHRSSST)
465 analysis fields inter-comparisons. Part 1: a GHRSSST multi-product ensemble (GMPE). *Deep*
466 *Sea Research Part II: Topical Studies in Oceanography*, **77-80**, 21–30.
- 467 Mead, A., and Coauthors, 2013: Human-mediated drivers of change — impacts on coastal ecosys-
468 tems and marine biota of South Africa. *African Journal of Marine Science*, **35 (3)**, 403–425.
- 469 Ohring, G., B. Wielicki, R. Spencer, B. Emery, and R. Datla, 2005: Satellite instrument calibration
470 for measuring global climate change. *Bulletin of the American Meteorological Society*, **86 (9)**,
471 1303–1313.
- 472 Pinheiro, J., and D. Bates, 2006: *Mixed-effects models in S and S-PLUS*. Springer Science &
473 Business Media.
- 474 Qiu, C., D. Wang, H. Kawamura, L. Guan, and H. Qin, 2009: Validation of AVHRR and TMI-
475 derived sea surface temperature in the northern South China Sea. *Continental Shelf Research*,
476 **29 (20)**, 2358–2366.
- 477 R Core Team, 2013: *R: A Language and Environment for Statistical Computing*. Vienna, Austria,
478 R Foundation for Statistical Computing, URL <http://www.r-project.org/>.
- 479 Reynolds, R. W., and T. M. Smith, 1994: Improved global sea surface temperature analyses using
480 optimum interpolation. *Journal of Climate*, **7 (6)**, 929–948.

481 Roberts, M. J., 2005: Chokka squid (*Loligo vulgaris reynaudii*) abundance linked to changes in
 482 South Africa's Agulhas Bank ecosystem during spawning and the early life cycle. *ICES Journal*
 483 *of Marine Science*, **62** (1), 33–55, doi:10.1016/j.icesjms.2004.10.002.

484 Rouault, M., P. Penven, and B. Pohl, 2009: Warming in the Agulhas Current system since the
 485 1980's. *Geophysical Research Letters*, **36** (12), 1–5.

486 Rouault, M., B. Pohl, and P. Penven, 2010: Coastal oceanic climate change and variability from
 487 1982 to 2009 around South Africa. *African Journal of Marine Science*, **32** (2), 237–246.

488 Santer, B. D., and Coauthors, 2008: Consistency of modelled and observed temperature trends in
 489 the tropical troposphere. *International Journal of Climatology*, **28** (13), 1703–1722.

490 Santos, F., M. Gomez-Gesteira, M. DeCastro, and I. Alvarez, 2012: Differences in coastal and
 491 oceanic SST trends due to the strengthening of coastal upwelling along the Benguela current
 492 system. *Continental Shelf Research*, **34**, 79–86.

493 Scherrer, D., and C. Körner, 2010: Infra-red thermometry of alpine landscapes challenges climatic
 494 warming projections. *Global Change Biology*, **16** (9), 2602–2613, doi:10.1111/j.1365-2486.
 495 2009.02122.x.

496 Scinocca, J., D. B. Stephenson, T. C. Bailey, J. Austin, and Coauthors, 2010: Estimates of past and
 497 future ozone trends from multimodel simulations using a flexible smoothing spline methodol-
 498 ogy. *Journal of Geophysical Research: Atmospheres*, **115** (D3).

499 Smit, A. J., M. Roberts, R. J. Anderson, F. Dufois, S. F. J. Dudley, T. G. Bornman, J. Olbers, and
 500 J. J. Bolton, 2013: A coastal seawater temperature dataset for biogeographical studies: Large
 501 biases between *in situ* and remotely-sensed data sets around the coast of South Africa. *PLoS*
 502 *ONE*, **8** (12), doi:10.1371/journal.pone.0081944.

503 Stocker, T., and Coauthors, 2013: *Climate change 2013: the physical science basis. Contribution*
504 *of working group I to the fifth assessment report of the intergovernmental panel on climate*
505 *change*. Cambridge University Press.

506 Straub, S. C., M. S. Thomsen, and T. Wernberg, 2016: The dynamic biogeography of the an-
507 thropocene: the speed of recent range shifts in seaweeds. *Seaweed Phylogeography*, Springer,
508 63–93.

509 Tittensor, D. P., C. Mora, W. Jetz, H. K. Lotze, D. Ricard, E. V. Berghe, and B. Worm, 2010:
510 Global patterns and predictors of marine biodiversity across taxa. *Nature*, **466 (7310)**, 1098–
511 1101, doi:10.1038/nature09329.

512 Tyberghein, L., H. Verbruggen, K. Pauly, C. Troupin, F. Mineur, and O. De Clerck, 2012: Bio-
513 ORACLE: a global environmental dataset for marine species distribution modelling. *Global*
514 *Ecology and Biogeography*, **21 (2)**, 272–281.

515 Vazquez-Cuervo, J., B. Dewitte, T. M. Chin, E. M. Armstrong, S. Purca, and E. Alburquerque,
516 2013: An analysis of SST gradients off the Peruvian coast: the impact of going to higher reso-
517 lution. *Remote Sensing of Environment*, **131**, 76–84, doi:10.1016/j.rse.2012.12.010.

518 Von Storch, H., 1999: Misuses of statistical analysis in climate research. *Analysis of Climate*
519 *Variability*, Springer, 11–26.

520 Wernberg, T., and Coauthors, 2016: Climate-driven regime shift of a temperate marine ecosystem.
521 *Science*, **353 (6295)**, 169–172.

522 Wilks, D. S., 2011: *Statistical methods in the atmospheric sciences*, Vol. 100. Academic press.

523 Wood, S., 2006: *Generalized additive models: an introduction with R*. CRC press.

524

LIST OF FIGURES

525	Fig. 1.	Map of South Africa indicating the location of the 129 time series currently comprising the South African Coastal Temperature Network. The location of the 84 time series used in this study are shown as solid white circles and the unused time series as opaque.	27
526			
527			
528	Fig. 2.	Box and whisker plot summarizing the 84 anomaly time series used in this study (<i>i.e.</i> after detrending) but before adding a decadal trend or rounding the data. The plot indicates the first and third quartile as the extremities of the boxes, the median is shown as the horizontal line within each box, the minima and maxima are indicated by the whiskers and the points are outliers.	28
529			
530			
531			
532			
533	Fig. 3.	The effect of time series length on the ability of the GLS model to accurately detect the trend added to each time series. The box and whisker plot shows the first and third quartile as the extremities of the boxes, the median is shown as the horizontal line within each box, and the minima and maxima are indicated by the whiskers. Points indicate the spread of the actual data and their size are scaled according to the length of the time series they represent.	29
534			
535			
536			
537			
538	Fig. 4.	The relationship between the length of a time series, the size of the modeled trend and its standard error (SE). Each individual line shows the modeled trend for 1 of the 84 sites used in this analysis to which a model was fitted iteratively as the time series length was ‘grown’ from 5 yr in length to the maximum duration available for the site. The panels show the progressive effect that decadal trend has on this relationship (indicated by the numeral above each panel), and the gray shade shown on the right, is mapped to the SE of the trend.	30
539			
540			
541			
542			
543			
544	Fig. 5.	The effect of the SD ($^{\circ}\text{C}$) of the anomaly time series before adding a decadal trend (initial SD), or rounding the data to the different levels of precision, on the significance of the modeled trend. The size of the symbols are scaled in direct proportion to the time series length and are shown at the top of the figure. Time series belonging to the three South African coastal sections are represented with shapes. The east coast (ec) typically has the most stable thermal regime of the three coasts, with the south coast (sc) having the greatest amount of variance and the west coast (wc) consisting of areas with both high and low variance. Linear models with 95% confidence intervals (indicated by gray ribbons) have been fitted separately for each coastal section, and illustrate the interaction between Initial SD in each group and the significance (p -value) of the GLS models. The panels, from top to bottom, show increasing decadal trends as indicated by numerals above the panels.	31
545			
546			
547			
548			
549			
550			
551			
552			
553			
554			
555	Fig. 6.	The relationship between the effect of initial SD ($^{\circ}\text{C}$) (<i>i.e.</i> the variance of the anomaly time series before adding artificial decadal trends; shown here in shades of gray), on the standard error (SE) of a modeled trend, controlled for by the length of the time series. The effect of the size of the added decadal trends on the relationship is imperceptible and therefore only a decadal trend of $0.20^{\circ}\text{C dec}^{-1}$ is presented.	32
556			
557			
558			
559			
560	Fig. 7.	The relationship between the amount of missing values (log %NA) and the significance of a modeled trend (p value). Each panel shows the effect of an increasingly larger amount of missing values indicated above each panel by numerals (from top left to bottom right): 1, 2.5, 5, 10, 12.5, and 15. The fitted black lines and 95% confidence intervals (shown as gray-shaded bands) represent each of the five decadal trends assessed ($^{\circ}\text{C decade}^{-1}$) shown using different black symbols: 0 (circles), 0.05 (triangles), 0.1 (squares), 0.15 (crosses), and 0.2 (squares with an \times symbol inside).	33
561			
562			
563			
564			
565			
566			
567	Fig. 8.	Plots representing correlations of the modeled trends acquired at different levels of rounding, which can be interpreted as representations of different measurement precisions. The effect	
568			

569 of rounding from 0.001 °C to 0.01 °C may be seen in the panel on the left. The panel on the
570 right shows that rounding from a precision of 0.001 °C to 0.5 °C has a visibly greater effect
571 on the deterioration of the correlation between the two sets of estimated trends. 34




FIG. 1. Map of South Africa indicating the location of the 129 time series currently comprising the South African Coastal Temperature Network. The location of the 84 time series used in this study are shown as solid white circles and the unused time series as opaque.




FIG. 2. Box and whisker plot summarizing the 84 anomaly time series used in this study (*i.e.* after detrending) but before adding a decadal trend or rounding the data. The plot indicates the first and third quartile as the extremities of the boxes, the median is shown as the horizontal line within each box, the minima and maxima are indicated by the whiskers and the points are outliers.




FIG. 3. The effect of time series length on the ability of the GLS model to accurately detect the trend added to each time series. The box and whisker plot shows the first and third quartile as the extremities of the boxes, the median is shown as the horizontal line within each box, and the minima and maxima are indicated by the whiskers. Points indicate the spread of the actual data and their size are scaled according to the length of the time series they represent.




FIG. 4. The relationship between the length of a time series, the size of the modeled trend and its standard error (SE). Each individual line shows the modeled trend for 1 of the 84 sites used in this analysis to which a model was fitted iteratively as the time series length was ‘grown’ from 5 yr in length to the maximum duration available for the site. The panels show the progressive effect that decadal trend has on this relationship (indicated by the numeral above each panel), and the gray shade shown on the right, is mapped to the SE of the trend.




FIG. 5. The effect of the SD (°C) of the anomaly time series before adding a decadal trend (initial SD), or rounding the data to the different levels of precision, on the significance of the modeled trend. The size of the symbols are scaled in direct proportion to the time series length and are shown at the top of the figure. Time series belonging to the three South African coastal sections are represented with shapes. The east coast (ec) typically has the most stable thermal regime of the three coasts, with the south coast (sc) having the greatest amount of variance and the west coast (wc) consisting of areas with both high and low variance. Linear models with 95% confidence intervals (indicated by gray ribbons) have been fitted separately for each coastal section, and illustrate the interaction between Initial SD in each group and the significance (p -value) of the GLS models. The panels, from top to bottom, show increasing decadal trends as indicated by numerals above the panels.




FIG. 6. The relationship between the effect of initial SD ($^{\circ}\text{C}$) (*i.e.* the variance of the anomaly time series before adding artificial decadal trends; shown here in shades of gray), on the standard error (SE) of a modeled trend, controlled for by the length of the time series. The effect of the size of the added decadal trends on the relationship is imperceptible and therefore only a decadal trend of $0.20^{\circ}\text{C dec}^{-1}$ is presented.




FIG. 7. The relationship between the amount of missing values (log %NA) and the significance of a modeled trend (p value). Each panel shows the effect of an increasingly larger amount of missing values indicated above each panel by numerals (from top left to bottom right): 1, 2.5, 5, 10, 12.5, and 15. The fitted black lines and 95% confidence intervals (shown as gray-shaded bands) represent each of the five decadal trends assessed ($^{\circ}\text{C decade}^{-1}$) shown using different black symbols: 0 (circles), 0.05 (triangles), 0.1 (squares), 0.15 (crosses), and 0.2 (squares with an \times symbol inside).




FIG. 8. Plots representing correlations of the modeled trends acquired at different levels of rounding, which can be interpreted as representations of different measurement precisions. The effect of rounding from 0.001 °C to 0.01 °C may be seen in the panel on the left. The panel on the right shows that rounding from a precision of 0.001 °C to 0.5 °C has a visibly greater effect on the deterioration of the correlation between the two sets of estimated trends.