

1 **Effects of Natural Variability of Seawater Temperature, Time Series Length,**
2 **Decadal Trend, and Instrument Precision on the Ability to Detect**
3 **Temperature Trends**

4 Robert Schlegel* and Albertus Smit

5 *Department of Biodiversity and Conservation Biology, University of the Western Cape, Bellville,*
6 *Republic of South Africa*

7 **Corresponding author address:* Robert Schlegel, Department of Biodiversity and Conservation
8 Biology, University of the Western Cape, Bellville, Republic of South Africa.
9 E-mail: 3503570@myuwc.ac.za

ABSTRACT

10 In South Africa 129 *in situ* temperature time series of 1 to 43 years are
11 used for investigations of the thermal characteristics of coastal seawater. They
12 are comprised of temperature recordings at precisions ranging from 0.5 °C to
13 0.001 °C and collected with handheld thermometers or underwater tempera-
14 ture recorders (UTRs). Using the naturally occurring range of seasonal sig-
15 nals, variability and temperature trends for 84 of these time series, the length,
16 decadal trend and data precision of each time series were systematically var-
17 ied before fitting generalized least squares (GLS) models to study the effect
18 these variables have on trend detection. We determined that low instrument
19 precision has less effect on the ability of a model to detect trends within a
20 time series than do the length, variance or the decadal trend itself, with length
21 contributing the most to trend detection. We found that time series at least 20
22 years in length may be used tentatively for climate change research, but that
23 time series >30 years in length are preferable. The implication is that long-
24 running thermometer time series in this dataset, and others around the world,
25 are more useful for decadal scale climate change studies than the shorter, more
26 precise UTR time series.

27 1. Introduction

28 The roughly 3,000 km of South Africa's coastline is bordered by the Benguela and Agulhas
29 currents (e.g. Roberts 2005; Hutchings et al. 2009), which, in combination with other nearshore
30 processes, affect the country's marine coastal ecosystems (Santos et al. 2012). A thorough under-
31 standing of these coastal processes is provided by several physical variables, of which temperature
32 is one of the main determinants (e.g. Blanchette et al. 2008; Tittensor et al. 2010; Couce et al.
33 2012). In order to ensure a true representation of organisms' biological thermal limits, nearshore
34 temperatures must be accurately recorded and monitored. Some sources warn of the pitfalls in
35 doing so *RWS: Add references here showing which sources say using SST for the coast is inappro-*
36 *priate*, and a study by Smit et al. (2013) showed that SST data have a warm bias as large as 6 °C
37 when compared to coastal *in situ* data. Nevertheless, a widespread approach in coastal ecolog-
38 ical research is to use satellite and/or model-generated temperature data as representation of the
39 sea surface temperature (SST) along coastlines (e.g. Blanchette et al. 2008; Broitman et al. 2008;
40 Tyberghein et al. 2012), because apparently the dangers of applying gridded SSTs to the coast
41 are not widely known or in many places in the world there simply are no suitable *in situ* coastal
42 temperature time series available. It is for this reason that we strongly recommended the use of *in*
43 *situ* data to support research conducted within 400 m from the shoreline.

44 Where records of *in situ* coastal seawater temperature do exist, the reliability of many of these
45 datasets that could be used in place of the remotely-sensed SST data remains to be verified. Users
46 of SST data benefit from it being refined through a number of well documented validation and
47 quality control processes (e.g. Reynolds and Smith 1994; Brown et al. 1999; Martin et al. 2012),
48 whereas the standards and methods with which local *in situ* data from a single dataset are collected
49 and refined may differ greatly. For example, there are currently seven organizations and/or govern-

50 mental departments (hereafter referred to as bodies) contributing coastal seawater temperature data
51 to the South African Coastal Temperature Network (SACTN). These bodies use different methods
52 and instruments to collect their data as no national standard has been set. One consequence of this
53 methodological disparity is that two thirds of the data were sampled with hand-held thermometers
54 that are manually recorded at a data precision of 0.5 °C, as opposed to the current generation of
55 Underwater Temperature Recorders (UTRs) with an instrument precision of up to 0.001 °C. If
56 these *in situ* data are to be used together *in lieu* of the satellite-based SST data, it is important that
57 the characteristics of the contributing data sources are understood in terms of their ability to yield
58 useful, reliable and accurate long-term measurements for use in climate change studies.

59 This prompted us to examine the 129 *in situ* time series that comprise the SACTN, whose loca-
60 tions and instrument types may be seen in Figure A1. The range of data precision and statistical
61 characteristics found within this dataset were used to guide a series of enquiry-driven analyses into
62 the suitability of the time series to yield statistically significant assessments of decadal tempera-
63 ture change. The length, decadal trend and data precision of each time series were adjusted in a
64 systematic manner, and forms the core of our analyses. Our aim was to assess the effect that each
65 of these variables has on the ability of a model to detect a decadal trend within time series that
66 differ in their decadal trends and variance properties. The effect gaps in the time series may have
67 on the fitting of models was also investigated as many of the time series used here have missing
68 data scattered throughout, which is unavoidable for a 20+ year time series that is sampled by hand
69 by a single technician at each site.

70 The study provides a better understanding of some of the determinants of a time series that are
71 influential in the detection success of decadal trends in coastal ocean temperature time series.

72 2. Methods

73 *a. Data Sources*

74 Our study lies within the political borders of South Africa's coastline. The location of each point
75 of collection may be found in Figure A1. Of the 129 time series used, 43 are recorded with UTRs
76 and the other 86 with hand-held mercury thermometers. The oldest currently running time series
77 began on January 1st, 1972; there are 11 total time series that started in the 70s, 53 more started in
78 the 80s, 34 began in the 90s, 18 in the 00s and 13 in the current decade.

79 The data are collected using two different methods and a variety of instruments. Hand-held
80 mercury thermometers (which are being phased out in favor of alcohol thermometers or electronic
81 instruments) are used in some instances at the shoreline, and represent seawater temperatures at
82 the surface. At other places, predominantly along the country's east coast, data are collected with
83 glass thermometers from a small boat at the location of shark nets along the coast (Cliff et al.
84 1988). Data at other localities are collected using delayed-mode instruments that are permanently
85 moored shallower than 10 m, but generally very close to the surface below the low-water spring
86 tide level.

87 Over the last 40+ years the electronic instruments used have changed and improved. The pre-
88 vious standard was the Onset Hobo UTR with a thermal precision of 0.01 °C. The meta-data
89 pertaining to these older temperature records and to those that came before, such as the instru-
90 mentation used and the motivation behind the levels of precision at which the data were recorded,
91 have over time been lost, highlighting the issues of staff rotation in government departments and
92 the importance of implementing meta-data standards at a very early stage in any monitoring pro-
93 gramme. The new standard currently being phased in is the Starmon Mini UTR. These devices
94 have a maximum thermal precision of $0.001\text{ °C} \pm 0.025\text{ °C}$ (<http://www.star-oddi.com/>). Of the 43

95 UTR time series in this dataset, 30 were recorded at a precision of 0.001 °C for their entirety, five
96 UTR time series include older data that were recorded at a precision of 0.01 °C or 0.1 °C and so
97 have been rounded down to match this level of precision. Eight additional UTR time series have
98 older data that were recorded at a precision of 0.1 °C.

99 The thermometer data are recorded manually and saved in an aggregated location at the head
100 offices of the collecting bodies. UTRs are installed and maintained by divers and data are retrieved
101 at least once annually. These data are digital and are downloaded to a hard drive at the respective
102 head offices of the collecting bodies.

103 *b. Data Management*

104 Each of the seven bodies contributing data to this study have their own method of data for-
105 matting. Steps are being taken towards a national standard as we move towards replacing all the
106 thermometer recordings with UTR devices; however, as of the writing of this article, one does not
107 yet exist. Data from each organization were formatted to a project-wide comma-separated values
108 (CSV) format with consistent column headers before any statistical analyses were performed. This
109 allowed for the same methodology to be used across the entire dataset, ensuring consistent analy-
110 sis. Before analysing the data they were scanned for any values above 35 °C or below 0 °C. These
111 data points were changed to NA, meaning ‘not available’, before including them in the SACTN
112 dataset.

113 All analyses and data management performed in this paper were conducted with R version 3.3.1
114 (2016-06-21) (R Core Team 2013). The script and data used to conduct the analyses and create
115 the tables and figures in this paper may be found at https://github.com/schrob040/Trend_Analysis.

116 Any time series with a temporal precision greater than one day were averaged into daily values
117 before being aggregated into the SACTN. A series of additional checks were then performed (e.g.

removing long stretches in the time series without associated temperature recordings) and time series shorter than five calendar years or collected at depths greater than 10 m were removed. At the time of this analysis, this daily dataset consisted of 84 time series, consisting of 819,499 days of data; these data were then binned further to the 26,924 monthly temperature values available for use in this study.

c. Systematic Analysis of Time Series

We used the 84 time series simply for their variance properties (comprised of seasonal, inter-annual, decadal and noise components), which reflect that of the thermal environment naturally present along the roughly 3,000 km of South African coastline. Unknown linear trends that may have been present in each time series were removed prior to the ensuing analysis by applying an ordinary least squares regression and keeping the detrended residuals. In doing so we avoided the need to simulate a series of synthetic time series, whose variance components may not have been fully representative of that naturally present in coastal waters. These detrended time series represent a range of time scales, from 72 to 519 months in duration.

To each of the 84 detrended time series we artificially added linear decadal trends of 0.00°C to $0.20^{\circ}\text{C dec}^{-1}$). In other words, we now had time series that captured the natural thermal variabilities around the coast, but with their decadal trends known *a priori*. The range of decadal trends was selected based around the global average of 0.124°C from Kennedy et al. (2011) and used in IPCC (2013). Furthermore, in order to replicate the instrumental precision of the instruments used to collect these time series, we rounded each of these (84 time series \times 5 decadal trends) to four levels of precision: 0.5°C , 0.1°C , 0.01°C and 0.001°C . Consequently, we had a pool of 1,680 time series with which to work.

140 To gain further insight into the effect of time series length on trend detection, each time series
141 was first shortened to a minimum length of 5 years, starting in January so that the timing of the
142 seasonal signal for each time series would be equitable. After fitting the model (see below) to
143 the all 1,680 of the shortened time series, the next year of data for each time series was added
144 and the models fitted again. This process was iterated until the full length of each time series
145 was attained. For example, if a time series consisted of 12 full years of data, it would require 160
146 models (8 iterations of increasing length plus 5 decadal trends plus 4 levels of precision); similarly,
147 720 models would be applied to a 40 year time series. Considering the 84 time series available,
148 the total number of individual models required to capture each combination of variables quickly
149 increased to 36,220.

150 In order to deal with missing values (NA) that were present in some of the time series, we initially
151 replaced these with linearly interpolated temperature values. It turned out that this was a terrible
152 idea because doing so resulted in artificially increasing the goodness of fit of the detected trend:
153 the degree to which this ‘improvement’ occurs is proportional to the amount of NA present and to
154 the magnitude of the linear decadal trend added (see Appendix A). The analysis presented here
155 therefore proceeded with non-interpolated data only.

156 Our approach of fitting models to each of the semi-artificial time series that we generated allowed
157 us to study the effect that the relevant variables (time series length, natural variability, added
158 slope and level of measurement precision) has on the ability of the time series model to faithfully
159 detect the decadal thermal trend, which was known *a priori*. The primary results of interest in
160 these analyses were the significance (*p*-value) of the model fit, the accuracy of the decadal trend
161 determined by the GLS model as well as the error associated with the trend estimate.

d. Time series model

The selection of the appropriate model can greatly influence the ability to detect trends Franzke (2012) and two broad approaches are widely used in climate change research (IPCC 2013). The first group of models estimates linear trends, and although linearity may not reflect reality (*i.e.* trends are very frequently non-linear), these models do provide the convenience of producing an easy to understand decadal trend (e.g. $0.10^{\circ}\text{C dec}^{-1}$) (Wilks 2006). The other group accommodates non-linear trajectories of temperature through time by the use of higher-degree polynomial terms or non-parametric smoothing splines, but the inconvenience comes from not being able to easily compare models among sites (insert refs here). Both groups of models can accommodate serially correlated error structures, which is often the cause for much criticism due to their effect on the uncertainty of the trend estimates (insert refs here). For example, Generalized Least Squares (GLS; yielding estimates of linear trends) and Generalized Additive Mixed Models (GAMM; non-linear fitting with no trend estimate provided) can both capture various degrees of serial autocorrelation (insert refs here). Although our exploratory analysis assessed two parameterizations of each of the model groups, we opted to proceed here with a GLS equipped with a second-order autoregressive AR(2) correlation structure (Wood 2006), which is similar to that used by the IPCC (IPCC 2013). The IPCC uses an AR(1) error term, but our analysis shows that AR(2) is better suited to our data. Another difference from the IPCC approach is that we nested the autoregressive component within year. This modeling approach allowed us to assess how various properties of the detrended data sets would affect the models ability to detect trends – in other words, by comparing the estimates of the trends themselves and how these deviate from the known trend.

AJS: I will insert the equation here...

3. Results

The range of statistical properties derived from the monthly dataset, which are relevant to the aims of this study, may be seen in Table A1. These *a priori* statistical properties of the time series greatly influence the GLS to produce reliable estimates for the known trends that we added to the data, at a wide range of significance levels (p -value), but these and other outcomes vary in a systematic manner. Important variables influential in affecting the GLS model include: the i) initial SD of the time series (after detrending but prior to adding artificial slopes); ii) time series length; iii) the magnitude of the added decadal trend; iv) the measurement precision; v) the amount of missing values in the time series (hereafter referred to as %NA); and vi) the type of instrument used to collect the data. *AJS: Would you please arrange the vars i-v, above, in order of decreasing importance? RWS: I will once they have all been quantified. Currently it is: i) length; ii) DT; iii) SD; iv) NA; v) type. But this may change.*

a. Factors affecting the detection of decadal trends by GLS

As would be expected, the magnitude of the decadal trend estimated by the GLS increases in direct proportion to the decadal trend which we added *a priori*. What is especially noteworthy in this analysis is that time series of longer duration more often result in trend estimates converging with the actual trend than those that are shorter (Figure A2). This effect is most evident from around 30 years. Furthermore, how well the model trend estimate converges with the actual trend is also very visible in the error estimate associated with the estimate of the trend (Figure A3): models fitted to short time series will always have greater trends with larger standard errors (SE) compared to longer ones. The strength of this correlation is $r = 0.56$ ($p < 0.001$) and it remains virtually unchanged as the added decadal trend increases. The p -value of the fitted models also vary in relation to time series duration and to the steepness of the decadal trend added to the data

207 (Figure A4). It is usually the longer time series equipped with steeper decadal trends that are
208 able to produce model fits with estimated trends that are statistically significant. Note, however,
209 that this p -value tests the null hypothesis that the estimated trend is no different from $0^{\circ}\text{C dec}^{-1}$
210 at $p \leq 0.05$, and *not* that the slope is not different from the added trend. Taken together, these
211 outcomes show that although our GLS model can very often result in trend estimates that *approach*
212 the true trend, it is seldom that those estimates are statistically significant in the sense that the
213 estimated trends differ statistically from $0^{\circ}\text{C dec}^{-1}$.

214 The variance of the detrended data is another variable that can affect model fitting, but its only
215 systematic influence concerns the SE of the trend estimate. Here, it acts in a manner that is entirely
216 consistent across all *a priori* trends (Figure A5). What we see is that as the variance of the data
217 increases (represented here as standard deviation, SD) the SE of the slope estimates increases too.
218 Moreover, it does so disproportionately more for time series of shorter duration. Again, as we
219 have seen with the estimated trend that converges to the true trend around 30 years, so too does
220 the initial SD of the data cease to be important in time series of around 3 decades in length.

221 The number of %NA permitted in any of our time series was limited to 15% per time series.
222 Twenty-five of the 84 time series have fewer than 1%NA. An additional 45 time series have up
223 to 5%NA, 10 have up to 10%NA and 4 have up to 15%NA. The mean number of NA for the data
224 is 2.65%. The relationship between %NA and the p -value of the models is shown in Figure A6.
225 At 2.5% or fewer NA their presence does not have any discernible effect on resultant p -values.
226 Progressively increasing the number of NAs above 5%, however, leads to a drastic improvement
227 of models fitted to series with no or gently increasing decadal trends (these generally have very
228 large p -values indicative of very poor fits), and a significant deterioration of models fitted to data
229 with steep decadal trends (for these data, the model generally fits better at low numbers of %NAs,
230 as suggested by the greater number p -values that approach 0.05). In other words, the inclusion of

missing values results in time series with no added decadal trend to veer away from $0^{\circ}\text{C dec}^{-1}$ towards a situation where they may erroneously appear to display a trend. On the other hand, time series that do indeed have decadal trends tend to produce fits that are not significantly different from $0^{\circ}\text{C dec}^{-1}$.

Regarding the effect that the level of measurement precision has on the GLS models, we see in Figure X (fig: correlations-new.pdf) that decreasing the precision from 0.001°C to 0.01°C has an undetectable effect on any differences in the modeled trends. The Root Mean Square Error (RMSE) between the slopes estimated from 0.001°C and 0.01°C data is 0.001. The correspondence between the slopes estimated for data reported at 0.5°C compared to that at 0.001°C decreases to a RMSE of 0.03.

The effect of decreasing data measurement precision from 0.001°C to 0.5°C has almost no appreciable effect on any of the measures of variance presented in this study. The effect of measurement precision on the accuracy of the modeled slope, however, becomes very pronounced going from 0.1°C to 0.5°C . This effect is exacerbated when a smaller decadal is added. For example, with a trend of $0.05^{\circ}\text{C dec}^{-1}$ added to the data, the mean accuracy of models fitted to data with a precision 0.1°C is only -0.14% different from the actual slope; however, the accuracy of the fitted trend at a precision of 0.5°C is 10.81% off, for a range of 10.95% between the most and least accurate mean values. This range decreases to 6.44% with $0.10^{\circ}\text{C dec}^{-1}$, 2.24% at $0.15^{\circ}\text{C dec}^{-1}$ and increases slightly to 2.30% at $0.20^{\circ}\text{C dec}^{-1}$. A precision of 0.5°C always provides clearly less accurate modeled trends than at higher precisions; however, the current analysis offered no one precision that consistently provides the most accurate estimate of the trends. This may however become determinable in an analysis of synthetic data with variance structures that are manipulated in a more consistent manner.

254 An analysis with a large number of variables as shown here is bound to have a medley of complex
255 interactions between the various statistics being measured; however, much of the range seen in the
256 results of the GLS models can be well explained by the influence of one independent variable, or
257 two operating in concert, as we have shown above.

258 **4. Discussion**

259 *RWS: This subsection contains three paragraphs from the "Time Series Analysis" discussion*
260 *subsection. RWS: None of this text has been edited. I must still do so.*

261 *a. Systematic Analysis of Time Series*

262 When the largest SD of 1.3 °C was applied to the time series in these data it increased the *p*-
263 value of the decadal trends detected above 0.05. This implies that if one is sampling seawater
264 temperature in a highly variable area, collecting an additional decade of data may not allow one
265 to establish a significant trend. Consequently, one must take into account the natural variability
266 of the seawater temperatures in a study area when determining how many months of data may be
267 required before climate change studies become feasible.

268 That there is a significant relationship between the significance of the detected trend and the
269 NA% of time series in this study. Interestingly, the amount of missing data in a time series affects
270 the behavior of the GLS model differently. Furthermore, if the missing data are linearly interpo-
271 lated this effect becomes even more pronounced. This means that linear interpolation is not an
272 acceptable method of filling in missing data in a time series analysis and should be avoided where
273 possible.

274 *RWS: This para must be altered to discuss how short time series affect the models ability to*
275 *detect the given DT. The time series analysis was applies to time series as short as 5 years at the*

276 risk of producing more erratic values in order to analyse as many of the time series in the dataset
 277 as possible. The smaller n for these time series allows any short-term phenomena to have a larger
 278 impact on the detected decadal change than in the longer time series, which may give the illusion
 279 that some of the shorter time series are showing very significant results in terms of decadal change.
 280 For example, the largest ΔT (R^2) values at either end of the spectrum for time series at or under 10
 281 years in length are $3.5\text{ }^{\circ}\text{C dec}^{-1}$ (0.69) and $-7.233\text{ }^{\circ}\text{C dec}^{-1}$ (0.81). The highest R^2 value for any
 282 time series at or under 10 years is 0.81. These extrapolated decadal trend values could not to be
 283 sustained over a long period as almost any point in the ocean would freeze over in only 40 years
 284 at $-7.233\text{ }^{\circ}\text{C dec}^{-1}$! Not all short time series have large ΔT (R^2) values; indeed, the lowest ΔT
 285 (R^2) value for time series under 10 years is $0.0\text{ }^{\circ}\text{C dec}^{-1}$ (0.00). The mean and median values for
 286 ΔT from time series under 10 years are $0.08\text{ }^{\circ}\text{C dec}^{-1}$ and $0.25\text{ }^{\circ}\text{C dec}^{-1}$. *AJS: Above the trend*
 287 *data are reported at two levels of precision... I think you need a statement somewhere near the*
 288 *start of the section here, or in the Results section, saying that the more precise reporting is for the*
 289 *UTR data while the lower precision trends were only provided for the other data. Or something*
 290 *like that... It looks weird when there is three decimal places some times, and only one other times.*
 291 The fact that the time series under 10 years in this study tend to have higher R^2 and lower p -
 292 values may seem to suggest that they would be at least as useful for climate change research as the
 293 longer time series; however this is not true. This is because the length variable has a non-linear
 294 influence on the R^2 and p -values of a time series. Whereas these values are almost always strongest
 295 within the first 10 years of a time series, the size of the decadal trends detected in these short time
 296 series are dubious in that such a strong signal would most likely not perpetuate for long in one
 297 direction and would therefore only be an artefact of the short length of the times series. This is
 298 an important consideration as many studies use *in situ* time series that are shorter than 10 years
 299 when relating temperature to other biotic or abiotic variables. It should go without saying that to

300 calculate a decadal trend in a time series, more than 10 years of data would be required. The size
301 of the outliers calculated from these shorter time series reaffirms that wisdom.

302 A bit of text here...

303 A bit of text here...

304 One of the original motivators for this paper was to investigate the effect instrument precision
305 had on a time series ability to assess long-term climate change in order to validate the use of the
306 low precision 0.5 °C thermometer data. Whereas the precision of these data is below the current
307 standard for climate change research, the length of the time series created with these instruments
308 are a valuable asset. The goodness of fit of linear models begins to increase once time series reach
309 30+ years in length. Forty-five of the 84 thermometer time series in this dataset are at or over 30
310 years whereas the longest UTR time series is less than 19 years long. By applying time series
311 analyses on the 20 new UTR time series with the high-precision of 0.001 °C and incrementally
312 decreasing this precision, we could deduce the hypothetical effect lower precisions would have
313 on a time series. The natural range of precisions within this dataset is 0.001 °C to 0.5 °C and
314 the almost complete lack of effect on the resultant R^2 and p values of these time series asserts
315 that the low precision thermometer data are as useful for climate change research as newer higher
316 resolution UTR data; assuming that one is willing to report ΔT values that have been rounded
317 to the nearest 0.1 °C in order to accurately reflect the level of precision at which the data were
318 collected.

319 *RWS: None of this text has been edited. I must still do so.*

320 5. Conclusion

321 We draw several key conclusions:

1. There is not a significant relationship between the goodness of fit (R^2) of a linear model to a time series and the NA% of that time series when the NAs are filled in via linear interpolation. This is an important finding as it means that, within reason, linear interpolation may be used to fill gaps in a time series before applying any time series analysis methods.
2. Length has the largest effect on the goodness of fit (R^2) of the decadal trend and natural variability (SD) has the largest effect on the significance (p) of the trend detected.
3. There is a predictable decrease in the goodness of fit (R^2) of a linear model to the trend line of a time series as it extends from 10 to 20 years in length. The goodness of fit (R^2) then begins to increase once the time series becomes roughly 30+ years long. Analyses of time series at or under 10 years in length should be interpreted with extreme caution in spite of them often having strong R^2 values.
4. Within the first decade of a time series, if the temperatures within the last few months move strongly in the opposite direction from the prevailing trend, the linear model used to detect the trendline may show an abrupt change in direction (i.e. a positive trend can become negative and *vice versa*).
5. After the first decade of data, the changes detected in almost all trends for all 105 time series become more gradual; however, many trend lines still change direction over the course of the following two decades.
6. It is at these changes in direction that the p -values for the time series plummet, though generally they tend to follow the same pattern of becoming weaker and then slowly stronger over time, as we see in the R^2 values.

7. There is a slight linear decrease in R^2 as the natural thermal variability (SD) of seawater increases; however, the decrease in p -values is larger and more rapid.
8. A precision greater than 0.5 °C is not required to confidently detect the long-term trend in a time series. This is an important consideration as many studies investigating the effects of climate change (e.g. Grant et al. 2010; Scherrer and Körner 2010; Lathlean and Minchinton 2012) do use lower precision 0.1 °C data. That being said, a precision of 0.001 °C or 0.01 °C is preferable over 0.5 °C. In fact, because the results from the higher precision of 0.001 °C were almost identical to the 0.5 °C tests, the higher precision is only necessary when one needs to identify trends at a precision of 0.01 °C or greater (Karl et al. 2015). This finding means that older, lower precision data may be combined with newer higher precision data within the same time series without concern that the reduced overall data precision will have a large negative impact on the time series ability to detect decadal trends. Indeed, extending time series in this way will only serve to make them more dependable as length is the primary criteria through which one should initially assess a time series ability to detect climate change before refining ones assumptions with any statistical analyses.
9. Decreasing the precision of measurements to greater than 0.1 °C has almost no appreciable effect on a time series ability to detect a long term trend, provided that the reported effect size matches the level of precision by the instruments.

We understand that time series of >30 years may be exceedingly rare. Therefore, while we move forward as a scientific community investigating the issues of climate change, the increasing length and continuity of any current and future time series must be ensured in order to construct and maintain a clear understanding of the trends in changing temperature that are occurring throughout Earth's oceans.

Acknowledgments. The authors would like to thank DAFF, DEA, EKZNW, KZNSB, SAWS and SAEON for contributing all of the raw data used in this study. Without it, this article and the SACTN would not be possible. This research was supported by NRF Grant (CPRR14072378735). The authors report no financial conflicts of interests. The data and analyses used in this paper may be found at https://github.com/schrob040/Trend_Analysis.

APPENDIX A

Effects of linear interpolation

Blurb here about linear interpolation effects.... Figure A7.

References

- Blanchette, C. A., C. Melissa Miner, P. T. Raimondi, D. Lohse, K. E. K. Heady, and B. R. Broitman, 2008: Biogeographical patterns of rocky intertidal communities along the Pacific coast of North America. *Journal of Biogeography*, **35** (9), 1593–1607.
- Broitman, B. R., N. Mieszkowska, B. Helmuth, and C. A. Blanchette, 2008: Climate and recruitment of rocky shore intertidal invertebrates in the eastern North Atlantic. *Ecology*, **89** (11 Suppl), S81–90.
- Brown, O. B., P. J. Minnett, R. Evans, E. Kearns, K. Kilpatrick, A. Kumar, R. Sikorski, and A. Závody, 1999: MODIS Infrared Sea Surface Temperature Algorithm Theoretical Basis Document Version 2.0. *University of Miami*, 31 098–33 149.
- Cliff, G., S. F. J. Dudley, and B. Davis, 1988: Sharks caught in the protective gill nets off Natal, South Africa. 1. The sandbar shark *Carcharhinus plumbeus* (Nardo). *South African Journal of Marine Science*, **7** (1), 255–265.

- 387 Couce, E., A. Ridgwell, and E. J. Hendy, 2012: Environmental controls on the global distribution
388 of shallow-water coral reefs. *Journal of Biogeography*, **39** (8), 1508–1523.
- 389 Franzke, C., 2012: Nonlinear trends, long-range dependence, and climate noise properties of sur-
390 face temperature. *Journal of Climate*, **25** (12), 4172–4183, doi:10.1175/JCLI-D-11-00293.1.
- 391 Grant, O. M., L. Tronina, J. C. Ramalho, C. Kurz Besson, R. Lobo-Do-Vale, J. Santos Pereira,
392 H. G. Jones, and M. M. Chaves, 2010: The impact of drought on leaf physiology of *Quercus*
393 *suber* L. trees: Comparison of an extreme drought event with chronic rainfall reduction. *Journal*
394 *of Experimental Botany*, **61** (15), 4361–4371, doi:10.1093/jxb/erq239.
- 395 Hutchings, L., and Coauthors, 2009: The Benguela Current: An ecosystem of four components.
396 *Progress in Oceanography*, **83** (1-4), 15–32, doi:10.1016/j.pocean.2009.07.046.
- 397 Karl, T. R., and Coauthors, 2015: Possible artifacts of data biases in the recent global surface
398 warming hiatus. *Science*, **348** (6242), 1469–1472, doi:10.1126/science.aaa5632.
- 399 Kennedy, J. J., N. A. Rayner, R. O. Smith, D. E. Parker, and M. Saunby, 2011: Reassessing biases
400 and other uncertainties in sea surface temperature observations measured in situ since 1850: 2.
401 Biases and homogenization. *Journal of Geophysical Research: Atmospheres*, **116**, D14104.
- 402 Lathlean, J. A., and T. E. Minchinton, 2012: Manipulating thermal stress on rocky shores to pre-
403 dict patterns of recruitment of marine invertebrates under a changing climate. *Marine Ecology*
404 *Progress Series*, **467**, 121–136, doi:10.3354/meps09996.
- 405 Martin, M., and Coauthors, 2012: Group for High Resolution Sea Surface temperature (GHRSSST)
406 analysis fields inter-comparisons. Part 1: A GHRSSST multi-product ensemble (GMPE). *Deep*
407 *Sea Research Part II: Topical Studies in Oceanography*, **77-80**, 21–30.

408 R Core Team, 2013: *R: A Language and Environment for Statistical Computing*. Vienna, Austria,
 409 R Foundation for Statistical Computing, URL <http://www.r-project.org/>.

410 Reynolds, R. W., and T. M. Smith, 1994: Improved global sea surface temperature analyses using
 411 optimum interpolation. *Journal of Climate*, **7** (6), 929–948.

412 Roberts, M. J., 2005: Chokka squid (*Loligo vulgaris reynaudii*) abundance linked to changes in
 413 South Africa’s Agulhas Bank ecosystem during spawning and the early life cycle. *ICES Journal*
 414 *of Marine Science*, **62** (1), 33–55, doi:10.1016/j.icesjms.2004.10.002.

415 Santos, F., M. Gomez-Gesteira, M. DeCastro, and I. Alvarez, 2012: Differences in coastal and
 416 oceanic SST trends due to the strengthening of coastal upwelling along the Benguela current
 417 system. *Continental Shelf Research*, **34**, 79–86.

418 Scherrer, D., and C. Körner, 2010: Infra-red thermometry of alpine landscapes challenges climatic
 419 warming projections. *Global Change Biology*, **16** (9), 2602–2613, doi:10.1111/j.1365-2486.
 420 2009.02122.x.

421 Smit, A. J., M. Roberts, R. J. Anderson, F. Dufois, S. F. J. Dudley, T. G. Bornman, J. Olbers, and
 422 J. J. Bolton, 2013: A coastal seawater temperature dataset for biogeographical studies: Large
 423 biases between *in situ* and remotely-sensed data sets around the coast of South Africa. *PLoS*
 424 *ONE*, **8** (12), doi:10.1371/journal.pone.0081944.

425 Stocker, T., D. Qin, G. K. Plattner, M. Tignor, S. K. Allen, J. Boschung, A. Nauels, Y. Xia, V. Bex,
 426 and P. M. Midgley (Eds.), 2013. Climate change 2013: the physical science basis: Working
 427 Group I contribution to the Fifth Assessment Report of the Intergovernmental Panel on Climate
 428 Change. Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA.
 429 doi:10.1017/CBO9781107415324.008

- 430 Tittensor, D. P., C. Mora, W. Jetz, H. K. Lotze, D. Ricard, E. V. Berghe, and B. Worm, 2010:
431 Global patterns and predictors of marine biodiversity across taxa. *Nature*, **466 (7310)**, 1098–
432 1101.
- 433 Tyberghein, L., H. Verbruggen, K. Pauly, C. Troupin, F. Mineur, and O. De Clerck, 2012: Bio-
434 ORACLE: A global environmental dataset for marine species distribution modelling. *Global*
435 *Ecology and Biogeography*, **21 (2)**, 272–281.
- 436 Wilks, D. S., 2006: Statistical Methods in the Atmospheric Sciences, 2nd edition. Elsevier,
437 Philadelphia, 627 pp.
- 438 Wood, S., 2006: Generalized Additive Models: An Introduction with R. Chapman and Hall/CRC,
439 Boca Raton, Florida, 410 pp.

440	LIST OF TABLES	
441	Table A1.	The mean values ($\pm sd$) of the length, initial sd of the time series, the significance (p) of the
442		detected trends, the percentage difference between the slope added <i>a priori</i> and the modelled
443		slope (% difference) as well as the standard error (se) around the detected trend from the 84
444		time series used in this study. Rows show the effect increasing size of added slopes have on
445		the results. Values shown here were derived only from the data rounded to a precision of
446		0.001 °C 23

447 TABLE A1. The mean values ($\pm sd$) of the length, initial sd of the time series, the significance (p) of the detected trends, the
448 percentage difference between the slope added *a priori* and the modelled slope (% difference) as well as the standard error (se)
449 around the detected trend from the 84 time series used in this study. Rows show the effect increasing size of added slopes have on
450 the results. Values shown here were derived only from the data rounded to a precision of 0.001 °C

added slope	p of trend	% difference	se of trend
0.00 °C dec ⁻¹	0.86 \pm 0.13	NA	2.51e ⁻³ \pm 2.40e ⁻³
0.05 °C dec ⁻¹	0.76 \pm 0.16	0.86 \pm 145.41	2.51e ⁻³ \pm 2.40e ⁻³
0.10 °C dec ⁻¹	0.62 \pm 0.23	0.43 \pm 72.71	2.51e ⁻³ \pm 2.40e ⁻³
0.15 °C dec ⁻¹	0.48 \pm 0.26	0.27 \pm 48.45	2.51e ⁻³ \pm 2.40e ⁻³
0.20 °C dec ⁻¹	0.37 \pm 0.28	0.20 \pm 36.34	2.51e ⁻³ \pm 2.40e ⁻³

451	LIST OF FIGURES	
452	Fig. A1. Location and instrument types used to sample each time series available for use in this study	25
453	Fig. A2. The effect of length on the ability of a model to accurately detect the trend within a time series	26
454	Fig. A3. The relationship between the length of a time series and the standard error (SE) of the modelled trend	27
455	Fig. A4. The effect the natural variation found within a time series has on the significance of the modelled trend	28
456	Fig. A5. The relationship between the effect of initial SD on the SE of a modelled trend	29
457	Fig. A6. The relationship between NA% amd the significance of a fitted trend	30
458	Fig. A7. Similar to Figure A6, this figure shows the effect missing data have on	31




FIG. A1. Location and instrument types used to sample each time series available for use in this study. In the legend, ‘new’ shows the underwater temperature recorder (UTR) time series that were recorded entirely with the newer UTRs that have a high precision of 0.001 °C, ‘old’ refers to UTR time series that were recorded, at least in part, with older UTRs and have data with precisions lower than 0.001 °C. The ‘thermo’ label shows the location of the thermometer time series.




FIG. A2. The effect of length on the ability of a model to accurately detect the trend within a time series.




FIG. A3. The relationship between the length of a time series and the standard error (SE) of the modelled trend.





FIG. A4. The effect the natural variation found within a time series has on the significance of the modelled trend. Also shown via size of dots is the interaction of length on these variables.



466 FIG. A5. The relationship between the effect of the initial SD of a time series on the SE of a modelled trend,
 467 controlled for by the length of the time series.




FIG. A6. The relationship between NA% and the significance of a fitted trend. Each panel shows the effect of an increasingly larger amount of missing values. The effect of missing data on a models ability to detect a larger decadal trend is initially negligible, but becomes significant as greater amounts of data are missing.




FIG. A7. Similar to Figure A6, this figure shows the effect missing data have on the significance of the slopes detected by GLS however; the missing values in the time series have been filled here via linear interpolation. The effect this has on the significance of the modeled trends is both immediate and dramatic. The behaviour of the quantity of interpolated data also differs from the effect of data left simply as NA. At lower levels of interpolation, missing data actually aid in the fitting of a more significant trend line. This phenomena reverses around 5%NA when the relationship becomes negative, meaning that as the amount of interpolated data increase, the significance of the fitted trend decreases.