The Statistical Value of Adaptive Feature Queries

Anonymous Authors[1]

Abstract

Questionnaires, laboratory panels and budgeted classifiers reveal at most b of m features per record. If each record's b questions must be fixed in advance instead of chosen from its earlier answers, how many more records does it take to estimate a known low-degree functional, such as the accuracy of a frozen screening rule? We show that pre-committed question sets simulate any adaptive protocol exactly, in law, with 2^{b-1} times the records in expectation, whatever the task and the population, and that this factor is attained when there are enough features. The worst case is robust to relaxations: with r rounds of adaptivity the simulation costs 2^{b-r} and a constant number of rounds can still lose $2^{b(1-o(1))}$; c extra pre-committed questions per record divide the worst-case price by at most $c+1$; and a mean budget instead of a hard cap by at most b, when there are exponentially many features. The price belongs to targeted collection: for all b-way interactions at once, uniform forms are optimal even among branching designs. A certified solver computes the optimal design of each kind for a given rule and population. On screening trees for several health outcomes, with the cap equal to the depth of the rule, optimal forms need 1.0–2.38 times the records of the adaptive optimum where the optimum was computed (at most 3.97 for the widest rules), and 1.53–1.93 at the median when the design is learned from a pilot charged to the same audit. Trained rules depend on few items beyond their depth, so one extra question per record lowers the median price from 1.47 to 1.14 and two to 1.02; counted in questions instead of records, however, branching keeps an advantage of about 1.39 over the best of the pre-committed designs we priced (forms of size b and $b+1$, and complete records of the rule's items).

[1]Anonymous Institution, Anonymous City, Anonymous Region, Anonymous Country. Correspondence to: Anonymous Author <anon.email@domain.com>.

Preliminary work. Under review by the International Conference on Machine Learning (ICML). Do not distribute.

1. Introduction

Population quantities are often estimated from records that each reveal only a few features. Questionnaires cap respondent burden, clinical protocols cap the tests per patient, and budgeted classifiers acquire a few inputs per case (Turney, 1995; Trapeznikov & Saligrama, 2013; Nan & Saligrama, 2017). The collector can pre-commit to each record's feature set, as in split questionnaires and laboratory panels ordered at once (Raghunathan & Grizzle, 1995; Chipperfield & Steel, 2009), or choose each question after seeing earlier answers, as in skip logic, adaptive matrix sampling (Gonzalez & Eltinge, 2008) and reflex testing. In machine-learning terms, the first is a static feature subset drawn per example and the second an adaptive feature-acquisition policy with a hard budget (Covert et al., 2023); here the goal is not a prediction for the example but an estimate for the population, such as the accuracy of a frozen rule or the accuracy gain of a model update. Two settings make the constraints concrete. In large assessments and health surveys the outcome is on the frame or in a registry, session length caps the items per respondent, and booklets are fixed before fieldwork; multistage adaptive forms were introduced to relax exactly this (Yamamoto et al., 2018). In on-device evaluation of a deployed rule the outcome is logged centrally, the features stay on the device, a privacy or communication budget caps how many a device discloses, and a device answers one pre-specified query, because interactive protocols need it to stay available across rounds (Joseph et al., 2019; Kairouz et al., 2021).

This paper asks what pre-commitment costs in records. That adaptivity within one input can be worth exponentially many queries is classical in decision-tree complexity, the huge-object model and interactive local privacy (Goldreich & Ron, 2022; Adar & Fischer, 2024; Joseph et al., 2019). Under a per-record cap the price is paid in records, and it is not known what it is exactly, how robust it is to small relaxations, or what it is for rules that are actually trained. We answer these three questions.

The exact price (Section 3). Pre-committed forms reproduce any adaptive protocol exactly in law, for every estimator and loss, with $K_{m,b} = \min\{2^{b-1}, \binom{m-1}{b-1}\}$ times the records in expectation. This is the folklore conversion of adaptive into non-adaptive queries made exact across records; the new part is that 2^{b-1} is attained, for a single

linear functional and whatever the accuracy of the rule, by an address contrast on $b - 1 + 2^{b-1}$ features.

Its robustness (Section 3). Neither a few rounds of adaptivity, nor a few extra questions, nor a mean budget removes the worst case. With r rounds the simulation costs 2^{b-r}, and a pointer tree loses at least $2^{b-1}/(1 + 2b/r)^r$, which is $2^{b(1-o(1))}$ for constant r; the two bounds do not match beyond that. Forms with $c < 2^{b-1}$ extra questions need between $2^{b-1}/(c+1)$ and $2^{b+1}/(c+3)$ times the records in the worst case, and in polynomial dimension forms twice as long as the rule's paths still lose $2^{b(1-o(1))}$.

The price at a population, computed with certificates (Sections 4–5). The minimax constant of each protocol class is a convex program, and a column-generation solver with exact pricing over trees finds the optimal design of every class. A working model for the unasked features leaves the simulation bound intact, removes the gap up to a factor two when the features are independent given the label, and does not remove it under strong couplings.

What it is for trained rules (Section 6). On screening trees for several health outcomes, with the cap equal to the depth of the rule, optimal forms need 1.0–2.38 times the records of branching, on the rules whose optimum was computed. This is far from the worst case, and in records it is easy to remove: the rules depend on a median of 3 items beyond their depth, and one extra pre-committed question lowers the median price from 1.47 to 1.14, two to 1.02, as does one follow-up round. In questions it is not removed: forms of size b, forms one question longer and complete records of the rule's items cost 1.47, 1.39 and 1.50 times the questions of branching at the median. Uniformly drawn forms cost 2.0–4.2 times the records of branching, so what matters first is optimizing the forms, which the solver does from a reference sample or a charged pilot.

The setting is deliberately simple: every sampled record's label is observed and charged separately, the target is one known contrast of degree at most b on binary features, and answers do not depend on question order. Section 8 discusses what this leaves out and what a practitioner can take from the results.

2. Estimand and Acquisition Protocols

Let X take values in $\mathcal{X} = \{-1, 1\}^m$ and let $Y \in \{-1, 1\}$ be a label. Freeze two rules $p_0, p_1 : \mathcal{X} \to [0, 1]$ that predict $+1$ with probabilities $p_j(x)$, put $d = p_1 - p_0$, $t(x, y) = yd(x)$, and

$$\Delta(P) = \mathbb{E}_P[t(X, Y)] = \mathbb{E}_P[Y d(X)]. \tag{1}$$

This is the accuracy difference of the two rules; $p_0 \equiv 1/2$ makes it affine in the accuracy of p_1 alone (sensitivity, specificity and other proper scores: Appendix G). The cap b bounds the feature queries per record and is *binding* when $m > b$.

Protocols. An interview ι maps a profile x to a transcript $\ell_\iota(x)$. A *fixed form* queries a set of coordinates chosen before the record is seen. An r*-round interview* queries batches $B_1, B_2(x_{B_1}), \ldots, B_r$, each chosen from the answers to earlier batches, with $\sum_s |B_s| \le b$. A *branching interview* is a decision tree of depth at most b. The menus are nested, $\mathcal{A}_{\text{fixed}} = \mathcal{A}_1 \subseteq \cdots \subseteq \mathcal{A}_b = \mathcal{A}_{\text{branch}}$. In machine-learning terms an interview is a per-example acquisition policy, a form is a static feature subset, and a collection of records under a design is an evaluation run, which we call an audit. In missing-data terms a fixed form is missingness completely at random by design, and the other interviews are randomized monotone coarsenings (Gill et al., 1997), which are ignorable. Across records a protocol may use all previous records to choose the next interview. $R_{\mathcal{A}}(d, n)$ is the minimax mean squared error over protocols and estimators with n records and interviews from $\mathcal{A}$.

Identification. The contrast is identifiable for all populations exactly when d is a sum of functions that are each constant on the transcript cells of one interview; for forms of size b and for depth-b trees this is the same condition, Fourier degree $\deg(d) \le b$ in the characters $\chi_S(x) = \prod_{j \in S} x_j$. It holds for every probability average of depth-b trees, such as the predicted probability of a depth-limited random forest, and fails for most thresholded ensembles at small caps.

Designs and constants. A *design* draws an interview ι with probability q_ι and scores its transcript with a table g_ι; the score $Y g_\iota(\ell_\iota(X))/q_\iota$ is unbiased for $Y d(X)$ at every profile exactly when $\sum_\iota g_\iota(\ell_\iota(x)) = d(x)$ for all x. Define the worst-case and population constants

$$V(d, \mathcal{A}) = \inf_{q, g:\, \sum_\iota g_\iota = d} \max_{x \in \mathcal{X}} \sum_\iota \frac{g_\iota(\ell_\iota(x))^2}{q_\iota}, \tag{2}$$

$$W_{\mathcal{A}}(P) = \inf_{q, H:\, \mathbb{E}[H|X,Y] = t(X,Y)} \mathrm{Var}_{P,q}(H), \tag{3}$$

where H is a score of the drawn interview's transcript and the label that may be tuned to P. Thus V is the worst-case and $W_{\mathcal{A}}(P)$ the population price of estimating Δ from interviews in $\mathcal{A}$, in units of variance per record; Theorem 3 shows that they are the first-order minimax risks, and ratios of constants are ratios of records needed.

A worked example. Let $m = 3$, $b = 2$ and $d(x) = [(1 + x_1)x_2 + (1 - x_1)x_3]/2$: the first answer selects which of two others is the target. Branching reads x_1 and then the selected coordinate, so it observes t. Under uniform profiles and an independent uniform label, the best pre-committed design draws $\{1, 2\}$ or $\{1, 3\}$ with probability

$1/2$ and doubles the score when the drawn form contains the selected coordinate: it needs twice the records.

3. The Price of Pre-Commitment

Theorem 1 (Exact simulation and the sharp gap)**.** *Let $1 \le b \le m$ and $K_{m,b} = \min\{2^{b-1}, \binom{m-1}{b-1}\}$.* (i) *Every protocol on n branching records, with any estimator, is reproduced with the same law at every population by a fixed-form protocol on a random number of records with mean $K_{m,b}n$.* (ii) *Consequently, for every contrast of degree at most b, $V_{\mathrm{fixed}} \le K_{m,b} V_{\mathrm{branch}}$, and at every law P of (X, Y),*

$$W_{\mathrm{branch}}(P) \le W_{\mathrm{fixed}}(P) \le K_{m,b}\, W_{\mathrm{branch}}(P). \qquad (4)$$

(iii) *The address contrast on $m = b - 1 + 2^{b-1}$ coordinates has $V_{\mathrm{fixed}}/V_{\mathrm{branch}} = 2^{b-1}$, and $W_{\mathrm{fixed}}(P)/W_{\mathrm{branch}}(P) = 2^{b-1}$ at uniform profiles whenever the label agrees with the rule with a probability in $(0, 1)$ that does not depend on the profile. With $m \le b^2$ there are examples with ratio at least $2^{b-b/(\lfloor \log_2 b \rfloor + 1) - \lfloor \log_2 b \rfloor}$.*

Proof. (i) Remove repeated questions from a tree τ and pad every root-to-leaf path, by a fixed rule, with unused coordinates to exactly b distinct ones. All path sets contain the root question, and the first $b - 1$ answers determine the set, so the collection $\mathcal{S}_\tau$ of distinct path sets has at most $K_{m,b}$ elements. To simulate one record of τ, draw one set of $\mathcal{S}_\tau$ uniformly and ask it as a form (Figure 1b). The answers reveal whether the drawn set is this record's own padded path set (follow the tree while its questions lie in the set), and for every profile exactly one set is; so the check succeeds with probability $1/|\mathcal{S}_\tau|$ whatever the record is. Accept a success with probability $|\mathcal{S}_\tau|/K_{m,b}$. An accepted record is then a draw from the population together with the transcript that τ would have produced, and the number of records used until acceptance is geometric with mean $K_{m,b}$ and independent of that transcript. The tree may depend on earlier transcripts, so the argument applies record by record to any protocol (Theorem C.1). *(ii)* For V, apply a near-optimal branching design to the records accepted from a fixed budget, whose number is binomial with success probability $1/K_{m,b}$. At a given P an explicit score suffices: draw a tree of the branching design and one of its path sets, and return $c + |\mathcal{S}_\tau|(H - c)$ if the set is the record's own and c otherwise, where H is the tree's score and $c = \Delta(P)$. This score has the conditional mean of H at every profile and at most $K_{m,b}$ times its variance (Appendix E.1). *(iii)* Let $\kappa = b - 1$ address bits select one of $D = 2^\kappa$ data bits, $d(A, Z) = Z_A$ (Figure 1a). Branching computes d. Under the uniform law, a form T with s address bits and $w \ge 1$ data bits has $s \le \kappa + 1 - w$ and explains the energy

$$\|\mathbb{E}[d \mid X_T]\|_2^2 = w\, 2^{s-2\kappa} \le 2^{-\kappa}, \qquad (5)$$

because each of its data bits is consistent with a fraction 2^{-s} of the partial addresses and is then the selected bit with conditional probability $2^{-(\kappa-s)}$. The dual of Theorem 3 with the uniform law and the witness $h = d$ compares $\|d\|_2^4 = 1$ with this energy, so $V_{\mathrm{fixed}} \ge D$. The same energy bounds the variance that a form explains when the label is a noisy copy of d (Proposition E.3), and products of address blocks give the example in polynomial dimension (Appendix C.6.1). □

Part (i) is the folklore conversion of an adaptive into a non-adaptive query procedure, carried out across records and made exact by the acceptance step; what the theorem adds is that the resulting constant is attained, already for one linear functional. Our sharp example uses $m = b - 1 + 2^{b-1}$ features; for $m \le b^2$ we only have the product examples, and for smaller m the extreme ratio is open (at $m = 4$, $b = 3$ it lies between 2.457 and $K_{4,3} = 3$; Proposition C.11). *One target or many.* The simulation involves no target, so the upper bounds hold for any family of targets and for rules chosen after collection, and the lower bound extends, for the worst member, to families that contain the address contrast and consist of functions of the address tree's transcript, such as all relabelings of its leaves. At the other extreme the price vanishes: for all monomials of degree b at once, uniform forms are minimax among all designs with unbiased scores, branching included, because every record yields exactly one question set however it is chosen (Proposition C.3). Adaptivity pays when the collector knows which questions the target needs; a multipurpose collection should use forms.

Theorem 2 (Rounds and extra questions)**.** *Let $K_r = \min\{2^{b-r}, \binom{m-r}{b-r}\}$ and let V_r, $W_r(P)$ be the constants of r-round interviews.* (a) *Part* (i) *of Theorem 1 holds for r-round interviews with K_r in place of $K_{m,b}$; hence $V_r \le K_r V_{\mathrm{branch}}$ and $W_r(P) \le K_r W_{\mathrm{branch}}(P)$ at every P.* (b) *The binary pointer tree on $m = 2^b - 1$ coordinates has $V_{\mathrm{branch}} = 1$ and $V_r \ge 2^{b-1}/(1 + 2b/r)^r$ for $1 \le r \le b$; a product of pointer trees on $m \le b^2$ coordinates has $\log_2 V_r \ge b - O(rb \log\log b/\log b)$.* (c) *For $b + c \le m$, forms of size $b + c$ reproduce every protocol on depth-b interviews with $K^+_{b,c} = 2^{b-1-j(c)}$ times the records in expectation, where $j(c)$ is the largest $j \le b - 1$ with $2^{j+1} - j - 2 \le c$, so that $K^+_{b,c} < 2^{b+1}/(c + 3)$ whenever $K^+_{b,c} > 1$. The address contrast needs exactly $2^{b-1}/(c + 1)$ times the records for $c < 2^{b-1}$, and a product of address blocks on $m \le b^2$ features needs $2^{b(1-o(1))}$ for every $c = O(b)$.*

For (a), follow the tree for $r - 1$ rounds and guess the rest of the path in the last batch.

Sketch of (b). By the dual of Theorem 3 with the uniform law and $h = d$, $V_r \ge 1/\max_\iota \mathbb{E}[\mathbb{E}[d \mid \ell_\iota]^2]$. Only a data bit that the interview has read carries signal, so the energy of ι




Figure 1. The three constructions behind Theorems 1 and 2. (a) Address contrast: $\kappa = b - 1$ address bits select one of $D = 2^\kappa$ data bits; branching reads the address and then the selected bit, while any fixed set of b coordinates explains at most a fraction $1/D$ of the target's variance under the uniform law. (b) Path guessing: a fixed form draws one of a tree's at most 2^{b-1} padded path sets and keeps the record when the path completes inside it. (c) Binary pointer tree: $L = b - 1$ levels of pointer bits select a child and the target is the data bit at the reached leaf; a round that reads a pointer bits raises the conditional probability of the most likely leaf by at most a factor $1 + 2a$ in expectation.

is at most $\mathbb{E}\pi_r$, where π_t is the conditional probability, given the transcript of t rounds, that the data bit of the selected leaf has been read (Figure 1c). Let M_t be 2^{b-1} times the largest conditional probability of a leaf given the transcript of t rounds, and $Q_t = M_t + 2^{b-1}\pi_t$, so that $Q_0 = 1$. A round that reads a pointer bits and λ data bits satisfies $\mathbb{E}[Q_t \mid \text{past}] \le (1 + 2a + \lambda)Q_{t-1}$: each pointer bit read adds at most $2M_{t-1}$ to M in expectation, because a chain of j newly read bits multiplies a leaf's probability by 2^j only if the first $j - 1$ point the right way, and each data bit adds at most the current largest leaf probability to π. With at most b questions in r rounds, $2^{b-1}\mathbb{E}\pi_r \le \mathbb{E}Q_r \le (1+2b/r)^r$. The potential is multiplicative over independent blocks, which gives the product statement (Appendix D.1).

The two rounds bounds do not match. They agree up to a polynomial in b for constant r; on the pointer tree a protocol within a polynomial factor of branching needs at least $0.37b - O(\log b)$ rounds; in polynomial dimension the lower bound covers only $r = o(\log b/\log\log b)$, since the product is evaluated exactly in $\lfloor \log_2 b \rfloor$ rounds; and at $b = 3, 4$ the pointer tree loses 1.80 and between 3.32 and 3.33 in two rounds against the upper bounds 2 and 4 (Appendix D.4), so we do not know a contrast that attains 2^{b-r} for $r \ge 2$.

For (c), guess only the first $b - 1 - j$ answers and put the whole subtree below the guessed node into the form, which costs $2^{j+1} - j - 2$ extra questions; for the address contrast, a form with all address bits and $c + 1$ data bits explains a fraction $(c+1)/2^{b-1}$ of the variance and no form explains more (Appendix C.2). Extra questions thus buy at most a factor linear in their number. The same holds for a mean budget: if records may ask up to $b + c$ features but only b on average, the worst-case price is between $\frac{b+c}{b} \cdot 2^{b-1}/(c+1)$ and $\frac{b+c}{b}K^+_{b,c}$, at least $2^{b-1}/b$ for every $c < 2^{b-1}$; under a pure mean budget it is at most m/b (ask all features with probability b/m), which is small in polynomial dimension and still about $2^{b-1}/b$ for the address contrast (Theorem C.2(d), Proposition C.9).

4. Constants and Their Computation

The simulation bounds the gap; the constants V and W of Section 2 measure it for a given contrast and population. Let $\sigma_P^2 = \mathrm{Var}_P(t)$; for a depth-$b$ tree τ that computes d, with padded path sets $S_\tau(x)$, let $\rho_P^2 = \max_{|T|=b} \mathrm{Var}_P(\mathbb{E}_P[t \mid X_T, Y])/\sigma_P^2$ be the largest share of the target's variance that one form explains, and $N_P(\tau) = (\sum_S v_S^{1/2})^2$ with $v_S = \mathbb{E}_P[(t-\Delta)^2 \mathbf{1}\{S_\tau(X) = S\}]/\sigma_P^2$ the effective number of question sets the tree uses at P.

Theorem 3 (Exact constants)**.** *Let $\mathcal{A}$ be a finite menu on which d is identifiable.* (a) *$nR_{\mathcal{A}}(d,n) \to V(d,\mathcal{A})$; for every $\epsilon > 0$ an iid design with an unbiased score has worst-case risk at most $(V + \epsilon)/n$ for every n, and the converse covers biased estimators, designs that adapt across records and interview choices that depend on the current label. Moreover $V(d,\mathcal{A}) = \sup_{\mu,h} \langle d, h\rangle_\mu^2 / \max_{\iota \in \mathcal{A}} \|\mathbb{E}_\mu[h \mid \ell_\iota(X)]\|_\mu^2$ over positive profile laws μ and witnesses h.* (b) *At a full-support law P, $W_{\mathcal{A}}(P)/n$ is the local asymptotic minimax risk over arbitrary estimators from the capped records, for interviews drawn without the record's label; drawing them with the label can only lower the constant (Proposition E.2).* (c) *If a depth-b tree τ computes d and $\sigma_P^2 > 0$, then $W_{\mathrm{branch}}(P) = \sigma_P^2$ and*

$$\frac{1}{\rho_P^2} \le \frac{W_{\mathrm{fixed}}(P)}{W_{\mathrm{branch}}(P)} \le N_P(\tau) \le K_{m,b},$$

with equality throughout for the address contrast under the uniform law. For every menu, constant c and wit-

ness h (with $0/0 = 0$), $W_{\mathcal{A}}(P) + (\Delta - c)^2 \geq \mathbb{E}_P[(t - c)h]^2 / \max_{\iota \in \mathcal{A}} \mathbb{E}_P[\mathbb{E}_P[h \mid \ell_\iota(X), Y]^2]$.

The ingredients are standard: minimax duality for linear functionals (Donoho, 1994; Juditsky & Nemirovski, 2009; Polyanskiy & Wu, 2026) and, for one menu, c-optimal design with the efficient augmented inverse-probability score under known coarsening (Elfving, 1952; Pukelsheim, 1993; Robins et al., 1994; Tsiatis, 2006). What is new is the comparison across menus and the exact optimization over the combinatorial menus of r-round and branching interviews. Ratios of W are ratios of records needed when n is large and each class uses its best design for P (Appendices B, C.7.2, E.1).

Computing every protocol class. Pad every transcript to exactly b coordinates, so that an interview is a partition of the cube into cells $\{x : x_T = a\}$ with $|T| = b$. Pointwise unbiasedness of the tables is then equivalent to matching the Fourier coefficients of t of degree at most b, and (3) over a list of interviews is a second-order cone program in the tables and the interview probabilities. Column generation solves a restricted master problem over a few interviews; for its dual witness, the most informative interview of any class $\mathcal{A}_r$ is found by dynamic programming over partial assignments. The witness certifies a lower bound through the last display of Theorem 3, and the master value is an upper bound. Pricing visits $O(\sum_{k \leq b} \binom{m}{k} 4^k)$ partial assignments per iteration and never touches the 2^m profiles (Appendix E.4).

The estimator in practice. For a rule given as a tree τ, the menu of τ's own padded path sets, at most 2^{b-1} forms whatever m is, gives the *path-set program*: a convex program with at most $2^{b-1} \cdot 2^{b+1}$ score variables whose optimum is a certified upper bound on $W_{\mathrm{fixed}}(P)$ (Proposition E.6). Its inputs are the cell masses of the path sets, estimated from reference data or from a pilot; its outputs are form probabilities q_S and tables g_S. The audit draws a form $S_i \sim q$ for record i, asks it, and reports the mean of $Y_i g_{S_i}(X_{i,S_i})/q_{S_i}$ with a standard error from the sample variance. The unbiasedness identities hold at every profile, so the estimate stays exactly unbiased when the reference is wrong; only its variance depends on it (Proposition E.9).

5. What a Working Model Changes

The constants so far assume nothing about the unasked features. If the law of (X, Y) is assumed to lie in a smooth parametric model with tangent space $\mathcal{T}$ at P, the constant of a menu *within the model* is $W^{\mathcal{T}}_{\mathcal{A}}(P) = \inf_q \sup_{h \in \mathcal{T}} \langle t - \Delta, h \rangle^2_P / \sum_\iota q_\iota J_\iota(h)$ with $J_\iota(h) = \mathbb{E}_P[\mathbb{E}_P[h \mid \ell_\iota(X), Y]^2]$, the c-optimal design value for the model's scores (Appendix E.2); it is a lower bound on the local minimax risk within the model, attained when the optimal design has nonsingular information, and the same solver computes it.

Theorem 4 (Adaptivity in a model)**.** (a) *In every model, at every law P and for $1 \leq r \leq b$: $W^{\mathcal{T}}_r(P) \leq K_r\, W^{\mathcal{T}}_{\mathrm{branch}}(P)$.* (b) *In the model $\mathcal{M}_{\mathrm{ci}}$ in which the features are independent given the label, at every law of the model and for every contrast: fixed forms, r-round and branching interviews have the same constant if the design may depend on the record's label, and $W^{\mathrm{ci}}_{\mathrm{fixed}}(P) \leq 2\, W^{\mathrm{ci}}_{\mathrm{branch}}(P)$ if it may not.* (c) *For every $b \geq 3$ and $\epsilon > 0$ there are full-support laws in the pairwise log-linear model, on the $b - 1 + 2^{b-1}$ features of the address contrast and 2^b auxiliary features, at which the constants of that contrast within the pairwise model, and within every log-linear model of higher order, satisfy $W^{(k)}_{\mathrm{fixed}}/W^{(k)}_{\mathrm{branch}} \geq (1 - \epsilon)\, 2^{b-1}/b$.*

Part (a) is the simulation of Theorem 1, which involves no model. Part (b) is the benign extreme, and it holds at rules of any accuracy: when the features are independent given the label, as in a naive-Bayes or latent-class model, a record informs the estimate only through *which* coordinates it reveals, and the constant of a design is $\mathrm{Var}(\mathbb{E}[t \mid Y]) + \sum_y P(y) \sum_j (\hat{d}^y_j)^2/\pi_j(y)$, where $\pi_j(y)$ is the probability that feature j is asked of a record with label y and $\hat{d}^y_j$ is the degree-one coefficient of the rule in class y. Whatever inclusion probabilities a tree realizes, a label-dependent mixture of forms realizes as well; without the label a tree can use early answers as a proxy for it, which is worth at most a factor two. Part (c) shows that parsimony alone is not enough: with a one-hot copy of the address and the products of its indicators with the data bits, tied by strong pairwise couplings, the address contrast is a polynomial of degree two on a set of probability nearly one; it is the address lower bound in another coding, and it shows that the order of a log-linear model measures parsimony only relative to the coding of the features. Between the two extremes we have bounds only at and near product laws, where the gap under a log-linear model of order k is at most $(b + 1)^k$ at uniform marginals, and none in between (Theorem E.5); Section 6 computes the constants there.

6. Experiments

The experiments ask three questions about trained rules. **Q1:** what does pre-commitment cost when the cap equals the depth of the rule and each class uses its best design? **Q2:** what removes that cost: optimizing the forms, a few extra questions, or one more round? **Q3:** what does a working model change, including when it is wrong? Every population P is a held-out or target empirical law mixed with a uniform mass of 1%; the constants are oracle computations at these laws, certified by primal and dual bounds (Appendix F), and only the simulated audits involve sam-

pling.

Q1: the price when the cap equals the depth. A depth-b screening tree asks at most b questions per person, so a validation study under the same burden must respect the cap. We train class-balanced CART trees on twelve yes/no items of the CDC Diabetes Health Indicators, derived from the Behavioral Risk Factor Surveillance System (BRFSS; 253,680 respondents, target diabetes or prediabetes) (Centers for Disease Control and Prevention, 2017), and on twelve binarized thyroid items (Quinlan, 1986), and validate them on a disjoint random half or on respondents of the other sex. Branching administers the rule and takes a plain mean; the question is what forms lose against it. On thyroid, and whenever a tree uses at most b items, one form asks everything the rule needs and nothing is lost. On BRFSS at caps four to six the best pre-committed design needs a median 1.35–1.58 times the respondents of branching (1.00–1.98 over runs; Table 1). Trees for four other BRFSS outcomes (21 variables) and for Spambase (57 features), with forms on the rule's items, give 1.07–2.38 in the 38 runs with a gain that were computed (median 1.46; Table 11). For the 5 rules with more than 13 items the optimum over all forms was not computed; the optimized design on the tree's own path sets, a convex program whose size does not grow with the number of items (Proposition E.6), certifies ratios of at most 2.20–3.97 for them, and it is within 14% of the optimum on the 52 rules where both are known (median 1.5%). Bootstrap resamples of the target records move a ratio by at most 0.6% on BRFSS and 6% on the 2,301 Spambase target records.

Ensembles. For an old and a new random forest of twenty depth-b trees (WDBC, Spambase, Ionosphere; $m \in \{6, 9, 12\}$ features) the contrast is the difference of predicted probabilities, which no single interview computes, so both classes face a design problem. At cap three the best forms need a median 1.51 times the records of the best branching design (range 1.14–1.85), and an optimal two-round design at most 1.09 times (Table 5). Per question, the best forms cost 0.96–1.16 times complete records and the best branching designs 0.76–0.78 times (medians by cap): at the median capped forms roughly break even with complete records, and the cap pays clearly only if interviews branch. On twenty Spambase features the picture is the same at $b = 3$: forms need 1.50–1.61 times the records of branching and two rounds at most 1.08 (Table 7).

Q2: what removes the price. *Optimization.* Much of what a naive comparison credits to adaptivity is design optimization: uniform randomization over all forms of the rule's own items, with optimal tables, needs a median 2.0–4.2 times the records of branching by cap, and uniform matrix sampling over all twelve items 9–23 times. The numbers above are for oracle designs, tuned to the target law. Forms fitted on 20,000 complete source records need a median 5% more and stay exactly unbiased; when such records exist and transport, imputation is a competitor (below). With no reference, the design is learned from a pilot charged to the audit (a quarter of $N = 4{,}000$ records, cross-fitted): with cell masses smoothed by a pairwise fit it costs a median 11–20% more than the oracle of the same run, and without smoothing 15–146% more, the pilot being too small at $b = 6$ (Tables 1 and 15).

Extra questions. With forms longer than the rule is deep, the price in records falls much faster than in the worst case of Theorem 2(c). Over the 52 rules with a gain for which both were computed, optimal forms need a median 1.47 times the records of branching at size b, 1.14 at size $b+1$ (at most 1.60) and 1.02 at size $b + 2$ (at most 1.29; Figure 2b, Table 12). The reason is the width of trained rules, not a failure of the theorem: these trees depend on a median of 3 and at most 7 items beyond their depth, whereas the address contrast depends on $2^{b-1} - 1$. *Counting questions.* Extra questions save records and hardly any questions. Per question asked, and relative to a branching interview charged b questions per record, forms of size b cost a median 1.47, forms of size $b + 1$ cost 1.39, and asking all K items of the rule in a fraction b/K of the records, a mean-budget design, costs 1.50 (1.17–2.40). Against each of these three pre-committed designs branching saves between a quarter and a third of the questions at the median (the best of the three, rule by rule, costs 1.39); a two-round interview with b questions saves them as well. Which design is best depends on whether records or questions are the scarce resource.

One more round. Of the 18 BRFSS diabetes trees with a gain, 11 are computed exactly by a two-round interview, 3 more are within a factor 1.040 of branching, and the remaining 4, at $b = 6$, within 1.32 (Figure 2a).

Q3: what a working model changes. A model buys efficiency with an assumption. In the model of Theorem 4(b), with features independent given the label, the constants are in closed form for every rule, including those whose design-based optimum was not computed: at the fit, forms drawn without the label need at most 1.5% more records than the branching optimum on every BRFSS rule and at most 14% on Spambase (60 rules; Table 21). At the pairwise log-linear fit of each target law, forms need 1.02–1.13 times the records of the best branching design on the diabetes trees (1.00–1.28 over all rules with at most ten items), where the design-based ratio at the same laws is 1.19–1.96 (Table 19); along the strongly coupled laws of Theorem 4(c) the pairwise ratio rises from one to 1.66 at $b = 3$ (Table 20).

The accuracy of a depth-b tree involves interactions of up to b items and the label, so the target laws do not satisfy

Table 1. BRFSS diabetes screening trees on twelve items (six runs per cap: two targets, three seeds): records needed relative to a branching interview that asks the tree's own b questions, which attains the full-information variance; medians over the six runs, range in brackets for the first column. *Pre-committed forms* with pointwise-unbiased scores: the optimal design with forms of size b at the target law, and with one or two extra questions per record (forms on the rule's items); uniform randomization over all forms of the rule's items with optimal tables; the design learned from a cross-fitted pilot of the audit itself with smoothed cell masses, all records charged ($N = 4{,}000$; a run in which one form holds all the rule's items needs no pilot and counts as 1). *Two rounds*: the optimal two-round design of size b (at $b = 6$ the median includes certified upper bounds from feasible two-round designs for 4 runs, Table 10). *Pairwise model*: $W_{\text{fixed}}/W_{\text{branch}}$ within the pairwise log-linear model at its fit. Bracket, reference-fitted designs and the three-way model: Table 8.

	optimal forms of size			two	uniform	pilot,	pairwise
b	b	$b+1$	$b+2$	rounds	forms	all-in	model
4	1.49 [1.00, 1.75]	1.11	1.00	1.00	2.30	1.62	1.09
5	1.35 [1.18, 1.54]	1.08	1.00	1.00	2.04	1.53	1.06
6	1.58 [1.42, 1.98]	1.24	1.09	$\leq$1.05	4.18	1.93	1.07




Figure 2. What removes the price of pre-commitment on trained rules. Records needed relative to the branching interview with b questions, without a model: (a) optimal r-round designs ($r = 1$: fixed forms; $r = b$: branching) for BRFSS diabetes trees (median over runs; at $b = 6$, $r = 2$ includes upper bounds) and forest updates (median over tasks); crosses mark the heuristic of following the root question; (b) optimal forms with c extra questions (thin lines: single rules; markers: medians over the rules computed at that size). (c) Simulated audits: mean squared error relative to the design-based mean under branching, for the design-based form score and for the pairwise EM estimate on three kinds of records (median over 12 trees; bands: quartiles).

the pairwise model, and the comparison is one of efficiency against robustness. At the target laws themselves we compute the limit of the pairwise maximum-likelihood estimate under each design and its sandwich variance (White, 1982; Golden et al., 2019) (17 runs; Table 24). The estimate is well defined essentially when the design observes every pair of items together, the classical condition for split questionnaires (Raghunathan & Grizzle, 1995; Rhemtulla & Little, 2012); the rule's own tree and the optimal design-based forms mostly do not. On pair-covering forms optimized for the model, the limit misses the true accuracy by at most 0.64 points, the squared bias equals the variance at a median of 69,000 records, and the sandwich variance is 0.57–1.15 times that of the branching design that is optimal for the model (median 1.04). In simulated audits of 500 to 32,000 respondents on 12 trees (Figure 2c, Table 22), the mean squared error of forms relative to branching is 1.40–1.55 for the design-based scores and 1.00–1.07 for the pairwise EM estimate up to 8,000 respondents; at 32,000 the bias shows, the pairwise estimate is less accurate than the design-based form score for 5 of the 12 trees, and its intervals cover 0.855–0.953 (Table 18).

Further checks. An optimized design is tailored to one rule: scoring another tree on its records needs 1.55 to 317 times the records of branching, whereas one design shared by three known trees costs each a median 10% more than its own optimum (Appendix F.6). Imputation from complete records of another population beats every capped design when it transports, and is biased in 3 of eight shift settings (Table 17).

7. Related Work

Adaptivity in query models. Within-input adaptivity is classical in decision-tree and property-testing complexity, with round hierarchies (Canonne & Gur, 2018; Li et al., 2026) and pointer chasing (Nisan & Wigderson, 1993). That a q-query adaptive procedure is simulated by 2^q non-adaptive queries to the same input is folklore there; the textbook non-adaptive estimator for a decision tree samples its Fourier coefficients (Kushilevitz & Mansour, 1993) and has sec-

ond moment $\|\hat{d}\|_1^2$, which is 4^{b-1} for the address contrast, against 2^{b-1} here. The locally bounded model of Adar & Fischer (2024), with its exponential separation from non-adaptive testers (Goldreich & Ron, 2022; Chakraborty et al., 2023), is the testing counterpart of our branching class. Adaptivity gaps are also studied in stochastic optimization (Dean et al., 2008; Golovin & Krause, 2011), batched bandits (Perchet et al., 2016; Gao et al., 2019) and adaptive sensing (Arias-Castro et al., 2013). *Information-constrained estimation* fixes a per-sample channel family (Acharya et al., 2019; Barnes et al., 2020; Acharya et al., 2022; Lau & Scarlett, 2026); pointer chasing separates rounds of interaction in local privacy (Joseph et al., 2019; 2020), and interactive protocols estimate some functionals at faster rates (Butucea et al., 2023). Ours is a noiseless analogue with exact constants. *Design and missing data by design.* For one menu our programs are c-optimal design and our solver a vertex-direction method (Fedorov, 1972; Wynn, 1970; Sagnol, 2011; Stenger, 1979; Gabler, 1990; Aronow & Lopatto, 2026). Split questionnaires and matrix sampling assign fixed forms (Raghunathan & Grizzle, 1995; Graham et al., 2006; Thomas et al., 2006; Chipperfield & Steel, 2009; Rhemtulla & Little, 2012), optimized under a model from a pilot (Adigüzel & Wedel, 2008; Jang et al., 2022); adaptive matrix sampling (Gonzalez & Eltinge, 2008), responsive design (Groves & Heeringa, 2006), two-phase designs (Reilly & Pepe, 1995; Gilbert et al., 2014; McIsaac & Cook, 2015; Tao et al., 2020; Chen & Lumley, 2020; 2022) and multistage adaptive tests (Yamamoto et al., 2018; Fink et al., 2024) are two-round interviews in our sense. These lines optimize one protocol within one design space; we compare the spaces. *Evaluation and acquisition in ML.* Restricted-focus-of-attention learning is our fixed-form class (Ben-David & Dichterman, 1998), and budgeted learning fits predictors from such views (Cesa-Bianchi et al., 2011; Hazan & Koren, 2012; Lizotte et al., 2003; Kapoor & Greiner, 2005). Active testing and prediction-powered inference allocate labels (Kossen et al., 2021; Zrnic & Candès, 2024; Angelopoulos et al., 2023), the reverse of our cost structure; active feature acquisition chooses features for individual predictions (Trapeznikov & Saligrama, 2013; Nan & Saligrama, 2017; Covert et al., 2023; von Kleist et al., 2025).

8. Discussion and Limitations

Within-record adaptivity is worth at most 2^{b-1} in records, for every task, model and population, sharply without a model, and it accrues round by round. A model with features independent given the label removes the gap, up to a factor two if forms are drawn without the label, and a pairwise model removes little of it when strong couplings encode the rule's structure. On the trained rules we examined, with the cap equal to the rule's depth, optimal forms needed between 1.0 and 2.38 times the records of branching where the optimum was computed; one extra question lowered the median from 1.47 to 1.14, and one adaptive round came within 1.32 of branching in every screening run; per question, the best of the three pre-committed designs we priced cost 1.39 times branching at the median.

In practice. If the rule uses at most b features, ask them all; if the interview can branch, administer the rule and take a plain mean. If questions must be pre-committed, do not randomize uniformly: solve the path-set program on a reference sample or on a pilot that is charged to the audit, draw one form per record from the resulting distribution, and average the scores, which are unbiased whatever the reference was. If records are scarce and the burden allows one or two questions more than the rule asks, or one batch of follow-ups, use them; if questions are scarce, no pre-committed design we priced helps, and only adaptivity, full branching or one follow-up round, saves them. A pairwise working model on pair-covering forms is nearly as efficient as branching for audits of a few thousand records, but its bias does not shrink, so claims that must not rest on a model should use the design-based scores.

Scope. For a single screening tree the adaptive optimum is the rule itself, so the comparison prices paper forms and panels ordered in one batch against computer-assisted administration of the rule, and it is made at the knife-edge where the cap equals the rule's depth; with one or two questions more, forms lose little on the rules we trained (Figure 2b); for ensembles the cap costs more than pre-commitment (Table 5). The target is one known low-degree functional fixed before collection: records collected for one rule can be nearly useless for another (Appendix F.6), for all monomials of degree b at once uniform forms are optimal (Proposition C.3), and thresholded ensembles exceed small caps. Labels are free and always observed; if labels are expensive, complete records can be cheaper (Appendix F.10). Answers are assumed not to depend on question order or mode; an illustrative flip of the last answer with probability 10% biases the scores by 1.54–2.17 standard errors at $n = 1000$ (Appendix F.9).

What the constants mean. Ratios of W compare designs tuned to the population in the limit of many records per cell, and the finite-sample bracket of Theorem B.1 carries a factor of 104. The populations are unweighted, smoothed empirical laws; no audit was run in the field. Pricing visits on the order of $\binom{m}{b}4^b$ partial assignments per iteration: adaptive classes were solved for $m \le 12$, $b \le 6$ and for $m = 20$ at $b = 3$ only.

Impact Statement

Shorter adaptive questionnaires and test sequences can reduce respondent and patient burden. Branching exposes

people to different questions; order effects, selective nonresponse and subgroup-specific burdens can bias estimates or hide unequal outcomes, and population-average gains do not imply subgroup gains, so deployments should monitor both.

References

Acharya, J., Canonne, C. L., and Tyagi, H. Inference under information constraints: Lower bounds from chi-square contraction. In *Proceedings of the Conference on Learning Theory*, volume 99 of *Proceedings of Machine Learning Research*, pp. 3–17, 2019. URL https://proceedings.mlr.press/v99/acharya19a.html.

Acharya, J., Canonne, C. L., Liu, Y., Sun, Z., and Tyagi, H. Interactive inference under information constraints. *IEEE Transactions on Information Theory*, 68(1):502–516, 2022.

Adar, T. and Fischer, E. Refining the adaptivity notion in the huge object model. In *Approximation, Randomization, and Combinatorial Optimization. Algorithms and Techniques (APPROX/RANDOM 2024)*, volume 317 of *Leibniz International Proceedings in Informatics*, pp. 45:1–45:16, 2024. doi: 10.4230/LIPIcs.APPROX/RANDOM.2024.45.

Adigüzel, F. and Wedel, M. Split questionnaire design for massive surveys. *Journal of Marketing Research*, 45(5):608–617, 2008. doi: 10.1509/jmkr.45.5.608.

Angelopoulos, A. N., Bates, S., Fannjiang, C., Jordan, M. I., and Zrnic, T. Prediction-powered inference. *Science*, 382(6671):669–674, 2023. doi: 10.1126/science.adi6000.

Arias-Castro, E., Candès, E. J., and Davenport, M. A. On the fundamental limits of adaptive sensing. *IEEE Transactions on Information Theory*, 59(1):472–481, 2013.

Aronow, P. M. and Lopatto, P. Minimax unbiased estimation for finite populations with bounded outcomes. arXiv:2605.20572, 2026.

Barnes, L. P., Han, Y., and Özgür, A. Lower bounds for learning distributions under communication constraints via Fisher information. *Journal of Machine Learning Research*, 21(236):1–30, 2020. URL https://jmlr.org/papers/v21/19-737.html.

Ben-David, S. and Dichterman, E. Learning with restricted focus of attention. *Journal of Computer and System Sciences*, 56(3):277–298, 1998. doi: 10.1006/jcss.1998.1569.

Blackwell, D. Equivalent comparisons of experiments. *The Annals of Mathematical Statistics*, 24(2):265–272, 1953. doi: 10.1214/aoms/1177729032.

Boros, E. and Hammer, P. L. Pseudo-Boolean optimization. *Discrete Applied Mathematics*, 123(1–3):155–225, 2002.

Butucea, C., Rohde, A., and Steinberger, L. Interactive versus noninteractive locally differentially private estimation: Two elbows for the quadratic functional. *The Annals of Statistics*, 51(2):464–486, 2023.

Canonne, C. L. and Gur, T. An adaptivity hierarchy theorem for property testing. *Computational Complexity*, 27(4):671–716, 2018. doi: 10.1007/s00037-018-0168-4.

Centers for Disease Control and Prevention. CDC diabetes health indicators. UCI Machine Learning Repository, https://doi.org/10.24432/C53919, 2017. Derived from the CDC Behavioral Risk Factor Surveillance System.

Cesa-Bianchi, N., Shalev-Shwartz, S., and Shamir, O. Efficient learning with partially observed attributes. *Journal of Machine Learning Research*, 12:2857–2878, 2011.

Chakraborty, S., Fischer, E., Ghosh, A., Mishra, G., and Sen, S. Testing of index-invariant properties in the huge object model. In *Proceedings of the 36th Conference on Learning Theory*, volume 195 of *Proceedings of Machine Learning Research*, pp. 3065–3136, 2023. URL https://proceedings.mlr.press/v195/chakraborty23a.html.

Chen, T. and Lumley, T. Optimal multiwave sampling for regression modeling in two-phase designs. *Statistics in Medicine*, 39(30):4912–4921, 2020.

Chen, T. and Lumley, T. Optimal sampling for design-based estimators of regression models. *Statistics in Medicine*, 41(8):1482–1497, 2022.

Chipperfield, J. O. and Steel, D. G. Design and estimation for split questionnaire surveys. *Journal of Official Statistics*, 25(2):227–244, 2009.

Cook, S., McKenzie, P., Wehr, D., Braverman, M., and Santhanam, R. Pebbles and branching programs for tree evaluation. *ACM Transactions on Computation Theory*, 3(2):4:1–4:43, 2012.

Covert, I., Qiu, W., Lu, M., Kim, N., White, N., and Lee, S.-I. Learning to maximize mutual information for dynamic feature selection. In *Proceedings of the 40th International Conference on Machine Learning*, volume 202 of *Proceedings of Machine Learning Research*, pp. 6424–6447, 2023. URL https://proceedings.mlr.press/v202/covert23a.html.

Dean, B. C., Goemans, M. X., and Vondrák, J. Approximating the stochastic knapsack problem: The benefit of adaptivity. *Mathematics of Operations Research*, 33(4):945–964, 2008.

Donoho, D. L. Statistical estimation and optimal recovery. *The Annals of Statistics*, 22(1):238–270, 1994. doi: 10.1214/aos/1176325367.

Elfving, G. Optimum allocation in linear regression theory. *The Annals of Mathematical Statistics*, 23(2):255–262, 1952. doi: 10.1214/aoms/1177729442.

Fedorov, V. V. *Theory of Optimal Experiments*. Academic Press, New York, 1972.

Fink, A., König, C., and Frey, A. Methodological aspects of the highly adaptive testing design for PISA. *Frontiers in Psychology*, 15:1446799, 2024.

Gabler, S. *Minimax Solutions in Sampling from Finite Populations*, volume 64 of *Lecture Notes in Statistics*. Springer, New York, 1990.

Gao, Z., Han, Y., Ren, Z., and Zhou, Z. Batched multi-armed bandits problem. In *Advances in Neural Information Processing Systems*, volume 32, 2019.

Gilbert, P. B., Yu, X., and Rotnitzky, A. Optimal auxiliary-covariate-based two-phase sampling design for semiparametric efficient estimation of a mean or mean difference, with application to clinical trials. *Statistics in Medicine*, 33(6):901–917, 2014. doi: 10.1002/sim.6006.

Gill, R. D. and Levit, B. Y. Applications of the van Trees inequality: a Bayesian Cramér–Rao bound. *Bernoulli*, 1:59–79, 1995.

Gill, R. D., van der Laan, M. J., and Robins, J. M. Coarsening at random: Characterizations, conjectures and counter-examples. In Lin, D.-Y. and Fleming, T. R. (eds.), *Proceedings of the First Seattle Symposium in Biostatistics: Survival Analysis*, pp. 255–294. Springer, 1997.

Golden, R. M., Henley, S. S., White, H., and Kashner, T. M. Consequences of model misspecification for maximum likelihood estimation with missing data. *Econometrics*, 7(3):37, 2019.

Goldreich, O. and Ron, D. Testing distributions of huge objects. In *Innovations in Theoretical Computer Science*, volume 215 of *LIPIcs*, pp. 78:1–78:19, 2022. doi: 10.4230/LIPIcs.ITCS.2022.78.

Golovin, D. and Krause, A. Adaptive submodularity: Theory and applications in active learning and stochastic optimization. *Journal of Artificial Intelligence Research*, 42:427–486, 2011.

Gonzalez, J. M. and Eltinge, J. L. Adaptive matrix sampling for the Consumer Expenditure Quarterly Interview Survey. In *Proceedings of the Section on Survey Research Methods*, pp. 2081–2088. American Statistical Association, 2008.

Graham, J. W., Taylor, B. J., Olchowski, A. E., and Cumsille, P. E. Planned missing data designs in psychological research. *Psychological Methods*, 11(4):323–343, 2006. doi: 10.1037/1082-989X.11.4.323.

Groves, R. M. and Heeringa, S. G. Responsive design for household surveys: Tools for actively controlling survey errors and costs. *Journal of the Royal Statistical Society: Series A*, 169(3):439–457, 2006. doi: 10.1111/j.1467-985X.2006.00423.x.

Hahn, J., Hirano, K., and Karlan, D. Adaptive experimental design using the propensity score. *Journal of Business & Economic Statistics*, 29(1):96–108, 2011. doi: 10.1198/jbes.2009.08161.

Hazan, E. and Koren, T. Linear regression with limited observation. In *Proceedings of the 29th International Conference on Machine Learning*, pp. 807–814, 2012.

Heitjan, D. F. and Rubin, D. B. Ignorability and coarse data. *The Annals of Statistics*, 19(4):2244–2253, 1991. doi: 10.1214/aos/1176348396.

Hopkins, M., Reeber, E., Forman, G., and Suermondt, J. Spambase. UCI Machine Learning Repository, 1999.

Jang, Zhu, and Yu. A-optimal split questionnaire designs for multivariate continuous variables. arXiv:2203.04883, 2022.

Joseph, M., Mao, J., Neel, S., and Roth, A. The role of interactivity in local differential privacy. In *IEEE Symposium on Foundations of Computer Science*, pp. 94–105, 2019. URL https://arxiv.org/abs/1904.03564.

Joseph, M., Mao, J., and Roth, A. Exponential separations in local differential privacy. In *Proceedings of the ACM-SIAM Symposium on Discrete Algorithms*, pp. 515–527, 2020. doi: 10.1137/1.9781611975994.31.

Juditsky, A. B. and Nemirovski, A. S. Nonparametric estimation by convex programming. *The Annals of Statistics*, 37:2278–2300, 2009. doi: 10.1214/08-AOS654.

Kairouz, P., McMahan, H. B., et al. Advances and open problems in federated learning. *Foundations and Trends in Machine Learning*, 14(1–2):1–210, 2021.

Kapoor, A. and Greiner, R. Learning and classifying under hard budgets. In *Machine Learning: ECML 2005*, volume 3720 of *Lecture Notes in Computer Science*, pp. 170–181. Springer, 2005.

Kossen, J., Farquhar, S., Gal, Y., and Rainforth, T. Active testing: Sample-efficient model evaluation. In *Proceedings of the 38th International Conference on Machine Learning*, volume 139 of *Proceedings of Machine Learning Research*, pp. 5753–5763, 2021.

Kushilevitz, E. and Mansour, Y. Learning decision trees using the Fourier spectrum. *SIAM Journal on Computing*, 22(6):1331–1348, 1993.

Lau, I. and Scarlett, J. Sequential 1-bit mean estimation with near-optimal sample complexity. In *Proceedings of the 29th International Conference on Artificial Intelligence and Statistics*, volume 300 of *Proceedings of Machine Learning Research*, pp. 217–225, 2026.

Li, R., Fu, H., Shi, Q., and Yao, P. Sharp two-round adaptivity and round hierarchies for semantic regular expressions. arXiv:2607.22799, 2026.

Lizotte, D., Madani, O., and Greiner, R. Budgeted learning of naive-Bayes classifiers. In *Proceedings of the Conference on Uncertainty in Artificial Intelligence*, 2003.

McIsaac, M. A. and Cook, R. J. Adaptive sampling in two-phase designs: a biomarker study for progression in arthritis. *Statistics in Medicine*, 34(21):2899–2912, 2015.

Nan, F. and Saligrama, V. Adaptive classification for prediction under a budget. In *Advances in Neural Information Processing Systems*, volume 30, 2017.

Nisan, N. and Wigderson, A. Rounds in communication complexity revisited. *SIAM Journal on Computing*, 22 (1):211–219, 1993. doi: 10.1137/0222016.

O'Donnell, R. *Analysis of Boolean Functions*. Cambridge University Press, 2014.

Perchet, V., Rigollet, P., Chassang, S., and Snowberg, E. Batched bandit problems. *The Annals of Statistics*, 44(2): 660–681, 2016.

Polyanskiy, Y. and Wu, Y. Dualizing Le Cam's method for functional estimation I: General theory. *The Annals of Statistics*, 54(1):1–24, 2026. doi: 10.1214/25-AOS2498.

Pukelsheim, F. *Optimal Design of Experiments*. Wiley, 1993.

Quinlan, R. Thyroid disease. UCI Machine Learning Repository, https://doi.org/10.24432/C5D010, 1986.

Raghunathan, T. E. and Grizzle, J. E. A split questionnaire survey design. *Journal of the American Statistical Association*, 90(429):54–63, 1995. doi: 10.1080/01621459.1995.10476488.

Reilly, M. and Pepe, M. S. A mean score method for missing and auxiliary covariate data in regression models. *Biometrika*, 82(2):299–314, 1995.

Rhemtulla, M. and Little, T. D. Planned missing data designs for research in cognitive development. *Journal of Cognition and Development*, 13(4):425–438, 2012.

Robins, J. M., Rotnitzky, A., and Zhao, L. P. Estimation of regression coefficients when some regressors are not always observed. *Journal of the American Statistical Association*, 89(427), 1994. doi: 10.1080/01621459.1994.10476818.

Romano, J. D., Le, T. T., La Cava, W., Gregg, J. T., Goldberg, D. J., Chakraborty, P., Ray, N. L., Himmelstein, D., Fu, W., and Moore, J. H. PMLB v1.0: an open-source dataset collection for benchmarking machine learning methods. *Bioinformatics*, 38(3):878–880, 2022.

Sagnol, G. Computing optimal designs of multiresponse experiments reduces to second-order cone programming. *Journal of Statistical Planning and Inference*, 141(5): 1684–1708, 2011. doi: 10.1016/j.jspi.2010.11.031.

Sigillito, V., Wing, S., Hutton, L., and Baker, K. Ionosphere. UCI Machine Learning Repository, 1989.

Sion, M. On general minimax theorems. *Pacific Journal of Mathematics*, 8:171–176, 1958. doi: 10.2140/pjm.1958.8.171.

Stenger, H. A minimax approach to randomization and estimation in survey sampling. *The Annals of Statistics*, 7 (2):395–399, 1979.

Tao, R., Zeng, D., and Lin, D.-Y. Optimal designs of two-phase studies. *Journal of the American Statistical Association*, 115(532):1946–1959, 2020. doi: 10.1080/01621459.2019.1671200.

Thomas, N., Raghunathan, T. E., Schenker, N., Katzoff, M. J., and Johnson, C. L. An evaluation of matrix sampling methods using data from the National Health and Nutrition Examination Survey. *Survey Methodology*, 32 (2):217–231, 2006.

Trapeznikov, K. and Saligrama, V. Supervised sequential classification under budget constraints. In *Proceedings of the Sixteenth International Conference on Artificial Intelligence and Statistics*, volume 31 of *Proceedings of Machine Learning Research*, pp. 581–589, 2013.

Tsiatis, A. A. *Semiparametric Theory and Missing Data*. Springer, 2006. doi: 10.1007/0-387-37345-4.

Turney, P. D. Cost-sensitive classification: Empirical evaluation of a hybrid genetic decision tree induction algorithm. *Journal of Artificial Intelligence Research*, 2:369–409, 1995. doi: 10.1613/jair.120.

von Kleist, H., Zamanian, A., Shpitser, I., and Ahmidi, N. Evaluation of active feature acquisition methods for time-varying feature settings. *Journal of Machine Learning Research*, 26(60):1–84, 2025.

Wentges, P. Weighted Dantzig–Wolfe decomposition for linear mixed-integer programming. *International Transactions in Operational Research*, 4(2):151–162, 1997.

White, H. Maximum likelihood estimation of misspecified models. *Econometrica*, 50(1):1–25, 1982.

Wolberg, W., Mangasarian, O., Street, N., and Street, W. Breast cancer wisconsin (diagnostic), 1993. URL https://archive.ics.uci.edu/dataset/17/breast-cancer-wisconsin-diagnostic.

Wynn, H. P. The sequential generation of D-optimum experimental designs. *The Annals of Mathematical Statistics*, 41(5):1655–1664, 1970.

Yamamoto, K., Shin, H. J., and Khorramdel, L. Multistage adaptive testing design in international large-scale assessments. *Educational Measurement: Issues and Practice*, 37(4):16–27, 2018.

Zrnic, T. and Candès, E. Active statistical inference. In *Proceedings of the 41st International Conference on Machine Learning*, volume 235 of *Proceedings of Machine Learning Research*, pp. 62993–63010, 2024. URL https://proceedings.mlr.press/v235/zrnic24a.html.

A. Guide, Notation and a Worked Example

Guide. Appendix B proves Theorem 3; Appendix C proves Theorem 1 and its extensions to finite samples, larger alphabets, polynomial dimension and mean budgets; Appendix D proves parts (a) and (b) of Theorem 2 and Appendix C.2 part (c) (Theorem C.2), including the product construction in polynomial dimension, and reports the round constants of the binary pointer tree at small caps; Appendix E proves the population statements (Theorems 1(ii), 2(a) and 3, Propositions E.2, E.3, E.6 and E.9), the statements about models (Theorem 4, Proposition E.4; Appendix E.2), and describes the solver and the form program; Appendix F gives every experimental detail; Appendix G extends the estimator to other proper scores; Appendix H describes code and checks.

Notation.

Symbol	Meaning
$d = p_1 - p_0$, $t(x, y) = yd(x)$	Contrast of the two frozen rules and the per-record target; $\Delta(P) = \mathbb{E}_P[t]$
ι, $\ell_\iota(x)$, $\mathcal{A}$	Interview, its transcript, and a finite menu of interviews
$\mathcal{A}_r(b)$	r-round interviews with at most b queries; $\mathcal{A}_1$ fixed forms, $\mathcal{A}_b$ branching trees
q_ι, g_ι	Interview probabilities and transcript score tables of a design
$V(d, \mathcal{A})$, V_r	Worst-case first-order constant (2); its value on $\mathcal{A}_r(b)$
$\Phi_n(d, \mathcal{A})$	Finite-sample program (B.1), within a factor 104 of the minimax risk (Theorem B.1)
$W_\mathcal{A}(P)$, $M_\mathcal{A}(P)$	Smallest variance and second moment of a pointwise-unbiased design at P; $M = W + \Delta^2$
σ_P^2, ρ_P^2	$\mathrm{Var}_P(t)$; largest share of it explained by one form, $\max_{\lvert T\rvert=b} \mathrm{Var}_P(\mathbb{E}_P[t \mid X_T, Y])/\sigma_P^2$
$N_P(\tau)$	Rényi-1/2 effective number of path sets, from the centered weights $v_S = \mathbb{E}_P[(t - \Delta)^2 \mathbf{1}\{S_\tau = S\}]/\sigma_P^2$
$W_{\mathrm{PS}}(P)$	Optimum over the padded path sets of a tree computing d (Proposition E.6)
$W_\mathcal{A}^{\mathrm{lab}}(P)$	Optimum when interview probabilities may depend on the label (Proposition E.2)
$\mathcal{T}$, $J_\iota(h)$, $W_\mathcal{A}^\mathcal{T}(P)$	Tangent space of a model at P, information of interview ι about $h \in \mathcal{T}$, and the local minimax constant within the model (E.1)
$\mathcal{M}_k$, $W_\mathcal{A}^{(k)}(P)$	Log-linear model with interactions of order at most k among the features and the label, and its constant
$K_{m,b}$	$\min\{2^{b-1}, \binom{m-1}{b-1}\}$
κ, $D = 2^\kappa$	Address bits and data bits of the address contrast; L levels of the binary pointer tree
U	Uniform law on the profile cube
B; n_1	Larger hard cap in the mixed-budget result; pilot size in Proposition E.9

The worked example in full. Let $m = 3$, $b = 2$ and $d(x) = [(1 + x_1)x_2 + (1 - x_1)x_3]/2$, and let P have uniform profiles and an independent uniform label, so that $\Delta(P) = 0$ and $\sigma_P^2 = \mathbb{E}[d^2] = 1$. The tree asks x_1 and then x_2 or x_3; it computes d, so $V_{\mathrm{branch}} = 1$ and $W_{\mathrm{branch}}(P) = 1$. The Fourier expansion is $d = \frac{1}{2}x_2 + \frac{1}{2}x_3 + \frac{1}{2}x_1x_2 - \frac{1}{2}x_1x_3$. Each coefficient carries a quarter of the variance, and every form of size two contains exactly two of the four sets: $\{1, 2\}$ captures x_2 and x_1x_2, $\{1, 3\}$ captures x_3 and x_1x_3, and $\{2, 3\}$ captures x_2 and x_3. Hence $\rho_P^2 = 1/2$ and $W_{\mathrm{fixed}}(P) \geq 2$. The two padded path sets $\{1, 2\}$ and $\{1, 3\}$ each carry weight $v_S = 1/2$, so $N_P = (2 \cdot 2^{-1/2})^2 = 2$, and drawing either form with probability $1/2$ and returning $2Yx_2\mathbf{1}\{x_1 = 1\}$ or $2Yx_3\mathbf{1}\{x_1 = -1\}$ attains second moment 2 at every profile. Thus $V_{\mathrm{fixed}} = W_{\mathrm{fixed}}(P) = 2 = K_{3,2}$, both ends of Theorem 3 coincide, and the optimized path-set design of Proposition E.6 equals the optimum. If instead the label agrees with $d(X)$ with probability 0.9, then $t = \pm 1$ with $\Delta = 0.8$ and $\sigma_P^2 = 0.36$; the centered weights are again $1/2$ each, so $N_P = 2$ and the fixed design that returns $0.8 + 2(t - 0.8)$ on a completed path and 0.8 otherwise has variance $2\sigma_P^2$, whereas the uncentered path-guessing score has variance $2 - 0.64 = 1.36 = 3.78\,\sigma_P^2$. Section 3 generalizes the example to κ address bits.

B. Identification and the Exact Characterization

B.1. General Statistical Guarantees

For a design distribution q on $\mathcal{A}$ and a table g_ι on each transcript alphabet, abbreviate $g_\iota(x) = g_\iota(\ell_\iota(x))$ and $s_g(x) = \sum_\iota g_\iota(x)$. Define

$$B_n(q, g; x) = (d(x) - s_g(x))^2 + \frac{1}{n} \sum_{\iota \in \mathcal{A}} \frac{g_\iota(x)^2}{q_\iota},$$
$$\Phi_n(d, \mathcal{A}) = \inf_{q,g} \max_{x \in \mathcal{X}} B_n(q, g; x). \tag{B.1}$$

A design of mass zero may only have a zero table, in which case its contribution is zero. The first term permits a controlled approximation of the update. The second accounts for observing only one of the tables per respondent.

Theorem B.1 (Finite-sample risk)**.** *For every finite $\mathcal{X}, \mathcal{A}$, every contrast d and every $n \geq 1$,*

$$\Phi_n(d, \mathcal{A})/104 \;\leq\; R_{\mathcal{A}}(d, n) \;\leq\; \Phi_n(d, \mathcal{A}).$$

The upper bound uses one iid design and the mean of $Y g_\iota(\ell_\iota(X))/q_\iota$. The lower bound allows the choice of interview to depend on all previous records.

The constant 104 is conservative. This is a convex quadratic-perspective program with a finite second-order cone formulation. Its size depends on enumerated profiles and interviews; Section E.5 instead uses a compact tree representation for a restricted menu.

Theorem B.2 (Exact leading variance)**.** *If $d \in \mathcal{H}_{\mathcal{A}}$, then $\lim_{n\to\infty} n R_{\mathcal{A}}(d, n) = V(d, \mathcal{A})$ for fixed $d, \mathcal{X}, \mathcal{A}$. An iid design and an unbiased score give risk at most $V(d, \mathcal{A})/n$ for every n.*

An optimal iid design already achieves the first-order worst-case coefficient. The lower bound even permits observing the current label before selecting an interview, so early labels do not improve this coefficient either. Within-person branching changes the available observation maps themselves. Population-specific performance can still benefit from learning a design from previous people.

Why the characterization holds. Conditioning on the complete record bounds the score mean's squared bias and variance by (B.1). For the converse, put $\Pi_\iota h = \mathbb{E}_\mu[h \mid \ell_\iota(X)]$ and $J_\mu(h) = \max_\iota \|\Pi_\iota h\|^2_{2,\mu}$. Convex duality gives

$$\Phi_n(d, \mathcal{A}) = \sup_{\mu, h} \frac{\langle d, h\rangle_\mu^2}{\|h\|^2_{2,\mu} + n J_\mu(h)}. \tag{B.2}$$

A witness perturbs the conditional label mean. Its numerator measures the change in the target, while $J_\mu(h)$ limits information from each interview. A change of measure and the KL chain rule handle sequential choices; a local van Trees argument gives the exact coefficient. Appendix B.3 gives the duality and information proofs. These are established tools for linear-functional estimation; the sharp comparison of query menus is the main result.

B.1.1. An Exact One-Feature Example

For an additive contrast $d(x) = c_0 + \sum_{j=1}^m c_j x_j$ on the unrestricted sign cube, put $M = \sum_{j=0}^m |c_j| = \|d\|_\infty$.

Proposition B.3. *For every $n \geq 1$ the minimax risk is $M^2/(1 + \sqrt{n})^2$ both with full profiles and with at most one feature query.*

Choose j with probability $|c_j|/M$, interpret $x_0 = 1$, and return the sign $Y \operatorname{sgn}(c_j) x_j$. Its mean is Δ/M. Estimating that mean reduces to a binomial problem. A population concentrated at an extremal profile gives the same binomial problem to an evaluator who sees all features (Appendix C.10). Equality concerns worst-case populations. The best length at equal monetary cost can differ for particular populations.

B.2. Identifiability and Irreducible Error

Let L_ι lift a transcript table to $\mathbb{R}^{\mathcal{X}}$, so $(L_\iota u)(x) = u(\ell_\iota(x))$. The space $\mathcal{H}_{\mathcal{A}} = \sum_\iota \operatorname{im}(L_\iota)$ is closed. Put $\delta = \inf_{v \in \mathcal{H}_{\mathcal{A}}} \|d - v\|_\infty$. In fact $\lim_{n\to\infty} R_{\mathcal{A}}(d, n) = \delta^2$. For the upper bound choose an approximant v attaining δ, decompose $v = \sum_\iota L_\iota g_\iota$,

and give every used interview positive probability. The mean score is uniformly within δ of Δ, and its variance tends to zero uniformly over P since all alphabets are finite.

If $\delta > 0$, finite-dimensional norm duality gives a signed vector w of ℓ_1 norm one annihilating $\mathcal{H}_{\mathcal{A}}$ with $\sum_x w(x)d(x) = \delta$. This follows by separating d from the open ℓ_∞ ball of radius δ around that subspace, or by duality of the linear program for uniform approximation. In particular $\sum_{x:\ell_\iota(x)=u} w(x) = 0$ for every ι, u. Give X the law $|w|$, and in the two populations set $Y = \pm \operatorname{sgn} w(X)$. For each y, each transcript cell has identical joint probability with $Y = y$ in the two populations. This remains true if y determines which interview is used. Sequentially conditioning on past records gives identical transcript laws for n interviews while the gains are $\pm\delta$. The average of the two squared risks is at least δ^2. The same reasoning gives necessity in the identification criterion.

For a binary coordinate mask T its cells span all characters χ_S with $S \subseteq T$ and no other characters, by the uniqueness of the multilinear expansion. A decision-tree leaf indicator is a product of up to b factors $(1 \pm x_i)/2$. Conversely the fixed query of S recovers χ_S for every $|S| \le b$. This proves the common identification threshold.

B.3. The General Minimax Characterization

Let $\|\cdot\|_\mu$ denote the norm in $L^2(\mu)$, identifying functions equal on the support of μ. All objects below are finite-dimensional. We can take q strictly positive when computing infima. If a feasible boundary design has $q_\iota = 0$, its table is zero; mixing q with a positive design and taking the mixture coefficient to zero recovers the same objective. Empty transcript cells can be discarded.

B.3.1. The Dual Form

For a fixed μ define, on the span of the transcript subspaces of $L^2(\mu)$,

$$N_\mu(v) = \inf_{\sum_\iota f_\iota = v} \sum_\iota \|f_\iota\|_\mu, \qquad f_\iota \in \operatorname{im}(L_\iota);$$

extend it by $+\infty$ outside that span. The dual seminorm on the full space is $N_\mu^*(h) = \max_\iota \|\Pi_\iota h\|_\mu$. Indeed $|\langle h, \sum_\iota f_\iota\rangle_\mu| \le \max_\iota \|\Pi_\iota h\|_\mu \sum_\iota \|f_\iota\|_\mu$, and a single normalized projection onto a maximizing subspace attains the bound. The closed unit ball of N_μ is the image of a compact product-space unit ball, so its squared norm is lower semicontinuous.

Maximizing over profiles in (B.1) is equivalent to maximizing its expectation over all μ. The objective is continuous affine in μ on the compact probability simplex and continuous convex in (q, g) for positive q. Sion's minimax theorem exchanges maximum and infimum (Sion, 1958). For fixed tables, Cauchy–Schwarz and $q_\iota \propto \|L_\iota g_\iota\|_\mu$ give

$$\inf_q \sum_\iota \|L_\iota g_\iota\|_\mu^2 / q_\iota = \Big(\sum_\iota \|L_\iota g_\iota\|_\mu \Big)^2$$

with zeros handled by limits. Thus at fixed μ the inner value is $\inf_v \{\|d - v\|_\mu^2 + N_\mu(v)^2/n\}$. By finite-dimensional Fenchel duality this equals

$$\sup_h \{2\langle d, h\rangle_\mu - \|h\|_\mu^2 - n N_\mu^*(h)^2\}.$$

To verify the factors, set $F(v) = \|d - v\|_\mu^2$ and $G(v) = N_\mu(v)^2/n$. Their conjugates satisfy

$$F^*(u) = \langle u, d\rangle_\mu + \|u\|_\mu^2/4, \qquad G^*(u) = n N_\mu^*(u)^2/4.$$

Fenchel duality gives $\inf_v (F + G)(v) = \sup_u [-F^*(-u) - G^*(u)]$; substituting $u = 2h$ gives the displayed expression. There is no duality gap because the first squared norm is finite and continuous everywhere and the second is proper convex. Optimizing over the scale of h proves (B.2).

Relation to fixed-channel constants. For a prescribed iid observation kernel, the χ^2-modulus bound of Polyanskiy & Wu (2026) brackets MSE between the squared modulus divided by $(1 + \sqrt{e})^2$ and the squared modulus, under its stated assumptions. Juditsky & Nemirovski (2009) instead formulate their main comparison using a confidence-level error criterion. Neither modulus is the joint-design objective in (B.1). Our factor 104 is a conservative conversion for that objective under sequential interview choice, not an improvement on fixed-channel constants. The additional argument is a witness whose information is bounded simultaneously for every eligible interview, including the two choices selected by the current label.

B.3.2. Finite-Sample Converse

The upper bound in Theorem B.1 follows from the conditional-score calculation in Section 4. For the lower bound fix μ, h and normalize $\|h\|_\mu^2 + nJ_\mu(h) = 1$. Write $C_\mu = 1 + \mathbb{E}_\mu|h| \le 2$ and set

$$\nu(x) = \mu(x)(1 + |h(x)|)/C_\mu, \qquad u(x) = \mu(x)h(x)/\nu(x)$$

on the support of μ, with $u = 0$ elsewhere. We have $\|u\|_\infty \le 2$, $\nu \ge \mu/2$, and $\mathbb{E}_\nu[du] = \langle d, h\rangle_\mu$. For every partition into transcript cells C,

$$\sum_C \nu(C)\,\mathbb{E}_\nu[u \mid C]^2 = \sum_{C:\nu(C)>0} \frac{\left(\sum_{x\in C}\mu(x)h(x)\right)^2}{\nu(C)} \le 2\|\Pi_\iota h\|_\mu^2.$$

For $\vartheta = 1/5$ define $P_\pm(x,y) = \nu(x)(1 \pm \vartheta y u(x))/2$. Their gains are $\pm\vartheta\langle d, h\rangle_\mu$. Condition on past interviews and on independent random coins. Under the null law with $X \sim \nu$ and Y fair and independent, the probability $p_0(z)$ of a leaf $z = (y, C)$ changes to $p_\pm(z) = p_0(z)(1 \pm \vartheta s_z)$ with $s_z = y\mathbb{E}_\nu[u \mid C]$. If the label selects interview ι_y, the null second moment is

$$\mathbb{E}_0 s_z^2 = \frac{1}{2}\sum_{y\in\{-1,1\}}\sum_{C\in\iota_y}\nu(C)\mathbb{E}_\nu[u \mid C]^2 \le \frac{1}{2}\sum_y 2\|\Pi_{\iota_y}h\|_\mu^2 \le 2J_\mu(h).$$

Also $|s_z| \le 2$. The conditional divergence satisfies

$$D_{\mathrm{KL}}(p_+\|p_-) \le \chi^2(p_+\|p_-) \le \frac{4\vartheta^2}{1-2\vartheta}\mathbb{E}_0 s_z^2 \le \frac{8\vartheta^2}{1-2\vartheta}J_\mu(h).$$

The bound is uniform in the realized past. Independent protocol randomness can be revealed at the start, then discarded by data processing. The KL chain rule bounds the divergence of the two full experiments by $8/15$. Pinsker's inequality bounds their total variation by $2/\sqrt{15}$. Classifying an estimate by the nearer of the two parameter values yields worst-case squared risk at least $\vartheta^2\langle d, h\rangle_\mu^2(1 - 2/\sqrt{15})/2 \ge \langle d, h\rangle_\mu^2/104$. Taking the supremum in (B.2) proves Theorem B.1. The numerical inequality is equivalent to $15 \cdot 27^2 > 104^2$. The losses come from the change of measure, the KL-to-total-variation conversion, and testing two alternatives. The same calculation applies when the current label cannot select the form; this constant is not caused solely by label-dependent selection.

B.3.3. Exact First-Order Constant

Assume $d \in \mathcal{H}_\mathcal{A}$. Apply the same minimax exchange to (2), retaining the affine pointwise constraint $s_g = d$. For strictly positive μ the pointwise constraint is equivalent to equality in $L^2(\mu)$, so the fixed-μ optimum is $N_\mu(d)^2$. Restricting the supremum to such μ preserves its value. Indeed mixing any μ with ϵ times a positive law gives an inner objective at least $(1 - \epsilon)$ times the original one for every feasible design. By duality,

$$V(d, \mathcal{A}) = \sup_{\mu>0,\, h:\, J_\mu(h)>0} \frac{\langle d, h\rangle_\mu^2}{J_\mu(h)}. \tag{B.3}$$

The zero contrast has value zero. Equation (B.2) and (B.3) also give $n\Phi_n \to V$: the upper bound is V by a zero-residual design, and every fixed witness in (B.3) gives the matching lower limit.

For a lower bound on $nR_\mathcal{A}$, fix positive μ, h and set $P_\theta(x,y) = \mu(x)(1 + \theta y h(x))/2$, $|\theta| \le \eta < 1/\|h\|_\infty$. Then $\Delta(P_\theta) = \theta c$, with $c = \langle d, h\rangle_\mu$. For each history and deterministic choice of interviews the conditional leaf law has the form $p_\theta(z) = p_0(z)(1 + \theta s_z)$. At the null $\mathbb{E}_0 s_z^2 \le J_\mu(h)$ and $|s_z| \le \|h\|_\infty$. Its Fisher information at θ is therefore at most $J_\mu(h)/(1 - \eta\|h\|_\infty)$. Conditional likelihood scores are martingale differences, so the full information is bounded by n times this quantity, including for a randomized protocol with its independent seed revealed.

Use a smooth prior π_η on $[-\eta, \eta]$ vanishing at both endpoints with finite information $I(\pi_\eta)$, for example $\eta^{-1}\cos^2(\pi\theta/(2\eta))$. Integration by parts in the joint data–parameter law gives $\mathbb{E}[(\widehat{\Delta} - \theta c)\partial_\theta \log(p_\theta\pi_\eta)] = c$. Cauchy–Schwarz gives the van Trees inequality (Gill & Levit, 1995), here a lower bound of

$$\frac{c^2}{nJ_\mu(h)/(1 - \eta\|h\|_\infty) + I(\pi_\eta)}$$

on the prior risk, and hence on worst-case risk. First take $n \to \infty$ after multiplying by n, then $\eta \downarrow 0$, then optimize (B.3). The means of the unbiased scores in (2) give the opposite inequality for every n, proving Theorem B.2.

B.3.4. When the Constant Is Attained in the Unbiased Class

For a fixed positive design law q such that the eligible cells span d, the analogous program $V_q(d)$ fixes q in (2). Its value divided by n is exactly the worst-case minimax MSE among globally unbiased estimators with iid design q. To see the lower bound, choose $X \sim \mu$ of full support and an independent fair label. For a symmetric unbiased statistic U_n of observations $O_i = (\iota_i, \ell_{\iota_i}, Y_i)$ put $H(O_1) = \mathbb{E}[U_n \mid O_1] - \mathbb{E}U_n$. Orthogonality of the one-observation projections gives $\operatorname{Var} U_n \geq n\mathbb{E}H^2$. Perturbing the joint law to $\mu(x)(1 + tyh(x))/2$ and differentiating unbiasedness gives $\langle d, h\rangle_\mu = n\mathbb{E}[H(O_1)Y_1 h(X_1)]$ for all h. Thus $d = \sum_\iota L_\iota g_\iota$ with $g_\iota(\ell) = nq_\iota \mathbb{E}[YH \mid \iota, \ell]$. Jensen bounds $\operatorname{Var} U_n$ below by $n^{-1} \inf_{s_g = d} \sum_\iota \|L_\iota g_\iota\|_\mu^2 / q_\iota$. Taking the supremum over full-support μ and exchanging extrema as above gives the claim. Symmetrization costs no risk with iid records.

C. The Sharp Gap and Its Extensions

C.1. The Sharp Universal Gap

We prove Theorem 1 and the program comparison

$$\Phi_n(d, \mathcal{A}_{\text{fixed}}(b)) \leq \Phi_{n/K_{m,b}}(d, \mathcal{A}_{\text{branch}}(b)). \tag{C.1}$$

Here the program is defined for any positive real sample-size argument. All scores in the construction below depend only on the queried feature values; the estimator subsequently multiplies them by Y. Theorems B.1 and B.2 already compare these constructions with the larger class of history-dependent protocols, including branching choices based on the current label.

C.1.1. Simulating One Tree without Branching

Let τ be a deterministic tree of depth at most b on $m \geq b$ coordinates, and let $H_\tau(x)$ be a score constant on each of its leaves. A repeated query on a path can be removed because its answer is already known. At every leaf with fewer than b distinct queries, append unused coordinates in increasing index order until the path length is b. Copy the original score to all descendants. For a tree with no queries, start with any coordinate. The padded tree has exactly b distinct queries on every path and the same score at every x.

Let $S_\tau(x)$ be its realized set of coordinates and let $\mathcal{S}_\tau$ be the collection of distinct such sets. Put $L_\tau = |\mathcal{S}_\tau|$. All sets contain the root, giving $L_\tau \leq \binom{m-1}{b-1}$. The first $b-1$ answers fix all queried coordinates, even though the final answer can change the leaf score. Thus $L_\tau \leq 2^{b-1}$, proving $L_\tau \leq K_{m,b}$.

Independently of X, Y, choose S uniformly from $\mathcal{S}_\tau$ and observe X_S. Run the padded tree until a queried coordinate is unavailable or a leaf is reached. Let $M_{\tau,S}(X_S)$ be the indicator of reaching a leaf. This event is measurable from the fixed-form transcript. For each full profile x,

$$M_{\tau,S}(x_S) = 1 \quad \Longleftrightarrow \quad S = S_\tau(x).$$

Indeed equality allows every successive query to be answered; conversely completion means that all b distinct coordinates of the actual path lie in the chosen set of size b. Define the simulated score to be $\widetilde{H}_{\tau,S} = L_\tau H_\tau$ on completion and zero otherwise. Then, pointwise in x,

$$\mathbb{E}_S[\widetilde{H}_{\tau,S}(x_S)] = H_\tau(x), \qquad \mathbb{E}_S[\widetilde{H}_{\tau,S}(x_S)^2] = L_\tau H_\tau(x)^2. \tag{C.2}$$

C.1.2. Mixtures and the Finite-Sample Program

For any feasible branching design (q, g) take $H_\tau = g_\tau / q_\tau$ on its positive-mass trees. Draw the tree with probability q_τ, then draw its path set as above. The resulting fixed-form procedure has conditional mean $s_g(x)$ and conditional second moment at most

$$K_{m,b} \sum_\tau g_\tau(x)^2 / q_\tau.$$

The same coordinate set can arise from several tree tags. To represent the construction using only the untagged fixed-mask menu, give S mass $q'_S = \sum_{\tau : S \in \mathcal{S}_\tau} q_\tau / L_\tau$ and replace its score by the weighted average of the tagged scores, with weights $(q_\tau / L_\tau)/q'_S$. The tag is drawn before the person, so these weights do not depend on the population or the answers. Writing

$p_{\tau,S} = q_\tau/L_\tau$ and $\overline{H}_S = \sum_\tau p_{\tau,S}\widetilde{H}_{\tau,S}/q'_S$, the reduction satisfies, pointwise,

$$\sum_S q'_S \overline{H}_S = \sum_{\tau,S} p_{\tau,S}\widetilde{H}_{\tau,S}, \qquad \sum_S q'_S \overline{H}_S^2 \le \sum_{\tau,S} p_{\tau,S}\widetilde{H}^2_{\tau,S}.$$

This is conditional Jensen averaging over the tag given the fixed set. Zero-mass forms are discarded.

For an unbiased branching design this gives $V_{\mathrm{fixed}} \le K_{m,b} V_{\mathrm{branch}}$ after taking infima. Fixed forms are available as branching trees, so the reverse comparison $V_{\mathrm{fixed}} \ge V_{\mathrm{branch}}$ holds. Moreover, Jensen gives $V_{\mathrm{branch}} \ge \|d\|^2_\infty > 0$. For a possibly biased design its residual $d - s_g$ is unchanged and its second moment is multiplied by at most $K_{m,b}$. Applying the same comparison to (B.1) gives (C.1). No attainment assumption is needed: apply the construction to designs approaching the infimum.

C.1.3. Path Guessing as an Exact Simulation

The construction above compares scores. The same guess gives a stronger statement, a comparison of experiments, which needs no score, no estimator and no asymptotics.

Theorem C.1 (Exact simulation)**.** *Let $1 \le r \le b \le m$ and $K_r = \min\{2^{b-r}, \binom{m-r}{b-r}\}$, so that $K_1 = K_{m,b}$. Let Π be any protocol that interviews n records with depth-b trees, choosing the tree of each record from the transcripts and labels of the earlier records and from the current label. There is a protocol Π' that uses only r-round interviews with at most b queries, chosen from the current label only if Π does so, interviews a random number N of records and outputs n transcripts with labels such that, for every law P of (X, Y):* (i) *the output of Π' has the same joint law as the n transcripts and labels of Π;* (ii) *N is a sum of n independent geometric variables with success probability $1/K_r$, independent of the output; in particular $\mathbb{E}N = K_r n$. With a fixed budget of n' records of r-round interviews, and a rule Π that is defined for every number of records, the same steps produce a number $B \sim \mathrm{Binomial}(n', 1/K_r)$ of transcripts such that, conditionally on $B = k$, they have the law of the first k transcripts of Π.*

Proof. Suppose that Π would interview its next record with the tree τ, padded as above. First let $r = 1$. Draw a fresh record, draw S uniformly from $\mathcal{S}_\tau$, independently of the features of the record given its label and the past, and ask the form S. The record is a *success* if $M_{\tau,S}(X_S) = 1$, that is, if the padded path of the record lies in S; the transcript $\ell_\tau(X)$ is then a function of X_S. For every profile x exactly one of the L_τ sets equals $S_\tau(x)$, so the conditional probability of success given $(X, Y) = (x, y)$ and the past is $1/L_\tau$ for the tree τ in use, which may depend on the past and, if Π allows it, on y. Accept a success with probability L_τ/K_1, by an independent coin, and otherwise discard the record. Acceptance then has probability exactly $1/K_1$ given (X, Y) and the past, so it is independent of (X, Y), and conditionally on acceptance (X, Y) has law P. Repeat with fresh records until one is accepted and output its transcript $\ell_\tau(X)$ and label. The output has the law of $(\ell_\tau(X), Y)$ under P given the past, which is the law of the next transcript of Π, and the number of records used is geometric with success probability $1/K_1$ and independent of the output. Induction over the n records of Π gives (i) and (ii). For $r \ge 2$ use the interviews of the proof of Theorem D.1(a): follow τ for $r - 1$ rounds to a node v, use for the last batch one of the $L_v \le K_r$ path sets of the subtree below v, drawn uniformly in advance for every node of depth $r - 1$, and call the record a success if the path completes in it; given (X, Y) this has probability $1/L_{v(X)}$, and accepting a success with probability $L_{v(X)}/K_r$ makes acceptance independent of (X, Y) with probability $1/K_r$. If Π chooses τ from the current label, Π' chooses its form or its last batch from the label as well, and not otherwise. For the last statement apply the same steps to n' records in turn; the acceptance indicators are independent Bernoulli variables with mean $1/K_r$ and independent of the transcripts produced. □

Consequences. Every estimator, test or decision rule that can be computed from n branching records can therefore be computed, with the same law under every P, from a random number of pre-committed records with mean $K_{m,b}n$, and from r-round records with mean $K_r n$. The inequality $V_r \le K_r V_{\mathrm{branch}}$ of Theorems 1(ii) and 2(a) follows to first order: run the iid design that attains V_{branch}/k for every sample size k on the B simulated records and return zero if $B = 0$; since $n'\mathbb{E}[B^{-1}; B \ge 1] \to K_r$ and $P(B = 0)$ is exponentially small, the risk is $(K_r + o(1))V_{\mathrm{branch}}/n'$. The same argument gives $W_r(P) \le K_r W_{\mathrm{branch}}(P)$ at full-support laws, where W is a local minimax risk (Proposition C.10); at every law the inequality is proved by the centered scores of Appendix E.1, and within a model by the information inequality of Theorem E.5(a), which is the same guess read as a statement about Fisher information. The score constructions of this appendix and the centered scores of Appendix E.1 remain of use because they give explicit pointwise-unbiased scores with a fixed sample size, which is what the designs of Section 6 implement; a pooled mean of t over the accepted records

is not unbiased with a fixed number of records because B may be zero. The simulation is the noiseless counterpart of the conversion of fully interactive into sequentially interactive locally private protocols, whose blow-up is linear in the compositionality of the protocol (Joseph et al., 2019); here the blow-up is the number of path sets, the simulation is exact, and for $r = 1$ and $m \geq b - 1 + 2^{b-1}$ Theorem 1(iii) shows that the factor cannot be improved even for estimating one linear functional. For a depth-b tree the classical non-adaptive baseline samples monomials of the Fourier expansion and has second moment $\|\hat{d}\|_1^2$, which equals 4^{b-1} for the address contrast; path guessing replaces it by 2^{b-1}.

C.1.4. The Address Function Attains the Bound

Let $\kappa = b - 1$, $D = 2^\kappa$ and $m = \kappa + D$. Identify the κ address bits with an index $A \in \{1, \dots, D\}$ in a fixed binary order, and let the remaining bits be $Z_1, \dots, Z_D$. Set $d = Z_A$. This Boolean contrast is realized by the two rules $p_1 = (1 + d)/2$, $p_0 = (1 - d)/2$. Its polynomial expression is

$$d(A, Z) = \sum_{j=1}^{D} Z_j \mathbf{1}\{A = j\}.$$

Each address indicator has degree κ, so $\deg(d) \leq \kappa + 1 = b$ and both menus identify the contrast.

Under the uniform profile law U, fix a mask with s address bits and a set R of r observed data indices. For a realized partial address α, let $C(\alpha)$ be its $2^{\kappa-s}$ consistent indices. Conditional independence and the centered unobserved data bits give

$$\mathbb{E}_{\mathsf{U}}[d \mid X_T] = 2^{-(\kappa-s)} \sum_{j \in R \cap C(\alpha)} Z_j.$$

The cross terms vanish on averaging over the observed independent signs. Each $j \in R$ is consistent with the random partial address with probability 2^{-s}. Consequently

$$\|\mathbb{E}_{\mathsf{U}}[d \mid X_T]\|_2^2 = 2^{-2(\kappa-s)} r 2^{-s} = r 2^{s-2\kappa}.$$

The value is zero if $r = 0$. Otherwise $s \leq \kappa + 1 - r$ and $r 2^{1-r} \leq 1$ for every integer $r \geq 1$, so it is at most $2^{-\kappa} = 1/D$. Querying all κ address bits and one data bit attains equality. Thus $J_{\mathsf{U}}(d) = 1/D$ for fixed forms, whereas $\langle d, d \rangle_{\mathsf{U}} = 1$. The dual formula (B.3) gives $V_{\mathrm{fixed}} \geq D$. This is a minimax lower bound over all protocols in the model, not only over unbiased score estimators, by Theorem B.2.

For the upper bound choose J uniformly from $\{1, \dots, D\}$ before the interview, and observe all address bits and Z_J. The feature score $H = D Z_J \mathbf{1}\{J = A\}$ satisfies $\mathbb{E}_J[H \mid X] = d(X)$ and $\mathbb{E}_J[H^2 \mid X] = D$ at every profile. Hence $V_{\mathrm{fixed}} = D$. Branching observes the address and then its data bit, giving feature score d with second moment one. Jensen gives the matching lower bound, so $V_{\mathrm{branch}} = 1$. In fact branching has the full-information risk $1/(1 + \sqrt{n})^2$ for every n by the binary-experiment calculation in Appendix C.10. The construction also covers $b = 1$, with no address bits and one data bit. It proves the asserted equality for every b.

Finally, for either menu write $C = D$ or $C = 1$ as appropriate. The uniform witness $h = d$ in (B.2) has denominator $1 + n/C$ and numerator one. Shrinking the preceding unbiased score by $\gamma = n/(n + C)$ gives residual $(1 - \gamma)d$ and constant second moment $\gamma^2 C$. The lower and upper bounds on the program therefore coincide:

$$\Phi_n = \frac{C}{n + C}.$$

The finite-sample minimax risk is bounded using Theorem B.1; identifying its exact ratio for fixed and branching forms is not part of Theorem 1.

C.2. Forms With Extra Questions

The gap of Theorem 1 compares forms and trees under the same cap. A validation study may be allowed a few more questions per record than the rule asks, and it is natural to ask how fast the price of pre-commitment falls when forms of size $b + c$ compete with trees of depth b. Write $V_{\mathrm{fixed}}(d, B)$ and $W^B_{\mathrm{fixed}}(P)$ for the constants of forms of size B, and keep $V_{\mathrm{branch}}(d, b)$, $W_{\mathrm{branch}}(P)$ for trees of depth b. For $c \geq 0$ let

$$j(c) = \max\{j \in \{0, \dots, b-1\} : \; 2^{j+1} - j - 2 \leq c\}, \qquad K^+_{b,c} = 2^{\,b-1-j(c)}.$$

Thus $j(0) = 0$, $j(1) = 1$, $j(4) = 2$, $j(11) = 3$ (for $b \geq 4$), and $2^{j(c)} > (c+3)/4$ as long as $j(c) < b-1$, so that $K^+_{b,c} < 2^{b+1}/(c+3)$.

Theorem C.2 (Extra questions)**.** *Let $1 \leq b \leq b+c \leq m$.*

(a) Simulation. *Every protocol on n records with interviews of depth at most b, with any estimator, is reproduced with the same law at every population by a protocol on forms of size $b+c$ that uses a random number of records with mean $K^+_{b,c}\, n$. Consequently $V_{\mathrm{fixed}}(d, b+c) \leq K^+_{b,c} V_{\mathrm{branch}}(d, b)$ for every contrast of degree at most b, $W^{b+c}_{\mathrm{fixed}}(P) \leq K^+_{b,c} W_{\mathrm{branch}}(P)$ at every law P, and the same holds for the constants within every model.*

(b) The address contrast. *For the address contrast with $\kappa = b-1$ address bits and $D = 2^\kappa$ data bits, and $0 \leq c \leq D-1$,*

$$\frac{V_{\mathrm{fixed}}(d, b+c)}{V_{\mathrm{branch}}(d, b)} = \frac{2^{b-1}}{c+1},$$

and $W^{b+c}_{\mathrm{fixed}}(P)/W_{\mathrm{branch}}(P) = 2^{b-1}/(c+1)$ at uniform profiles whenever the label agrees with the rule with a probability in $(0,1)$ that does not depend on the profile. Hence the largest price of pre-commitment with c extra questions lies between $2^{b-1}/(c+1)$ and $K^+_{b,c}$, two bounds that agree for $c \in \{0,1\}$ and differ by a factor below four for every such c.

(c) Polynomial dimension. *Let d be the product of k address blocks with κ address bits each (Appendix C.6.1), so that $b = k(\kappa+1)$, $m = k(\kappa + 2^\kappa)$ and $V_{\mathrm{branch}}(d, b) = 1$. Then for $0 \leq c \leq k(2^\kappa - 1)$,*

$$V_{\mathrm{fixed}}(d, b+c) \;\geq\; \frac{2^{\,b-k}}{(1+c/k)^k},$$

with equality when k divides c. If $2^\kappa \leq b$, so that $m \leq b^2$, and $c \leq \alpha b$, the ratio is at least $2^{\,b(1-\eta)}$ with $\eta = (1 + \log_2(1 + \alpha(\kappa+1)))/(\kappa+1)$, which tends to zero as κ grows; caps b that are not multiples of $\kappa+1$ are handled by adding blocks with $\kappa = 0$, as in Appendix C.6.1.

(d) Mean budgets. *Let $b \geq 2$ and consider fixed-form protocols whose forms have at most $b+c$ features and whose expected size, given the past, is at most b (the hybrid class of Appendix C.7.1 with $B = b+c$). For every contrast of degree at most b their constants are at most $\frac{b+c}{b} K^+_{b,c}$ times those of trees of depth b, and for the address contrast with $0 \leq c \leq D-1$ the worst-case constant of this class is exactly $\frac{b+c}{b} \cdot \frac{2^{b-1}}{c+1} \geq \frac{2^{b-1}}{b}$ times $V_{\mathrm{branch}}(d, b)$.*

Extra questions therefore buy a factor that is linear in their number and no more: one extra question halves the worst-case price, a constant number divides it by a constant, and in polynomial dimension even forms twice as long as the rule's paths leave a gap of $2^{b(1-o(1))}$. The price disappears only when the forms can hold a constant fraction of all the questions the tree might ask, which is the regime of the mean-budget remark of Appendix C.7.1. On trained rules the same quantity behaves very differently, because such rules depend on few items beyond their depth (Appendix F.3).

Proof. (a) Let τ be a padded tree of depth b and $j = j(c)$. Index the nodes of τ by the answers that lead to them. For a sequence g of $b-1-j$ answers, let F_g consist of the $b-j$ coordinates asked along the path that follows g from the root, together with all coordinates asked in the subtree below the last of these nodes; that subtree has $2+4+\cdots+2^j = 2^{j+1}-2$ nodes, so $|F_g| \leq b - j + 2^{j+1} - 2 \leq b+c$, and F_g is padded to size $b+c$ by a fixed rule. To simulate one record of τ, draw g uniformly and ask the form F_g. The answers to the first $b-1-j$ coordinates of the path show whether the record follows g; for every profile exactly one g is followed, so this happens with probability $2^{-(b-1-j)}$ whatever the record is, and then the whole path of the record lies in F_g and the transcript $\ell_\tau(X)$ is a function of the answers. The rest is the proof of Theorem C.1: an accepted record is a draw from the population with its transcript under τ, the number of records until acceptance is geometric with mean 2^{b-1-j} and independent of the transcript, and the tree may depend on earlier transcripts and on the label. The inequality for V follows from the simulation with a fixed budget, and those for W and for the model constants from the centred score $c_0 + K^+_{b,c}(H - c_0)\mathbf{1}\{\text{the record follows } g\}$ with $c_0 = \Delta(P)$ and from the information inequality $\sum_g J_{F_g}(h) \geq J_\tau(h)$, exactly as in Theorems E.1(a) and E.5(a).

(b) Under the uniform law a form with $s \leq \kappa$ address bits and w data bits explains the energy $w\, 2^{s-2\kappa}$ (Appendix C.1), and $s + w \leq b + c$. With $\delta = \kappa - s \geq 0$ we have $w \leq c + 1 + \delta \leq (c+1)2^\delta$, so $w\, 2^{s-2\kappa} \leq (c+1)2^{-\kappa}$, with equality at $s = \kappa$, $w = c+1 \leq D$; the largest energy is $(c+1)/D$. The dual of Theorem 3 with the uniform law and the witness $h = d$ gives $V_{\mathrm{fixed}}(d, b+c) \geq D/(c+1)$, and Proposition E.3 gives the same bound for $W^{b+c}_{\mathrm{fixed}}(P)/\sigma^2_P$ under label noise.

For the upper bounds ask all address bits and a uniformly drawn set of $c+1$ data bits: the addressed bit is in the set with probability $(c+1)/D$ at every profile, the score $DYd/(c+1)$ on that event is pointwise unbiased with second moment $D/(c+1)$, and its centred version has variance $D\sigma_P^2/(c+1)$. The comparison of the two bounds uses $2^{j(c)} > (c+3)/4$.

(c) A form that observes s_j address bits and r_j data bits of block j explains the energy $\prod_j r_j 2^{s_j-2\kappa}$ (Proposition C.7), with $\sum_j (s_j + r_j) \le b + c$. Put $u_j = s_j + r_j - \kappa$ and $\delta_j = \kappa - s_j \ge 0$, so that $r_j 2^{s_j} = 2^{\kappa}(u_j + \delta_j)2^{-\delta_j}$. If $u_j \ge 1$ then $u_j + \delta_j \le u_j 2^{\delta_j}$, and the factor is at most $2^{\kappa}u_j$; if $u_j \le 0$ then $r_j = u_j + \delta_j \ge 1$ and the factor is $2^{\kappa} r_j 2^{u_j - r_j} \le 2^{\kappa + u_j - 1}$. The function $\phi(u) = u$ for $u \ge 1$ and $\phi(u) = 2^{u-1}$ for $u \le 0$ has nonincreasing ratios $\phi(u+1)/\phi(u)$ on the integers, so under $\sum_j u_j \le k + c$ the product $\prod_j \phi(u_j)$ is largest for a balanced allocation with all $u_j \ge 1$, where it is at most $(1 + c/k)^k$ by the inequality of arithmetic and geometric means. Hence every form explains at most $(1 + c/k)^k 2^{-k\kappa}$, and the dual gives the bound. If k divides c, asking all address bits and $1 + c/k$ uniformly drawn data bits in every block attains it. For the last statement, $b - k - k\log_2(1 + c/k) \ge b - k(1 + \log_2(1 + \alpha(\kappa+1)))$ and $k = b/(\kappa+1)$.

(d) Ask a form F_g of part (a) with probability $b/(b+c)$ and nothing otherwise; the expected number of questions is at most b, a record is usable with probability $\frac{b}{b+c}2^{-(b-1-j)}$ whatever its profile, and the arguments of (a) apply with this probability. For the address contrast the value is Proposition C.9 with $B = b + c$, where $B - \kappa = c + 1$; it is at least $2^{b-1}/b$ because $(b+c)/(c+1) \ge 1$. □

C.3. Families of Targets

Theorem 1 is about one target fixed before collection. Two remarks and one proposition describe what happens for a family of targets served by one set of records, with the target chosen after collection.

The simulation of Theorem C.1 involves no target, so its factor bounds the price of pre-commitment for every family of targets and every criterion. The lower bound extends, for the worst member, to every family that contains the address contrast and whose members are functions, bounded by one, of the transcript of the address tree, for example all relabelings of its leaves by signs: a design that asks this one tree on every record observes every member exactly, so every member has branching constant at most one, while the address member needs 2^{b-1} under forms. At the other extreme the price disappears.

Proposition C.3 (All monomials of degree b at once)**.** *Let $1 \le b \le m$ and consider the family of monomials χ_A, $|A| = b$. Let a design draw a padded interview ι of depth b with probability q_ι, without knowledge of the target (padding is without loss of generality, since a cell with fewer than b questions carries no information about a monomial of degree b), and let it have, for every A, tables g_ι^A with $\sum_\iota g_\iota^A(\ell_\iota(x)) = \chi_A(x)$ for all x. Then*

$$\max_{|A|=b} \max_x \sum_\iota \frac{g_\iota^A(\ell_\iota(x))^2}{q_\iota} \;\ge\; \binom{m}{b},$$

with equality for the uniform distribution over the forms of size b and the tables $g_S^A = \chi_A \mathbf{1}\{S = A\}$, that is, the scores $\binom{m}{b} Y \chi_A \mathbf{1}\{S = A\}$. Thus fixed forms, r-round and branching interviews have the same simultaneous worst-case constant $\binom{m}{b}$ for this family among designs with pointwise-unbiased scores, and uniform matrix sampling attains it. Under uniform forms the same constant $\binom{m}{b}\|d\|_2^2$ holds at every profile for every homogeneous target $d = \sum_{|S|=b} \hat{d}(S)\chi_S$, with the tables $g_S = \hat{d}(S)\chi_S$.

Proof. Fix A and write $g_\iota = g_\iota^A$. A transcript of ι is a cell $c = \{x_{T_c} = u_c\}$ with $|T_c| = b$, and under the uniform law $\mathbb{E}[\mathbf{1}\{X \in c\}\chi_A(X)]$ is $2^{-b}\chi_A(u_c)$ if $T_c = A$ and zero otherwise, because a coordinate of A outside T_c is free in c. Taking the Fourier coefficient at A of the unbiasedness identity therefore gives

$$1 = \sum_\iota \sum_{c:\, T_c = A} g_\iota(c)\chi_A(u_c)2^{-b} \le \Big(\sum_\iota \sum_{c:\, T_c = A} \frac{g_\iota(c)^2}{q_\iota} 2^{-b}\Big)^{1/2} \Big(\sum_\iota q_\iota\, p_\iota(A)\Big)^{1/2},$$

by the Cauchy–Schwarz inequality, where $p_\iota(A) = 2^{-b}|\{c : T_c = A\}|$ is the probability, under the uniform law, that the question set of ι equals A. The first factor is at most the square root of the average over x of the second moment $\sum_\iota g_\iota(\ell_\iota(x))^2/q_\iota$, hence of its maximum. So the maximum is at least $1/\pi(A)$ with $\pi(A) = \sum_\iota q_\iota p_\iota(A)$. Every record has exactly one question set, so $\sum_{|A|=b} \pi(A) = 1$ and some A has $\pi(A) \le 1/\binom{m}{b}$. For uniform forms $\pi(A) = 1/\binom{m}{b}$ and the

stated tables have second moment $\binom{m}{b}$ at every profile. The bound is the dual bound of Theorem 3 for the witness $h = \chi_A$ under the uniform law with the design held fixed, so it is not specific to the tables, and it applies at each value of the label when the design may depend on the label. □

The price of pre-commitment is thus a property of targeted collection. A branching interview spends its questions where one target needs them, and every profile still yields a single question set; when all question sets matter equally, nothing is gained by choosing them from the answers. Between one tree and all monomials, for instance for the trees of a forest or for candidate rules on the same items, we have only the computations of Appendix F.6.

C.4. Additional Sharpness Results

Proposition C.4 (Finite-sample address gap)**.** *For the address contrast with $D = 2^{b-1}$, every $n \geq 1$ satisfies*

$$D\left(\frac{\sqrt{n}+1}{\sqrt{n}+\pi\sqrt{D}}\right)^2 \leq \frac{R_{\text{fixed}}(b,n)}{R_{\text{branch}}(b,n)} \leq D.$$

In particular, $n \geq 60D$ suffices for a ratio at least $D/2$.

Appendix C.5.1 proves the result by a finite-width van Trees bound and a constant-risk shrinkage estimator.

Corollary C.5 (Finite answer alphabets)**.** *For features with at most $s \geq 2$ answers and nonzero identifiable d, the coefficient ratio is at most $\min\{s^{b-1}, \binom{m-1}{b-1}\}$. Its supremum over dimensions is s^{b-1}, attained at $m = b - 1 + s^{b-1}$.*

The proof uses an s-ary address and the same simulation (Appendix C.5.2); ordinal features therefore need not be dichotomized for the theory.

Consider the four-bit Boolean contrast

$$f(x) = \tfrac{1}{2}(x_1x_2 + x_1x_3 + x_4x_2 - x_4x_3). \tag{C.3}$$

If $x_2 = x_3$ then $f = x_1x_2$, and otherwise $f = x_4x_2$. This is the four-bit sortedness function up to sign and permutation (O'Donnell, 2014, Chapter 1). Indeed, in the order (x_3, x_1, x_2, x_4) it is positive exactly when the sequence changes sign at most once. On k disjoint blocks let $d_k = \prod_{r=1}^{k} f(x^{(r)})$. The difference is realized by $p_1 = (1+d_k)/2$ and $p_0 = (1-d_k)/2$. Mixing a common rule into both predictors scales the contrast and its risks. The 4^k Fourier terms of d_k all have degree $2k$ and coefficient magnitude 2^{-k}.

Theorem C.6 (Complete query-budget curve)**.** *For d_k, both risks equal one for $b < 2k$. For $2k \leq b \leq 4k$ define*

$$C_{\text{fixed}} = 2^{4k-b}, \qquad C_{\text{branch}} = 4^{(3k-b)_+}.$$

In either class the minimax risk is $\Theta(\min\{1, C/n\})$, with absolute constants, and the exact first-order coefficient V is C. At $b \geq 3k$, branching attains the full-profile risk $1/(1+\sqrt{n})^2$ at every n.

At $b = 2k$ both interview types identify the gain and both pay a 4^k sample factor for fixed small error. At $b = 3k$ branching needs only the full-data number of people, while fixed forms still pay 2^k. These comparisons allow a different randomly chosen fixed form for every respondent. The penalty comes from committing to this person's questions before seeing their answers.

Figure 3 in Appendix C.10.1 plots this exact synthetic budget curve.

Where the exponents come from. In each block, two, three and four fixed coordinates capture at most $1/4$, $1/2$ and all of its uniform Fourier mass. The masses multiply across blocks. Randomized signed-term scores attain the reciprocal second moments. A branching path with nonzero conditional mean must query at least two coordinates per block; at budget $2k + r$, at least $k - r$ blocks receive only two and each contributes at most $1/2$ in absolute conditional mean. These bounds yield the matching witnesses and scores (Appendix C.10.1).

Two rounds suffice for these extremal examples. The address construction asks all address bits in one batch and the selected data bit in a second. The block family also attains its entire branching curve in two rounds. Consequently the sharp supremum of V_{fixed}/V_R is already 2^{b-1} for every allowed round count $R \geq 2$ (Appendix D.2). This concerns fixed versus two-round interviews; a general separation between two rounds and unrestricted branching is a different question.

An instance-specific comparison. If a depth-b tree τ computes d, then $V_{\text{branch}} = M^2$ for $M = \|d\|_\infty$. Its number L_τ of padded path sets improves the universal upper bound. Write $E_b = \max_{|T|\le b} \sum_{S\subseteq T} \widehat{d}(S)^2$ using uniform Fourier coefficients. Then

$$\max\left\{1, \frac{\|d\|_{2,\mathrm{U}}^4}{M^2 E_b}\right\} \le \frac{V_{\text{fixed}}}{M^2} \le \min\{L_\tau, K_{m,b}\}.$$

Appendix C.8 proves the interval.

C.5. Finite-Sample and Nonbinary Extensions

C.5.1. When the Address Gap Becomes Visible

Consider the address contrast with D data coordinates, either the binary construction of Theorem 1 or the s-ary construction below. The data coordinates take values in $\{-1, 1\}$. The fixed score $H = DYZ_J\mathbf{1}\{J = A\}$ has $\mathbb{E}H = \Delta$ and $\mathbb{E}H^2 = D$ under every population. With $\gamma = \sqrt{n}/(1+\sqrt{n})$, its shrunken sample mean has risk

$$\frac{\gamma^2 D}{n} + \left((1-\gamma)^2 - \frac{\gamma^2}{n}\right)\Delta^2 = \frac{D}{(1+\sqrt{n})^2}.$$

The zero estimator also has risk at most one. Branching recovers the sign Yd and has exact minimax risk $(1+\sqrt{n})^{-2}$ by Appendix C.10.

For a converse take independent uniform address coordinates and independent fair data bits, with profile law μ. On the subfamily $P_\theta(x, y) = \mu(x)(1 + \theta y d(x))/2$, the gain is θ. For a fixed mask T, put $u_T = \mathbb{E}_\mu[d \mid X_T]$. A transcript (x_T, y) has density proportional to $1 + \theta y u_T(x_T)$, so its Fisher information is

$$I_T(\theta) = \mathbb{E}_\mu\left[\frac{u_T^2}{1-\theta^2 u_T^2}\right] \le \frac{1}{D(1-\eta^2)}, \qquad |\theta| \le \eta < 1.$$

Here $|u_T| \le 1$ and the address projection calculation gives $\mathbb{E}u_T^2 \le 1/D$. The bound is uniform in the chosen mask. Conditional scores are martingale differences, so choosing masks from earlier respondents cannot increase total information beyond $n/[D(1-\eta^2)]$. The same bound holds if the current label selects the mask: for either selected mask the law of u_T is symmetric under flipping all data bits, and $\mathbb{E}[u_T^2/(1 \pm \theta u_T)] = \mathbb{E}[u_T^2/(1-\theta^2 u_T^2)]$.

The prior $\eta^{-1}\cos^2(\pi\theta/(2\eta))$ on $[-\eta, \eta]$ has Fisher information π^2/η^2. The van Trees argument in Appendix B.3 therefore gives

$$R_{\text{fixed}}(b, n) \ge \frac{1}{n/[D(1-\eta^2)] + \pi^2/\eta^2}.$$

Minimizing the denominator over $\eta^2 \in (0, 1)$ gives $(\sqrt{n/D} + \pi)^2$. Consequently

$$\frac{D}{(\sqrt{n} + \pi\sqrt{D})^2} \le R_{\text{fixed}}(b, n) \le \min\left\{1, \frac{D}{(1+\sqrt{n})^2}\right\}.$$

This proves Proposition C.4. The ratio is at least $D/2$ whenever $n \ge 60D$, since $\sqrt{60}/(\sqrt{60}+\pi) > 1/\sqrt{2}$. This threshold is sufficient, not necessary; the result does not identify the exact finite-sample fixed-form risk.

C.5.2. Features with More Than Two Answers

Let each of the m coordinates have at most $s \ge 2$ possible answers. Pad a depth-b tree to b distinct coordinates as in Appendix C.1. Its first $b-1$ answers determine the entire query set, giving at most s^{b-1} sets. Every set contains the root coordinate, so there are also at most $\binom{m-1}{b-1}$. The same simulation and the same conditional second-moment argument give Corollary C.5 and its finite-program counterpart.

For sharpness put $\kappa = b - 1$ and $D = s^\kappa$. Use κ independent uniform s-valued address coordinates and D independent fair data bits; features with fewer than s possible answers are permitted. The contrast is $d = Z_A$. A mask with u address coordinates and r data coordinates has

$$\big\|\mathbb{E}[d \mid X_T]\big\|_2^2 = r s^{u-2\kappa} \le s^{-\kappa}, \qquad u + r \le \kappa + 1,$$

because $rs^{1-r} \le 1$ for integers $r \ge 1$. Thus the variance dual gives $V_{\mathrm{fixed}} \ge D$. Querying the complete address and one independent uniform data coordinate yields a pointwise unbiased score with second moment D, while branching evaluates d with second moment one. The coordinate count is $\kappa + D$. If a common alphabet of exactly s symbols is required, map each data symbol to a sign and use a lower-bound population supported on two symbols of opposite signs. The bounds allow arbitrary population laws.

Both complete menus span sums of functions of at most b coordinates: each adaptive leaf indicator depends on at most b coordinates, and fixed forms span every such function. Thus the identification spaces still coincide. The binary Fourier-degree language is replaced by interaction order in a product basis.

C.5.3. Approaching an Identification Floor

For any d, define $\delta = \inf_{v \in \mathcal{H}_{\mathcal{A}}} \|d - v\|_\infty$. The indistinguishable pair in Appendix B.2 proves $R_{\mathcal{A}}(d,n) \ge \delta^2$ for every n. For each $v \in \mathcal{H}_{\mathcal{A}}$, use a design with conditional score mean $Yv(X)$ and maximal second moment approaching $V(v,\mathcal{A})$. Relative to the original target $Yd(X)$, its mean has absolute bias at most $\|d - v\|_\infty$. Hence

$$\delta^2 \le R_{\mathcal{A}}(d,n) \le \inf_{v \in \mathcal{H}_{\mathcal{A}}} \left\{ \|d - v\|_\infty^2 + \frac{V(v,\mathcal{A})}{n} \right\}. \tag{C.4}$$

A best uniform approximant exists in this finite-dimensional subspace and has finite V. For any such $v_\star$, the excess risk is at most $V(v_\star,\mathcal{A})/n$. On a common binary cap, the two menus have the same δ; applying Theorem 1 to each approximant compares their variance terms by $K_{m,b}$. This is an upper bound on approach to the floor, not a matching excess-risk rate.

For example, on twenty binary features let $d(x) = 0.8x_1 + 0.2\prod_{j=1}^{20} x_j$ and set $b = 5$. Its degree exceeds the cap, but $v = 0.8x_1$ has uniform approximation error 0.2. The uniform signed witness proportional to $\prod_{j=1}^{20} x_j$ annihilates every degree-five function, proving that this error is optimal. Both menus satisfy

$$0.04 \le R_{\mathcal{A}}(d,n) \le 0.04 + 0.64/n.$$

For a general dense logistic contrast, determining an accurate approximant can itself be difficult. More respondents alone cannot remove a positive identification floor. Additional structural assumptions or a larger per-person cap are required for consistency.

C.6. Products of Addresses and Equality Certificates

C.6.1. Nearly Maximal Gaps in Polynomial Dimension

For $\kappa \ge 0$, an address block consists of κ address bits and 2^κ data bits, and returns the addressed data bit. At $\kappa = 0$ it is just one data bit. Let f_j be address blocks on disjoint coordinates, with address lengths $\kappa_1, \dots, \kappa_k$, and put

$$d = \prod_{j=1}^{k} f_j, \qquad b = \sum_{j=1}^{k} (\kappa_j + 1), \qquad m = \sum_{j=1}^{k} (\kappa_j + 2^{\kappa_j}), \qquad D = 2^{\sum_j \kappa_j}.$$

Proposition C.7 (Product address coefficient)**.** *For this contrast,* $V_{\mathrm{branch}}(d,b) = 1$ *and* $V_{\mathrm{fixed}}(d,b) = D = 2^{b-k}$*. Two rounds attain the branching coefficient.*

Proof. Query all address bits first, and the k selected data bits second. This evaluates d within budget b, giving coefficient one by Jensen's inequality. For a fixed design, independently guess one data index in each block, query those k bits and all address bits, and return Dd if every guess matches, zero otherwise. Completion has probability $1/D$ at every profile, so the conditional mean is d and second moment is D.

For the converse let μ be uniform on the full cube and use $h = d$ in the variance dual. An arbitrary fixed form can allocate its queries among blocks in any way. If it observes s_j address bits and r_j data bits in block j, independence gives

$$\|\mathbb{E}_\mu[d \mid X_T]\|_2^2 = \prod_{j=1}^{k} r_j 2^{s_j - 2\kappa_j}.$$

If some $r_j = 0$, this is zero. Otherwise $r_j \le 2^{r_j-1}$ and $\sum_j (s_j + r_j) \le b$, whence

$$\prod_j r_j 2^{s_j - 2\kappa_j} \le 2^{\sum_j (s_j+r_j) - k - 2\sum_j \kappa_j} \le 2^{-\sum_j \kappa_j} = D^{-1}.$$

Since $\mathbb{E}_\mu[dh] = 1$, the variance dual gives $V_{\mathrm{fixed}} \ge D$. This treats all fixed forms, including those that redistribute queries across blocks. The general minimax theorem then covers arbitrary biased estimators and history-dependent form selection. □

For any fixed $\kappa \ge 1$, equal blocks give

$$m = \frac{\kappa + 2^\kappa}{\kappa + 1} b, \qquad \frac{V_{\mathrm{fixed}}}{V_{\mathrm{branch}}} = 2^{\kappa b/(\kappa+1)} \quad \text{when } \kappa + 1 \text{ divides } b.$$

In particular $m = 3b/2$ permits $2^{b/2}$. More generally, for any fixed $\zeta > 0$ one can choose a constant κ so that the exponent is at least $(1-\zeta)b$ while $m = O_\zeta(b)$; the hidden dimension multiplier grows with $1/\zeta$. This does not solve the sharp frontier for a prescribed ratio m/b, such as $4/3$.

For every $b \ge 2$, choose $\kappa = \lfloor \log_2 b \rfloor$, $k_0 = \lfloor b/(\kappa+1) \rfloor$, and $\rho_0 = b - k_0(\kappa+1)$. Use k_0 blocks of address length κ and ρ_0 blocks of address length zero. Then $m = k_0(\kappa + 2^\kappa) + \rho_0 \le b^2$, and

$$\log_2(V_{\mathrm{fixed}}/V_{\mathrm{branch}}) = k_0\kappa = b - k_0 - \rho_0 \ge b - \frac{b}{\kappa+1} - \kappa.$$

Thus polynomial dimension already permits $2^{b - O(b/\log b)}$. The construction uses the standard product method; the content is the exact coefficient and the lower bound uniform over all query allocations.

C.6.2. When the Full-Information Coefficient Is Attained

Write $M = \|d\|_\infty > 0$ and let $\mathcal{H}_\iota$ be the space of functions observable under interview ι. Define the bounded-decomposition quantity

$$\mathcal{C}_{\mathcal{A}}(d) = \inf\left\{ \sum_\iota \|g_\iota\|_\infty : g_\iota \in \mathcal{H}_\iota,\ \sum_\iota g_\iota = d \right\}.$$

It is computed by a linear program: minimize $\sum_\iota u_\iota$ subject to $\sum_\iota g_\iota = d$ and $-u_\iota \le g_\iota(c) \le u_\iota$ on every cell. Its size is that of the enumerated profiles and menu, not polynomial in a general compact representation.

Proposition C.8 (Equality and a bounded-score certificate)**.** *For any identifiable d, $M \le \mathcal{C}_{\mathcal{A}}(d)$ and*

$$M^2 \le V(d, \mathcal{A}) \le \mathcal{C}_{\mathcal{A}}(d)^2.$$

Consequently $\mathcal{C}_{\mathcal{A}}(d) = M$ suffices for $V(d, \mathcal{A}) = M^2$. Equivalently, it suffices that d/M be a convex combination of observable functions with values in $[-1, 1]$. If $|d(x)| = M$ at every profile, then $V(d, \mathcal{A}) = M^2$ if and only if one interview in $\mathcal{A}$ computes d exactly. In particular, a Boolean contrast on the full cube has $V_{\mathrm{fixed}}(d, b) = 1$ if and only if it depends on at most b coordinates.

Proof. The triangle inequality gives $\mathcal{C}_{\mathcal{A}}(d) \ge M$. For a decomposition of total norm C, choose $q_\iota = \|g_\iota\|_\infty / C$, dropping zero terms. Each returned score g_ι/q_ι is bounded by C, giving $V \le C^2$. Jensen gives $V \ge M^2$.

For the last assertion the finite program attains its optimum: a bounded objective bounds every g_ι, the simplex is compact, and the quadratic perspectives are closed at $q_\iota = 0$. At an optimum with value M^2, for each profile,

$$d(x)^2 = \left(\sum_\iota q_\iota H_\iota(x) \right)^2 \le \sum_\iota q_\iota H_\iota(x)^2 \le M^2.$$

If $|d(x)| = M$ everywhere, equality in Jensen forces $H_\iota(x) = d(x)$ for every positive-weight interview and every profile. Any such interview computes d. The converse follows by using that interview alone. A fixed coordinate set computes d precisely when it contains all of its relevant coordinates. □

For general real-valued d, equality in Jensen is forced only on the extremal set $E = \{x : |d(x)| = M\}$. Every used form must distinguish opposite-sign extremal profiles. That condition is necessary but does not by itself control second moments away from E. The bounded-decomposition test is a sufficient certificate; it is not asserted to be necessary for general d. Deterministic classifier differences take values in $\{-1, 0, 1\}$, so they are not covered by the constant-magnitude equivalence when zero contrasts occur.

Few extremal profiles alone do not imply a small gap. Take an even number $k \geq 2$ of equal address blocks of length $\kappa \geq 1$, let f be their product, and put $\ell(x) = m^{-1} \sum_{i=1}^{m} x_i$. For $0 < \epsilon < 1$ define $d_\epsilon = (1 - \epsilon)f + \epsilon\ell$. The only profile with $|d_\epsilon| = 1$ is the all-positive profile: $\ell = \pm 1$ only at the two constant profiles, and $f = 1$ at both when k is even. A mixture of the two-round exact score for f and a uniformly sampled one-coordinate score for ℓ has magnitude one and conditional mean d_ϵ, so $V_{\mathrm{branch}} = 1$. Under the uniform law, $\mathbb{E}[f\ell] = 0$ because any coordinate lies in only one block and another independent block has zero mean. The witness $h = f$ from Proposition C.7 therefore gives

$$V_{\mathrm{fixed}}(d_\epsilon, b) \geq (1 - \epsilon)^2 2^{k\kappa}.$$

An exponentially large gap is therefore compatible with a non-Boolean contrast having a unique absolute maximum. Empirical near-equality should be established by certificates rather than inferred from extremal-profile counts.

C.6.3. Optimizing a Root Augmentation

With $U = (X_{\mathrm{root}}, Y)$ and $w = q_S^{-1} - 1$, the variance identity in Appendix E.7 implies that, for fixed q, the population-optimal root function is

$$c^*(U) = \frac{\mathbb{E}[wt \mid U]}{\mathbb{E}[w \mid U]},$$

with any value when the denominator is zero. This follows by minimizing the conditional quadratic $\mathbb{E}[w(t - c)^2 \mid U]$. The weighted calibration counterpart uses the same form law as the original comparator and a single pseudo-observation at the calibration grand mean, weighted by the mean calibration weight. It remains globally unbiased. A reduction in its calibration criterion does not guarantee a reduction under the held-out law.

C.7. Expected Budgets and a Local Efficiency Comparison

C.7.1. The Expected-Budget and Hybrid Address Coefficients

A mean feature budget b bounds the conditional expected number of queries on each new respondent, for every past history and every input profile. A fixed form has no input-dependent length, so its distribution satisfies $\sum_T q_T |T| \leq b$. In the hybrid model it also obeys $|T| \leq B$ almost surely; branching procedures obey the corresponding expected-length and depth constraints. The label is observed in both models and is charged separately. These conventions impose a uniform design constraint, not a budget that holds only under an unknown favorable population. Write $R^{b,B}_{\mathrm{fixed}}(n)$ and $R^{b,B}_{\mathrm{branch}}(n)$ for the minimax risks with these constraints.

Proposition C.9 (Exact address interpolation)**.** *Let $d(A, Z) = Z_A$ have $\kappa \geq 1$ address bits and $D = 2^\kappa$ data bits, so $m = \kappa + D$. For $\kappa + 1 \leq b \leq B \leq m$, with integer B,*

$$\lim_{n\to\infty} nR^{b,B}_{\mathrm{fixed}}(n) = \frac{DB}{b(B - \kappa)}, \qquad \lim_{n\to\infty} nR^{b,B}_{\mathrm{branch}}(n) = 1.$$

In particular the mean-budget comparison ($B = m$) has exact ratio m/b. For an arbitrary nonzero contrast on m binary features and $0 < b \leq m$, the mean-budget fixed-to-branching first-order ratio is at most m/b. If $m \geq 3$ and $\lfloor \log_2 m \rfloor + 1 \leq b \leq m$, the supremum over contrasts is between $m/(2b)$ and m/b.

Proof. For the upper construction, with probability b/B ask all κ address bits and a uniformly sampled set of $B - \kappa$ data bits; otherwise ask no features. Return $DB/[b(B - \kappa)]$ times Yd when the selected set contains the addressed bit, and zero otherwise. The nonzero-score probability is $b(B - \kappa)/(BD)$ at every profile. Hence the score is pointwise unbiased, its second moment is $DB/[b(B - \kappa)]$, its maximum query count is B, and its expected count is b. Branching evaluates the address in $\kappa + 1$ queries. The usual full-information Bernoulli submodel gives its matching coefficient one.

For the fixed-form lower bound use uniform independent feature bits and the label submodel $P_\theta(x,y) = 2^{-m}(1+\theta y d(x))/2$. A form observing s address bits and r data bits has information at zero $J_T = r2^{s-2\kappa}$. Put $c = s + r$. We claim

$$J_T \le \frac{c(B-\kappa)}{BD} \quad (c \le B). \tag{C.5}$$

The empty form has zero information. If $1 \le c \le \kappa+1$, the inequality $r2^{1-r} \le 1$ gives $J_T \le 2^{c-1-2\kappa}$; the sequence $2^{c-1}/c$ is nondecreasing, so $J_T/c \le 1/[D(\kappa+1)] \le (B-\kappa)/(BD)$. If $c \ge \kappa+1$, write $j = \kappa - s \ge 0$ and $r_0 = c-\kappa \ge 1$. Then $(r_0+j)2^{-j} \le r_0$, giving $J_T \le (c-\kappa)/D$. The ratio $(c-\kappa)/c$ is increasing in c, proving (C.5) in this case as well.

Conditionally on any previous records, the randomized form therefore has information at zero at most $b(B-\kappa)/(BD)$. For $|\theta| \le \eta < 1$, its information is at most this quantity divided by $1-\eta$, since the conditional score is bounded by one. The sequential likelihood score is a martingale sum. Applying the scalar van Trees argument of Appendix B.3, then taking $n \to \infty$ and $\eta \downarrow 0$, proves the converse for arbitrary estimators. The same argument allows a label-dependent form distribution when its conditional expected cost is bounded for each label. Thus the bound does not rely on unbiasedness or an iid allocation of forms.

For general d, query the entire profile with probability b/m and return $(m/b)Yd$ on that event, zero otherwise. Its second moment is at most $(m/b)\|d\|_\infty^2$, whereas even full information has first-order minimax coefficient $\|d\|_\infty^2$. For the stated factor-two lower bound, choose the largest $\kappa \ge 1$ such that $m' = \kappa + 2^\kappa \le m$, and ignore the remaining coordinates. Then $m' > m/2$ and $\kappa + 1 \le b$. If $b \le m'$, the address ratio is m'/b; if $b > m'$, the universal lower bound one is already at least $m/(2b)$. This proves both assertions. Existence of the general first-order coefficients also follows by the same finite-menu program with its compact, linear expected-cost constraints. □

The interpolation does not claim a sharp hybrid supremum over all contrasts. It identifies an exact family and the order of the unrestricted mean-budget supremum in the indicated budget range. For example, $\kappa = 3$ gives $m = 11$: at mean budget $b = 4$, increasing the permitted maximum from $B = 4$ to 11 reduces the exact address coefficient from 8 to $11/4$. Paying for labels does not invalidate either experiment. If one label and each feature cost one and the hard total-answer cap is b_{total}, replace the feature cap by $b_{\text{total}} - 1$. With a label cost c_Y, a padded b-feature protocol costs $c_Y + b$ per person; that common cost is positive and need not be small.

C.7.2. Why Population-Specific Efficiency Also Constrains Biased Estimators

The result below is a finite-state application of local asymptotic efficiency and optimal design, rather than a new general efficiency principle (Robins et al., 1994; Tsiatis, 2006; Hahn et al., 2011; Barnes et al., 2020). It supplies the local interpretation missing from a comparison based only on globally unbiased estimators.

Write $Z = (X,Y)$, $t(Z) = Yd(X)$, and let P assign positive probability to every state in a specified finite space $\mathcal{Z}$. A fixed interview ι reveals $O_\iota = (\ell_\iota(X), Y)$. The menu is finite and the interview for the current respondent is chosen before their answers, although all previous respondents may be used. Put

$$W_{\mathcal{A}}(P) = \inf_{q,H:\,\sum_\iota q_\iota H_\iota(O_\iota(z))=t(z)\ \forall z} \operatorname{Var}_{P,q}(H_A(O_A)).$$

For $c > 0$ define the local risk, with the center P available to the procedure,

$$L_{n,c}(P) = \inf_{\pi,\widehat{\Delta}} \sup_{\substack{Q \text{ a probability law} \\ \|Q-P\|_2 \le c/\sqrt{n}}} n\mathbb{E}_Q[(\widehat{\Delta} - \mathbb{E}_Q t)^2].$$

Proposition C.10 (Local minimax counterpart)**.** *If t is identifiable from the menu, then*

$$\lim_{c\to\infty} \liminf_{n\to\infty} L_{n,c}(P) = \lim_{c\to\infty} \limsup_{n\to\infty} L_{n,c}(P) = W_{\mathcal{A}}(P).$$

The infimum permits biased and nonlinear estimators and history-dependent form choices. If one allowed branching interview computes t, its local constant is $\operatorname{Var}_P(t)$; the local fixed-to-branching ratio is therefore $W_{\text{fixed}}(P)/\operatorname{Var}_P(t)$ when the denominator is nonzero.

Proof. Choose an orthonormal matrix U spanning the zero-sum tangent space of the probability simplex and parameterize $P_\theta = P + U\theta$. The derivative of the target is the vector $v = U^\mathsf{T} t$. If A_ι is the incidence matrix of the observed cells, the information matrix of one observation is

$$I_\iota(P) = U^\mathsf{T} A_\iota^\mathsf{T} \operatorname{diag}(A_\iota P)^{-1} A_\iota U.$$

Every nonempty observed cell has positive mass. For a design q, elementary least squares for its score functions gives

$$\inf_{H:\,\sum_\iota q_\iota H_\iota = t} \operatorname{Var}_{P,q}(H) = v^\mathsf{T} \left(\sum_\iota q_\iota I_\iota(P) \right)^\dagger v,$$

with the value $+\infty$ when v is outside the matrix range. Indeed, a centered observation function ϕ must satisfy $\mathbb{E}[\phi\, s] = v$, where s is the observation score vector. Its minimum squared norm is the displayed quadratic form. Because P has full support, these moment constraints and $\mathbb{E}\phi = 0$ are equivalent to $\sum_\iota q_\iota \phi_\iota(O_\iota(z)) = t(z) - \mathbb{E}_P t$ for every z.

For the converse place a smooth product prior on a cube of radius $c/(\sqrt{\dim(\theta)}\sqrt{n})$ in tangent coordinates, vanishing at its boundary. Its information matrix is nC_0/c^2 for a fixed positive-definite C_0. The prior is inside the required Euclidean ball and, for sufficiently large n, inside the probability simplex. Uniformly on it, $I_\iota(P_\theta) = I_\iota(P) + O(c/\sqrt{n})$ in operator norm. Under any sequential protocol the likelihood scores form martingale differences. After averaging over both the prior and histories, their total information divided by n is $\sum_\iota \bar{q}_\iota I_\iota(P) + O(c/\sqrt{n})$ for some $\bar{q}$ in the simplex. Integration by parts and matrix Cauchy–Schwarz (the multivariate van Trees inequality) bound the normalized Bayes risk below by

$$v^\mathsf{T} \left(\sum_\iota \bar{q}_\iota I_\iota(P) + C_0/c^2 + \epsilon_n I \right)^{-1} v, \qquad \epsilon_n \to 0$$

for fixed c, with a nonnegative uniform error bound ϵ_n. This uses no unbiasedness or regularity of the estimator. Minimize over $\bar{q}$, take $n \to \infty$, then $c \to \infty$. Compactness of the design simplex and the variational identity $v^\mathsf{T} M^\dagger v = \sup_u (2v^\mathsf{T} u - u^\mathsf{T} M u)$, with the same infinite-value convention, show that this limit is $\min_q v^\mathsf{T} (\sum_\iota q_\iota I_\iota(P))^\dagger v = W_{\mathcal{A}}(P)$.

For the upper bound choose a minimizing fixed design and its globally unbiased score at the center P. Its iid mean has normalized risk $\operatorname{Var}_{Q,q}(H)$, which converges uniformly to $W_{\mathcal{A}}(P)$ on each shrinking ball. A procedure approaching the infimum gives the same argument if desired. Full observation has constant $\operatorname{Var}_P(t)$ and cannot be improved by coarsening; a branching interview computing t attains it. □

The center-dependent local design is an oracle, not a calibration algorithm. The full-support qualification is essential to this regular argument; this is why every population in Section 6 mixes the empirical law with a uniform mass, whose size is varied in Appendix F.1. The finite-sample globally unbiased bound applies at boundary empirical laws. Neither result says that a biased estimator cannot improve at a single distribution or at a small finite sample size.

C.8. Instance Certificates

C.8.1. Instance-Specific Coefficients

Suppose a depth-b tree τ computes d exactly, let $M = \|d\|_\infty > 0$, and let L_τ be its number of distinct padded path sets. Under the uniform law on the binary cube, put

$$E_b(d) = \max_{|T| \le b} \sum_{S \subseteq T} \widehat{d}(S)^2.$$

Then

$$\max\left\{1, \frac{\|d\|_{2,\mathsf{U}}^4}{M^2 E_b(d)}\right\} \le \frac{V_{\mathrm{fixed}}(d,b)}{V_{\mathrm{branch}}(d,b)} \le \min\{L_\tau, K_{m,b}\}, \qquad V_{\mathrm{branch}}(d,b) = M^2. \tag{C.6}$$

Indeed the tree score d has maximum second moment M^2, while Jensen gives that lower bound for every unbiased representation. The variance dual with $\mu = \mathsf{U}, h = d$ gives the first inequality; Parseval identifies its denominator as $E_b(d)$. Simulating τ gives the L_τ bound, and the universal theorem supplies $K_{m,b}$. This certificate is instance-specific but still

concerns worst-case populations. It does not bound the ratio $W(P)/\operatorname{Var}_P(Yd)$ at an individual population by the same instance coefficient ratio: the denominator of that latter ratio is a population variance, not M^2.

The path-set bound need not be attainable for a given tree. On four coordinates with cap three, take the chain querying X_4, stopping on its negative branch, then querying X_3, stopping on its negative branch, and otherwise querying X_2. Padding early leaves with the smallest unused coordinates yields the three sets $\{1,2,4\}$, $\{1,3,4\}$ and $\{2,3,4\}$, so $L_\tau = 3$. Nevertheless every assignment of values to the original leaves depends only on X_2, X_3, X_4. One fixed triple computes it, and the coefficient ratio is one for every nonzero assignment. Thus L_τ is a simulation upper bound and can depend on padding; it is not a sharp invariant of every underlying tree.

For four features, our coefficient solver minimizes v subject to $\sum_\iota M_\iota g_\iota = d$, $\sum_\iota q_\iota = 1$, and $\sum_\iota g_\iota(x)^2/q_\iota \le v$ on all sixteen profiles. Here M_ι lifts a triple-score table. An implementable score supplies an upper bound. A nonnegative probability vector μ and arbitrary vector h supply the lower bound

$$\frac{(h^T d)^2}{\max_\iota \sum_C (\sum_{x\in C} h_x)^2/\mu(C)},$$

where C ranges over form ι's cells. This follows directly from Cauchy–Schwarz applied to every unbiased decomposition. We normalize the nonnegative state dual, add 10^{-12} mass per state, and evaluate this expression; no solver-reported objective is treated as a certificate without reconstruction. The returned primal tables are projected onto the exact observation-span constraint. Both bounds and the numerical equality residual are recorded with the code.

C.9. An Exact Four-Feature Example Beyond the Block Gap

The ratio two in the four-bit block is not extremal at $(m, b) = (4, 3)$.

Proposition C.11. *There is a Boolean contrast on four features, computable by a depth-three tree, with*

$$V_{\mathrm{branch}}(d,3) = 1, \qquad V_{\mathrm{fixed}}(d,3) = \frac{7+2\sqrt{2}}{4} = 2.457106\ldots.$$

Proof. Write $\zeta = \sqrt{2}$ and $\alpha = (7+2\zeta)/4$. Define

$$f_-(x) = \frac{(1-x_2)x_1 - (1+x_2)x_3}{2}, \qquad f_+(x) = -\frac{(1-x_3)x_2 + (1+x_3)x_1}{2},$$

$$d(x) = \frac{(1-x_4)f_-(x) + (1+x_4)f_+(x)}{2}.$$

Query x_4 first. If its value is -1, query x_2 and then x_1 or x_3; if its value is $+1$, query x_3 and then x_2 or x_1. This computes $d \in \{-1, 1\}$ with three queries. Jensen's inequality and the exact tree score give $V_{\mathrm{branch}} = 1$.

For the fixed upper bound use only the forms 124, 134 and 234, with respective probabilities $(2-\zeta)/2$, $(2-\zeta)/2$ and $\zeta - 1$. In a form $T = (i, j, k)$ index its answers by $c = (1+x_i)/2 + 2(1+x_j)/2 + 4(1+x_k)/2$. The following table specifies twice the unbiased decomposition table g_T; the returned score is $Y g_T/q_T$.

c	$2g_{124}$	$2g_{134}$	$2g_{234}$
0	$\zeta - 3$	$4 - 3\zeta$	$2\zeta - 3$
1	$3 - \zeta$	$2 - \zeta$	$2\zeta - 1$
2	$\zeta - 1$	$\zeta - 2$	$3 - 2\zeta$
3	$1 - \zeta$	$3\zeta - 4$	$1 - 2\zeta$
4	$2 - \zeta$	$1 - \zeta$	$2\zeta - 1$
5	$4 - 3\zeta$	$\zeta - 1$	$1 - 2\zeta$
6	$3\zeta - 4$	$3 - \zeta$	$2\zeta - 3$
7	$\zeta - 2$	$\zeta - 3$	$3 - 2\zeta$

Number full profiles lexicographically by $i = 8(1+x_1)/2 + 4(1+x_2)/2 + 2(1+x_3)/2 + (1+x_4)/2$, and put $E = \{1, 2, 3, 6, 8, 12, 13, 15\}$. Substitution gives $\sum_T g_T = d$ on all sixteen profiles. The conditional second moment equals α on E, and $(23 - 10\zeta)/4$ outside E. The latter is smaller because their difference is $3\zeta - 4 > 0$. Thus $V_{\mathrm{fixed}} \le \alpha$.

For the matching lower bound, let μ be uniform on E. Set $\xi = 3 + 2\zeta$ and define a signed vector η, in the same full-profile order, by

$$32\eta = (-\xi, 3, -5, 5, \xi, -\xi, -3, \xi, 5, \xi, \xi, -\xi, 3, -3, -\xi, -5).$$

For a form T let $I_T = \sum_C \eta(C)^2/\mu(C)$ over its cells, with $0/0 = 0$. Every zero-mass cell in this example also has zero signed mass. Direct summation yields

$$\eta^T d = \alpha, \qquad I_{123} = \alpha - \frac{1}{2}, \qquad I_{124} = I_{134} = I_{234} = \alpha.$$

The Cauchy–Schwarz certificate above therefore gives $V_{\text{fixed}} \geq (\eta^T d)^2 / \max_T I_T = \alpha$. Forms with fewer coordinates are coarsenings of triples and cannot increase I_T. All displayed identities are exact in $\mathbb{Q}(\sqrt{2})$; the supplied symbolic script verifies them without a numerical optimizer. □

Consequently the worst ratio at $(4, 3)$ lies in $[(7 + 2\sqrt{2})/4, 3]$. This excludes two as its value, but neither proves that three is attainable nor determines the exact supremum over all identifiable real contrasts.

Corollary C.12. *For the same contrast and budget three, $V_2(d, 3) \geq 4/3 > V_{\text{branch}}(d, 3) = 1$.*

Proof. Use the uniform law and witness $h = d$. A deterministic two-round protocol can be represented by a first set S of size one or two, followed by a set of $3 - |S|$ remaining coordinates depending on the first answers. Padding covers protocols using fewer queries; a single fixed triple is included by splitting it into a first pair and a final fixed coordinate. For a given S, its largest possible projection energy is

$$J(S) = 2^{-|S|} \sum_{u \in \{-1,1\}^S} \max_{\substack{T \subseteq [4] \setminus S \\ |T| = 3 - |S|}} \mathbb{E}\big[\mathbb{E}[d \mid X_S = u, X_T]^2 \mid X_S = u\big] .$$

Substitution of the displayed contrast gives the complete table:

First set S	$J(S)$
$\{1\}, \{4\}, \{1, 2\}, \{1, 3\}, \{1, 4\}, \{2, 3\}$	$1/2$
$\{2\}, \{3\}, \{2, 4\}, \{3, 4\}$	$3/4$

Thus every allowed two-round observation has projection energy at most $3/4$. The variance dual with $\mathbb{E}[d^2] = 1$ gives $V_2 \geq 4/3$, including mixtures of these protocols. The symbolic script also verifies all ten entries using rational arithmetic. This is a strict constant separation at the smallest cap where one exists; Theorem D.3 gives the exponential separation for large caps. □

C.10. The Binomial Value and a Synthetic Budget Curve

The binomial minimax value. For any nonzero d set $M = \|d\|_\infty$. With full profiles a randomized sign with conditional mean $Yd(X)/M$ has population mean Δ/M. For n iid signs W_i with unknown mean μ_0, the estimate $\sqrt{n}\,\overline{W}/(1 + \sqrt{n})$ has variance $(1 - \mu_0^2)/(1 + \sqrt{n})^2$ and squared bias $\mu_0^2/(1 + \sqrt{n})^2$. For the lower bound fix a profile at which $|d| = M$ and vary the sign label. A $\text{Beta}(\sqrt{n}/2, \sqrt{n}/2)$ prior on the success probability of the sign makes this constant-risk rule its posterior mean on the mean-sign scale. The Bayes risk equals the displayed frequentist risk, proving minimaxity. The full-profile value is therefore $M^2/(1 + \sqrt{n})^2$. The one-feature construction in Proposition B.3 simulates these same signs, and the constant-profile lower bound applies. For $M = 0$ return zero.

C.10.1. Proof of Theorem C.6

If $b < 2k$, under uniform X the laws $Y = \pm d_k(X)$ give the same observations on every allowed leaf. The risk is at least one and zero attains it. For $b \geq 2k$ we show that $J_{\mathsf{U}}(d_k) = 1/C$ and construct a per-person score H with $\mathbb{E}[H \mid X, Y] = Y d_k(X)$ and $\mathbb{E}[H^2 \mid X, Y] = C$ for each design type.

Fixed forms. Under the uniform law, projecting a multilinear expansion onto a mask retains exactly its contained Fourier terms. In a block of size four a mask with $j = 2, 3, 4$ coordinates captures at most $1/4, 1/2, 1$ of the total squared coefficient mass, respectively. The first maximum is achieved by a supported pair and the other two by any mask of the indicated size. A block with fewer than two coordinates captures zero mass. Independence of blocks gives total mass at most 2^{b-4k}, and choosing a supported pair in every two-coordinate block attains it for any budget allocation assigning between two and four coordinates to every block. For two coordinates sample one of four supported pairs uniformly and return twice its signed product. For three, choose $\{1, 2, 3\}$ or $\{2, 3, 4\}$ with equal probability. Return $2f$ when the first form is chosen and $x_2 = x_3$, or when the second is chosen and $x_2 \neq x_3$; return zero otherwise. For four the score is f itself. The scores have conditional means f and constant second moments $4, 2, 1$. Multiply independently randomized block scores and Y to obtain the desired H.

Branching forms. When $b = 2k + r$, $0 \leq r \leq k$, evaluate r blocks via branching, using three queries each; use a random signed pair score of magnitude two in every other block. Then H has constant absolute value 2^{k-r} and square $C = 4^{k-r}$. When $b \geq 3k$, evaluate every block. For the lower witness, at any deterministic leaf under uniform X, an unqueried coordinate remains uniform independent of the others. Nonzero conditional mean for the product requires that every block has at least two exposed coordinates. At most r blocks have more than two, so $|\mathbb{E}_{\mathsf{U}}[d_k \mid \ell]| \leq 2^{-(k-r)}$. Averaging yields the bound on J. It is attained by evaluating r blocks exactly and querying one predetermined supported pair in each other block. The absolute conditional mean is then $2^{-(k-r)}$ at every leaf.

Exact program and leading coefficients. The witness $\mu = \mathsf{U}, h = d_k$ in (B.2) gives $\Phi_n \geq 1/(1 + n/C)$. Randomized query procedures can be represented as mixtures over deterministic interviews by drawing all their random coins in advance. Taking their unbiased transcript scores with shrinkage $\gamma = n/(n + C)$ gives $s_g = \gamma d_k$ and a second-moment contribution at most $\gamma^2 C$ at every profile. Thus $\Phi_n \leq (1 - \gamma)^2 + \gamma^2 C/n = C/(n + C)$. The same witness in (B.3) and the unshrunk scores give $V = C$. Theorem B.1 yields the nonasymptotic rates. At $b \geq 3k$, evaluating d_k yields a sign of mean Δ and the binomial minimax value above.

Exact binary comparison for branching. For $2k \leq b \leq 3k$ put $\lambda = 2^{b-3k}$. The branching score λH is a sign with mean $\lambda\Delta$ for every P. Conversely consider the subfamily of uniform X with conditional label law $(1 + \theta y d_k(x))/2$. Then $\Delta = \theta \in [-1, 1]$. At each leaf, conditionally on the history and independent random coins, $p_\theta(z) = p_0(z)(1 + \theta s_z)$ with $|s_z| \leq \lambda$ and $\sum_z p_0(z) s_z = 0$. A sign w of mean $\lambda\theta$ can generate this leaf via the probability kernel $K(z \mid w) = p_0(z)(1 + w s_z/\lambda)$. Simulating the leaves successively reproduces the complete protocol. This is an explicit experiment comparison in the sense of Blackwell (1953): on the indicated subfamily an arbitrary interview can be simulated using the scalar sign. Together with the construction for every P, it gives exact equality of minimax risks with the binary-mean experiment, even for biased estimators.

Why the population supremum matters. A uniform-law calculation can be uninformative even when a contrast is identifiable. For example, take an extra switch $z \in \{-1, 1\}$ and set $\widetilde{d} = 1 + \mathbf{1}\{z = 1\}(d_k - 1)$. This remains Boolean, has degree $2k + 1$, and has uniform mean $1/2$, so even an empty form captures at least $1/4$ of the squared mean signal in the uniform witness $h = \widetilde{d}$. At the identifiable width $2k + 1$, however, a population supported on $z = 1$ contains the original hard experiment at that width. An auditor given the switch for free still has branching risk of order $\min\{1, 4^{k-1}/n\}$ for $k \geq 1$. Optimizing over profile laws and perturbations in (B.2) accounts for this possibility.

D. Rounds of Adaptivity

D.1. Rounds of Adaptivity

Interviews with r rounds. An r-round interview chooses coordinate batches $B_1,\ B_2(x_{B_1}),\ \ldots,\ B_r(x_{B_1 \cup \cdots \cup B_{r-1}})$. Each batch is a function of the answers to earlier batches only, and $\sum_s |B_s| \leq b$ on every input. Let $\mathcal{A}_r(b)$ be the finite menu of deterministic r-round interviews and $V_r(d, b) = V(d, \mathcal{A}_r(b))$; randomized interviews are mixtures and are already represented by the design probabilities in (2). Fixed forms are $\mathcal{A}_1(b)$. Sequentializing each batch shows that every r-round interview is a depth-b tree, and a depth-b tree is a b-round interview with one query per round. Hence $\mathcal{A}_1(b) \subseteq \mathcal{A}_2(b) \subseteq \cdots \subseteq \mathcal{A}_b(b) = \mathcal{A}_{\mathrm{branch}}(b)$ and $V_1 \geq V_2 \geq \cdots \geq V_b = V_{\mathrm{branch}}$. All menus identify the same contrasts, $\deg(d) \leq b$, because the two extreme menus do (Appendix B.2). Theorems B.1 and B.2 apply to each finite menu $\mathcal{A}_r(b)$, so the lower bounds below hold against biased, sequential and label-dependent procedures.




Figure 3. Exact synthetic budget curve for the same contrast on 24 features ($k = 6$). The fixed coefficient is 64 and the branching coefficient is one at 18 queries.

Theorem D.1 (Rounds of adaptivity, full statement of Theorem 2)**.** (a) *For $1 \le r \le b \le m$ and every nonzero contrast of degree at most b on $\{-1, 1\}^m$,*

$$V_{\mathrm{branch}}(d, b) \le V_r(d, b) \le \min\left\{2^{b-r}, \binom{m-r}{b-r}\right\} V_{\mathrm{branch}}(d, b).$$

(b) *For integers $r, \kappa \ge 1$ put $D = 2^\kappa$ and $b = r\kappa + 1$. The pointer-chasing contrast $N_{r,\kappa}$ defined below satisfies*

$$V_{r+1}(N_{r,\kappa}, b) = V_{\mathrm{branch}}(N_{r,\kappa}, b) = 1, \qquad \frac{D}{b} \le V_r(N_{r,\kappa}, b) \le D.$$

Corollary D.2 (Two rounds can be exponentially worse than branching)**.** *For every $b \ge 3$,*

$$\sup_{m,d} \frac{V_2(d, b)}{V_{\mathrm{branch}}(d, b)} \ge \frac{2^{\lfloor (b-1)/2 \rfloor}}{b}, \qquad \sup_{m,d} \frac{V_r(d, b)}{V_{r+1}(d, b)} \ge \frac{2^{\lfloor (b-1)/r \rfloor}}{b} \quad (1 \le r < b).$$

Proof. Take $\kappa = \lfloor (b-1)/r \rfloor \ge 1$ and $N_{r,\kappa}$, which uses $r\kappa + 1 \le b$ queries along every path. Branching and $(r+1)$-round interviews still compute it exactly under the larger cap b, while the hit bound in the proof of part (b) uses only the total number b of queries. For the first display use $r = 2$ and $V_{\mathrm{branch}} \le V_3$. □

The constant separation $V_2 \ge 4/3$ of Corollary C.12 is therefore not the end of the hierarchy. Part (b) and the corollary place the exponent for $2 \le r < b$ between $\lfloor (b-1)/r \rfloor - \log_2 b$ and $b - r$; the binary pointer tree (Theorem D.3) raises the lower end to $b - 1 - r \log_2(1 + 2b/r)$, its products do so in polynomial dimension, and Appendix D.4 computes the constants of the tree at small caps.

Proof of part (a). The left inequality is menu inclusion. For the right inequality pad a depth-b tree τ as in Appendix C.1, so that every path has b distinct coordinates. In rounds $1, \dots, r-1$ ask one question per round and follow τ exactly. This reaches a node v of depth $r-1$, whose padded subtree has depth $b - r + 1$. Its path sets all contain the subtree root and are determined by the first $b - r$ answers below v; hence their number L_v is at most 2^{b-r}, and also at most $\binom{m-r}{b-r}$ because the remaining $b - r$ coordinates are chosen from the $m - r$ coordinates not on the followed prefix or at v. In round r choose one of these L_v sets uniformly, independently of the person, and attempt to complete the path. On completion return L_v times the tree score and otherwise zero. Conditionally on x the mean equals the tree score and the second moment is L_v times its square. The mixture and merging argument of Appendix C.1 applies verbatim to randomized branching designs, so every feasible branching design is simulated with second moment inflated by at most $\max_v L_v$.

The pointer-chasing contrast. Levels $\ell = 1, \dots, r$ consist of registers. Level ℓ has $D^{\ell-1}$ registers $R_\ell(i_1, \dots, i_{\ell-1})$, indexed by $(i_1, \dots, i_{\ell-1}) \in [D]^{\ell-1}$, and each register stores κ binary coordinates encoding an element of $[D]$. There are D^r data coordinates $Z_{i_1 \cdots i_r}$. The pointer path is $j_1 = R_1$ and $j_\ell = R_\ell(j_1, \dots, j_{\ell-1})$, and

$$N_{r,\kappa}(x) = Z_{j_1 \cdots j_r} = \sum_{i \in [D]^r} Z_i \prod_{\ell=1}^{r} \mathbf{1}\{R_\ell(i_{<\ell}) = i_\ell\}.$$

Each register indicator has degree κ, so $\deg N_{r,\kappa} = r\kappa + 1 = b$. Reading the κ coordinates of the level-ℓ register on the pointer path in round ℓ, and the selected data coordinate in round $r+1$, computes the contrast with b queries in $r+1$ rounds. With Jensen's inequality this gives $V_{r+1} = V_{\mathrm{branch}} = 1$. For the upper bound $V_r \le D$, read the pointer registers of levels $1, \dots, r-1$ in the first $r-1$ rounds. In round r, read the level-r register together with one data coordinate $Z_{j_1 \cdots j_{r-1} J}$, where J is uniform on $[D]$ and drawn in advance. The score $D Z_{j_1 \cdots j_{r-1} J} \mathbf{1}\{J = j_r\}$ is unbiased with second moment D at every profile.

The lower bound $V_r(N_{r,\kappa}, b) \ge D/b$. Let U be the uniform law and write $c^\star = (j_1, \dots, j_r)$ for the selected data cell. Since $\mathbb{E}_{\mathsf{U}}[N^2] = 1$, the dual formula (B.3) with witness $h = N$ gives $V_r \ge 1/\max_{\iota \in \mathcal{A}_r(b)} E_\iota$, where $E_\iota = \mathbb{E}_{\mathsf{U}}[\mathbb{E}_{\mathsf{U}}[N \mid \ell_\iota(X)]^2]$. Fix a deterministic r-round interview ι.

Step 1: a generous transcript. Let ι^+ ask the same questions as ι, but at the end of every round additionally reveal the full content of every register touched in that round. The transcript of ι is a function of that of ι^+, and conditional expectation onto a larger σ-field cannot decrease the second moment, so $E_\iota \le E_{\iota^+}$.

Step 2: only the selected data coordinate carries signal. Let Obs be the set of data coordinates queried by ι^+. For a data cell c, on the transcript event $\{c \notin \mathrm{Obs}\}$ the transcript and the pointer path are functions of the coordinates other than Z_c, because the interview never reads Z_c. Since Z_c is an independent uniform sign, $\mathbb{E}[Z_c \mathbf{1}\{c^\star = c\} \mid \ell_{\iota^+}]$ vanishes there. Hence

$$\mathbb{E}[N \mid \ell_{\iota^+}] = \sum_{c \in \mathrm{Obs}} Z_c \, \mathbb{P}(c^\star = c \mid \ell_{\iota^+}),$$

so $|\mathbb{E}[N \mid \ell_{\iota^+}]| \le \mathbb{P}(c^\star \in \mathrm{Obs} \mid \ell_{\iota^+}) \le 1$ and

$$E_{\iota^+} \le \mathbb{E}\big[\mathbb{P}(c^\star \in \mathrm{Obs} \mid \ell_{\iota^+})\big] = \mathbb{P}(c^\star \in \mathrm{Obs}).$$

Step 3: observing $c^\star$ in r rounds requires a lucky guess. Let $\mathcal{G}_t$ be the σ-field of the ι^+ transcript before round t, and let K_t be the largest k such that the pointer registers $R_1, R_2(j_1), \dots, R_k(j_{<k})$ are all revealed before round t; thus $K_1 = 0$ and K_t is nondecreasing. A query of round t is *deep* if it touches a register of level at least $K_t + 2$ or a data coordinate while $K_t \le r - 1$; it is a *hit* if, in addition, the touched register or data coordinate lies on the pointer path. Every deep location is indexed by a tuple whose (K_t+1)st entry is $\mathcal{G}_t$-measurable, and lies on the pointer path only if this entry equals j_{K_t+1}. The latter is the content of the register $R_{K_t+1}(j_{\le K_t})$, which is unrevealed before round t. For an adaptive sequence of queries to independent uniform coordinates, the unrevealed coordinates remain independent and uniform given the transcript. Hence j_{K_t+1} is uniform on $[D]$ given $\mathcal{G}_t$, and every deep query is a hit with conditional probability at most $1/D$.

If no hit occurs in rounds $1, \dots, t$, then $K_{t+1} \le K_t + 1$. Indeed, $K_{t+1} \ge K_t + 2$ requires the pointer register of level $K_t + 2$ to be revealed by the end of round t, say in round $t' \le t$. Since $K_{t'} \le K_t$, that query was deep in round t' and lay on the pointer path, so it was a hit. By induction, without hits $K_t \le t - 1 \le r - 1$ for every $t \le r$. Then every query of a data coordinate in rounds $1, \dots, r$ is deep, and $c^\star$ can be queried only by a hit. Since the total number of queries is at most b,

$$\mathbb{P}(c^\star \in \mathrm{Obs}) \le \mathbb{P}(\text{some hit}) \le \mathbb{E}\Big[\sum_{t=1}^{r} \sum_{\text{deep queries of round } t} \mathbb{P}(\text{hit} \mid \mathcal{G}_t)\Big] \le \frac{b}{D}.$$

Combining the three steps gives $E_\iota \le b/D$ for every $\iota \in \mathcal{A}_r(b)$, and therefore $V_r \ge D/b$. □

A binary pointer tree: few rounds lose almost the whole exponent. The pointer-chasing family separates consecutive numbers of rounds, but its lower bound $2^{\lfloor (b-1)/r \rfloor}/b$ leaves the exponent for a fixed small number of rounds far below the upper bound $b - r$ of part (a). The following contrast closes most of this gap. Let $L \ge 1$ and $\beta = L + 1$. Arrange $2^L - 1$ *pointer coordinates* as the internal nodes u of a complete binary tree with levels $0, \dots, L-1$ and attach 2^L *data coordinates*

Z_v to its leaves v, so that a block has $2^\beta - 1$ coordinates. Encode the pointer coordinate of node u as a bit $x_u \in \{0,1\}$ that selects one of the two children of u. The *pointer path* starts at the root and descends through the selected children; let $c^\star(x)$ be the leaf it reaches, and

$$B_L(x) = Z_{c^\star(x)} = \sum_v Z_v \prod_{u \in \mathrm{path}(v)} \mathbf{1}\{x_u \to v\},$$

where $\mathrm{path}(v)$ is the set of the L internal nodes above v and $\{x_u \to v\}$ is the event that the bit of u selects the child on the way to v. Each indicator has degree one, so $\deg B_L = L + 1 = \beta$, and following the path with one question per round computes B_L with β queries. The function is the tree evaluation problem of Cook et al. (2012), and round hierarchies for the query cost of oracle evaluation under bounded adaptivity are known (Canonne & Gur, 2018; Li et al., 2026). Here the total number of queries is capped, exact evaluation in fewer than β rounds is impossible for every strategy, and we quantify the variance that the missing rounds cost. To reach polynomial dimension we also take products: for $k \geq 1$ let

$$B_L^{\otimes k}(x^{(1)}, \ldots, x^{(k)}) = \prod_{i=1}^{k} B_L(x^{(i)})$$

on k disjoint blocks, a contrast of degree $k\beta$ on $k(2^\beta - 1)$ coordinates.

Theorem D.3 (Binary pointer trees and their products)**.** *Let $L \geq 1$, $\beta = L + 1$, $k \geq 1$ and $b \geq k\beta$, and let $d = B_L^{\otimes k}$ on $m = k(2^\beta - 1)$ coordinates.* (a) *$V_{\mathrm{branch}}(d, b) = 1$ and $V_r(d, b) \leq 2^{b-r}$ for $1 \leq r \leq b$.* (b) *For every $1 \leq r \leq b$,*

$$V_r(d, b) \;\geq\; \frac{2^{k(\beta-1)}}{\left(1 + \frac{2b}{rk}\right)^{rk}}.$$

In particular the single tree B_{b-1} on $m = 2^b - 1$ coordinates ($k = 1$, $\beta = b$) satisfies $V_r(B_{b-1}, b) \geq 2^{b-1}/(1 + 2b/r)^r$, and $V_1(B_{b-1}, b) \geq 2^{b-1}/b$.

Corollary D.4 (Few rounds lose almost the whole exponent)**.** (i) *For every $b \geq 2$ and $1 \leq r \leq b$,*

$$\frac{2^{b-1}}{(1 + 2b/r)^r} \;\leq\; \sup_{m,d} \frac{V_r(d, b)}{V_{\mathrm{branch}}(d, b)} \;\leq\; 2^{b-r}, \qquad \textit{in particular} \quad \frac{2^{b-1}}{(b+1)^2} \;\leq\; \sup_{m,d} \frac{V_2(d, b)}{V_{\mathrm{branch}}(d, b)} \;\leq\; 2^{b-2}.$$

Hence for every fixed r the exponent of the r-round gap is $b - O(r \log b)$. (ii) *An r-round protocol that is within a factor b^c of full branching on every contrast of degree at most b needs $r \log_2(1 + 2b/r) \geq b - 1 - c \log_2 b$, and therefore $r \geq 0.37\, b - (1 + c \log_2 b)$ rounds: a number of rounds linear in the cap. The witnessing contrast is the single tree B_{b-1} on $2^b - 1$ coordinates.* (iii) *Polynomial dimension. For every $b \geq 4$ and $r \geq 1$, with $\beta = \lfloor \log_2 b \rfloor$ and $k = \lfloor b/\beta \rfloor$, the contrast $B_{\beta-1}^{\otimes k}$ has degree at most b, lives on $m \leq b^2$ coordinates, and satisfies $V_{\mathrm{branch}} = 1$ and*

$$\log_2 V_r \;\geq\; \Big(\frac{b}{\beta} - 1\Big)(\beta - 1) - \frac{rb}{\beta} \log_2 \Big(1 + \frac{4\beta}{r}\Big) \;=\; b - O\Big(\frac{r\, b \log\log b}{\log b}\Big) \quad (b \to \infty,\ r \textit{ fixed}).$$

Proof. (i) The upper bound is Theorem D.1(a) and the lower bound is Theorem D.3(b) with $k = 1$. (ii) Combine $V_r \leq b^c$ with (i) and take logarithms. The function $f(r) = r \log_2(1 + 2b/r)$ is increasing and concave, and $f(\alpha^\star b) = b$ for the solution $\alpha^\star \approx 0.376$ of $\alpha \log_2(1 + 2/\alpha) = 1$, where $f'(\alpha^\star b) = \log_2(1 + 2/\alpha^\star) - 2/((\alpha^\star + 2)\ln 2) > 1.44$. By concavity $f(r) \leq b - 1.44(\alpha^\star b - r)$ for $r \leq \alpha^\star b$, so $f(r) \geq b - 1 - c \log_2 b$ forces $r \geq \alpha^\star b - (1 + c \log_2 b)/1.44$. (iii) Here $2^\beta \leq b$, so $m = k(2^\beta - 1) < kb \leq b^2$, and $k\beta \leq b < (k+1)\beta \leq 2k\beta$ gives $b/k < 2\beta$. Theorem D.3(b) and $k \geq b/\beta - 1$, $k \leq b/\beta$ give the display; with $\beta = \lfloor \log_2 b \rfloor$ the subtracted terms are $O(b/\log b)$ and $O(rb \log\log b/\log b)$. □

For two rounds the bound of (i) exceeds one once $2^{b-1} > (b+1)^2$, that is for $b \geq 8$; for smaller caps Appendix D.4 computes the round constants of B_{b-1} numerically (two rounds lose at least 1.80 at $b = 3$ and 3.32 at $b = 4$). The single tree uses a number of coordinates exponential in b, like the address contrast; part (iii) removes this at the price of the lower-order term in the exponent, exactly as products of address blocks do for fixed forms (Proposition C.7). The product is, however, evaluated exactly in $\beta = \lfloor \log_2 b \rfloor$ rounds within the cap: reading level j of every block in round j uses $k\beta \leq b$ questions. Part (iii) therefore gives a lower bound only for $r = o(\log b/\log\log b)$, and the linear number of rounds in (ii) is proved only for the single tree, with exponentially many coordinates; whether more than logarithmically many rounds can be necessary on a number of coordinates polynomial in b is open. Whether the factor $(1 + 2b/r)^r$ can be removed, so that the r-round gap is 2^{b-r} up to constants, is open.

Proof of Theorem D.3. (a) Following the pointer path of every block returns d itself with $k\beta \le b$ queries, so $V_{\mathrm{branch}} \le 1$; every unbiased design has $\sum_\iota g_\iota(x) = d(x) = \pm 1$, and Cauchy–Schwarz gives $\sum_\iota g_\iota(x)^2/q_\iota \ge 1$ at every x, so $V_{\mathrm{branch}} \ge 1$. The remaining inequality is Theorem D.1(a).

(b) Let U be the uniform law on all coordinates. Under U the blocks are independent, $\mathbb{E}_{\mathsf{U}}[d^2] = 1$, and the pointer path of each block is a uniformly distributed root-to-leaf path. By the dual formula (B.3) with witness $h = d$, $V_r(d,b) \ge 1/\max_{\iota \in \mathcal{A}_r(b)} E_\iota$ with $E_\iota = \mathbb{E}_{\mathsf{U}}[\mathbb{E}_{\mathsf{U}}[d \mid \ell_\iota(X)]^2]$. Fix a deterministic r-round interview ι. Let S_t be the set of coordinates it reads in round t; S_t is a function of the transcript T_{t-1} of rounds $1,\dots,t-1$, and we may assume that no coordinate is read twice. For block i write $a_{t,i}$ and $\lambda_{t,i}$ for the numbers of its pointer and data coordinates in S_t and $n_{t,i} = a_{t,i} + \lambda_{t,i}$, so that $\sum_{t,i} n_{t,i} \le b$; let $I_{t,i}$ be the set of pointer coordinates of block i read in rounds $1,\dots,t$, $\mathrm{Obs}_{t,i}$ the set of its leaves whose data coordinate is read in rounds $1,\dots,t$, and $c_i^\star$ the leaf selected in block i.

Step 1: only selected leaves carry signal. The event $\{T_r = \tau\}$ fixes the values of the coordinates read under τ and nothing else, so conditionally on the transcript the unread coordinates remain independent and uniform and the blocks remain independent. Hence $\mathbb{E}_{\mathsf{U}}[d \mid \ell_\iota] = \prod_i \mathbb{E}_{\mathsf{U}}[B_L(X^{(i)}) \mid \ell_\iota]$. Within a block, the data coordinate of an unread leaf v is independent of the transcript and of the pointer path, so $\mathbb{E}_{\mathsf{U}}[Z_v \mathbf{1}\{c_i^\star = v\} \mid \ell_\iota] = 0$, and

$$\big|\mathbb{E}_{\mathsf{U}}[B_L(X^{(i)}) \mid \ell_\iota]\big| = \Big|\sum_{v \in \mathrm{Obs}_{r,i}} Z_v\, \mathbb{P}(c_i^\star = v \mid T_r)\Big| \le \pi_{r,i}, \qquad \pi_{t,i} := \mathbb{P}(c_i^\star \in \mathrm{Obs}_{t,i} \mid T_t).$$

Since $0 \le \pi_{r,i} \le 1$,

$$E_\iota \le \mathbb{E}\Big[\prod_{i=1}^k \pi_{r,i}^2\Big] \le \mathbb{E}\Big[\prod_{i=1}^k \pi_{r,i}\Big]. \tag{D.1}$$

Step 2: the conditional law of a selected leaf. For a leaf v of block i let $R_t(v) = |\mathrm{path}(v) \cap I_{t,i}|$ be the number of pointer coordinates on its path read by the end of round t, and let $C_t(v)$ be the event that every one of them selects the child towards v. Since $c_i^\star = v$ exactly when all L pointer coordinates on $\mathrm{path}(v)$ select the child towards v, and the unread ones are uniform given T_t,

$$\mathbb{P}(c_i^\star = v \mid T_t) = \mathbf{1}\{C_t(v)\}\, 2^{R_t(v) - L}. \tag{D.2}$$

Step 3: a potential for one block. Put $M_{t,i} = \max_v \mathbf{1}\{C_t(v)\} 2^{R_t(v)}$ over the leaves of block i, so that $M_{0,i} = 1$ and, by (D.2), $\max_v \mathbb{P}(c_i^\star = v \mid T_t) = M_{t,i} 2^{-L}$, and put

$$Q_{t,i} = M_{t,i} + 2^L \pi_{t,i}, \qquad Q_{0,i} = 1.$$

We claim

$$\mathbb{E}[Q_{t,i} \mid T_{t-1}] \le (1 + 2a_{t,i} + \lambda_{t,i})\, Q_{t-1,i} \le (1 + 2n_{t,i})\, Q_{t-1,i}. \tag{D.3}$$

The most likely leaf. We show $\mathbb{E}[M_{t,i} \mid T_{t-1}] \le (1 + 2a_{t,i}) M_{t-1,i}$. Condition on T_{t-1}, which fixes S_t, drop the block index, and let $v^\star$ attain M_t. If no coordinate of S_t lies on $\mathrm{path}(v^\star)$, then $R_t(v^\star) = R_{t-1}(v^\star)$ and $C_t(v^\star) = C_{t-1}(v^\star)$, so $M_t \le M_{t-1}$. Otherwise let u be the deepest node of $\mathrm{path}(v^\star) \cap S_t$, write $\mathrm{path}(u)$ for the nodes above u together with u itself, and define

$$G(u) = \mathbf{1}\{\text{every node of } (\mathrm{path}(u) \setminus \{u\}) \cap I_t \text{ selects towards } u\}\, 2^{|\mathrm{path}(u) \cap I_t|},$$

$$F(u) = \max_{w \text{ below } u} \mathbf{1}\{\text{every node of } (\mathrm{path}(w) \setminus \mathrm{path}(u)) \cap I_{t-1} \text{ selects towards } w\}\, 2^{|(\mathrm{path}(w) \setminus \mathrm{path}(u)) \cap I_{t-1}|}.$$

No node of $\mathrm{path}(v^\star)$ strictly below u is read in round t, so every read node there belongs to I_{t-1} and $\mathbf{1}\{C_t(v^\star)\} 2^{R_t(v^\star)} \le G(u)F(u)$: the factor $G(u)$ accounts for the nodes of $\mathrm{path}(u)$ other than u and for the count including u, the bound $\mathbf{1}\{x_u \to v^\star\} \le 1$ drops u's own selection, and $F(u)$ dominates the contribution of the nodes below u because $w = v^\star$ is admissible in its maximum. Consequently $M_t \le M_{t-1} + \sum_{u \in S_t \cap U} G(u)F(u)$, the sum being over the pointer coordinates of the block in S_t. The quantity $F(u)$ is T_{t-1}-measurable. Given T_{t-1}, the $|\mathrm{path}(u) \cap S_t| - 1$ newly read nodes strictly above u select towards u independently with probability $1/2$ each, so

$$\mathbb{E}[G(u) \mid T_{t-1}] = \mathbf{1}\{\text{every node of } \mathrm{path}(u) \cap I_{t-1} \text{ selects towards } u\}\, 2^{|\mathrm{path}(u) \cap I_{t-1}|} \cdot 2.$$

Multiplying by $F(u)$ reassembles, for the leaf w below u attaining $F(u)$, exactly $2 \cdot \mathbf{1}\{C_{t-1}(w)\} 2^{R_{t-1}(w)} \le 2M_{t-1}$, because $u \notin I_{t-1}$. Summing over the a_t pointer coordinates of the block in S_t proves the claim.

The leaves already read. The sets $\mathrm{Obs}_{t-1,i}$ and S_t are T_{t-1}-measurable and disjoint, so $\pi_{t,i} = \mathbb{P}(c_i^\star \in \mathrm{Obs}_{t-1,i} \mid T_t) + \mathbb{P}(c_i^\star \in S_t \mid T_t)$ (we write $c_i^\star \in S_t$ when the data coordinate of that leaf is read in round t), and by the tower property and (D.2),

$$\mathbb{E}[\pi_{t,i} \mid T_{t-1}] = \pi_{t-1,i} + \mathbb{P}(c_i^\star \in S_t \mid T_{t-1}) \le \pi_{t-1,i} + \lambda_{t,i}\, M_{t-1,i}\, 2^{-L}.$$

Adding 2^L times this inequality to the bound for $M_{t,i}$ gives $\mathbb{E}[Q_{t,i} \mid T_{t-1}] \le (1 + 2a_{t,i} + \lambda_{t,i}) M_{t-1,i} + 2^L \pi_{t-1,i}$, which is (D.3).

Step 4: blocks multiply. Given T_{t-1} the batch S_t is fixed, the answers of round t in different blocks are independent, and $Q_{t,i}$ is a function of T_{t-1} and of the round-t answers in block i. Hence $\mathbb{E}[\prod_i Q_{t,i} \mid T_{t-1}] = \prod_i \mathbb{E}[Q_{t,i} \mid T_{t-1}] \le \prod_i (1+2n_{t,i}) \prod_i Q_{t-1,i}$. Let $\Psi_t = \prod_i Q_{t,i} \prod_{s\le t} \prod_i (1+2n_{s,i})^{-1}$. As the $n_{t,i}$ are T_{t-1}-measurable, $\mathbb{E}[\Psi_t \mid T_{t-1}] \le \Psi_{t-1}$ and $\mathbb{E}[\Psi_r] \le \Psi_0 = 1$.

Step 5: combination. The pathwise constraint $\sum_{t\le r} \sum_i n_{t,i} \le b$ and the inequality between arithmetic and geometric means over the rk pairs (t,i) give $\prod_{t\le r} \prod_i (1 + 2n_{t,i}) \le (1 + 2b/(rk))^{rk}$. Since $2^L \pi_{r,i} \le Q_{r,i}$,

$$\mathbb{E}\Big[\prod_{i=1}^{k} \pi_{r,i}\Big] \le 2^{-Lk}\, \mathbb{E}\Big[\prod_i Q_{r,i}\Big] = 2^{-Lk}\, \mathbb{E}\Big[\Psi_r \prod_{t\le r} \prod_i (1+2n_{t,i})\Big] \le 2^{-k(\beta-1)} \Big(1 + \frac{2b}{rk}\Big)^{rk}.$$

With (D.1) and the dual formula this proves the first statement of (b). For $r = 1$ and $k = 1$, (D.2) at $t = 0$ gives directly $\mathbb{P}(c^\star \in \mathrm{Obs}) \le \lambda_1 2^{-L} \le b 2^{-L}$, which is the last statement. □

The argument is a potential-function version of the lucky-hit bound: a round can raise the conditional probability of the most likely leaf only by a factor linear in the number of pointer coordinates it reads, whereas following the path raises it by a factor two per question; the potential Q adds the probability that a leaf already read is the selected one, which makes the bound multiplicative across blocks even though an interview may split its questions among blocks adaptively. As a numerical check, the released code verifies (D.3) exhaustively on the tree with $L = 2$ (all 4,752 pairs of a state and a nonempty batch; the largest ratio of the two sides is exactly one) and on random states and batches for $L = 3$ and $L = 4$ (largest ratio one and 0.83).

Interpretation. Each round can advance the known part of the pointer path by at most one level unless a question lands on an unknown pointer target, which happens with probability at most $1/D$ per question. Additional questions within a round therefore do not substitute for a round. Practical counterparts of rounds are questionnaire pages, clinic visits and batches of laboratory orders. The address contrast of Theorem 1 is the case $r = 1$, where two rounds already attain the branching coefficient. The pointer-chasing family shows that this is not a general property of two-stage designs.

D.2. Two Rounds Already Attain the Universal Extremum

An R-round interview uses at most R batches, choosing each batch of coordinates using only the preceding batches' answers, with at most b coordinates in total. Let V_R use all such interviews. One round is the fixed-form menu and b rounds suffice for full branching. For any $R \ge 2$,

$$1 \le V_{\mathrm{fixed}}/V_R \le K_{m,b}, \qquad \sup_{m,d\ne 0} \frac{V_{\mathrm{fixed}}(d,b)}{V_R(d,b)} = 2^{b-1},$$

where the supremum is over identifiable binary contrasts. The upper bound follows from menu inclusion. The address construction attains equality using two batches: all address bits, then the addressed data bit. Thus the universal extremum measures the value of permitting a second batch; it does not require b sequential rounds.

The block construction also needs at most two rounds throughout its identifiable budget curve. At $b = 2k + R$, $0 \le R \le k$, use the paper's three-query strategy on R blocks and two-query strategy on the others. Ask the first two queries in every block together; then ask the selected third coordinate in the R adaptive blocks. At $b \ge 3k$ this already evaluates the contrast exactly. The address and block examples do not separate two rounds from unrestricted branching. Corollary C.12 supplies a constant separation with a four-feature contrast, and the pointer constructions of Appendix D.1 supply the exponential hierarchy.

D.3. Labels and Across-Person Adaptation

Let R_{pre} require selecting the interview independently of the current label, and let R_{label} allow observing that label first, while both permit adaptation across people. For an identifiable contrast, Theorem B.2 and the iid upper construction imply

$$\lim_{n\to\infty} nR_{\mathrm{pre}}(d,n) = \lim_{n\to\infty} nR_{\mathrm{label}}(d,n) = V(d,\mathcal{A}).$$

Thus neither access to earlier people nor early access to the current label reduces the first-order worst-case coefficient relative to an optimal iid design.

There is also an exact statement for the *finite program*. If one extends Φ_n to separate form laws and score tables for $y=-1,+1$, its objective is the maximum over these labels of their individual worst-profile quadratic objectives. Multiplying the negative-label tables by -1 turns both subproblems into the definition of $\Phi_n(d,\mathcal{A})$. Each has value at least Φ_n, and using the same form law and sign-adjusted tables attains it. Early labels therefore do not improve this program. This is not an equality claim for the exact finite-n minimax risks, which the program bounds only within a constant.

D.4. Round Constants of the Binary Pointer Tree at Small Caps

The two-round bound of Theorem 2(b) exceeds one only for $b \geq 8$. For small caps we compute the round constants of B_{b-1} directly. Under the uniform law U with a label independent of the profile, $M_r(\mathsf{U})$, the optimal second moment over r-round designs, is a lower bound on V_r, because $V_r = \sup_P M_r(P)$ by Sion's theorem applied to (2). For $b \leq 4$ the cube is small enough to evaluate the uniform-optimal design at every profile; its worst profile is an upper bound on V_r. The solver of Appendix E.4 computes $M_r(\mathsf{U})$ with certified bounds that agree to 10^{-7} or better. At $b=3$ and $b=4$ the two bounds

Table 2. Round constants of the binary pointer tree B_{b-1} ($m = 2^b - 1$). Entries are $M_r(\mathsf{U})$, a lower bound on V_r; the brackets give the worst profile of the uniform-optimal design, an upper bound on V_r, when it differs. The last columns are the upper bound 2^{b-2} of Theorem 2(a) and the lower bound $2^{b-1}/(b+1)^2$ of Theorem 2(b) for $r=2$.

b	m	fixed	$r=2$	$r=3$	$r=4$	2^{b-2}	$2^{b-1}/(b+1)^2$
3	7	4.000	1.800	1.000		2	0.25
4	15	8.000	3.324 [3.334]	1.800	1.000	4	0.32

agree to the reported precision except for two rounds at $b=4$, where they differ by 0.3%. Two rounds therefore lose almost the factor 2^{b-2} allowed by Theorem 2(a) already at these caps, and each missing round costs a factor close to two. The cap $b=5$ ($m=31$, 169,911 forms) was not completed.

E. Population Constants and Their Computation

E.1. The Value of Adaptivity at a Given Population

Fix a menu $\mathcal{A}$ and a law P on $\mathcal{X}\times\{-1,1\}$. A *design* is a probability vector (q_ι) on $\mathcal{A}$ with tables $g_\iota(\ell,y)$ that satisfy $\sum_\iota g_\iota(\ell_\iota(x),y) = t(x,y) = yd(x)$ for all (x,y). Its score is $H = g_\iota(\ell_\iota(X),Y)/q_\iota$ with $\iota\sim q$ drawn independently of the record, which is unbiased for t at every profile and label; conversely every function H of $(\iota,\ell_\iota(X),Y)$ with $\mathbb{E}[H\mid X,Y]=t$ arises in this way. $W_{\mathcal{A}}(P)$ is the infimum of $\mathrm{Var}_{P,q}(H)$ over designs, $M_{\mathcal{A}}(P) = W_{\mathcal{A}}(P) + \Delta(P)^2$ the infimum of the second moment, and $\sigma_P^2 = \mathrm{Var}_P(t)$. At a full-support law, $W_{\mathcal{A}}(P)$ is also the local asymptotic minimax risk over arbitrary estimators (Proposition C.10), so ratios of W are ratios of the numbers of records that two menus need. For a deterministic depth-b tree τ that computes d, $S_\tau(x)$ is its padded path set and $\mathcal{S}_\tau$ the collection of these sets (Appendix C.1).

Theorem E.1 (Records needed at a population, full statement of Theorems 1(ii), 2(a) and 3)**.** *Let d be a nonzero contrast of degree at most b on $\{-1,1\}^m$ and P any law on $\mathcal{X}\times\{-1,1\}$.*

(a) Uniform gap. *For $1\leq r\leq b$,*

$$W_{\mathrm{branch}}(P) \;\leq\; W_r(P) \;\leq\; \min\left\{2^{b-r}, \binom{m-r}{b-r}\right\} W_{\mathrm{branch}}(P),$$

and in particular $W_{\mathrm{fixed}}(P) \leq K_{m,b} W_{\mathrm{branch}}(P)$.

(b) Witness bound. *For every menu $\mathcal{A}$, every constant c and every function h on $\mathcal{X} \times \{-1, 1\}$ with $\mathbb{E}_P[(t-c)h] \neq 0$,*

$$W_{\mathcal{A}}(P) + (\Delta(P) - c)^2 \;\geq\; \frac{\mathbb{E}_P[(t-c)h]^2}{\max_{\iota \in \mathcal{A}} \mathbb{E}_P\big[\mathbb{E}_P[h \mid \ell_\iota(X), Y]^2\big]}.$$

With $c = 0$ this bounds the second moment $M_{\mathcal{A}}(P)$, and with $c = \Delta(P)$ it bounds the variance $W_{\mathcal{A}}(P)$.

(c) Bracket. *If a depth-b tree τ computes d and $\sigma_P^2 > 0$, then $W_{\mathrm{branch}}(P) = \sigma_P^2$ and*

$$\frac{1}{\rho_P^2} \;\leq\; \frac{W_{\mathrm{fixed}}(P)}{W_{\mathrm{branch}}(P)} \;\leq\; N_P(\tau) \;\leq\; |\mathcal{S}_\tau| \;\leq\; K_{m,b},$$

where

$$\rho_P^2 = \max_{|T|=b} \frac{\mathrm{Var}_P\big(\mathbb{E}_P[t \mid X_T, Y]\big)}{\mathrm{Var}_P(t)}, \qquad N_P(\tau) = \Big(\sum_{S \in \mathcal{S}_\tau} v_S^{1/2}\Big)^2, \quad v_S = \frac{\mathbb{E}_P\big[(t - \Delta(P))^2 \mathbf{1}\{S_\tau(X) = S\}\big]}{\sigma_P^2}.$$

More generally $W_{\mathrm{fixed}}(P) \leq \big(\sum_S \mathbb{E}_P[(t-c)^2 \mathbf{1}\{S_\tau(X) = S\}]^{1/2}\big)^2 - (\Delta(P) - c)^2$ for every constant c.

(d) *For the address contrast with $\kappa = b - 1$ under the uniform law with independent uniform labels, $1/\rho_P^2 = N_P(\tau) = K_{m,b} = D$, so all inequalities in* (c) *are equalities.*

The quantity ρ_P^2 is the largest fraction of the variance of the target that a single form explains, a population R^2. The weights v_S sum to one, and $N_P(\tau) = \exp H_{1/2}(v)$ is the exponential of their Rényi entropy of order $1/2$: an effective number of question sets that the tree uses at P. Adaptivity can be valuable only when the tree's questions diverge over many well-populated paths, and it is valuable when in addition no fixed set of b questions explains most of the variance of the target; correlated features and shared coordinates raise ρ_P^2.

Proof. (a) The left inequality is menu inclusion. For the right one fix a branching design: trees τ with probabilities $q_\tau > 0$ and tables $g_\tau(\ell, y)$ with $\sum_\tau g_\tau(\ell_\tau(x), y) = t(x, y)$, and write $H_\tau(x, y) = g_\tau(\ell_\tau(x), y)/q_\tau$, so that $\sum_\tau q_\tau H_\tau = t$ pointwise. Pad every tree as in Appendix C.1; padding refines the transcript, so H_τ is still a function of the padded leaf and the label. For a node v of τ at depth $r-1$ (the root if $r = 1$), the padded subtree below v has depth $b - r + 1$; let $\mathcal{S}_{\tau,v}$ be the collection of coordinate sets of its root-to-leaf paths and $L_{\tau,v} = |\mathcal{S}_{\tau,v}|$. These sets contain the coordinate asked at v and are determined by the first $b - r$ answers below v, so $L_{\tau,v} \leq \min\{2^{b-r}, \binom{m-r}{b-r}\} =: K_r$. Fix a constant c and consider the following randomized r-round interview with at most b queries. Draw $\tau \sim q$ and, independently of the record, one set S_v uniformly from $\mathcal{S}_{\tau,v}$ for every node v of depth $r-1$. In rounds $1, \ldots, r-1$ ask one question per round and follow τ to a node v; in round r ask the coordinates of S_v. The path of x below v can be followed to a leaf exactly when its coordinate set equals S_v, which has conditional probability $1/L_{\tau,v}$ given (x, y, τ), and this event is measurable from the transcript. Return

$$\widetilde{H} = c + L_{\tau,v}\big(H_\tau(x, y) - c\big) \text{ on completion}, \qquad \widetilde{H} = c \text{ otherwise}.$$

The pair $(\tau, (S_v)_v)$ indexes a deterministic r-round interview, so $\widetilde{H}$ is a score of a design on $\mathcal{A}_r(b)$. Conditionally on (x, y) and τ, its mean is $H_\tau(x, y)$ and $\mathbb{E}[(\widetilde{H} - c)^2 \mid x, y, \tau] = L_{\tau,v}(H_\tau(x, y) - c)^2 \leq K_r(H_\tau(x, y) - c)^2$. Hence $\mathbb{E}[\widetilde{H} \mid X, Y] = t$, and with $H = H_\tau$ for $\tau \sim q$,

$$\mathrm{Var}(\widetilde{H}) = \mathbb{E}[(\widetilde{H} - c)^2] - (\Delta(P) - c)^2 \leq K_r\, \mathbb{E}[(H - c)^2] - (\Delta(P) - c)^2.$$

Choose $c = \Delta(P)$: then $\mathbb{E}[(H - c)^2] = \mathrm{Var}(H)$ and $\mathrm{Var}(\widetilde{H}) \leq K_r\, \mathrm{Var}(H)$. The constant c is part of the score, not an estimate: the design is pointwise unbiased for every value of c, and $W_r(P)$ is by definition an infimum over designs that may be tuned to P. Taking the infimum over branching designs proves (a).

(b) Let (q, g) be a design on $\mathcal{A}$ with score H. The tables $g_\iota - c q_\iota$ sum to $t - c$, and $\mathbb{E}[(H - c)^2] = \sum_\iota \|g_\iota - c q_\iota\|_P^2 / q_\iota \geq (\sum_\iota \|g_\iota - c q_\iota\|_P)^2$ by Cauchy–Schwarz in q. For every h,

$$\mathbb{E}_P[(t-c)h] = \sum_\iota \mathbb{E}_P\big[(g_\iota - c q_\iota)\, \mathbb{E}_P[h \mid \ell_\iota(X), Y]\big] \leq \Big(\sum_\iota \|g_\iota - c q_\iota\|_P\Big) \max_\iota \big\|\mathbb{E}_P[h \mid \ell_\iota(X), Y]\big\|_P,$$

because $g_\iota - cq_\iota$ is a function of $(\ell_\iota(X), Y)$. Square, use $\mathbb{E}[(H-c)^2] = \mathrm{Var}(H) + (\Delta(P)-c)^2$ and take the infimum over designs.

(c) Following τ returns t itself, and every unbiased score has $\mathrm{Var}(H) \geq \mathrm{Var}(\mathbb{E}[H \mid X, Y]) = \sigma_P^2$, so $W_{\mathrm{branch}}(P) = \sigma_P^2$. For the lower bound apply (b) to the fixed forms with $c = \Delta(P)$ and $h = t - \Delta(P)$: the numerator is σ_P^4 and $\mathbb{E}_P[\mathbb{E}_P[t-\Delta \mid X_T, Y]^2] = \mathrm{Var}_P(\mathbb{E}_P[t \mid X_T, Y])$ because $\mathbb{E}_P[t \mid X_T, Y]$ has mean $\Delta(P)$. For the upper bound choose $q_S > 0$ on $\mathcal{S}_\tau$, draw S before the person answers and return $c + (t(X,Y) - c)/q_S$ if the tree's path completes inside S and c otherwise. Completion occurs exactly when $S = S_\tau(X)$ (Appendix C.1), so the score is unbiased at every profile and label, and $\mathbb{E}[(H-c)^2] = \sum_S q_S^{-1} \mathbb{E}_P[(t-c)^2 \mathbf{1}\{S_\tau(X) = S\}]$. Minimizing over the simplex gives $q_S \propto \mathbb{E}_P[(t-c)^2 \mathbf{1}\{S_\tau = S\}]^{1/2}$ and the value $(\sum_S \mathbb{E}_P[(t-c)^2 \mathbf{1}\{S_\tau = S\}]^{1/2})^2$; sets of zero weight are handled by letting their probability tend to zero. Subtracting $(\Delta(P)-c)^2$ gives the general statement, and $c = \Delta(P)$ gives $N_P(\tau)\sigma_P^2$. Finally $(\sum_S v_S^{1/2})^2 \leq |\mathcal{S}_\tau| \sum_S v_S = |\mathcal{S}_\tau|$ by Cauchy–Schwarz, and $|\mathcal{S}_\tau| \leq K_{m,b}$.

(d) Here $\Delta = 0$, $t^2 = 1$ and the D path sets are equally likely, so $N_P = D = K_{m,b}$. With independent labels, $\mathbb{E}[t \mid X_T, Y] = Y\mathbb{E}[d \mid X_T]$, whose second moment is at most $2^{-\kappa} = 1/D$ by (5), so $1/\rho_P^2 \geq D$. □

Relation to the uncentered bracket. Setting $c = 0$ in (b) and (c) gives the second-moment bracket $\mathbb{E}_P[t^2]/\Gamma_P \leq M_{\mathrm{fixed}}(P) \leq N_P^0(\tau)\mathbb{E}_P[t^2]$ with $\Gamma_P = \max_T \mathbb{E}_P[\mathbb{E}_P[t \mid X_T, Y]^2]/\mathbb{E}_P[t^2]$ and N_P^0 computed from the weights $\mathbb{E}_P[t^2 \mathbf{1}\{S_\tau = S\}]$. Subtracting Δ^2 turns it into a bracket for $W_{\mathrm{fixed}}/W_{\mathrm{branch}}$ with lower end $(\mathbb{E}_P[t^2]/\Gamma_P - \Delta^2)/\sigma_P^2$ and upper end $(N_P^0\mathbb{E}_P[t^2] - \Delta^2)/\sigma_P^2$. The centered lower end is never smaller: writing $s = \mathbb{E}_P[t^2]$ and $e = \Gamma_P s$, the claim $(s-\Delta^2)^2 \geq (s^2/e - \Delta^2)(e - \Delta^2)$ reduces to $s^2/e + e \geq 2s$. The uncentered upper end is unbounded as $\Delta^2 \to \mathbb{E}_P[t^2]$, that is for accurate rules, whereas the centered one never exceeds $K_{m,b}$; for the worked example of Section 2 with uniform profiles and a label that agrees with d with probability 0.9, the centered upper end is 2 and the uncentered one is $(2 - 0.8^2)/0.36 \approx 3.78$. Centering is what makes part (a) hold at every population: without it, path guessing inflates the second moment by K_r and the variance by more.

Label-dependent designs. Labels are observed before the questions are asked, so a collector may let the interview probabilities depend on the label, as in outcome-dependent two-phase sampling. Write $\mathcal{A}^{\mathrm{lab}}$ for this larger class of designs on a menu $\mathcal{A}$: probability vectors $q_{\cdot|y}$ for $y = \pm 1$ with tables that sum to $t(\cdot, y)$ for each label.

Proposition E.2 (Label-dependent designs)**.** (i) *$V(d, \mathcal{A})$ is unchanged if interview probabilities may depend on the current label (Theorem B.2).* (ii) *For every law P, $M_{\mathcal{A}}^{\mathrm{lab}}(P) = \sum_{y=\pm 1} M_{\mathcal{A}}(P^{(y)})$, where $P^{(y)} = P(\cdot, y)$ is the sub-probability law of the profiles with label y and $M_{\mathcal{A}}(P^{(y)})$ is the optimum of the program for the target yd under that mass; hence $W_{\mathcal{A}}^{\mathrm{lab}}(P) = \sum_y M_{\mathcal{A}}(P^{(y)}) - \Delta(P)^2 \leq W_{\mathcal{A}}(P)$.* (iii) *Part* (a) *of Theorem E.1 holds for label-dependent designs: $W_r^{\mathrm{lab}}(P) \leq K_r W_{\mathrm{branch}}^{\mathrm{lab}}(P)$, and $W_{\mathrm{branch}}^{\mathrm{lab}}(P) = \sigma_P^2$ when a depth-b tree computes d.*

Proof. (i) The converse of Theorem B.2 allows the interview to depend on the current label, and label-independent designs attain V. (ii) The second moment of a label-dependent design is $\sum_y \sum_\iota \mathbb{E}_P[g_{\iota,y}(\ell_\iota(X))^2 \mathbf{1}\{Y = y\}]/q_{\iota|y}$, and neither the constraints nor the probability vectors couple the two labels. (iii) The simulation in the proof of Theorem E.1(a) is carried out separately for each label with the same constant $c = \Delta(P)$; the conditional bound $\mathbb{E}[(\widetilde{H}-c)^2 \mid x, y] \leq K_r \mathbb{E}[(H-c)^2 \mid x, y]$ holds for each (x, y). The last statement is the Jensen bound. □

Each $M_{\mathcal{A}}(P^{(y)})$ is computed by the solver of Appendix E.4 with the same certificates. Appendix F.3 reports that label-dependent form probabilities lower the records ratio of the screening trees by at most 0.5%, and by at most 0.8% when they are also stratified by sex.

Accurate populations. The populations that attain the worst-case constants in Appendices C and D have labels independent of the profile, so that the rule under audit is no better than chance. The gaps do not depend on this.

Proposition E.3 (The gaps persist at every accuracy)**.** *Let d take values in $\{-1, 1\}$, let $X \sim \mu$ for an arbitrary profile law μ, and let $Y = d(X)\xi$, where $\xi \in \{-1, 1\}$ is independent of X with $\mathbb{P}(\xi = 1) = p \in (0, 1)$. Then $t = \xi$, $\Delta(P) = 2p - 1$, $\sigma_P^2 = 1 - \Delta^2$, and for every interview ι, with $e_\iota = \mathbb{E}_\mu[d \mid \ell_\iota(X)]$,*

$$\mathrm{Var}_P\left(\mathbb{E}_P[t \mid \ell_\iota(X), Y]\right) = \sigma_P^4 \, \mathbb{E}_\mu\left[\frac{e_\iota^2}{1 - \Delta^2 e_\iota^2}\right] \;\leq\; \sigma_P^2 \, \mathbb{E}_\mu[e_\iota^2].$$

Consequently, for every menu $\mathcal{A}$,

$$\frac{W_{\mathcal{A}}(P)}{\sigma_P^2} \;\geq\; \frac{1}{\max_{\iota\in\mathcal{A}} \|\mathbb{E}_\mu[d \mid \ell_\iota(X)]\|_\mu^2},$$

which is the bound that the dual of Theorem 3 gives for $V(d,\mathcal{A})$ with the witness $h = d$ at μ. In particular: (i) *for the address contrast with $\kappa = b-1$ and uniform profiles, $W_{\mathrm{fixed}}(P)/W_{\mathrm{branch}}(P) = 2^{b-1}$ for every $p \in (0,1)$, that is, at every accuracy $(1+\Delta)/2$ of the rule;* (ii) *for the pointer-tree contrasts of Theorem D.3 with uniform profiles, $W_r(P)/W_{\mathrm{branch}}(P) \geq 2^{k(\beta-1)}(1+2b/(rk))^{-rk}$ for every $p \in (0,1)$.*

Proof. Since $d^2 = 1$, $t = Yd(X) = \xi$. Fix ι and a transcript, and write e for e_ι at that transcript, so that $\mathbb{P}(d = \pm 1 \mid \ell_\iota) = (1 \pm e)/2$. For $y \in \{-1,1\}$, $\mathbb{P}(Y = y \mid \ell_\iota) = \frac{1}{2}(1+e)\mathbb{P}(\xi = y) + \frac{1}{2}(1-e)\mathbb{P}(\xi = -y) = \frac{1}{2}(1+y\Delta e)$, and by Bayes' rule $\mathbb{P}(d = y \mid \ell_\iota, Y = y) = p(1+ye)/(1+y\Delta e)$. Hence

$$\mathbb{E}_P[t \mid \ell_\iota, Y = y] = \frac{p(1+ye) - (1-p)(1-ye)}{1+y\Delta e} = \frac{\Delta + ye}{1+y\Delta e}, \qquad \mathbb{E}_P[t \mid \ell_\iota, Y = y] - \Delta = \frac{ye\,\sigma_P^2}{1+y\Delta e}.$$

Averaging the square over y with the weights $\frac{1}{2}(1+y\Delta e)$ gives $\frac{1}{2}e^2\sigma_P^4[(1+\Delta e)^{-1} + (1-\Delta e)^{-1}] = e^2\sigma_P^4/(1-\Delta^2 e^2)$, and taking the expectation over the transcript proves the identity. The inequality uses $e^2 \leq 1$, so that $1 - \Delta^2 e^2 \geq \sigma_P^2$. The bound on $W_{\mathcal{A}}$ is Theorem E.1(b) with $c = \Delta$ and $h = t - \Delta$, whose numerator is σ_P^4. (i) Branching computes d, so $W_{\mathrm{branch}}(P) = \sigma_P^2$; the energy of every form is at most $2^{-\kappa}$ by (5), so the ratio is at least 2^κ, and Theorem E.1(a) bounds it by $K_{m,b} = 2^\kappa$. (ii) The proof of Theorem D.3(b) bounds $\mathbb{E}_{\mathsf{U}}[\mathbb{E}_{\mathsf{U}}[d \mid \ell_\iota]^2]$ for every r-round interview, and branching computes d. □

For $b = 2, 3$ the solver confirms (i) numerically: $W_{\mathrm{fixed}}/W_{\mathrm{branch}} = 2.000000$ and 4.000000 at $p \in \{0.5, 0.6, 0.75, 0.9, 0.97\}$. The identity also shows that the explained share ρ_P^2 of an accurate rule is at most the label-free energy $\mathbb{E}_\mu[e^2]$ and equals it for forms that either determine d or carry no information about it.

Several contrasts. A collector may want the same records to serve several rules. For contrasts $d_1, \ldots, d_J$ of degree at most b and weights $\omega_j > 0$, let a design share one interview distribution and carry one table per contrast, and let $W_{\mathcal{A}}^{\Sigma}(P)$ be the smallest value of $\sum_j \omega_j \operatorname{Var}(H_j)$. The simulation in the proof of Theorem E.1(a) acts on the interview and not on the table: with one centring constant $c_j = \Delta_j(P)$ per contrast and the same drawn tree and path set, it returns $c_j + L_{\tau,v}(H_{\tau,j} - c_j)$ on completion and c_j otherwise, and every coordinate keeps its mean while its variance grows by at most K_r. Hence $W_r^{\Sigma}(P) \leq K_r W_{\mathrm{branch}}^{\Sigma}(P)$ at every population, and the fixed-form program is the same second-order cone program with stacked tables. Appendix F.6 computes it for pairs and triples of screening rules.

E.2. The Value of Adaptivity in a Structured Model

The constants V and $W_{\mathcal{A}}(P)$ are for an analyst who assumes nothing about the law of the unasked features. Here the law of (X,Y) is known to lie in a model, and we ask what remains of the gap between pre-committed and branching questions. The answer is the content of Theorem E.5: the simulation bound survives in every model; when the features are independent given the label a design matters only through its inclusion probabilities, and the gap is at most a factor two; in a model with interactions of order at most k the exponential gap becomes polynomial in the cap at product laws and near them, and it disappears for the address contrast when $k = 2$; but parsimony alone does not help, because pairwise laws with strong couplings can encode the address structure and bring the exponential gap back.

Constants in a model. Let P have full support and let $\mathcal{T} \subseteq L_0^2(P)$ be a linear space of mean-zero functions of (X,Y), the tangent space at P of a smooth parametric model $\mathcal{M} \ni P$. For an interview ι and $h \in L^2(P)$ write $J_\iota(h) = \mathbb{E}_P\big[\mathbb{E}_P[h \mid \ell_\iota(X), Y]^2\big]$, the Fisher information that the records of ι carry about the direction h, and for a design q on a menu $\mathcal{A}$ let $J_q = \sum_\iota q_\iota J_\iota$. Define

$$W_{\mathcal{A}}^{\mathcal{T}}(P) = \inf_q \sup_{h\in\mathcal{T}} \frac{\langle t - \Delta(P), h\rangle_P^2}{J_q(h)}, \tag{E.1}$$

with $0/0 = 0$ and the value $+\infty$ if $\langle t - \Delta, h\rangle_P \neq 0 = J_q(h)$ for some $h \in \mathcal{T}$. If $s = (s_1, \ldots, s_p)$ is a basis of $\mathcal{T}$, $v = \mathbb{E}_P[(t-\Delta)s]$ and $I_\iota = \mathbb{E}_P\big[\mathbb{E}_P[s \mid \ell_\iota, Y]\mathbb{E}_P[s \mid \ell_\iota, Y]^{\mathsf{T}}\big]$, then $W_{\mathcal{A}}^{\mathcal{T}}(P) = \min_q v^{\mathsf{T}}(\sum_\iota q_\iota I_\iota)^\dagger v$, with the value $+\infty$

when v is outside the range of the matrix, a c-optimal design problem. For $\mathcal{T} = L_0^2(P)$ this is the constant $W_{\mathcal{A}}(P)$ of (3) (proof of Proposition C.10), and $\mathcal{T} \subseteq \mathcal{T}'$ implies $W_{\mathcal{A}}^{\mathcal{T}}(P) \le W_{\mathcal{A}}^{\mathcal{T}'}(P) \le W_{\mathcal{A}}(P)$: a model can only lower the constant of a menu, and the question is whether it lowers the constant of fixed forms by more than that of branching.

Proposition E.4 (Local minimax risk in a model)**.** *Let $\{P_\theta : \theta \in \Theta \subseteq \mathbb{R}^p\}$ be a model on the finite space $\mathcal{X} \times \{-1, 1\}$ with θ_0 in the interior of Θ, $P_{\theta_0} = P$ of full support, $\theta \mapsto P_\theta$ continuously differentiable with a Jacobian of rank p at θ_0, and let $\mathcal{T}$ be the span of its scores at θ_0. For every sequence of protocols that choose each record's interview from $\mathcal{A}$ using the previous records, and every sequence of estimators,*

$$\lim_{c\to\infty} \liminf_{n\to\infty} \sup_{\|\theta-\theta_0\|\le c/\sqrt{n}} n\,\mathbb{E}_\theta\big[(\widehat{\Delta} - \Delta(P_\theta))^2\big] \;\ge\; W_{\mathcal{A}}^{\mathcal{T}}(P).$$

Proof. This is the converse half of the proof of Proposition C.10 with the tangent coordinates $P + U\theta$ of the simplex replaced by the model's own parameter, and with two changes. The cube of radius $c/\sqrt{pn}$ around θ_0 that carries the smooth product prior lies in Θ for large n because θ_0 is interior. On this cube the information matrix of one record of interview ι is $I_\iota(\theta) = I_\iota + o(1)$ and the gradient of $\theta \mapsto \Delta(P_\theta)$, which is no longer linear, is $v + o(1)$, uniformly, by continuous differentiability. The multivariate van Trees inequality then bounds the normalized Bayes risk below by $(v + o(1))^{\mathsf{T}}(\sum_\iota \bar{q}_\iota I_\iota + C_0/c^2 + \epsilon_n I)^{-1}(v + o(1))$ for some $\bar{q}$ in the simplex; the matrix is bounded below by C_0/c^2 uniformly in $\bar{q}$, so the $o(1)$ terms vanish in the limit for fixed c. Minimizing over $\bar{q}$ and letting $n \to \infty$ and then $c \to \infty$ gives $\min_q v^{\mathsf{T}}(\sum_\iota q_\iota I_\iota)^{\dagger} v$. □

Like $W_{\mathcal{A}}(P)$ in Proposition C.10, the constant is local, with the center P available to the procedure: when $\sum_\iota q_\iota I_\iota$ is nonsingular at a minimizing design, the maximum-likelihood estimator of $\Delta(P_\theta)$ under that design attains it for regular models such as the log-linear models below. A minimizing design may have singular information, and then $\Delta(P_\theta)$ need not be identified from its records away from P; this happens for the address contrast under the pairwise model below, whose optimal designs read only data bits. The value is then approached, up to a factor $1/(1-\epsilon)$, by mixing the design with a mass ϵ of interviews that identify the model. We use the lower bound, the comparison of menus, and the simulations of Appendix F.8.

Models of bounded interaction order. Number the label as coordinate $m+1$. The *order-k model* $\mathcal{M}_k$ is the hierarchical log-linear model $\log p(x, y) = \sum_{A:\,|A|\le k} \theta_A \chi_A(x, y)$ over $A \subseteq [m+1]$; $k = 2$ has main effects and pairwise interactions and $k = m+1$ is saturated. Its tangent space at any of its laws is $\mathcal{T}_k = \operatorname{span}\{\chi_A - \mathbb{E}_P \chi_A : 1 \le |A| \le k\}$, and we write $W_{\mathcal{A}}^{(k)}(P)$ for (E.1) with $\mathcal{T} = \mathcal{T}_k$. A *product law* makes $X_1, \dots, X_m, Y$ independent; it lies in every $\mathcal{M}_k$.

Theorem E.5 (Adaptivity in a model, full statement of Theorem 4)**.** *Let $1 \le b \le m$ and let d be a contrast of degree at most b.*

(a) Every model. *For every depth-b tree τ with padded path sets $\mathcal{S}_\tau$ and every law P, the uniform design on $\mathcal{S}_\tau$ satisfies*

$$\frac{1}{|\mathcal{S}_\tau|} \sum_{S \in \mathcal{S}_\tau} J_S(h) \;\ge\; \frac{J_\tau(h)}{|\mathcal{S}_\tau|} \qquad \text{for every } h \in L^2(P), \tag{E.2}$$

where J_S is the information of the form S. Consequently, for every tangent space $\mathcal{T}$ and $1 \le r \le b$,

$$W_{\mathrm{branch}}^{\mathcal{T}}(P) \;\le\; W_r^{\mathcal{T}}(P) \;\le\; \min\left\{2^{b-r}, \binom{m-r}{b-r}\right\} W_{\mathrm{branch}}^{\mathcal{T}}(P).$$

(b) Features independent given the label. *Let $\mathcal{M}_{\mathrm{ci}}$ be the model of all full-support laws under which $X_1, \dots, X_m$ are independent given Y, with tangent space $\mathcal{T}_{\mathrm{ci}}$ and constants $W_{\mathcal{A}}^{\mathrm{ci}}$; for designs whose interview probabilities $q_\iota(y)$ may depend on the record's label, $W_{\mathcal{A}}^{\mathrm{ci,lab}}$ is (E.1) with $J_q(h) = \sum_y P(y) \sum_\iota q_\iota(y)\,\mathbb{E}[\mathbb{E}[h \mid \ell_\iota(X), Y]^2 \mid Y = y]$ (Proposition E.4 applies to such designs with the menu of pairs of interviews, one for each label). For $P \in \mathcal{M}_{\mathrm{ci}}$ let ϕ_j^y be X_j standardized under $P(\cdot \mid Y = y)$, $\hat{d}_j^{\,y} = \mathbb{E}_P[d(X)\phi_j^y(X_j) \mid Y = y]$, and*

$$M_b(c) = \min\Big\{ \sum_{j=1}^m \frac{c_j^2}{\pi_j} :\; 0 \le \pi_j \le 1,\; \sum_{j=1}^m \pi_j \le b \Big\} \qquad (0/0 = 0).$$

Then, for every contrast d and every law $P \in \mathcal{M}_{\mathrm{ci}}$:

(i) *if the design may depend on the record's label, fixed forms, r-round interviews for every r and branching interviews have the same constant,* $W^{\mathrm{ci,lab}}(P) = \mathrm{Var}_P(\mathbb{E}_P[t \mid Y]) + \sum_y P(y)\, M_b(\hat{d}^{\,y})$*;*

(ii) *for designs chosen without the label,* $W^{\mathrm{ci,lab}}(P) \le W^{\mathrm{ci}}_{\mathrm{branch}}(P) \le W^{\mathrm{ci}}_{r}(P) \le W^{\mathrm{ci}}_{\mathrm{fixed}}(P) = \mathrm{Var}_P(\mathbb{E}_P[t \mid Y]) + M_b(\bar{c}) \le 2\, W^{\mathrm{ci,lab}}(P)$*, where* $\bar{c}_j^{\,2} = \sum_y P(y)(\hat{d}_j^{\,y})^2$*;*

(iii) *if P is a product law with $P(Y=1) = 1/2$, then also in the pairwise model* $W^{(2)}_{\mathrm{fixed}}(P) = W^{(2)}_{r}(P) = W^{(2)}_{\mathrm{branch}}(P) = (\mathbb{E}_P d)^2 + M_b(\hat{d})$ *for every r, with* $\hat{d}_j = \mathbb{E}_P[d(X)\phi_j(X_j)]$*.*

The minimum is attained by $\pi_j = \min\{1, |c_j|/\lambda\}$ *for the λ that makes the sum equal to b (and by* $\pi_j = \mathbf{1}\{c_j \neq 0\}$ *if at most b of the c_j are nonzero); it equals* $(\sum_j |c_j|)^2/b$ *when* $b \max_j |c_j| \le \sum_j |c_j|$*.*

(c) Bounded interaction order at product laws. *Let P be a product law whose coordinates and label take each value with probability at least $\lambda \in (0, 1/2]$, and let $2 \le k \le b+1$. For every depth-b tree τ, the fixed design* $q^\tau_S = P(S_\tau(X) = S)$ *satisfies* $\sum_S q^\tau_S J_S(h) \ge J_\tau(h)/\nu_k$ *for every $h \in \mathcal{T}_k$, with*

$$\nu_k = \nu_k(b, \lambda) = \sum_{j=1}^{k} \binom{b+1}{j} \Big(\frac{1-\lambda}{\lambda}\Big)^j,$$

and therefore

$$W^{(k)}_{\mathrm{fixed}}(P) \;\le\; \min\{K_{m,b},\ \nu_k(b,\lambda)\}\, W^{(k)}_{\mathrm{branch}}(P).$$

At uniform laws $\nu_k = \sum_{j=1}^{k} \binom{b+1}{j} \le (b+1)^k$*. The bound is stable: if $k \le b$ and P_0 is such a product law, then for every $\epsilon > 0$ there is a neighbourhood of P_0 in $\mathcal{M}_k$ on which* $W^{(k)}_{\mathrm{fixed}}(P) \le (1+\epsilon)\, \nu_k(b,\lambda)\, W^{(k)}_{\mathrm{branch}}(P)$*.*

(d) The address contrast. *Let d be the address contrast with $\kappa = b-1$ address bits and $D = 2^\kappa$ data bits, and P the uniform law with an independent uniform label, so that $\sigma_P^2 = 1$ and* $W_{\mathrm{fixed}}(P)/W_{\mathrm{branch}}(P) = 2^{b-1}$*. Write* $N(s,j) = \sum_{i=0}^{j} \binom{s}{i}$*. Then*

$$W^{(2)}_{\mathrm{fixed}}(P) = W^{(2)}_{\mathrm{branch}}(P) = \frac{1}{b},$$

and for $3 \le k \le b+1$,

$$W^{(k)}_{\mathrm{branch}}(P) \le 1, \qquad W^{(k)}_{\mathrm{fixed}}(P) \;\ge\; L_k(b) = \frac{N(b-1, k-2)^2}{\max_{0 \le s \le b-1}(b-s)\, N(s, k-2)} \;\ge\; \frac{1}{b}\binom{b-1}{k-2}.$$

Here $L_3(b) = b^2/\lfloor (b+1)^2/4 \rfloor < 4$ *and* $L_{b+1}(b) = 2^{b-1}$*. For $k = 3$, moreover,* $W^{(3)}_{\mathrm{fixed}}(P) < 4$ *for every $b \ge 2$, and already two-round interviews give*

$$W^{(3)}_{\mathrm{branch}}(P) \;\le\; U_3(b) = \min_{\substack{s+r=b,\ s,r \ge 1 \\ r \le 2^{\kappa-s}}} \Big(\frac{\kappa}{s\sqrt{r}} + \frac{\kappa - s}{s\sqrt{b}}\Big)^2 = \frac{9 + o(1)}{b},$$

so that $W^{(3)}_{\mathrm{fixed}}(P)/W^{(3)}_{\mathrm{branch}}(P) \ge L_3(b)/U_3(b) = (4/9 + o(1))\, b$*.*

(e) Strong dependence. *Let $b \ge 3$, $\kappa = b-1$ and $D = 2^\kappa$. Add to the $\kappa + D$ coordinates of the address contrast $2D$ auxiliary binary features $s, p \in \{0,1\}^D$, which d does not depend on; write z_j for the indicator of $Z_j = 1$. (Coding a feature by $\{0,1\}$ is an affine change of the ± 1 coordinate and leaves every $\mathcal{M}_k$ unchanged.) For every $\epsilon > 0$ there is a full-support law P in the pairwise model $\mathcal{M}_2$ on these $m = b - 1 + 3 \cdot 2^{b-1}$ features and the label, under which, with probability at least $1 - \epsilon$, s is the one-hot code of the address and $p_j = s_j z_j$ for all j, such that for every $k \ge 2$,*

$$\frac{W^{(k)}_{\mathrm{fixed}}(P)}{W^{(k)}_{\mathrm{branch}}(P)} \;\ge\; (1-\epsilon)\, \frac{2^{b-1}}{b}.$$

For $b = 2$ the same construction gives only the trivial bound one.

Part (a) is the model-level form of Theorem C.1: with probability $1/|\mathcal{S}_\tau|$ a record of the uniform path-set design contains the transcript that τ would have produced, whatever the analyst intends to do with it, so the factor 2^{b-1} bounds the value of branching for maximum likelihood in any model, for Bayesian analysis and for imputation alike. Part (b) is the benign extreme: when the features are independent given the label, whatever their association with the label and whatever the accuracy of the rule, a record contributes to the estimate only through which coordinates it reveals, and inclusion probabilities are all that a design controls. Any pair of inclusion vectors that a tree realizes is realized by a label-dependent mixture of forms, so adaptivity within the record has no first-order value once the label may be used. Without the label, a tree can use early answers as a proxy for the label and realize inclusion vectors that differ between the classes, which no label-blind mixture of forms does; the factor two in (ii) bounds this advantage, and it is nearly attained: if one feature is a 1%-noisy copy of the label and the rule averages one of two groups of fair coins according to that feature, label-blind forms need 1.31, 1.47 and 1.57 times the records of the tree that asks the copy first, at $b = 3, 4, 5$, about $2(b-1)/b$. The model is the one behind naive-Bayes classifiers and latent-class analyses with the outcome as the class; it is a submodel of $\mathcal{M}_2$, and (iii) shows that at balanced product laws the feature-by-feature interactions of $\mathcal{M}_2$ change nothing. Away from those laws they do: for random depth-b rules at product laws with an unbalanced label, and at laws of $\mathcal{M}_{\mathrm{ci}}$ with label-dependent features, the pairwise constants computed by the solver differ by up to 15% between forms and branching (a numerical exploration with the solver of Section 4). Parts (c) and (d) say that, at laws without dependence, the bound is far from attained when the model is parsimonious. Under the order-k model at uniform laws the price of pre-commitment is at most $(b+1)^k$ for every contrast, and for the address contrast, which attains 2^{b-1} without a model, it is exactly one under the pairwise model, a case of (b)(iii), and 2^{b-1} again at $k = b+1$, when the model contains the interaction of all b queried items with the label. In between it is polynomial. Under the order-k model forms need at least $L_k(b) \geq \binom{b-1}{k-2}/b$ times the records of the rule's own tree, a polynomial of degree $k-3$. Under the three-way model this factor stays below four, but the rule's own tree is no longer the best branching design once $b \geq 6$: an interview that reads some address bits and then several data bits consistent with them needs only $U_3(b) = O(1/b)$, so the gap grows at least linearly in the cap ($U_3 = 1$ for $b \leq 5$; the bound L_3/U_3 is 3.09 at $b = 6$, 4.10 at $b = 8$ and 9.28 at $b = 20$). At product laws the exponential separation is thus a statement about interactions of order growing with the cap. Part (e) shows that this reading needs weak dependence and not only a parsimonious model. In the law of part (e) the address bits, their one-hot copy and the product indicators are tied by strong pairwise couplings, as dummy-coded categorical items and derived indicators are in a real questionnaire. On the set that carries almost all of the mass the target is a polynomial of degree two, so it is nearly in the tangent space of the pairwise model, the model is of no help, and forms need $2^{b-1}/b$ times the records of the rule's tree, within a factor b of the bound of part (a). The order of a log-linear model therefore measures parsimony only relative to the coding of the features. The values of the constants along a path from the uniform law to this law, and at pairwise fits of noisy address populations, are in Table 20. The exact worst-case exponent at product laws, between $\max\{k-3, 1\}$ and k for $k \geq 3$, is open, and so is a bound of type (c) at laws with dependence among the features, where conditional expectations of low-order monomials are no longer low-order; Appendix F.8 computes the constants there.

Proof. *(a)* Fix $S \in \mathcal{S}_\tau$ and let $G_S = \{x : S_\tau(x) = S\}$. The event G_S is determined by x_S: starting at the root, follow τ on the coordinates in S; the path of x stays inside S and is padded to S exactly when $S_\tau(x) = S$. On G_S the transcript $\ell_\tau(x)$ is a function of x_S. Hence, by Jensen's inequality for the conditional expectation given the coarser information,

$$J_S(h) \;\geq\; \mathbb{E}\big[\mathbb{E}[h \mid X_S, Y]^2 \mathbf{1}_{G_S}\big] \;\geq\; \mathbb{E}\big[\mathbb{E}[h \mid \ell_\tau(X), Y]^2 \mathbf{1}_{G_S}\big],$$

and summing over $S \in \mathcal{S}_\tau$, whose events G_S partition $\mathcal{X}$, gives $\sum_S J_S(h) \geq J_\tau(h)$, which is (E.2). A tree that is not padded, or repeats a question, is refined by a padded tree, whose information J is at least as large by the same Jensen step, so an optimal branching design may be taken to consist of padded trees. If q is a design on trees that attains $W^{\mathcal{T}}_{\mathrm{branch}}(P)$, the fixed design $\sum_\tau q_\tau \,\mathrm{Unif}(\mathcal{S}_\tau)$ has $J(h) \geq J_q(h)/\max_\tau |\mathcal{S}_\tau|$ for every h, so the supremum in (E.1) grows by at most $\max_\tau |\mathcal{S}_\tau| \leq K_{m,b}$ (Appendix C.1). For r rounds, replace the forms by the r-round interviews of the proof of Theorem D.1(a), which follow τ for $r-1$ rounds and ask one of at most $\min\{2^{b-r}, \binom{m-r}{b-r}\}$ guessed completions in the last batch. For every profile exactly one completion is correct, the event that a given completion is correct is determined by that interview's transcript, on it the transcript determines ℓ_τ, and the completions are drawn uniformly, so the two inequalities above apply with these events in place of the G_S.

(b) The scores of $\mathcal{M}_{\mathrm{ci}}$ at P are the functions $h(x, y) = c(y) + \sum_j g_j(x_j, y)$ with $\mathbb{E}c(Y) = 0$ and $\mathbb{E}[g_j(X_j, Y) \mid Y] = 0$, that is, $g_j(x_j, y) = \beta_{y,j}\phi^y_j(x_j)$; these three kinds of terms are orthogonal in $L^2(P)$. Let ι be a padded interview of any class, chosen possibly as a function of y, and let $S(x)$ be the set of coordinates it queries on the profile x. A transcript is a

cylinder event $\{X_S = u\}$, and given $Y = y$ the coordinates outside S keep their conditional laws, so

$$\mathbb{E}[h \mid \ell_\iota(X), Y] = c(Y) + \sum_{j \in S(X)} g_j(X_j, Y).$$

The event that ι queries j is determined by the answers to the coordinates queried before j (in earlier batches, for an r-round interview), which are independent of X_j given Y. Hence $\mathbb{E}[g_j \mathbf{1}\{j \in S(X)\} \mid Y] = 0$ and $\mathbb{E}[g_j^2 \mathbf{1}\{j \in S(X)\} \mid Y = y] = \pi_j^\iota(y)\beta_{y,j}^2$ with $\pi_j^\iota(y) = P(j \in S(X) \mid Y = y)$. For $i \neq j$, split the event $\{i, j \in S(X)\}$ according to which of the two is queried later (in an arbitrary order if they are in the same batch), say j at a node v of the interview: the event of reaching v and the value g_i depend on coordinates other than j, so $\mathbb{E}[g_i g_j \mathbf{1}\{j \text{ is queried at } v \text{ after } i\} \mid Y] = 0$. Therefore

$$J_\iota(h) = \mathbb{E}[c(Y)^2] + \sum_y P(y) \sum_j \pi_j^\iota(y)\, \beta_{y,j}^2,$$

and for a design the same holds with $\pi_j(y) = \sum_\iota q_\iota(y)\pi_j^\iota(y)$. The projection of $t - \Delta$ on $\mathcal{T}_{\mathrm{ci}}$ has the components $c^*(y) = \mathbb{E}[t \mid Y = y] - \Delta$ and $\beta_{y,j}^* = \mathbb{E}[t\phi_j^y \mid Y = y] = \pm \hat{d}_j^y$, so by the Cauchy–Schwarz inequality in these orthogonal coordinates the supremum in (E.1) equals

$$\mathrm{Var}_P(\mathbb{E}[t \mid Y]) + \sum_y P(y) \sum_j \frac{(\hat{d}_j^y)^2}{\pi_j(y)}.$$

A design enters only through the inclusion vectors $\pi(y)$. Every padded interview asks exactly b coordinates of every record, so each $\pi(y)$ lies in the hypersimplex $H_b = \{\pi \in [0,1]^m : \sum_j \pi_j = b\}$, whose extreme points are the indicator vectors of the forms; the objective is nonincreasing in π, so the constraint $\sum_j \pi_j \le b$ of M_b may be taken with equality. Label-dependent mixtures of forms therefore realize every pair $(\pi(-1), \pi(1)) \in H_b^2$, which is all that any design of any class can realize; minimizing over each $\pi(y)$ separately gives (i). Without the label, mixtures of forms realize exactly the pairs with $\pi(-1) = \pi(1)$, which gives the formula for $W_{\mathrm{fixed}}^{\mathrm{ci}}$, and the other classes lie between it and (i) because the menus are nested. If $\pi^*(y)$ are optimal in (i), the common vector $\pi = (\pi^*(-1) + \pi^*(1))/2 \ge \pi^*(y)/2$ at most doubles every term, which gives the last inequality of (ii). The form of the minimizer of M_b follows from the Lagrange conditions of the convex program. For (iii), at a product law with a balanced label the standardized monomials ϕ_A, $A \subseteq [m+1]$, are orthonormal and $\langle t, \phi_A \rangle = 0$ unless A contains the label, because $\mathbb{E}Y = 0$; the gradient in $\mathcal{T}_2$ is supported on $\{m+1\}$ and the pairs $\{j, m+1\}$, where it equals $\mathbb{E}d$ and $\hat{d}_j$. For an interview chosen without the label, the information matrix in the basis $\{\phi_A\}$ has no entries between sets with and without the label, since $\mathbb{E}\phi_{m+1} = 0$ and ϕ_{m+1}^2 has mean one, and on the sets with the label it is the diagonal matrix with entries one and π_j^ι by the computation above. The value $v^\mathsf{T} I^\dagger v$ is then $(\mathbb{E}d)^2 + \sum_j \hat{d}_j^2/\pi_j$ for every design, and the inclusion vectors of all classes fill H_b.

(c) Let ϕ_j be the standardized jth variable and $\phi_A = \prod_{j \in A} \phi_j$; at a product law the ϕ_A, $A \subseteq [m+1]$, are orthonormal, and $\{\phi_A : 1 \le |A| \le k\}$ is a basis of $\mathcal{T}_k$. Write $h = \sum_A a_A \phi_A$. A transcript of a padded interview is a cylinder event $\{X_S = u\}$, so the coordinates outside S keep their marginal laws given the transcript and $\mathbb{E}[\phi_A \mid \ell, Y] = \phi_A \mathbf{1}\{A \subseteq S \cup \{m+1\}\}$. For a form this gives $J_S(h) = \sum_{A \subseteq S \cup \{m+1\}} a_A^2$. For the tree,

$$J_\tau(h) = \mathbb{E}\Big[\Big(\sum_{A \in \mathcal{A}(X)} a_A \phi_A\Big)^2\Big] \le \mathbb{E}\Big[\Big(\sum_{A \in \mathcal{A}(X)} \phi_A^2\Big)\Big(\sum_{A \in \mathcal{A}(X)} a_A^2\Big)\Big] \le \nu_k \sum_A a_A^2\, P\big(A \setminus \{m+1\} \subseteq S_\tau(X)\big),$$

where $\mathcal{A}(x)$ is the collection of sets $A \subseteq S_\tau(x) \cup \{m+1\}$ with $1 \le |A| \le k$, because $\phi_j^2 \le (1-\lambda)/\lambda$ and at most $\binom{b+1}{j}$ sets of size j lie in $S_\tau(X) \cup \{m+1\}$. The last sum is $\sum_S q_S^\tau J_S(h)$. Mixing over the trees of an optimal branching design gives the bound with ν_k, and (a) gives the bound with $K_{m,b}$.

(d) Here $\Delta = 0$, $t = Yd$ and $\phi_A = \chi_A$. The gradient is $v_A = \mathbb{E}[t\chi_A]$, which vanishes unless $A = B \cup \{j\} \cup \{m+1\}$ with B a set of address bits and j a data bit, and then $|v_A| = 2^{-\kappa}$ since $\hat{d}(B \cup \{j\}) = \mathbb{E}[\mathbf{1}\{\text{address} = j\}\chi_B] = \pm 2^{-\kappa}$. In $\mathcal{T}_k$ there are $D\,N(\kappa, k-2)$ such sets. By the computation in (c), the information matrix of a fixed design q in the basis χ_A is diagonal with entries $c_A(q) = P_q(A \setminus \{m+1\} \subseteq S)$, so

$$W_{\mathrm{fixed}}^{(k)}(P) = \min_q \sum_A \frac{v_A^2}{c_A(q)} \;\ge\; \min_q \frac{(\sum_A |v_A|)^2}{\sum_A c_A(q)} \;\ge\; \frac{N(\kappa, k-2)^2}{\max_s (b-s) N(s, k-2)},$$

by Cauchy–Schwarz, since a form with s address bits and $b-s$ data bits contains $(b-s)N(s,k-2)$ of the sets $B\cup\{j\}$. The maximum is at most $bN(\kappa,k-2)$, which gives $L_k(b)\ge N(\kappa,k-2)/b\ge\binom{b-1}{k-2}/b$. For $k=3$, $N(s,1)=1+s$ and $\max_s(b-s)(1+s)=\lfloor(b+1)^2/4\rfloor$; for $k=b+1$, $N(s,\kappa)=2^s$ and the maximum is 2^κ at $s=\kappa$. The tree that reads the address and the selected bit observes t, so $W^{(k)}_{\mathrm{branch}}\le W_{\mathrm{branch}}=\sigma_P^2=1$. For the upper bound at $k=3$, draw $s=\lfloor b/2\rfloor$ address bits and $b-s$ data bits uniformly. Then $c_A=(b-s)/D$ for the D sets with $B=\emptyset$ and $c_A=s(b-s)/(\kappa D)$ for the κD sets with $|B|=1$, so

$$W^{(3)}_{\mathrm{fixed}}(P)\le\frac{1}{b-s}+\frac{\kappa^2}{s(b-s)}\le\frac{2}{b}+\frac{4(b-1)^2}{b^2-1}=\frac{2}{b}+4-\frac{8}{b+1}<4,$$

using $s(b-s)\ge(b^2-1)/4$ and $b-s\ge b/2$. At $b=3$ and $b=4$ the first bound equals 2.50 and 2.75, the values of Table 3.

For the bound on branching at $k=3$ we use weak duality in (E.1): if a design q has scores $H_\iota(\ell_\iota,Y)$ with $\sum_\iota q_\iota\mathbb{E}[H_\iota h]=\langle t-\Delta,h\rangle_P$ for all $h\in\mathcal{T}$, then $\langle t-\Delta,h\rangle_P^2=(\sum_\iota q_\iota\mathbb{E}[H_\iota\mathbb{E}[h\mid\ell_\iota,Y]])^2\le\sum_\iota q_\iota\mathbb{E}[H_\iota^2]\,J_q(h)$ by Cauchy–Schwarz, so $W^{\mathcal{T}}_{\mathcal{A}}(P)\le\sum_\iota q_\iota\mathbb{E}[H_\iota^2]$. Fix $s,r\ge1$ with $s+r=b$ and $r\le2^{\kappa-s}$. With probability $1-\epsilon$, draw a set S_a of s address bits uniformly, read them, and then read a set R of r data bits drawn uniformly among the $2^{\kappa-s}$ data bits whose index agrees with the answers on S_a; the score is $cY\sum_{j\in R}Z_j/(1-\epsilon)$ with $c=\kappa/(rs)$. With probability ϵ, read a uniform set R' of b data bits; the score is $-\delta Y\sum_{j\in R'}Z_j/\epsilon$ with $\delta=(\kappa-s)/(sb)$. This is a mixture of two-round interviews. Given the profile, data bit j belongs to R with probability $r2^{s-\kappa}\mathbf{1}\{j$ agrees with the address on $S_a\}$, and averaging the indicator over S_a gives $2^{-s}\sum_B P(B\subseteq S_a)\chi_B(\text{address})\chi_B(j)$ over sets B of address bits, with $P(B\subseteq S_a)=1$ for $B=\emptyset$ and s/κ for $|B|=1$. The conditional mean of the score given (X,Y) is therefore $Y\sum_j Z_j2^{-\kappa}\sum_B w_B\chi_B(\text{address})\chi_B(j)$ with $w_\emptyset=cr-\delta b=1$ and $w_B=crs/\kappa=1$ for $|B|=1$, whereas $t=Y\sum_j Z_j2^{-\kappa}\sum_B\chi_B(\text{address})\chi_B(j)$. The two functions have the same coefficients on all sets of at most three coordinates, hence the same inner product with every $h\in\mathcal{T}_3$, and

$$W^{(3)}_{\mathrm{branch}}(P)\le\frac{c^2r}{1-\epsilon}+\frac{\delta^2b}{\epsilon},\qquad\min_\epsilon=\big(c\sqrt{r}+\delta\sqrt{b}\big)^2=\Big(\frac{\kappa}{s\sqrt{r}}+\frac{\kappa-s}{s\sqrt{b}}\Big)^2,$$

since the data bits are independent of the address and of each other. The choice $s=\kappa$, $r=1$ is the rule's own tree and gives one. With $s=\lceil3b/4\rceil$, which is admissible for large b, the two terms are $(8/3+o(1))/\sqrt{b}$ and $(1/3+o(1))/\sqrt{b}$, so $U_3(b)=(9+o(1))/b$, and $L_3(b)\to4$.

For $k=2$ the gradient is supported on the D pairs $\{j,m+1\}$ with j a data bit. Forms of b data bits drawn uniformly have $c_{\{j,m+1\}}=b/D$ and value $D\,2^{-2\kappa}D/b=1/b$, and the display gives $W^{(2)}_{\mathrm{fixed}}\ge D^2\,2^{-2\kappa}/b=1/b$. For a branching design, let I be its information matrix and I_{vv} the block of the D pairs. Then $v^{\mathsf T}I^\dagger v=\sup_u(2v^{\mathsf T}u-u^{\mathsf T}Iu)\ge\sup_{u\text{ supported on the block}}(2v^{\mathsf T}u-u^{\mathsf T}I_{vv}u)=v^{\mathsf T}I_{vv}^\dagger v$, with the value $+\infty$ if some c_j below vanishes. For data bits $j\ne l$ the entry of a tree is $\mathbb{E}[X_jX_l\mathbf{1}\{j,l\in S_\tau(X)\}]$, which is zero: on the event that l is read after j, the indicator is determined by the answers read before l, and X_l is independent of them with mean zero; the other order is symmetric. So I_{vv} is diagonal with entries $c_j=P(j\text{ is read})$, $\sum_j c_j\le b$, and $v^{\mathsf T}I_{vv}^\dagger v=\sum_j 2^{-2\kappa}/c_j\ge1/b$. Mixtures of trees have the same diagonal block structure by linearity.

(c), stability. Let $k\le b$, let $h_a=\sum_A a_A(\chi_A-\mathbb{E}_P\chi_A)$ for a coefficient vector a of Euclidean norm one over the sets with $1\le|A|\le k$, and write J^P for information at the law P. The menus of forms and of padded depth-b trees are finite, and for full-support laws $(P,a)\mapsto J^P_\iota(h_a)$ is continuous, so $\omega(\delta)=\sup\{|J^P_\iota(h_a)-J^{P_0}_\iota(h_a)|:\|P-P_0\|\le\delta,\ \|a\|=1,\ \iota\}$ tends to zero with δ. Every set A with $|A|\le k\le b$ lies in $S\cup\{m+1\}$ for some form S, so the uniform design u on all forms has $c_0=\min_{\|a\|=1}J^{P_0}_u(h_a)>0$. For a tree τ and $\eta\in(0,1)$ put $q^{\tau,\eta}=(1-\eta)q^\tau+\eta u$ with q^τ from (c) at P_0. Then $J^{P_0}_{q^{\tau,\eta}}(h_a)\ge\max\{(1-\eta)J^{P_0}_\tau(h_a)/\nu_k,\ \eta c_0\}$, and for $\|P-P_0\|\le\delta$ with $\omega(\delta)\le\eta c_0/2$,

$$J^P_\tau(h_a)\le J^{P_0}_\tau(h_a)+\omega\le\frac{\nu_k}{1-\eta}\big(J^P_{q^{\tau,\eta}}(h_a)+\omega\big)+\omega\le\frac{\nu_k}{1-\eta}\Big(1+\frac{4\omega}{\eta c_0}\Big)J^P_{q^{\tau,\eta}}(h_a),$$

using $J^P_{q^{\tau,\eta}}(h_a)\ge\eta c_0/2$ and $\nu_k\ge1$. The tangent space $\mathcal{T}_k$ at P consists of the multiples of the h_a, so the fixed design $\sum_\tau q_\tau q^{\tau,\eta}$ built from an optimal branching design q at P has information at least $(1-\eta)(1+4\omega/(\eta c_0))^{-1}/\nu_k$ times that of q in every direction of $\mathcal{T}_k$. Choosing η and then δ small gives the factor $(1+\epsilon)\nu_k$. The radius δ depends on m, b, k, λ and ϵ and is not explicit.

(e) Write the address as $a \in \{-1,1\}^\kappa$, identify a data index j with its sign vector, and let $\langle a, j\rangle \in \{-\kappa, \dots, \kappa\}$ be their inner product, which equals κ exactly when j is the index selected by a. With $z_j = (1+Z_j)/2$ define the quadratic function

$$\mathcal{Q}(a,s,z,p) = \sum_j s_j\langle a,j\rangle - (\kappa+1)\Big(\sum_j s_j - 1\Big)^2 - \sum_j \big(3p_j + s_j z_j - 2p_j s_j - 2p_j z_j\big),$$

and let P_β be the law with $P_\beta(x) \propto e^{\beta\mathcal{Q}(x)}$ and an independent uniform label. $\mathcal{Q}$ is a polynomial of degree two in the binary features, so $P_\beta \in \mathcal{M}_2$ for every $\beta \geq 0$, with full support. The last sum is the standard penalty for a product of two binary variables (Boros & Hammer, 2002): each term is nonnegative and vanishes exactly when $p_j = s_j z_j$. If exactly one s_j equals one, the first two terms equal $\langle a, j\rangle \leq \kappa$, with equality exactly when j is the selected index; if none does they equal $-(\kappa+1)$; and if $n \geq 2$ do, they are at most $n\kappa - (\kappa+1)(n-1)^2 < \kappa$. So $\mathcal{Q} \leq \kappa$ with equality exactly on the set where s is the one-hot code of the address and $p_j = s_j z_j$ for all j, and as $\beta \to \infty$ the law P_β converges to the law P_∞ under which the address, the data bits and the label are independent and uniform and s and p are these functions of them.

On the support of P_∞, $d = Z_a = \sum_j s_j(2z_j - 1) = 2\sum_j p_j - 1$, so $t = h_0$ almost surely for the polynomial $h_0 = Y(2\sum_j p_j - 1)$ of degree two in (p, Y), whose centered version lies in $\mathcal{T}_2 \subseteq \mathcal{T}_k$ at every P_β. By (E.1),

$$W^{(k)}_{\text{fixed}}(P_\beta) \;\geq\; \frac{\mathrm{Cov}_{P_\beta}(t,h_0)^2}{\max_S J_S^{P_\beta}(h_0 - \mathbb{E}_{P_\beta}h_0)}.$$

For a bounded function h and an interview ι, $J^P_\iota(h) = \sum_c \mathbb{E}_P[h\mathbf{1}_c]^2/P(c)$ over the transcript cells, and a cell contributes at most $\|h\|^2_\infty P(c)$, so $J_S^{P_\beta} \to J_S^{P_\infty}$ and the right-hand side converges to $1/\max_S J_S^{P_\infty}(t)$, because $\mathrm{Var}_{P_\infty}(t) = 1$. Fix a form S of b of the m features, with data bits R_z and product bits R_p. Under P_∞ and given that the selected index is j, the features in S other than Z_j and p_j are functions of the address and of data bits independent of Z_j, while $p_j = z_j$; hence the conditional mean of Z_j given X_S and the selected index j is Z_j or $2p_j - 1$ if $j \in R_z \cup R_p$ and zero otherwise. Therefore $|\mathbb{E}[d \mid X_S]| \leq P(\text{selected index} \in R_z \cup R_p \mid X_S)$ and

$$J_S^{P_\infty}(t) = \mathbb{E}\big[\mathbb{E}[d \mid X_S]^2\big] \leq P(\text{selected index} \in R_z \cup R_p) \leq \frac{b}{D}.$$

So $\liminf_\beta W^{(k)}_{\text{fixed}}(P_\beta) \geq D/b$. The tree that reads the address and the selected data bit computes d at every profile, so $W^{(k)}_{\text{branch}}(P_\beta) \leq W_{\text{branch}}(P_\beta) = \mathrm{Var}_{P_\beta}(t) = 1$, because $t = \pm 1$ and the independent uniform label gives $\Delta(P_\beta) = 0$. Taking β large gives the claim, and P_β puts mass tending to one on the support of P_∞. For $b = 2$ the bound D/b equals one. □

Computed constants for the address contrast. Table 3 lists the constants of Theorem E.5(d) for $b = 3$ and $b = 4$, computed by the model version of the solver (Appendix F.8). Under the three-way model the gap is 2.50 and 2.75 against the lower bounds $L_3(3) = 2.25$ and $L_3(4) = 2.67$, and at these caps the optimal branching design is the rule's own tree ($U_3 = 1$ for $b \leq 5$); under the pairwise model both classes need $1/b$ of the records that the design-based branching optimum needs.

Table 3. The address contrast at the uniform law within the order-k log-linear model: constants relative to $\sigma_P^2 = 1$, computed by the solver (certified gaps below 10^{-5}). Complete: bound for complete records. $L_k(b)$: the lower bound of Theorem E.5(d) on the fixed constant. Without a model the ratio is 2^{b-1}.

b	m	k	Complete	Fixed	Branch	Ratio	$L_k(b)$
3	6	2	0.250	0.333	0.333	1.00	0.333
3	6	3	0.750	2.500	1.000	2.50	2.250
3	6	4	1.000	4.000	1.000	4.00	4.000
4	11	2	0.125	0.250	0.250	1.00	0.250
4	11	3	0.500	2.750	1.000	2.75	2.667

Contrasts that branching cannot compute. For an ensemble whose contrast is not computed by one depth-b tree, part (b) of Theorem E.1 applies to both menus. Every depth-b branching interview is a coarsening of a complete depth-b tree,

and the energy of a tree is a sum over its leaves. Hence $\max_\iota \mathbb{E}_P[\mathbb{E}_P[h \mid \ell_\iota, Y]^2]$ over all depth-b trees is computed exactly by dynamic programming over partial assignments, in time $O(m^b 2^b)$. This yields certified lower bounds on $M_{\mathrm{branch}}(P)$ for every dual witness h. Appendix F.2 uses these bounds inside column generation to compute the branching coefficient over all depth-three trees.

E.3. A Computable Upper End: the Optimized Path-Set Design

The upper end $N_P(\tau)$ of Theorem 3 is the variance of centered completion-only path guessing, which discards every answer on a form except a completed path. Optimizing the full score program on the same small menu tightens it without enumerating the $\binom{m}{b}$ forms.

Proposition E.6 (Optimized path-set design)**.** *Let a depth-b tree τ compute d, let $S_1, \dots, S_L$ be its distinct padded path sets ($L \le K_{m,b}$), and let $W_{\mathrm{PS}}(P)$ be the optimum of the population program (3) over the fixed-form menu $\{S_1, \dots, S_L\}$, with all pointwise-unbiased tables. Then*

$$W_{\mathrm{fixed}}(P) \;\le\; W_{\mathrm{PS}}(P) \;\le\; N_P(\tau)\, W_{\mathrm{branch}}(P).$$

Proof. The menu $\{S_1, \dots, S_L\}$ is a set of fixed forms, so its optimum is at least the optimum over all forms. The menu also identifies d: every leaf indicator of τ depends only on that leaf's path coordinates, which lie in its padded set, so d is a sum of tables on the S_j. Centered path guessing with $q_j \propto v_j^{1/2}$ is a feasible design in this program, because the constant $c = \Delta(P)$ is a table on every form, and its variance is $N_P(\tau)\sigma_P^2$ by the proof of Theorem E.1(c), where $W_{\mathrm{branch}}(P) = \sigma_P^2$. □

The program has at most $2L2^b$ score variables and is solved from reference marginals; Appendix F.3 compares W_{PS} with the all-forms optimum on every screening run. The two coincide whenever an optimal fixed design uses only path sets of τ; in general the gap measures the value of forms that no path of the tree reads.

E.4. Computing Optimal Designs for Every Protocol Class

All population constants in Section 6 are computed by one column-generation solver whose cost depends on $\binom{m}{b}2^b$, not on the 2^m profiles.

Cells and masses. Pad every transcript of an interview to exactly b coordinates. An interview of $\mathcal{A}_r$ is then a partition of the cube into cells $C(T,u) = \{x : x_T = u\}$ with $|T| = b$; for a fixed form all cells share T, for an r-round interview T varies across cells. Padding refines a transcript and therefore never increases the optimal second moment. For a law P that mixes records with a uniform mass α over profiles and labels, the masses $P(T,u,z)$ of all $\binom{m}{b}2^{b+1}$ cells are computed once from the records.

Master problem. For a list of interviews, minimize $\sum_{\iota,c,z} P(c,z) g_{\iota,z}(c)^2/q_\iota$ subject to $\sum_\iota q_\iota = 1$ and the Fourier identities $\sum_{\iota,c} \widehat{\mathbf{1}}_c(S) g_{\iota,z}(c) = (2z-1)\widehat{d}(S)$ for all $|S| \le b$, where $\widehat{\mathbf{1}}_{C(T,u)}(S) = 2^{-b}\chi_S(u)$ for $S \subseteq T$ and zero otherwise. Because cell indicators and d have degree at most b, the identities are equivalent to pointwise unbiasedness at every profile. The problem is a second-order cone program; we solve it with Clarabel through CVXPY at tolerance 10^{-10}. Inside column generation the solver's solution steers the iteration; the final design is repaired to exact feasibility by the least-norm correction weighted by the interview probabilities, which adds the least second moment, so every reported upper bound is the second moment of an exactly pointwise-unbiased design. Looser solver tolerances produced visibly suboptimal masters in our tests, which the consistency checks of Appendix F (the fixed optimum may not exceed the path-set or fitted designs) detect.

Pricing. Let $\eta_z(S)$ be the multipliers of the identities. The information of a cell is

$$e(T,u) = \sum_z \frac{\psi_z(T,u)^2}{P(T,u,z)}, \qquad \psi_z(T,u) = 2^{-b}\sum_{S \subseteq T} \eta_z(S)\chi_S(u),$$

a Walsh transform of η restricted to T; all cells cost $O(\binom{m}{b} b 2^b)$ operations. The information of an interview is the sum over its cells. The most informative interview in $\mathcal{A}_r$ is found by dynamic programming over partial assignments (A,a), $|A| < b$:

$$V_1(A,a) = \max_{T \supseteq A, |T|=b} \sum_{u:\, u_A = a} e(T,u), \qquad V_\rho(A,a) = \max_{B \supsetneq A,\, |B| \le b} \sum_{a' :\, a'_A = a} V_{\rho-1}(B,a'),$$

with $V_\rho(T,u) = e(T,u)$ for $|T| = b$; the best r-round interview has information $V_r(\emptyset,\emptyset)$, the best tree is the special case of singleton batches, and the best form is $\max_T \sum_u e(T,u)$. The program visits $O(\sum_{k\le b}\binom{m}{k}4^k)$ entries, and an optimal interview is reconstructed by following the maximizers.

Certificates and stabilization. Normalize η so that $\sum_{z,S}\eta_z(S)(2z-1)\widehat{d}(S) = 1$. By Theorem E.1(b) with $c = 0$, $1/\max_{\iota\in\mathcal{A}_r} e(\iota)$ is a lower bound on $M_{\mathcal{A}_r}(P) = W_{\mathcal{A}_r}(P) + \Delta(P)^2$ for every such η, and the master value is an upper bound. Every pointwise-unbiased design also satisfies $M_{\mathcal{A}}(P) \ge \mathbb{E}_P[t^2]$ by Jensen's inequality, which certifies designs that attain full information. Each iteration adds up to six (twelve for the forests and the 21-item study) interviews whose information exceeds the master's maximum, and the solver stops when no interview of the class improves the master or when the bounds agree to a relative error of 10^{-7} (10^{-4} for the forest tasks at $b \ge 4$). Master duals of column generation are degenerate, and unstabilized pricing can stall for many iterations; we price at a convex combination, with equal weights, of the current dual and the dual with the best dual bound so far (Wentges, 1997), and fall back to the current dual when the smoothed point finds no improving interview. Removing unused columns from the master made the iteration cycle in our tests, so the master keeps every generated interview. The master starts from the support of the optimal fixed design (and, for the forests, the padded component trees), so it stays small.

Validation. On stored tasks the solver reproduces the fixed constants, the explained energy of the best form and the certified branching constants of an independent dense implementation (released with the code) to six significant digits, and every generated interview is checked to partition the cube. Branching and r-round designs at $m \le 12$ converge in tens to a few hundred iterations; pricing is no longer the bottleneck, and the master problem dominates the running time (Appendix F.2).

E.5. The Form Program and the Path-Set Menu

A form can reveal useful coordinates even when it fails to complete a tree path. For a frozen depth-b tree, take the coordinate set on each root-to-leaf path, extend it to size b with fixed unused coordinates, and retain the L distinct sets as a tractable menu. Let $g_{T,y}(u)$ be a table on *all* answers u to form T, and let $\widehat{P}_T(u,y)$ be calibrated marginal frequencies. For binary features solve

$$\begin{aligned} \min_{q,g} \quad & \sum_{T,u,y} \widehat{P}_T(u,y)\frac{g_{T,y}(u)^2}{q_T}, \\ \text{subject to} \quad & \sum_{T\supseteq S} \widehat{g}_{T,y}(S) = y\widehat{d}(S) \quad \text{for every } S, y, \\ & \sum_T q_T = 1, \qquad q_T \ge \beta/L. \end{aligned} \tag{E.3}$$

Here $\widehat{g}_{T,y}(S)$ is the Fourier coefficient of the table on its own form; it is a linear transform of g. Fitted designs in Section 6 use $\beta = 0.05$. These constraints are exactly equivalent to $\sum_T g_{T,y}(x_T) = y d(x)$. Returning $g_{T,Y}(X_T)/q_T$ is therefore globally unbiased even when the reference frequencies are inaccurate; the reference affects only the variance.

There are at most $2L2^b$ score entries and coefficient constraints, and the program avoids enumerating the 2^m full profiles. Pairwise shared-coordinate corrections span the complete unbiased score space on the supplied menu over the full binary cube (Proposition E.7). A correction adds $f(X_{T\cap T'}, Y)/q_T$ on form T and subtracts it divided by $q_{T'}$ on form T'. The optimization therefore includes every such control variate. In the language of coarsening at random, (E.3) chooses the optimal augmentation of the inverse-probability-weighted score under the reference law, jointly with the form probabilities. For ordinal data the same program applies to tensor contrasts on the categories defined by the frozen tree's thresholds.

For an existing randomized archive, keep its known q fixed and optimize only the tables. Form-specific marginal frequencies can be estimated from incomplete training records; they need not be assembled into a joint reference distribution. Identification then requires coverage by the actual form menu, not only the degree bound for the menu of all size-b forms.

A *completion-only* score returns an inverse-probability weighted target on a matching padded path and zero otherwise; the experiments call its uniform version path guessing.

Proper scoring rules. For Brier improvement, with $z = (Y+1)/2$,

$$(p_0 - z)^2 - (p_1 - z)^2 = d(1 - p_0 - p_1) + Yd.$$

In the form program, replace Yd by this known target and impose separate coefficient identities for each label. The same unbiasedness guarantee holds. For ensembles of degree r, Brier loss can require degree $2r$; log loss can introduce still higher interactions. Appendix G gives the identification conditions and explains why AUC requires a separate pairwise design.

E.6. Shared-Coordinate Corrections and the Complete Score Space

Fix a menu of coordinate sets $\mathcal{T}$. For binary features write $g_{T,y}(x_T) = \sum_{S \subseteq T} \gamma_{T,y,S} \chi_S(x)$. Uniqueness of the Fourier expansion gives the following elementary representation.

Proposition E.7 (All shared-coordinate corrections)**.** *The score tables satisfy $\sum_T g_{T,y}(x_T) = yd(x)$ on the full binary cube if and only if*

$$\sum_{T \supseteq S} \gamma_{T,y,S} = y\widehat{d}(S) \quad \textit{for every } S.$$

For a fixed positive design q, every change to a globally unbiased score is a sum of pairwise transfers of functions on shared coordinates: add $f(X_{T \cap T'}, Y)/q_T$ to form T and subtract the same function divided by $q_{T'}$ from form T'.

Proof. Equate the coefficients of each χ_S. For the second claim, any score difference has, for each S and each label, coefficients summing to zero over forms containing S. Choose one of those forms as an anchor and transfer every other coefficient to or from the anchor. Each transfer is observable under both forms. Terms appearing in only one form must have zero difference. Summing the transfers proves the statement. □

Thus including all such terms uses the entire unbiased score space on the supplied menu. This is a product-basis implementation of standard control variates and known-coarsening influence-function optimization, not a new general missing-data efficiency bound.

The implementation works in cell values. A block of the constraint matrix is the normalized Walsh transform of a form's table, placed in the rows of its contained coordinate subsets. If F is the resulting matrix, the constraints are $Fg_y = y\widehat{d}$. Its rows are orthogonal: FF^{T} is diagonal, with entry $\sum_{T \supseteq S} 2^{-|T|}$ for subset S. We use this identity to repair numerical equality residuals explicitly. Rotated second-order cones represent $g_{T,y}(u)^2 \le q_T v_{T,y}(u)$.

For a given population, marginal form–label frequencies are sufficient for the objective. No full-profile enumeration is needed. With L forms of size at most b, there are at most $2L2^b$ score entries and $2L2^b$ coefficient constraints. A tree provides the target coefficients through its leaves, or by Fourier transforming its known completion-score decomposition. Choosing an unrestricted menu can still require $\binom{m}{b}$ forms; this implementation does not solve that combinatorial design problem.

For ordinal features one splits each coordinate at all thresholds used for it in the frozen tree, with a fixed middle cut for otherwise unused padding coordinates. We use orthonormal Helmert contrasts on those categories and their tensor products. The same argument is exact on the resulting product space. This retains every observed tree-threshold category, but not within-category distinctions between raw ordinal answers. Its cell count is $\sum_T \prod_{j \in T} s_j$, where s_j is the category count.

E.7. Efficiency and Augmented Fixed Forms

E.7.1. Relation to the Missing-Data Efficiency Bound

Fix a known form law q and a full-support law P on the finite complete-record space. The observed record is $O = (A, o_A(Z))$. Its score along a centered complete-data perturbation h is $\mathbb{E}_P[h(Z) \mid O]$. A mean-zero observed influence function $\phi(O)$ for $\Delta = \mathbb{E}_P t$ must therefore satisfy

$$\mathbb{E}[\phi(O) \mid Z] = t(Z) - \Delta.$$

Conversely this equation gives the required pairing with every complete-data score, and hence an influence function in this finite regular model. Minimizing $\mathbb{E}\phi^2$ is the efficiency bound. Substituting $\phi = H - \Delta$ identifies it with the fixed-design population variance $W_q(P)$. This is the usual known-coarsening influence-function geometry (Robins et al., 1994; Tsiatis, 2006), not a new asymptotic efficiency formula. The proposition's additional statement is finite-n pointwise optimality over

globally unbiased nonlinear estimators and sequential form assignments, together with its extension to boundary laws. It does not assert a finite-sample bound for all locally regular biased estimators.

Fixed random forms implement planned missingness. A tree's terminal cell is generated using observed answers, so its known acquisition mechanism is a designed coarsening-at-random mechanism (Heitjan & Rubin, 1991; Gill et al., 1997). This description does not cover arbitrary unrecorded selection or nonresponse.

E.8. Learning a Design from a Compact Tree

Fix a tree that recovers $t(z) = yd(x)$ within b distinct queries. Pad each path deterministically to b coordinates. Enumerate its distinct path sets $\mathcal{S} = \{S_1, \dots, S_L\}$ and let $S(z)$ denote the reached set. If an assigned fixed form S_j permits following the padded tree to completion, then $S(z) = S_j$, as in Appendix C.1. For any strictly positive form law q, the score

$$H_q(z, J) = \frac{t(z)}{q_J} \mathbf{1}\{S(z) = S_J\}, \qquad J \sim q,$$

is computable from the selected form and label. It is unbiased conditional on every full record. With $w_j = \mathbb{E}_P[t(Z)^2 \mathbf{1}\{S(Z) = S_j\}]$,

$$\operatorname{Var}_P H_q = \sum_{j=1}^{L} \frac{w_j}{q_j} - \Delta(P)^2. \tag{E.4}$$

Within this importance-score family the unconstrained optimum is $q_j \propto \sqrt{w_j}$, with infimal second moment $(\sum_j \sqrt{w_j})^2$. Zero-mass cells are handled by limits. We enforce $q_j \geq \beta/L$, ensuring global unbiasedness even on path sets absent from calibration. If some $w_j > 0$, the constrained optimum satisfies $q_j = \max\{\beta/L, c\sqrt{w_j}\}$, with c chosen to normalize the probabilities. If all weights vanish, every feasible q is optimal. Iteratively fixing cells below the floor computes the positive-weight solution without profile enumeration. This oracle optimizes a specified score family, not all fixed-form estimators.

E.8.1. A Worked Two-Question Example

Let $d(x) = [(1 + x_1)x_2 + (1 - x_1)x_3]/2$. Branching queries x_1 and then x_2 when $x_1 = 1$, or x_3 when $x_1 = -1$. The two fixed forms are

Form	Probability	Observed score
$\{1, 2\}$	$1/2$	$2Yx_2\mathbf{1}\{x_1 = 1\}$
$\{1, 3\}$	$1/2$	$2Yx_3\mathbf{1}\{x_1 = -1\}$

At every full record the mean score is Yd and its second moment is two. The branching score has second moment one. For an independent fair label both score means vanish, so their sample-mean variances are exactly $2/n$ and $1/n$. This describes these concrete estimators; the minimax results additionally justify their worst-case comparison against all protocols.

E.8.2. A Finite Calibration Guarantee

Suppose the tree is fitted independently of r calibration records and $|t| \leq M$. Full calibration records reveal $W_j = t^2 \mathbf{1}\{S = S_j\}$ directly. Alternatively, assign each calibration person a uniformly random fixed path set. On successful completion return $W_j = Lt^2 \mathbf{1}\{J = j, S = S_j\}$ and return zero on failure. This alternative obeys the same fixed per-person cap and uses only observable quantities. In both cases $\mathbb{E}W = w$. Let $\widehat{w}$ be the mean and let $\widehat{q}$ minimize $\sum_j \widehat{w}_j/q_j$ subject to $q_j \geq \beta/L$.

Proposition E.8 (Learning path-set probabilities)**.** *Let $q^\star_\beta$ be the population optimum with the same probability floor. Set $B = M^2$ for full calibration or $B = LM^2$ for uniformly assigned fixed calibration. With probability at least $1 - \alpha$ over the r independent calibration records,*

$$\operatorname{Var}_P H_{\widehat{q}} - \operatorname{Var}_P H_{q^\star_\beta} \leq \frac{2LB}{\beta} \sqrt{\frac{2(L \log 2 + \log(1/\alpha))}{r}}.$$

The resulting audit mean on fresh respondents remains globally unbiased for every realized calibration sample.

Proof. Each calibration vector has at most one nonzero entry and $\|W\|_1 \le B$. For any sign vector $\sigma \in \{-1,1\}^L$, $\sigma^T W \in [-B, B]$. Hoeffding's inequality and a union bound over 2^L vectors give

$$\|\widehat{w} - w\|_1 = \max_{\sigma} \sigma^T(\widehat{w} - w) \le B\sqrt{2(L\log 2 + \log(1/\alpha))/r}.$$

The difference between the empirical and population second-moment objective is uniformly at most $(L/\beta)\|\widehat{w} - w\|_1$. Empirical minimization costs at most twice this uniform deviation. Subtracting the common Δ^2 proves the bound. Pointwise conditional unbiasedness holds for each positive q, hence also conditional on the entire calibration sample. □

The bound is conservative and becomes informative only with sufficiently large calibration samples. It proves consistency relative to a constrained family oracle, not a calibration-free optimality claim. Form enumeration takes time polynomial in the given tree's size and m; that tree may itself have exponentially many leaves in b. A deployment needs only its form probabilities and the tree, not a table over all s^m profiles.

E.8.3. Confidence Intervals and Costs

Conditional on calibration, each path score has magnitude at most $M/q_{\min}$. Thus the ordinary score mean has a distribution-free $(1-\alpha)$ confidence interval with half-width

$$\frac{M}{q_{\min}}\sqrt{\frac{2\log(2/\alpha)}{n}},$$

intersected with $[-M, M]$. The branching score has the same bound with $q_{\min} = 1$. These Hoeffding intervals are valid but conservative; they are not claims of optimal interval width or power. The paper's risk comparisons concern MSE and do not order intervals for biased estimators.

For a menu with known per-person costs $c_\iota > 0$, let $c(q) = \sum_\iota q_\iota c_\iota$. An iid unbiased score mean with n respondents uses expected cost $nc(q)$, so its worst-case MSE-times-cost upper criterion is $c(q)\max_x \sum_\iota g_\iota(x)^2/q_\iota$. Set $r_\iota = q_\iota/c(q)$. Joint cost-weighted design becomes the convex program

$$U_c(d, \mathcal{A}) = \inf_{\substack{r,g:\ \sum_\iota c_\iota r_\iota = 1 \\ \sum_\iota g_\iota = d}} \max_x \sum_\iota \frac{g_\iota(x)^2}{r_\iota}.$$

Conversely $q_\iota = r_\iota / \sum_j r_j$ recovers the design, proving equivalence of the two criteria. For fixed forms, c_ι may include recruitment, the label and item-specific costs. For a branching tree, taking c_ι as its maximum path cost gives a conservative criterion. This uses expected total expenditure and continuous sample-size scaling; it does not impose an almost-sure stopping budget or establish a sharp adaptivity ratio for arbitrary costs.

E.9. Learning the Design from the Audit Itself

The population-specific constants $W_{\mathcal{A}}(P)$ are attained by designs tuned to P. Section 6 tunes them on separate reference records. No reference is needed asymptotically: because every cell mass of every interview is estimable from the records assigned that interview, a pilot phase of the audit itself can supply the design, and pointwise unbiasedness keeps every intermediate design exactly unbiased.

For a finite menu $\mathcal{A}$, a law P on $\mathcal{X} \times \{-1,1\}$ and a design (q, g) feasible for (3), write $J_P(q,g) = \sum_\iota \sum_{c,y} P_\iota(c,y) g_\iota(c,y)^2/q_\iota$ for its second moment, where $P_\iota(c,y) = P(\ell_\iota(X) = c, Y = y)$ are the cell masses, so that $M_{\mathcal{A}}(P) = \inf_{q,g} J_P(q,g)$. For $\beta \in (0,1]$ let $M_{\mathcal{A}}^{\beta}(P)$ denote the infimum over designs with $q_\iota \ge \beta/|\mathcal{A}|$ for all ι.

Proposition E.9 (Two-phase self-calibration)**.** *Let $n_1 = n_1(n) \to \infty$ with $n_1/n \to 0$, and let $\alpha_n, \beta_n \in (0,1]$ tend to zero with $\alpha_n n_1^{1/2} \to \infty$. Phase one interviews n_1 records with the uniform design on $\mathcal{A}$ and forms, for every interview ι, the Horvitz–Thompson estimates $\widehat{P}_\iota(c,y) = (|\mathcal{A}|/n_1)\sum_{i \le n_1} \mathbf{1}\{\iota_i = \iota,\ \ell_\iota(X_i) = c,\ Y_i = y\}$ and the smoothed masses $\widetilde{P}_\iota = (1-\alpha_n)\widehat{P}_\iota + \alpha_n u_\iota$, where u_ι is uniform on the cells and labels of ι. Let $(\hat{q}, \hat{g})$ minimize $J_{\widetilde{P}}$ over feasible designs with $q \ge \beta_n/|\mathcal{A}|$, and interview the remaining $n_2 = n - n_1$ records with $(\hat{q}, \hat{g})$. Then the mean $\widehat{\Delta}$ of their scores is unbiased for $\Delta(P)$ conditionally on phase one, for every P, and*

$$n_2 \operatorname{Var}(\widehat{\Delta} \mid \textit{phase one}) = J_P(\hat{q}, \hat{g}) - \Delta(P)^2, \qquad J_P(\hat{q}, \hat{g}) \xrightarrow{\mathbb{P}} M_{\mathcal{A}}(P), \qquad n \operatorname{Var}(\widehat{\Delta} \mid \textit{phase one}) \xrightarrow{\mathbb{P}} W_{\mathcal{A}}(P).$$

Proof. Unbiasedness holds because $(\hat{q}, \hat{g})$ satisfies the pointwise constraints $\sum_\iota \hat{g}_\iota(\ell_\iota(x), y) = t(x, y)$ for every (x, y), whatever the phase-one data. Conditionally on phase one the phase-two scores are iid with mean $\Delta(P)$ and second moment $J_P(\hat{q}, \hat{g})$, which gives the variance identity. For the limit put $P^\alpha_\iota = (1 - \alpha_n)P_\iota + \alpha_n u_\iota$. Each $\widehat{P}_\iota(c, y)$ is an average of n_1 iid bounded terms with mean $P_\iota(c, y)$, so $\max_{\iota,c,y} |\widehat{P}_\iota(c, y) - P_\iota(c, y)| = O_{\mathbb{P}}(n_1^{-1/2})$ over the finitely many cells. Since $\widetilde{P}_\iota - P^\alpha_\iota = (1 - \alpha_n)(\widehat{P}_\iota - P_\iota)$ and $P^\alpha_\iota \geq \alpha_n u_\iota$,

$$\rho_n := \max_{\iota,c,y} \Big| \frac{\widetilde{P}_\iota(c, y)}{P^\alpha_\iota(c, y)} - 1 \Big| \leq \frac{\max_{\iota,c,y} |\widehat{P}_\iota(c, y) - P_\iota(c, y)|}{\alpha_n \min_\iota u_\iota} = O_{\mathbb{P}}\big((\alpha_n n_1^{1/2})^{-1}\big) \to 0.$$

Let (q^α, g^α) be optimal for J_{P^α} under the floor β_n. Because J is linear in the masses,

$$(1 - \alpha_n) J_P(\hat{q}, \hat{g}) \leq J_{P^\alpha}(\hat{q}, \hat{g}) \leq \frac{J_{\widetilde{P}}(\hat{q}, \hat{g})}{1 - \rho_n} \leq \frac{J_{\widetilde{P}}(q^\alpha, g^\alpha)}{1 - \rho_n} \leq \frac{1 + \rho_n}{1 - \rho_n} M^{\beta_n}_{\mathcal{A}}(P^\alpha).$$

Finally $M^\beta_{\mathcal{A}}(P^\alpha) \to M_{\mathcal{A}}(P)$ as $\alpha, \beta \to 0$: for any feasible (q, g), mixing q with the uniform design at weight β keeps the tables feasible and multiplies J by at most $(1 - \beta)^{-1}$, so $M^\beta_{\mathcal{A}}(P^\alpha) \leq (1 - \beta)^{-1} J_{P^\alpha}(q, g) \leq (1 - \beta)^{-1}[(1 - \alpha)J_P(q, g) + \alpha J_u(q, g)]$, whose infimum over (q, g) tends to $M_{\mathcal{A}}(P)$; conversely $M^\beta_{\mathcal{A}}(P^\alpha) \geq (1 - \alpha)M_{\mathcal{A}}(P)$. Combining the displays gives $\limsup J_P(\hat{q}, \hat{g}) \leq M_{\mathcal{A}}(P)$ in probability, and $J_P(\hat{q}, \hat{g}) \geq M_{\mathcal{A}}(P)$ holds for every feasible design. Since $n_2/n \to 1$, the last claim follows from $W_{\mathcal{A}}(P) = M_{\mathcal{A}}(P) - \Delta(P)^2$. □

The proposition is an asymptotic statement about a finite menu: phase one must estimate the masses of all $\sum_\iota |\mathrm{cells}(\iota)|$ cells, which requires n_1 large compared with $|\mathcal{A}|2^b$, and in the experiments of Section 6 reference records play the role of phase one. A sequential version that refits the design after every batch of records inherits exact unbiasedness by the same argument; its rate is not analyzed here. Learning the design across records is standard adaptive importance sampling; the point here is that the pointwise constraints make every fitted design exactly unbiased, so that the price of learning is paid in variance only.

F. Experimental Details

Every experiment of Section 6 is described here.

F.1. Data, Populations and Software

Data. WDBC (569 records, 30 continuous features) comes from scikit-learn (Wolberg et al., 1993); Spambase (4,601 records) and Ionosphere (351 records) from the PMLB mirror of the UCI files (Hopkins et al., 1999; Sigillito et al., 1989; Romano et al., 2022); the CDC Diabetes Health Indicators (253,680 respondents of the 2015 Behavioral Risk Factor Surveillance System) from a public mirror of the Kaggle release referenced by the UCI entry (Centers for Disease Control and Prevention, 2017); and the thyroid files (7,200 patients) from public mirrors of the UCI release (Quinlan, 1986). Each mirror was checked by reproducing values computed from the original downloads to nine decimals, which also confirms the record order.

Populations. Every population P is an empirical law of held-out or target records mixed with a uniform mass α over all profiles and both labels, $\alpha = 1\%$ unless stated. Full support makes $W_{\mathcal{A}}(P)/n$ a local minimax risk (Proposition C.10) and keeps every cell mass positive. Oracle designs are optimized at P; fitted designs are optimized on a reference law (training or source records, smoothed with $\alpha = 10\%$) with a floor of $5\%/L$ on each of the L interview probabilities of the final master problem, and evaluated at P. Every design is pointwise unbiased, so oracle and fitted designs differ only in variance.

Software. Python 3.13.5, NumPy 2.5.4, SciPy 1.18.1, scikit-learn 1.9.1, pandas 3.1.1, CVXPY 1.9.3 with Clarabel 0.11.1, single-threaded. All constants are computed by the solver of Appendix E.4; it reproduces the values of an independent dense implementation on every overlapping task.

F.2. Forest Updates

Protocol. For each dataset and seed, a random permutation assigns n_{old} records to train the old forest, the next n_{new} to train the new forest, and the remaining n_{eval} records to the evaluation law: $(n_{\mathrm{old}}, n_{\mathrm{new}}, n_{\mathrm{eval}}) = (100, 200, 269)$ for

Table 4. Independent forest updates, cap $b = 3$, 20 splits per cell. Exact fixed/branching variance ratio over all forms and all depth-3 trees (median and 10th–90th percentiles). Restr.: branching restricted to the 40 component trees and all triples. Fix/full: fixed-form variance relative to the full-information variance. Fitted: designs optimized on training records, evaluated held-out ($m = 12$, five splits).

Data	m	Exact ratio	Restr.	Fix/full	Fitted
WDBC	6	1.36 [1.27, 1.48]	1.08	2.11	
	9	1.57 [1.43, 1.69]	1.09	3.91	
	12	1.66 [1.53, 1.74]	1.08	5.55	1.67
Spambase	6	1.41 [1.27, 1.52]	1.03	2.31	
	9	1.45 [1.36, 1.59]	1.03	3.09	
	12	1.58 [1.49, 1.66]	1.04	4.75	1.60
Ionosphere	6	1.36 [1.20, 1.46]	1.07	2.06	
	9	1.61 [1.45, 1.77]	1.10	4.19	
	12	1.73 [1.58, 1.78]	1.10	5.94	1.79

Table 5. Forest updates of depth b under cap b (WDBC, Spambase, Ionosphere; binding tasks). Fixed/branch: exact $W_{\mathrm{fixed}}/W_{\mathrm{branch}}$, median [minimum, maximum]. Fit: median excess of the ratio realized by designs optimized on training records over the exact ratio of the same task (†the 15 tasks with $m = 12$). Own: share of the log-gap captured when branching may use only the forests' own trees. Two: exact optimal two-round design, W_2/W_{branch} (‡five splits per cell: 45 and 15 tasks; at $b = 2$ two rounds are branching). Cap: W_{fixed} and W_{branch} relative to the full-information variance.

						Cap: /full	
b	tasks	Fixed/branch	Fit	Own	Two	fixed	branch
2	45	1.31 [1.12, 1.49]	+1%	49%	1	4.1	3.3
3	180	1.51 [1.14, 1.85]	+2%†	17%	1.04‡	3.5	2.3
4	15	1.24 [1.13, 1.35]	+3%	26%	1.03‡	1.4	1.1

WDBC, $(500, 1000, 3101)$ for Spambase and $(60, 120, 171)$ for Ionosphere. Continuous features are binarized at their training medians; m features are chosen at equally spaced positions of the eligible feature list, without labels. Each forest has twenty trees of depth b, $\sqrt{m}$ features per split and minimum leaf size $\max\{3, n_{\mathrm{old}}/30\}$. The contrast d is the difference of the two forests' predicted probabilities on the cube; it has degree at most b. The cap is binding for $m > b$. At $b = 3$ twenty splits per $(m, \text{dataset})$ cell are used; at $b = 2$ and $b = 4$ five splits. The cap $b = 4$ was completed for $m = 6$ only: the branching program over all depth-four trees with forty padded component trees did not finish for $m \geq 9$ within our compute budget, and $b = 5$ was not run. At $m = 12$ the held-out records occupy a median of 171, 388 and 121 of the 4,096 profiles for WDBC, Spambase and Ionosphere (ranges 164–180, 373–405 and 113–130 over twenty splits). Spambase, whose 3,101 held-out records cover few profiles densely, gives a median exact ratio of 1.58 at $b = 3$, $m = 12$, close to WDBC (1.66) and Ionosphere (1.73), so the gains are not an artifact of sparse held-out sets; Appendix F.1 also varies the smoothing mass and resamples the held-out records.

Designs and certificates. The fixed constant optimizes over all $\binom{m}{b}$ forms; the branching constant over all depth-b trees by column generation. The restricted menu adds only the forty component trees to the forms; at $b \geq 4$ their short paths are padded to depth b, a refinement within the cap, so the restricted-menu ratios at $b \geq 4$ are slightly more favourable to branching than at $b \leq 3$. At $b = 4$ column generation stops when the certified bounds agree to relative error 10^{-4} or no tree improves the master; over the 15 tasks the largest relative gap of the reported ratio is 2e-08, the median number of iterations 28 and the median time 40 seconds per task on one core. At $b \in \{2, 3\}$ an independent dense implementation certified every task to 10^{-6}.

Two-round designs. For five splits per dataset at $(m, b) \in \{(6, 3), (9, 3), (12, 3), (6, 4)\}$ the solver computes the optimal fixed, two-round (and at $b = 4$ three-round) and branching designs with certified bounds (relative gap at most 1e-04). Of the 45 tasks that the dense implementation had also solved, the fixed-to-branching ratios agree to a relative error of 2×10^{-4} on 44; on the remaining 1 the retrained forests differ slightly from the stored ones (Δ differs in the third decimal) and the ratios differ by at most 0.2%. The share of the log-gap that a class recovers is $1 - \log(W_r/W_{\mathrm{branch}})/\log(W_{\mathrm{fixed}}/W_{\mathrm{branch}})$. Two rounds recover a median 89% (range 80–95%); at $b = 4$ three rounds recover a median 98% (Table 6).

Twenty features. To test the forest findings beyond $m = 12$ we repeat the Spambase protocol with $m = 20$ features (the quantile-spaced subset of the eligible features), where the cube has 2^{20} profiles and the adaptive classes cannot be solved

Table 6. Forest updates by cap and dimension (medians over splits). Fixed/branch at $b = 3$ uses twenty splits per dataset (Table 4); the two-round columns use five splits per dataset and the solver of Appendix E.4: W_2/W_{branch}, the share of the log-gap that the second round recovers (median and minimum over tasks), and the largest relative gap between the certified bounds of the branching constant.

b	m	WDBC	Spambase	Ionosphere	fitted	W_2/W_{branch}	recovered	min	width
2	6	1.31	1.35	1.31	1.31	–	–	–	–
2	9	1.23	1.25	1.44	1.26	–	–	–	–
2	12	1.27	1.35	1.42	1.35	–	–	–	–
3	6	1.36	1.41	1.36	–	1.023	92%	85%	6e-07
3	9	1.57	1.45	1.61	–	1.041	90%	86%	1e-04
3	12	1.66	1.58	1.73	1.67	1.057	89%	86%	8e-05
4	6	1.23	1.28	1.20	1.27	1.031	85%	80%	9e-05

to optimality in reasonable time. The fixed optimum is certified over all $\binom{20}{b}$ forms. Column generation over two-round interviews (at $b = 3$) and over branching interviews is given a wall-clock budget (15 and 25 minutes at $b = 3$; 25 minutes for branching at $b = 4$); when it is stopped by the budget, the certified lower and upper bounds reached give intervals for the ratios. At $b = 3$ column generation over branching interviews closed within its budget on all three splits, so the ratios

Table 7. Forest updates on $m = 20$ median-binarized Spambase features (twenty-tree forests of depth b, 3,101 held-out records, 1% smoothing). The fixed optimum is certified over all forms. Column generation over two-round and branching interviews was stopped after a wall-clock budget, so their constants are certified intervals and the ratios to branching are intervals: fixed / branch $\in [W_{\mathrm{fixed}}/\overline{W}_{\mathrm{branch}},\ W_{\mathrm{fixed}}/\underline{W}_{\mathrm{branch}}]$. Used: features on which the forests split. Iter.: column-generation iterations for branching.

b	seed	forms	used	fixed / complete	fixed / branch	two rounds / branch	iter.
3	0	1,140	17	6.9	[1.54, 1.54]	[1.00, 1.08]	130
3	1	1,140	18	3.7	[1.50, 1.50]	[1.00, 1.03]	121
3	2	1,140	17	5.1	[1.61, 1.61]	[1.04, 1.05]	114
4	0	4,845	18	5.8	[1.03, 5.76]	–	66

1.50–1.61 are exact to the certified gap; they lie in the range of the twelve-feature tasks. The two-round programs did not close in 15 minutes, and their certified intervals put two rounds between 1.00 and 1.08 times the records of branching. At $b = 4$ (4,845 forms) the fixed optimum was certified in seven minutes, but 25 minutes of column generation over branching interviews left only the trivial lower bound for the one split that was run, so its interval is uninformative; the other two splits were not run. Exact adaptive designs at $m = 20$ and $b \geq 4$ are beyond this implementation.

Smoothing mass and resampling. For the fifteen $b = 3$ tasks with $m = 12$ (five splits per dataset) the exact constants are recomputed with $\alpha \in \{0.1\%, 1\%, 5\%, 10\%\}$ and, at $\alpha = 1\%$, on bootstrap resamples of the held-out records with the forests fixed.

Variation	tasks	median ratio	median rel. change	max rel. change	median $\lvert \log \rvert$ change
$\alpha = 0.1\%$	15	1.654	+0.0%	2.0%	0.003
$\alpha = 5\%$	15	1.660	+0.8%	2.7%	0.008
$\alpha = 10\%$	15	1.690	+1.8%	5.3%	0.018
bootstrap of held-out records ($\alpha = 1\%$)	27	1.602	-1.1%	7.5%	0.012

Screening for a gain from reference data. For every forest task at $b \in \{2, 3, 4\}$ and every screening run, the uncentered lower end $(\mathbb{E}_P[t^2]/\Gamma_P - \Delta^2)/\sigma_P^2$ of Appendix E.1 is recomputed on the reference law alone and compared with the realized exact ratio.

Tasks	n	Spearman($\widehat{L}$, realized)	median realized/$\widehat{L}$	realized$< \widehat{L}$	rule: $\widehat{L} \leq 1.05 \Rightarrow$ realized ≤ 1.25
forests	240	0.52	1.08	75	2/2
screening trees	48	0.78	1.00	0	31/31

For a screening tree this quantity is a lower bound on the records ratio at the reference law. For a forest update it is not: no single tree computes the contrast, branching itself needs more records than complete data, and the quantity bounds W_{fixed} relative to the full-information variance; it exceeds the realized fixed-to-branching ratio in 75 of the 240 forest tasks. We

therefore use it only as a screen. In these data the screen *at most 1.05 ⇒ do not plan for a gain* has no misses: all 17 screening runs with a realized gain above 1.25 have a reference value above 1.05, all 31 runs with a value at most 1.05 realize at most 1.25, and none of the 214 forest tasks with a realized gain above 1.25 has a value at most 1.05 (only two forest tasks have a value at most 1.05).

F.3. Screening Questionnaires

Items and targets. BRFSS items: high blood pressure, high cholesterol, body-mass index at least 30, smoking, stroke, heart disease, physical activity, heavy drinking, fair or poor general health, difficulty walking, age 60 or older, and income in the five lowest categories; the target is the diabetes-or-prediabetes indicator. Thyroid items: age above its upper quartile, sex, thyroxine treatment, thyroid surgery, query hypothyroid, goitre, TSH above its upper quartile, T3, TT4 and FTI below their lower quartiles, T4U above its upper quartile, and sickness, with quartiles from the training file and no labels; the target is any abnormal class.

Splits and trees. For the random-half target, a random half of the records is the source; for the shifted target, the source is one sex and the target the other. Up to 20,000 source records train a class-balanced CART tree of depth b (minimum leaf size 50 for BRFSS, 10 for thyroid); up to 20,000 further source records are the reference for fitted designs. The estimand is the target accuracy of the tree's decision, $d \in \{-1, 1\}$, so $M_{\mathrm{branch}} = \mathbb{E}[t^2] = 1$ and branching attains the full-information variance. Seeds $0, 1, 2$ vary the split and the tree.

Designs. Best: the optimal fixed design over all $\binom{12}{b}$ forms. Fitted: the same program on the source reference. Label: the optimal fixed design whose form probabilities may depend on the label, $W^{\mathrm{lab}}_{\mathrm{fixed}}(P)$ of Proposition E.2, computed as two certified programs, one per label. Strat.: the same with strata defined by the label and sex (random-half BRFSS targets, where both sexes occur). PS: the optimized path-set design of Proposition E.6, at the target law and fitted. H2: follow the root question, then use the best fixed form of the remaining size in the reached subcube. r-round (exact): the certified optimum over all r-round interviews. Two fit.: column generation over two-round interviews on the source reference with the 5% floor, started from the oracle two-round design, evaluated at the target. U-p: the uniform design over the tree's padded path sets with the tables that minimize the variance for that design. U-i: the uniform design over all b-subsets of the items the decision depends on (a single form if it depends on at most b items), with the best tables for that design. U-HT: the uniform design over all forms with the monomial Horvitz–Thompson tables $g_T = \sum_{S \subseteq T} \widehat{t}(S)\chi_S / N(S)$, where $N(S) = \binom{m-|S|}{b-|S|}$ is the number of forms that contain S. U-opt: the uniform design with the tables that minimize the variance for that design, a weighted least-norm problem. $1/\rho_P^2$ and N_P: the ends of Theorem 3.

Table 8. Screening trees on twelve items, all designs without a model: records needed relative to branching, which asks the tree's own questions and attains the full-information variance (medians over two targets and three seeds; the range is over runs). Best: optimal fixed design over all forms at the target law. Fitted: the same program on source records, evaluated at the target. Label: optimal fixed design whose form probabilities may depend on the label. Two: exact optimal two-round design (at $b = 6$ the median includes certified upper bounds from feasible two-round designs for 4 runs, Table 10). Two fit.: two-round design fitted on source records. Naive pre-committed designs with the best scores for each: U-paths, uniform over the tree's question paths; U-items, uniform over all b-subsets of the items the rule depends on; U-all, uniform over all $\binom{12}{b}$ forms. $1/\rho_P^2$, N_P: the ends of Theorem 3. – not computed.

Data, b	runs	Best	[min, max]	Fitted	Label	Two	Two fit.	U-paths	U-items	U-all	$1/\rho_P^2$	N_P
BRFSS, 3	6	1.00	[1.00, 1.34]	1.04	1.00	1.00	1.00	1.03	1.00	15.8	1.00	1.5
BRFSS, 4	6	1.49	[1.00, 1.75]	1.57	1.48	1.00	1.01	1.80	2.30	21.7	1.18	3.3
BRFSS, 5	6	1.35	[1.18, 1.54]	1.42	1.35	1.00	1.01	1.72	2.04	12.9	1.08	5.4
BRFSS, 6	6	1.58	[1.42, 1.98]	1.68	–	≤1.05	–	2.30	4.18	11.1	1.13	10.5
Thyroid, 3–6	24	1.00	[1.00, 1.00]	1.04	1.00	1.00	1.04	1.20	1.00	4.9	1.00	3.6

Label-dependent and stratified forms. Over the 30 runs, letting the form probabilities depend on the label lowers the records ratio by at most 0.5% (certified gap between the bounds at most 1e-04), and stratifying in addition by sex by at most 0.8% over 9 runs. The prevalence of the target is about 14%, so the two labels do weight the tree's paths differently, but the forms that serve one label serve the other almost as well.

Two-round designs. A two-round interview that asks a first batch B_1 computes the decision exactly if and only if, for every answer pattern on B_1, the restricted decision depends on at most $b - |B_1|$ further items. If such a batch exists, the

Table 9. All screening settings (medians over three seeds). Ratios are records needed relative to branching. Best/Fitted: optimal and fitted fixed designs; Label, Strat.: optimal fixed designs whose form probabilities depend on the label, and on the label and sex; PS: optimized path-set design (oracle/fitted); H2: heuristic two-round design; r=2, 3: exact optimal r-round designs; 2 fit.: fitted two-round design; U-p, U-i: uniform designs over the tree's path sets and over the b-subsets of the items the rule depends on, with optimal scores; U-HT, U-opt: uniform design over all forms with monomial Horvitz–Thompson scores and with optimal scores; $1/\rho_P^2$, N_P: ends of Theorem 3.

Data	b	Target	Acc.	Paths	Best	Fitted	Label	Strat.	PS	PS fit	H2	r=2	r=3	2 fit.	U-p	U-i	U-HT	U-opt	$1/\rho_P^2$	N_P
BRFSS	3	random half	0.629	2	1.00	1.04	1.00	1.00	1.00	1.00	1.00	1.00	–	1.00	1.06	1.00	20.4	11.2	1.00	1.5
BRFSS	3	other sex	0.741	1	1.00	1.04	1.00	–	1.00	1.00	1.00	1.00	–	1.00	1.00	1.00	44.6	23.8	1.00	1.0
BRFSS	4	random half	0.664	4	1.46	1.52	1.45	1.45	1.50	1.52	1.14	1.00	1.00	1.01	1.77	2.30	43.2	21.5	1.18	3.2
BRFSS	4	other sex	0.681	4	1.52	1.61	1.51	–	1.57	1.62	1.13	1.00	1.00	1.01	1.97	2.29	47.6	22.6	1.19	3.7
BRFSS	5	random half	0.675	7	1.28	1.32	1.28	1.27	1.28	1.29	1.15	1.02	1.00	1.02	1.66	1.89	26.9	11.9	1.04	5.2
BRFSS	5	other sex	0.734	6	1.43	1.52	1.43	–	1.44	1.50	1.27	1.00	1.00	1.01	1.88	2.06	30.5	14.5	1.13	5.5
BRFSS	6	random half	0.719	14	1.53	1.58	–	–	1.53	1.56	1.40	1.06	–	–	2.41	3.09	29.7	11.6	1.13	11.0
BRFSS	6	other sex	0.722	13	1.60	1.70	–	–	1.65	1.70	1.39	1.01	1.00	–	2.21	4.30	25.3	10.2	1.13	9.8
Thyroid	3	random half	0.823	3	1.00	1.04	1.00	–	1.00	1.00	1.00	1.00	–	1.04	1.19	1.00	32.4	10.8	1.00	2.4
Thyroid	3	other sex	0.819	3	1.00	1.04	1.00	–	1.00	1.01	1.00	1.00	–	1.00	1.37	1.00	31.9	10.7	1.00	2.8
Thyroid	4	random half	0.823	3	1.00	1.04	1.00	–	1.00	1.00	1.00	1.00	1.00	1.04	1.15	1.00	11.3	4.9	1.00	2.4
Thyroid	4	other sex	0.848	4	1.00	1.04	1.00	–	1.00	1.01	1.00	1.00	1.00	1.00	1.60	1.00	30.3	10.4	1.00	3.3
Thyroid	5	random half	0.823	6	1.00	1.03	–	–	1.00	1.00	1.00	1.00	1.00	–	1.11	1.00	6.1	3.2	1.00	5.5
Thyroid	5	other sex	0.848	5	1.00	1.04	–	–	1.00	1.01	1.00	1.00	1.00	–	1.27	1.00	10.7	4.9	1.00	4.0
Thyroid	6	random half	0.823	7	1.00	1.03	–	–	1.00	1.00	1.00	1.00	1.00	–	1.09	1.00	4.0	2.4	1.00	6.1
Thyroid	6	other sex	0.848	5	1.00	1.04	–	–	1.00	1.01	1.00	1.00	1.00	–	1.27	1.00	5.7	3.1	1.00	4.0

interview observes t and $W_2 = W_{\text{branch}}$; at a full-support law the converse holds as well, because a design with variance σ_P^2 must return t at every profile. We enumerate all first batches inside the items the decision depends on. Of the 24 BRFSS runs, 17 are two-round computable, and the enumeration agrees with the solver on all 43 runs for which the solver also computed the two-round optimum (ratio one exactly when computable). The trees that are not two-round computable are the larger ones: two of the six at $b = 5$ and five of the six at $b = 6$; class-balanced CART produces unbalanced trees with few distinct paths at the smaller caps. Exact two-round optima were computed by the solver for every run at $b \leq 5$. At $b = 6$ one run is two-round computable. For the other five, column generation was given one hour per run on one core (about fifty to sixty iterations and five to seven hundred generated interviews). One run closed its bracket to a relative gap of 1.3×10^{-4} and is counted as solved; for the other 4 Table 10 reports the bracket reached. Its upper end is the variance of a feasible, exactly unbiased two-round design. Its lower end is one where the dual bound of the stopped run was not yet above the full-information bound; for these runs the ratio is nevertheless known to exceed one, since the trees are not two-round computable. The forest updates are not two-round computable either (all sixty exact two-round ratios exceed one), and neither is the pointer tree.

Further rules and bootstrap intervals. To see whether the constants of the diabetes trees are typical, we trained trees for four other outcomes in the same survey (heart disease or attack, stroke, high blood pressure, difficulty walking) on the 21 binarized variables of Appendix F.4, with diabetes status as a feature and the outcome removed from the features, and for Spambase on its 57 median-binarized features. Each run trains a class-balanced tree of depth b on at most 20,000 records of a random half (2,300 for Spambase), takes the other half as the target law (with 1% uniform mass on the items the rule depends on and the label), and solves the fixed program over all forms of the items the rule depends on. Forms that include other variables could only lower the fixed constant, so the reported ratios are upper bounds on the ratio for the best forms on all variables; for the diabetes trees of Table 1 the program is over all forms of the twelve items. The adaptive optimum is the rule's tree, with constant $\mathrm{Var}_P(t)$. Of the 45 runs, 38 have a gain and were computed, 2 are trivial, and 5 were not computed (5 rules depend on more than 13 items; 0 did not finish). For 28 of the computed runs the bootstrap, the uniform design and the pairwise fit were computed as well; for the other 10 only the optimal forms. The ratios lie in the range of the diabetes trees of Table 1, with a few larger values (up to 2.38). For the runs that were not computed, the lower end of the bracket is 1.14–1.43, within the range of the computed ratios, and the upper end 8–20 is not informative; the largest rules are thus not known to behave differently, but their ratios are not known either. Resampling the target records with the tree fixed moves a ratio by at most 0.6% of its value on BRFSS, where a target half has more than 10^5 records, and by at most 6% on Spambase (2,301 target records; 5th to 95th percentile of 20 resamples, 26 runs in all). The variation in the tables is therefore between rules and training samples, not sampling error of the target law; the bootstrap does not cover the uncertainty of training the rule. Uniform randomization over the forms of the rule's items needs 1.3–16.8 times the records of branching (median 2.1), and the ratio within the pairwise model is 1.00–1.28 (median 1.09, 26 runs).

Table 10. Every BRFSS screening run (twelve items; R: random-half target, S: other-sex target; seeds 0–2) and every 21-item run: records needed relative to branching. Items: number of items the decision depends on. Two: exact optimal two-round design; a bracket gives the certified lower and upper bounds reached within the time budget. – not computed.

Run	b	Acc.	Items	Paths	Best	Fitted	Label	Two	Two fit.	PS	$1/\rho_P^2$
R0	3	0.604	2	2	1.000	1.035	1.000	1.000	1.035	1.000	1.000
R1	3	0.736	3	4	1.000	1.040	1.000	1.000	1.000	1.000	1.000
R2	3	0.629	3	2	1.000	1.037	1.000	1.000	1.000	1.000	1.000
S0	3	0.742	3	1	1.000	1.040	1.000	1.000	1.000	1.000	1.000
S1	3	0.720	3	1	1.000	1.040	1.000	1.000	1.000	1.000	1.000
S2	3	0.741	4	2	1.338	1.409	1.338	1.000	1.008	1.338	1.070
R0	4	0.664	6	4	1.457	1.524	1.451	1.000	1.009	1.496	1.153
R1	4	0.659	5	4	1.406	1.461	1.404	1.000	1.009	1.458	1.176
R2	4	0.687	6	3	1.550	1.611	1.543	1.000	1.010	1.608	1.192
S0	4	0.664	4	2	1.000	1.035	1.000	1.000	1.035	1.000	1.000
S1	4	0.684	7	5	1.749	1.867	1.740	1.000	1.011	1.817	1.213
S2	4	0.681	6	4	1.517	1.610	1.511	1.000	1.006	1.570	1.188
R0	5	0.675	6	7	1.184	1.231	1.183	1.018	1.020	1.189	1.031
R1	5	0.678	7	8	1.279	1.323	1.275	1.000	1.007	1.281	1.042
R2	5	0.663	8	5	1.513	1.581	1.508	1.040	1.046	1.544	1.099
S0	5	0.692	7	6	1.260	1.322	1.259	1.000	1.006	1.261	1.070
S1	5	0.746	8	6	1.537	1.645	1.535	1.000	1.011	1.564	1.144
S2	5	0.734	7	9	1.430	1.522	1.425	1.000	1.011	1.439	1.132
R0	6	0.729	10	14	1.982	2.059	–	[1.000, 1.320]	–	2.022	1.264
R1	6	0.719	9	17	1.530	1.580	–	[1.000, 1.063]	–	1.534	1.132
R2	6	0.686	9	13	1.417	1.475	–	[1.012, 1.031]	–	1.438	1.066
S0	6	0.690	10	12	1.604	1.700	–	1.000	–	1.645	1.119
S1	6	0.725	10	13	1.714	1.827	–	1.013	–	1.767	1.150
S2	6	0.722	10	20	1.561	1.664	–	[1.000, 1.109]	–	1.570	1.132
21 items, R0	3	0.604	2	2	1.000	1.000	–	1.000	–	1.000	1.000
21 items, R1	3	0.736	3	4	1.000	1.016	–	1.000	–	1.000	1.000
21 items, R2	3	0.629	3	2	1.000	1.002	–	1.000	–	1.000	1.000
21 items, R0	4	0.664	6	5	1.457	1.535	–	1.000	–	1.496	1.153
21 items, R1	4	0.659	5	4	1.406	1.471	–	1.000	–	1.458	1.176
21 items, R2	4	0.687	6	3	1.550	1.623	–	1.000	–	1.608	1.192

Extra pre-committed questions. For every rule of Tables 1 and 11 we solve the fixed program over all forms of size $b + c$ on the items the rule depends on, $c = 0, \dots, 4$, as long as the program has at most 4.5×10^5 cells, and the path-set program at size b. Of the 59 rules that depend on more than b items and were run (1 further rule, with eleven items at cap six, was stopped unfinished), 52 have the optima at sizes b and $b + 1$ and 51 also at $b + 2$; the others have more items than the solver handles at those sizes. At size $b + 1$ the forms need a median 1.14 times the records of branching (at most 1.60), against 1.47 at size b: a median 64% and at least 35% of the log-gap is removed. At size $b + 2$ the median is 1.02 (at most 1.29; 92% removed in the median). Theorem C.2 allows much less in the worst case, a factor two for one extra question at any cap. The difference is the number of items beyond the depth, $K - b$, which is 3 in the median and at most 7 for the rules computed at size $b + 1$ (at most 10 over all rules) and $2^{b-1} - 1$ for the address contrast. The comparison is with the branching interview of depth b, which already attains the full-information variance for a single tree, so extra questions cannot help branching. The optimized path-set design of size b needs a median 1.5% and at most 14% more records than the optimal forms on the 52 rules where both were computed; for the 5 rules whose optimum at size b was not computed it certifies that forms need at most 2.20–3.97 times the records of branching, against the lower ends 1.14–1.43 of the bracket. The values in this table are primal values of the program over all forms of the given size, accurate to the tolerance of the cone solver (about 10^{-3} in relative terms).

F.4. All Twenty-One BRFSS Variables

All 21 variables of the CDC file are binarized at pre-specified cut points: high blood pressure, high cholesterol, cholesterol check, body-mass index at least 30, smoking, stroke, heart disease, physical activity, fruit, vegetables, heavy drinking, any health coverage, cost barrier to a doctor visit, fair or poor general health, at least 14 days of poor mental health, at least 14 days of poor physical health, difficulty walking, sex, age 60 or older, no college education, and income in the five lowest categories. Splits, trees and references are as for the twelve-item study (random-half target). The form program has $\binom{21}{b}$ forms, 1,330 at $b = 3$ and 5,985 at $b = 4$, and the profile cube has 2^{21} elements; the solver never enumerates the cube, and at $b = 4$ one master problem over all forms certifies the optimum. In all three seeds at $b = 4$ the tree selects only items of the twelve-item study and coincides with the tree of that study (runs R0–R2 of Table 10), so its constants coincide as well; the purpose of this study is the size of the program, whose optimum must be certified against all 5,985 forms. All three trees at $b = 4$ are two-round computable, so their two-round ratio is exactly one. Label-dependent designs and the cap $b = 5$

Table 11. Further rules: class-balanced depth-b trees for four other BRFSS outcomes (trained on 20,000 respondents with the 21 binarized variables, diabetes status included as a feature) and for Spambase (57 binarized features, 2,300 training records), three seeds each; target law on a disjoint half. Runs: with a gain / trivial (the rule depends on at most b items, ratio 1) / not computed (5 rules with more than 13 items; 0 runs not finished). For 10 of the runs with a gain only the optimal forms were computed, without the bootstrap, the uniform design and the model fit. Last column: for the runs that were not computed, the bracket $[1/\rho_P^2, N_P]$ of Theorem 3 over forms on the rule's items. For the runs with a gain: number K of items the rule depends on; records needed relative to branching by the optimal forms on these items, median [range]; the largest distance, over the runs, from this ratio to the 5th or the 95th percentile of 20 bootstrap resamples of the target records with the tree fixed, relative to the ratio (runs with $K \le 11$); uniform randomization over all forms of the rule's items with optimal tables; and the ratio within the pairwise model at the pairwise fit (runs with $K \le 10$).

target	b	runs	K	optimal forms	bootstrap	uniform, items	pairwise	bracket if not computed
BRFSS, heart disease	4	3 / 0 / 0	5–7	1.29 [1.16, 1.53]	±0.5%	2.15 [1.44, 3.22]	1.10 [1.06, 1.18]	–
	5	3 / 0 / 0	8–10	1.67 [1.57, 2.23]	±0.4%	3.73 [2.93, 8.92]	1.19 [1.15, 1.24]	–
	6	1 / 0 / 2	13–13	2.38 [2.38, 2.38]	–	16.76 [16.76, 16.76]	–	[1.16, 16.5], [1.14, 17.4]
BRFSS, stroke	4	3 / 0 / 0	6–6	1.37 [1.30, 1.44]	±0.3%	1.95 [1.91, 2.05]	1.12 [1.12, 1.23]	–
	5	3 / 0 / 0	8–9	1.32 [1.30, 1.79]	±0.6%	3.24 [2.19, 3.74]	1.10 [1.08, 1.28]	–
	6	1 / 0 / 2	11–11	1.74 [1.74, 1.74]	–	–	–	[1.43, 20.0], [1.20, 15.9]
BRFSS, high blood pressure	4	3 / 0 / 0	5–5	1.25 [1.16, 1.33]	±0.3%	1.57 [1.51, 1.59]	1.01 [1.01, 1.01]	–
	5	3 / 0 / 0	6–6	1.13 [1.13, 1.13]	±0.3%	1.41 [1.41, 1.41]	1.01 [1.00, 1.01]	–
	6	3 / 0 / 0	7–10	1.21 [1.07, 1.38]	±0.2%	1.31 [1.31, 1.31]	1.00 [1.00, 1.00]	–
BRFSS, difficulty walking	4	1 / 2 / 0	5–5	1.24 [1.24, 1.24]	±0.2%	1.40 [1.40, 1.40]	1.06 [1.06, 1.06]	–
	5	3 / 0 / 0	7–8	1.48 [1.32, 1.56]	±0.6%	1.99 [1.93, 2.81]	1.08 [1.04, 1.09]	–
	6	3 / 0 / 0	10–10	1.75 [1.61, 1.78]	–	–	–	–
Spambase, spam	4	3 / 0 / 0	6–8	1.95 [1.40, 2.06]	±5.7%	4.46 [2.21, 5.13]	1.18 [1.03, 1.25]	–
	5	3 / 0 / 0	10–12	1.99 [1.92, 2.04]	–	8.38 [8.38, 8.38]	–	–
	6	2 / 0 / 1	11–13	1.94 [1.76, 2.13]	–	–	–	[1.27, 8.4]

Table 12. Extra pre-committed questions on trained rules. For every rule that depends on $K > b$ items: the records that optimal forms of size $b + c$ on the rule's items need relative to the branching interview with b questions, medians over rules [range]; a form of size K asks every item of the rule and has ratio one. Rules: with $c = 0$ and $c = 1$ computed / all with a gain (the others have more items than the solver handles at that size); the columns $c = 2$ and $c = 3$ are over the rules computed at that size. Excess: $K - b$, over all rules with a gain. Path sets: the optimized path-set design of size b relative to the optimal forms of size b, where both were computed.

target	rules	excess	$c = 0$	$c = 1$	$c = 2$	$c = 3$	path sets
BRFSS, diabetes, $b = 4$	5 / 5	2 [1, 3]	1.52 [1.41, 1.75]	1.12 [1.00, 1.24]	1.00 [1.00, 1.05]	1.00 [1.00, 1.00]	+3.7%
BRFSS, diabetes, $b = 5$	6 / 6	2 [1, 3]	1.35 [1.18, 1.54]	1.08 [1.00, 1.18]	1.00 [1.00, 1.04]	1.00 [1.00, 1.00]	+0.5%
BRFSS, diabetes, $b = 6$	6 / 6	4 [3, 4]	1.58 [1.42, 1.98]	1.24 [1.12, 1.43]	1.09 [1.03, 1.18]	1.02 [1.00, 1.06]	+1.7%
BRFSS, heart disease	6 / 9	3 [1, 8]	1.55 [1.16, 2.23]	1.17 [1.00, 1.60]	1.03 [1.00, 1.28]	1.00 [1.00, 1.12]	+1.0%
BRFSS, stroke	7 / 9	4 [2, 10]	1.37 [1.30, 1.79]	1.13 [1.08, 1.39]	1.02 [1.00, 1.19]	1.00 [1.00, 1.05]	+1.6%
BRFSS, high blood pressure	9 / 9	1 [1, 4]	1.16 [1.07, 1.38]	1.00 [1.00, 1.17]	1.00 [1.00, 1.06]	1.00 [1.00, 1.00]	+0.0%
BRFSS, difficulty walking	7 / 7	3 [1, 4]	1.56 [1.24, 1.78]	1.17 [1.00, 1.37]	1.08 [1.00, 1.15]	1.00 [1.00, 1.00]	+3.7%
Spambase, spam	6 / 8	5 [2, 8]	1.97 [1.40, 2.06]	1.48 [1.15, 1.56]	1.20 [1.00, 1.29]	1.06 [1.00, 1.10]	+5.3%

(20,349 forms) were not completed at $m = 21$ within our compute budget.

Table 13. All 21 BRFSS variables (random-half target; medians over three seeds). Forms: $\binom{21}{b}$. Time: median minutes for the certified fixed design on one core.

b	forms	Acc.	items used	Best	Fitted	PS	$1/\rho_P^2$	N_P	time
3	1330	0.629	4	1.00	1.00	1.00	1.00	1.5	0.0
4	5985	0.664	6	1.46	1.54	1.50	1.18	3.5	6.1

F.5. Learning the Design from a Pilot

Proposition E.9 is asymptotic and discards the pilot. To measure the all-in cost of learning at finite sizes we use the BRFSS runs with a gain and the tree's path-set menu, whose optimum is within a few percent of the all-forms optimum (Table 9). A pilot of n_1 target respondents is interviewed with the uniform design on the path sets. The pilot is split into two halves at random; the masses of the cells of each path-set form are estimated from one half, mixed with a uniform mass of 10%, and used to compute the tables for the uniform design that score the *other* half. These scores are pointwise unbiased and independent of the records they score. The main sample of $N - n_1$ respondents uses the path-set design learned from the whole pilot with the 5% floor. The final estimate averages all N scores; its variance is $[n_1 w_{\text{pilot}} + (N - n_1) w_{\text{learned}}]/N^2$ for the per-record variances w, which are computed exactly at the target law for 30 independent pilots per run and size. With a quarter of the records in the pilot, the all-in price exceeds the oracle all-forms price by a median 15% and 35% at

Table 14. Learning the design from a pilot of the audit itself, with the pilot charged (BRFSS runs with a gain; medians over runs of the median over 30 pilots). Pilot: per-record variance of the cross-fitted scores of the n_1 pilot respondents (uniform design on the path sets). Learned: per-record variance of the path-set design learned from the pilot. All-in: variance of the estimator that uses all $N = 4n_1$ records with weights proportional to sample sizes, $[n_1 \cdot \text{pilot} + 3n_1 \cdot \text{learned}]/N$. All entries are relative to branching; the last columns give the oracle path-set design and the oracle design over all forms. Cells: number of (form, pattern, label) masses to estimate.

b	runs	cells	n_1	pilot	learned	all-in	oracle PS	oracle all forms
3	1	32	250	1.56	1.37	1.42	1.34	1.34
			500	1.53	1.36	1.40		
			1000	1.52	1.35	1.39		
			2000	1.51	1.35	1.39		
4	5	128	250	2.43	1.80	1.96	1.57	1.52
			500	2.13	1.71	1.75		
			1000	1.99	1.64	1.67		
			2000	1.90	1.61	1.65		
5	6	416	250	3.26	2.46	2.61	1.36	1.35
			500	2.95	1.90	2.09		
			1000	2.31	1.66	1.80		
			2000	1.99	1.54	1.63		
6	6	1728	250	4.53	4.15	4.24	1.61	1.58
			500	5.57	4.21	4.59		
			1000	5.40	3.71	4.13		
			2000	4.79	2.76	3.30		

$b = 4, 5$ for $N = 4{,}000$ and by 13% and 20% for $N = 8{,}000$. The excess falls roughly in proportion to the number of cells per pilot respondent; at $b = 6$ pilots of this size are too small. A design fitted on a large reference does better at every cap (Table 10), so a pilot is the tool for the case in which no reference exists, and the all-forms menu, with $\binom{12}{b}2^{b+1}$ cells, is out of reach of pilots of this size. A non-asymptotic guarantee for the two-phase procedure is not proved here.

Smoothing the pilot with a parsimonious model. The tables and form probabilities of a design are unbiased for any law used to choose them, so the pilot's cell masses may come from a model without any cost in validity. We fit the pairwise log-linear model for the label and the items of the path-set forms to the pilot's incomplete records by EM and take every form's cell masses from the fitted model, mixed with a uniform mass of 1%; everything else is as above (12 pilots per run and size). Smoothing lowers the all-in price at every cap and most where the pilot is thinnest: for $N = 4{,}000$ from 1.67 to

Table 15. The pilot of Table 14 with its cell masses smoothed by a pairwise log-linear fit (BRFSS runs with a gain; medians over runs of the median over 12 pilots; $N = 4n_1$). Empirical: cell masses from the pilot's frequencies, as in Table 14. Pairwise fit: cell masses from the pairwise model fitted to the pilot by EM. Pilot, learned and all-in are per-record variances relative to branching, as in Table 14; over oracle is the median over runs of the all-in price divided by that run's oracle optimum over all forms.

			empirical			pairwise fit				oracle	
b	runs	n_1	pilot	learned	all-in	pilot	learned	all-in	over oracle	PS	all forms
4	5	1,000	1.97	1.64	1.67	1.86	1.59	1.65	+11%	1.57	1.52
		2,000	1.89	1.61	1.65	1.84	1.59	1.64	+11%		
5	6	1,000	2.33	1.70	1.82	1.78	1.40	1.53	+12%	1.36	1.35
		2,000	1.98	1.54	1.64	1.75	1.38	1.50	+10%		
6	6	1,000	5.38	3.79	4.33	2.66	1.70	1.93	+20%	1.61	1.58
		2,000	4.91	2.82	3.37	2.42	1.66	1.84	+15%		

1.65 at $b = 4$, from 1.82 to 1.53 at $b = 5$ and from 4.33 to 1.93 at $b = 6$ (medians), which is 11%, 12% and 20% above the oracle optimum over all forms (median of per-run ratios); for $N = 8{,}000$ the excess is 11%, 10% and 15%. The percentages quoted in this subsection are medians over runs of the per-run ratio, not ratios of the tabulated medians.

F.6. One Set of Records for Several Rules

An optimized design concentrates on the forms that serve one contrast. To measure what this costs when the records are later used for another rule, we take the other-sex BRFSS target, whose population is the same for every seed, and treat the three trees of seeds 0–2 at each cap as three rules. For every ordered pair we compute the best pointwise-unbiased scores for

the second rule on records collected under the first rule's optimal design, mixed with a 5% uniform floor over all forms so that every contrast of degree at most b stays identifiable. For every pair and for the triple we also solve the fixed-form program for several contrasts of Appendix E.1, which minimizes the sum of the rules' records ratios over one shared form distribution. Reuse can be nearly free or nearly useless: the reused records need between 1.55 and 317 times the records of

Table 16. Using one set of records for several rules (BRFSS, other-sex target; the three trees of seeds 0–2 at each cap are rules A, B, C). Entries are records needed relative to branching for that rule alone. Own: the rule's own optimal fixed design. Reuse: best scores for the rule on records collected under another rule's optimal design mixed with a 5% uniform floor (minimum and maximum over the two other designs). Joint: the fixed design that minimizes the sum of the records ratios of the rules in the group.

b	Rule	Own	Reuse min	Reuse max	Joint A,B	Joint A,C	Joint B,C	Joint A,B,C
4	A	1.00	1.97	2.14	1.40	1.34		1.51
	B	1.75	30.6	316.7	1.94		1.76	1.87
	C	1.52	1.69	286.6		1.67	1.60	1.67
5	A	1.26	20.4	24.5	1.36	1.40		1.43
	B	1.54	10.2	42.2	1.59		1.54	1.56
	C	1.43	1.55	41.2		1.59	1.48	1.58

branching, depending on how many of the second rule's paths the first design happens to cover. A joint design for rules known in advance costs each rule a median 10% (at most 51%) more than its own optimum. For comparison, giving each of three rules its own third of the respondents and administering its tree costs each rule a factor of three.

F.7. Model-Based Alternatives

Simulated audits. Each replication draws $n = 1000$ target respondents with replacement. Branching averages t. The fixed-form score draws a form from the fitted design and averages its score. Truth is the contrast on the full target sample. The trees and fitted designs are those of seed 0, so the random-half and sex settings are runs R0 and S0 of Table 10; in particular the exact records ratio of the sex-shift tree at $b = 4$ is 1.000 (fitted 1.006), so the fitted form and branching have nearly equal standard deviations in the simulation, whereas Table 9 reports medians over three seeds.

Imputation from complete source records. Regression imputation uses the same observed forms and answers and replaces t by its conditional mean given the asked items and the label among the complete source records, with a naive standard error. Multiple imputation draws, for each of ten imputations, Bayesian-bootstrap weights over the source records, imputes the unasked items of each respondent from a donor that agrees on the asked items and the label, and combines by Rubin's rules. Both use information that the capped audit does not collect: complete records whose conditional law of unasked items is assumed to transport to the target. Before any result was computed we fixed four shifts defined by variables outside the twelve items: sex (female source, male target), education (college graduates, target without education beyond high school), cost barrier (no cost barrier to a doctor visit, target with a cost barrier) and physical-health burden (no poor physical-health days in the past month, target with at least 14). Source and target do not overlap in the shift variable, so an imputation model cannot condition on it. All four are reported in Table 17 (400 replications).

Under the random-half target, MI intervals over-cover (0.975–0.993), so Rubin's variance is conservative there. For a frozen imputation bias $\mathfrak{b}_I$ and per-record variances σ_I^2 and σ_U^2 of the imputation and design-based estimators, imputation has the smaller mean squared error exactly when $n\mathfrak{b}_I^2 < \sigma_U^2 - \sigma_I^2$, so a biased imputation can win at small n and lose as the audit grows.

A working model on the capped records. Planned-missing data are usually analysed by maximum likelihood or multiple imputation under a working model for the items, fitted to the incomplete records themselves; missingness is by design, so no transport assumption is involved. We fit the log-linear model with all main effects and two-way interactions among the label and the items that the screening rule uses (items the rule ignores carry no information about the target) by EM, with one iterative-proportional-fitting sweep per M-step, and estimate Δ by $\sum_v p(v)t(v)$. Intervals use a delete-a-group jackknife with eight groups and a Student quantile. The estimator is applied to the records of the fitted fixed-form design and to records of balanced random forms (uniform over all $\binom{12}{b}$ forms), in the same simulated audits (400 replications). At $n = 1000$ the working model on the fitted-form records has the smallest RMSE of the four estimators in most settings, 0.90–1.01 times that of branching: the model shares strength across paths that the design-based scores treat separately. Its bias is between 0.01 and 0.25 branching standard deviations and its coverage 0.920–0.968 (below 0.925 in 1 of 10 settings). The bias is a

Table 17. Imputation from complete source records: simulated BRFSS audits with $n = 1000$ target respondents and 400 replications, for the random-half target (shift: none) and the four pre-specified shifts. The trees are those of seed 0; the first two blocks are runs R0 and S0 of Table 10. Bias, SD and RMSE are multiplied by 100; coverage of nominal 95% intervals has Monte Carlo standard error about 0.011. TV: total variation distance between source and target item profiles. Regression imputation: conditional mean of t given the asked items and the label among the source records, naive standard error. Multiple imputation: ten proper hot-deck imputations from the same source records, Rubin's rules.

Shift	b	TV	Estimator	Bias	SD	RMSE	Coverage	Width
none	4	0	branching	-0.14	2.96	2.96	0.955	11.70
			fitted fixed form	-0.11	3.60	3.60	0.965	14.42
			regression imputation	-0.45	2.59	2.63	0.955	10.27
			multiple imputation	-0.42	2.58	2.61	0.993	13.21
none	5	0	branching	-0.05	2.92	2.92	0.950	11.60
			fitted fixed form	-0.06	3.19	3.18	0.955	12.85
			regression imputation	-0.06	2.67	2.67	0.958	10.84
			multiple imputation	-0.06	2.68	2.68	0.975	12.41
sex	4	0.21	branching	0.05	3.06	3.05	0.953	11.69
			fitted fixed form	0.06	3.12	3.11	0.948	11.89
			regression imputation	0.32	3.01	3.03	0.940	11.50
			multiple imputation	0.33	3.02	3.03	0.955	11.87
sex	5	0.21	branching	0.08	2.89	2.89	0.965	11.42
			fitted fixed form	0.11	3.44	3.44	0.938	13.12
			regression imputation	-0.45	2.80	2.84	0.948	10.72
			multiple imputation	-0.47	2.81	2.85	0.983	12.22
education	4	0.47	branching	0.08	3.12	3.12	0.948	12.28
			fitted fixed form	-0.06	3.60	3.60	0.960	14.20
			regression imputation	2.39	2.79	3.67	0.863	11.28
			multiple imputation	2.38	2.83	3.69	0.920	13.27
education	5	0.47	branching	0.04	3.11	3.11	0.950	12.25
			fitted fixed form	-0.03	3.39	3.38	0.953	13.28
			regression imputation	1.67	2.96	3.39	0.902	11.76
			multiple imputation	1.66	2.97	3.39	0.935	12.73
cost barrier	4	0.41	branching	-0.09	3.23	3.22	0.945	12.01
			fitted fixed form	-0.18	3.80	3.80	0.943	14.75
			regression imputation	0.39	2.82	2.85	0.930	10.76
			multiple imputation	0.39	2.84	2.86	0.980	13.31
cost barrier	5	0.41	branching	0.01	3.13	3.12	0.935	11.75
			fitted fixed form	-0.10	3.89	3.89	0.938	14.45
			regression imputation	-0.72	2.84	2.93	0.925	10.58
			multiple imputation	-0.70	2.92	3.00	0.960	13.01
poor health	4	0.68	branching	0.21	3.17	3.17	0.950	12.39
			fitted fixed form	0.23	3.23	3.23	0.950	12.59
			regression imputation	0.65	3.15	3.21	0.960	12.23
			multiple imputation	0.65	3.15	3.22	0.965	12.57
poor health	5	0.68	branching	0.09	3.17	3.16	0.945	12.37
			fitted fixed form	0.14	3.94	3.94	0.940	15.04
			regression imputation	5.49	2.96	6.24	0.515	11.44
			multiple imputation	5.48	2.96	6.23	0.667	13.23

property of the model, not of the sample size: the accuracy of a depth-b tree depends on interactions of up to b items and the label, which a two-way model does not represent. At $n = 16,000$ the same bias is up to 1.02 of the estimator's own standard deviations, coverage falls to 0.855–0.953 (below 0.90 in 4 of 10 settings), and the RMSE is 0.91–1.42 times that of branching and above that of the design-based form score in 5 settings; the design-based scores keep coverage 0.935–0.958. A richer model would reduce the bias and raise the variance; with all interactions it is the design-based estimator. On balanced random forms the model-based estimate is unbiased to a similar degree but 0.99–1.78 times less accurate at $n = 1000$ than the design-based score on optimized forms, so the choice of forms matters for model-based analysis as well.

F.8. Constants and Audits Within Log-Linear Models

Computation. The constants $W_{\mathcal{A}}^{(k)}(P)$ of Appendix E.2 are computed on the K items that the rule depends on and the label; items the rule ignores are left out of the model and carry no information about the target. The reference law P_k is the maximum-likelihood fit of the order-k model to the target law (mixed with 1% uniform mass), obtained by iterative proportional fitting, so that P_k lies in the model. The basis of the tangent space is $s_A = \chi_A - \mathbb{E}_{P_k}\chi_A$, $1 \le |A| \le k$, and an interview's information matrix is $I_\iota = G_\iota G_\iota^{\mathsf{T}}$ with one column of G_ι per transcript cell and label, $G_\iota[A, c] = \mathbb{E}[s_A \mathbf{1}_c]/P_k(c)^{1/2}$. The weights of a menu solve the Elfving dual $\min\{s : v^{\mathsf{T}}u = 1,\ \|G_\iota^{\mathsf{T}}u\| \le s\ \forall\iota\}$, a second-order cone program whose multipliers are the interview probabilities. Fixed forms are the $\binom{K}{b}$ subsets of the rule's items. Branching designs are found by column generation: for the current u, the measure $\eta = \sum_A u_A s_A P_k$ is priced by the tree dynamic program of Appendix E.4, which returns the depth-b tree that maximizes $u^{\mathsf{T}} I_\tau u$ over all trees on the K items.

Table 18. A working model on the capped records: simulated BRFSS audits with $n = 1000$ and $n = 16,000$ target respondents (400 replications each). Design-based scores (branching; fitted fixed form) against the EM estimate (early-stopped; maximum likelihood where an interior maximiser exists) under a two-way log-linear model for the label and the items the rule uses, fitted by EM to the fitted-form records (EM, fitted forms) and to balanced random forms (EM, uniform forms). Bias, SD and RMSE are multiplied by 100; coverage of nominal 95% intervals (delete-a-group jackknife for EM) has Monte Carlo standard error about 0.011. – not computed.

				$n = 1000$				$n = 16,000$			
Shift	b	items	Estimator	Bias	SD	RMSE	Cov.	Bias	SD	RMSE	Cov.
none	4	6	branching	-0.01	3.05	3.05	0.950	-0.01	0.76	0.76	0.943
			fitted fixed form	-0.07	3.71	3.70	0.950	-0.03	0.88	0.88	0.953
			EM, fitted forms	0.74	2.88	2.97	0.935	0.77	0.76	1.08	0.855
			EM, uniform forms	0.20	4.91	4.91	0.953	0.50	1.19	1.29	0.922
none	5	6	branching	0.00	3.06	3.06	0.940	-0.03	0.75	0.75	0.940
			fitted fixed form	0.04	3.37	3.37	0.950	-0.03	0.84	0.84	0.938
			EM, fitted forms	0.20	2.75	2.75	0.948	0.14	0.71	0.72	0.932
			EM, uniform forms	0.15	4.25	4.25	0.960	0.18	1.13	1.14	0.932
sex	4	4	branching	0.15	2.96	2.96	0.958	-0.05	0.77	0.77	0.935
			fitted fixed form	0.18	3.00	3.00	0.955	-0.04	0.78	0.78	0.943
			EM, fitted forms	0.51	2.79	2.84	0.955	0.37	0.69	0.78	0.920
			EM, uniform forms	-0.11	5.33	5.33	0.925	0.35	1.22	1.27	0.943
sex	5	7	branching	0.13	2.87	2.87	0.955	-0.08	0.74	0.74	0.940
			fitted fixed form	0.02	3.29	3.29	0.960	-0.04	0.84	0.84	0.950
			EM, fitted forms	0.03	2.75	2.75	0.953	0.02	0.68	0.68	0.930
			EM, uniform forms	-0.46	4.16	4.19	0.963	-0.01	1.00	1.00	0.938
education	4	5	branching	-0.15	2.85	2.86	0.970	0.01	0.77	0.77	0.953
			fitted fixed form	-0.25	3.42	3.42	0.973	0.01	0.86	0.86	0.958
			EM, fitted forms	-0.63	2.81	2.88	0.968	-0.45	0.74	0.87	0.920
			EM, uniform forms	-0.84	5.00	5.06	0.958	-0.44	1.25	1.32	0.925
education	5	6	branching	-0.15	2.83	2.83	0.970	0.01	0.77	0.77	0.948
			fitted fixed form	-0.04	3.02	3.01	0.978	-0.00	0.86	0.86	0.938
			EM, fitted forms	-0.36	2.51	2.54	0.963	-0.17	0.72	0.74	0.953
			EM, uniform forms	-0.65	4.14	4.18	0.978	-0.20	1.03	1.05	0.950
cost barrier	4	5	branching	0.13	3.18	3.18	0.935	-0.04	0.74	0.74	0.953
			fitted fixed form	-0.07	3.85	3.85	0.960	-0.02	0.94	0.94	0.943
			EM, fitted forms	0.79	2.99	3.09	0.920	0.68	0.70	0.98	0.870
			EM, uniform forms	0.45	4.85	4.87	0.930	0.69	1.13	1.32	0.922
cost barrier	5	8	branching	0.01	3.08	3.08	0.938	-0.03	0.74	0.74	0.953
			fitted fixed form	-0.11	3.79	3.78	0.943	0.01	0.93	0.93	0.953
			EM, fitted forms	0.48	3.01	3.04	0.932	0.33	0.68	0.75	0.935
			EM, uniform forms	0.41	3.71	3.73	0.963	0.45	0.89	0.99	0.945
poor health	4	4	branching	-0.21	3.21	3.21	0.948	-0.00	0.77	0.77	0.958
			fitted fixed form	-0.19	3.23	3.23	0.960	-0.00	0.79	0.79	0.955
			EM, fitted forms	-0.72	3.07	3.15	0.943	-0.60	0.74	0.95	0.885
			EM, uniform forms	-0.75	4.84	4.90	0.958	-0.69	1.19	1.38	0.940
poor health	5	7	branching	-0.28	3.26	3.27	0.948	-0.03	0.77	0.76	0.958
			fitted fixed form	-0.37	4.00	4.02	0.953	-0.07	0.99	0.99	0.948
			EM, fitted forms	-0.83	2.97	3.08	0.950	-0.69	0.74	1.01	0.882
			EM, uniform forms	-0.73	4.20	4.26	0.948	-0.62	1.05	1.21	0.890

Every value has the certificate $(v^{\mathsf{T}}u)^2/\max_\iota u^{\mathsf{T}}I_\iota u$, with the maximum exact over the class. With $k = K + 1$ the model is saturated and the program reproduces the design-based constants of the main solver (checked to 10^{-6} on a random law).

Reading. At the pairwise fits, pre-committed forms need 1.02–1.13 times the records of the best branching design (median 1.07), against 1.19–1.96 (median 1.50) without a model at the same laws; at the three-way fits, 1.08–1.40 (median 1.22). The optimal branching design within a model is not the rule's own tree: administering the rule attains the design-based constant $\mathrm{Var}(t)$ whatever the analysis, whereas the optimal design mixes trees and attains 0.71–0.84 of it under the pairwise model. Efficient pairwise analysis of the forms that are optimal for the design-based score needs a median 3% more records than the forms optimized for the model. For the runs in which the rule depends on at most b items, one form of these items is a complete record for the model and all classes coincide. The fitted laws are not the target laws: these constants describe a population that satisfies the model, and the audits below measure what happens on the actual target.

Laws with dependence. Theorem E.5 gives a polynomial bound at and near product laws and an exponential lower bound at pairwise laws with strong couplings. Table 20 computes the constants on two paths in between. On the first, the address contrast with $b = 3$ is embedded with its auxiliary features (a one-hot copy of the address and the four products; 14 features) and the coupling strength β of the pairwise law of part (e) is raised from zero. On the second, the address populations of Proposition E.3, in which the label agrees with the rule with a given probability, are projected on the pairwise and three-way models. Along the strongly coupled path the pairwise constants of both classes rise from $1/3$ towards their values without a model, and their ratio rises from 1.00 at $\beta = 0$ to 1.66 at $\beta = 6$, above the limit $2^{b-1}/b = 1.33$ guaranteed by the theorem

Table 19. Constants within log-linear models for the BRFSS screening trees whose rule depends on more than b items (18 of 24 runs; in the others one form of the rule's items is complete data). P_k is the maximum-likelihood fit of the order-k model to the target law on the label and the rule's K items. All entries are relative to $\mathrm{Var}_{P_k}(t)$, the constant of design-based branching. Design: $W_{\mathrm{fixed}}(P_k)$ without a model. Complete: efficiency bound for complete records in the model. Fixed, Branch: $W^{(k)}$ for forms of b of the K items and for depth-b trees, certified to a relative gap of 10^{-3} or better, except the three-way branching constants at $b = 6$ (*), where column generation was stopped after 15 minutes with a certified gap of at most 2.9%; the value shown is the upper bound, so the ratio is a lower bound that is at most that much too small. Ratio: fixed over branch.

				pairwise model, $k = 2$					three-way model, $k = 3$				
b	target	seed	K	Design	Complete	Fixed	Branch	Ratio	Design	Complete	Fixed	Branch	Ratio
3	other sex	2	4	1.34	0.79	0.92	0.84	1.09	1.34	0.95	1.21	0.99	1.23
4	random half	0	6	1.43	0.76	0.87	0.81	1.08	1.45	0.90	1.17	0.94	1.25
4	random half	1	5	1.40	0.76	0.87	0.81	1.07	1.40	0.91	1.15	0.94	1.22
4	random half	2	6	1.55	0.74	0.90	0.81	1.12	1.55	0.90	1.24	0.95	1.32
4	other sex	1	7	1.75	0.73	0.92	0.82	1.13	1.74	0.90	1.35	0.96	1.40
4	other sex	2	6	1.52	0.74	0.88	0.80	1.10	1.51	0.90	1.22	0.94	1.30
5	random half	0	6	1.19	0.76	0.81	0.77	1.05	1.18	0.90	0.99	0.91	1.08
5	random half	1	7	1.27	0.78	0.80	0.78	1.02	1.27	0.91	1.02	0.93	1.10
5	random half	2	8	1.49	0.75	0.87	0.79	1.10	1.51	0.89	1.17	0.93	1.26
5	other sex	0	7	1.25	0.76	0.81	0.78	1.04	1.26	0.88	1.02	0.90	1.14
5	other sex	1	8	1.51	0.70	0.80	0.75	1.07	1.53	0.84	1.13	0.89	1.28
5	other sex	2	7	1.40	0.70	0.79	0.74	1.07	1.43	0.86	1.08	0.89	1.21
6	random half	0	10	1.96	0.67	0.83	0.73	1.13	1.97	0.82	1.18	0.89*	1.33
6	random half	1	9	1.52	0.68	0.76	0.71	1.06	1.52	0.82	1.02	0.86*	1.19
6	random half	2	9	1.40	0.72	0.79	0.75	1.06	1.41	0.83	0.98	0.86*	1.15
6	other sex	0	10	1.59	0.74	0.80	0.76	1.05	1.60	0.85	1.05	0.90*	1.17
6	other sex	1	10	1.70	0.67	0.77	0.71	1.07	1.71	0.81	1.06	0.87*	1.22
6	other sex	2	10	1.55	0.68	0.77	0.72	1.07	1.55	0.83	1.04	0.88*	1.18

and below the factor 4 of part (a). At moderate coupling the gap is still small (1.02 at $\beta = 1$ and 1.10 at $\beta = 2$); it opens where the auxiliary features become nearly deterministic functions of the address and the data. At the pairwise fits of the noisy address populations the ratio stays at 1.00 at every accuracy, although the ratio without a model is 2^{b-1} at each of these populations: the association of the label with the data bits, which is what accuracy adds, does not bring the address structure into the pairwise tangent space. At the three-way fits the ratio is 2.46–3.35.

The conditional-independence model. The closed forms of Theorem E.5(b) need only the class-conditional degree-one coefficients of the rule, so they are available for every rule: the 60 rules of Tables 1 and 11 that depend on more than b items. At the conditional-independence fit of the target law, forms drawn without the label need a median 0.2% and at most 1.5% more records than the optimum with label-dependent forms on the BRFSS rules, and at most 14% on Spambase, where the classes differ more; the branching optimum without the label lies between the two. Under this working model, then, pre-commitment costs nothing that matters on any rule we trained, at caps four to six and for rules with up to 16 items. The model is cruder than the pairwise model: it ignores the associations among the items, which the target laws have. Its constants for forms are 0.58–0.72 of the variance of t on the diabetes trees (0.51–0.78 over all rules), and its bias for the true accuracy was not computed; we report it as the case in which the value of adaptivity is known in closed form, not as a recommended analysis.

Audits under three collection designs. For the BRFSS trees at caps four and five (two targets, three seeds) and audit sizes from 500 to 32,000, n target respondents are drawn with replacement and interviewed under three designs with the same cap: the fitted forms of Table 10, optimized for the design-based score on the source reference; forms optimized for the pairwise model at the pairwise fit of the source reference, over the b-subsets of the rule's items, with a 5% uniform floor; and branching, which administers a minimum-depth tree for the rule on its items and pads every path to b questions with the rule's lowest-index unasked items. Estimators are the design-based scores (the fitted tables on the fitted forms; the mean of t under branching) and the EM estimate under the pairwise and the three-way log-linear model for the label and the rule's items: EM from the uniform law on the records of each design, stopped when the estimate of the contrast changes by less than 10^{-5} or after 500 iterations. Where the observed-data likelihood has an interior maximizer this is the maximum-likelihood estimate; on designs that never observe some pairs of items together, such as the rule's tree and most fitted forms, it is an early-stopped iterate (see the paragraph on misspecification below) (400 replications per tree and size; no intervals; coverage at $n = 1{,}000$ and $16{,}000$ is in Table 18). Missingness is ignorable under all three designs, so the same likelihood applies. For the design-based scores the ratio of mean squared errors stays near the population ratio at every size (medians 1.40–1.55). For the three-way EM estimate the ratio between forms and branching records is 1.26–1.50

Table 20. Constants within log-linear models at laws with dependence, relative to $\mathrm{Var}_P(t)$. Left: the address contrast with $b = 3$ and the auxiliary features of Theorem E.5(e) (14 features), along the pairwise laws $P_\beta \propto e^{\beta \mathcal{Q}}$ from the uniform law ($\beta = 0$) towards the law on which the one-hot copy and the products are exact; the theorem gives a limit of at least $2^{b-1}/b = 1.33$ for the ratio, and the bound of part (a) is 4. Right: the address contrast with label $Y = d(X)\xi$, $P(\xi = 1) = \mathrm{acc}$, projected on the pairwise and three-way models by maximum likelihood; without a model the ratio is 2^{b-1} at each of these populations (Proposition E.3). Values certified to 10^{-3} or better, except entries marked *, where column generation over branching interviews was stopped by its budget and the ratio is a lower bound.

strong couplings, $k = 2$				pairwise and three-way fits					
β	Fixed	Branch	Ratio	b	acc	$k = 2$ ratio	$k = 3$ fixed	$k = 3$ branch	$k = 3$ ratio
0	0.333	0.333	1.00	3	0.5	1.00	2.500	1.000	2.50
0.5	0.350	0.350	1.00	3	0.6	1.00	2.516	1.000	2.52
1	0.432	0.425	1.02	3	0.7	1.00	2.564	1.000	2.56
1.5	0.618	0.595	1.04	3	0.8	1.00	2.636	1.000	2.64
2	0.839	0.761	1.10	3	0.9	1.00	2.675	1.000	2.68
3	1.223	0.892	1.37	3	0.97	1.00	2.463	1.000	2.46
4	1.465	0.934	1.57	4	0.5	1.00	2.750	1.000	2.75
6	1.635	0.987*	1.66	4	0.7	1.00	2.857	1.000	2.86
				4	0.9	1.00	3.187	1.000	3.19
				4	0.97	1.00	3.352	1.000	3.35

Table 21. The model in which the features are independent given the label (Theorem E.5(b)), at its fit to each target law: closed-form constants for every rule that depends on more than b items, including the rules whose design-based optimum was not computed. Ratio: label-blind forms over the optimum with label-dependent forms, which equals the branching optimum with the label and bounds the branching optimum without it from below; the ratio therefore bounds $W^{\mathrm{ci}}_{\mathrm{fixed}}/W^{\mathrm{ci}}_{\mathrm{branch}}$ from above. The other columns are relative to the variance of t under the fit: the constants of label-blind forms and of complete records within the model. Medians over runs [range].

target	runs	forms / branching, at most	forms	complete records
BRFSS, diabetes (12 items)	17	1.002 [1.000, 1.007]	0.62 [0.58, 0.72]	0.56 [0.47, 0.61]
BRFSS, heart disease	9	1.003 [1.000, 1.011]	0.64 [0.56, 0.71]	0.50 [0.40, 0.67]
BRFSS, stroke	9	1.002 [1.001, 1.015]	0.67 [0.56, 0.70]	0.59 [0.36, 0.65]
BRFSS, high blood pressure	9	1.001 [1.000, 1.004]	0.66 [0.58, 0.68]	0.61 [0.54, 0.64]
BRFSS, difficulty walking	7	1.004 [1.001, 1.007]	0.61 [0.56, 0.65]	0.55 [0.49, 0.61]
Spambase, spam	9	1.050 [1.021, 1.143]	0.61 [0.51, 0.78]	0.50 [0.35, 0.55]

on the fitted forms and 1.27–1.43 on the forms optimized for the pairwise model; for the pairwise model it is 1.00–1.07 and 0.94–1.01 up to 8,000 respondents (at most 2.25 and 2.30 for a single tree). These orders of magnitude agree with the population constants of Table 19, although the target laws do not satisfy the models. Branching here is the rule's own tree, not the branching design that is optimal within the model, which Table 19 puts a few percent below the optimal forms. At large sizes the comparison is governed by the bias of the model, on either kind of record: at 32,000 respondents the pairwise fit has a median absolute bias of 0.40 of its standard deviation on forms (0.17 at 1,000) and 0.33 on branching records (0.05 at 1,000), the pairwise fit to branching records is less accurate than the plain mean of t for 3 and 6 of the 12 trees at 16,000 and 32,000, and the ratio between the two designs ranges from 0.74 to 4.97 over trees. Against the design-based form score on the same fitted-form records, the pairwise EM estimate has the smaller mean squared error for every tree up to $n = 2,000$ and the larger for 3 and 5 of the 12 trees at 16,000 and 32,000 (RMSE up to 1.48 and 1.95 times larger); which trees these are cannot be read from the capped records. A model-assisted score, which uses a working model only to choose the tables of a pointwise-unbiased score, as in the smoothed pilot of Appendix F.5, cannot have a smaller variance than the oracle tables for the same form distribution; the advantage of the model-based estimate at small n is bought with the model's bias.

A model-assisted estimator on the audit's own records. The design-based form score of the audits takes its tables from 20,000 reference records. To see what a design-unbiased analysis achieves with the audit's records alone, the n records on the fitted forms are split into two folds; on each fold the pairwise model for the label and the rule's items is fitted by EM, the fitted law (with the items the rule ignores taken as independent and uniform, and 1% uniform mass) gives cell masses for every form, and the variance-optimal tables for the given form probabilities at these masses, $g_z = W_z F^{\mathsf{T}}(F W_z F^{\mathsf{T}})^{-1}(\pm\hat{t})$ with $W_z = \mathrm{diag}(q/\mathrm{mass}_z)$ and F the Fourier constraint matrix of the form program, score the records of the other fold. The tables satisfy the unbiasedness identities exactly and are independent of the records they score, so the estimator is unbiased at every population whatever the model. The cross-fitted estimator has the mean squared error of the reference-table score

Table 22. Simulated audits under three collection designs with the same cap (12 BRFSS trees: caps 4 and 5, two targets, three seeds; 400 replications per tree and size). Entries are ratios of mean squared errors, the finite-sample analogue of a records ratio: median over trees [minimum, maximum]. Design-based: the fitted form score against the mean of t under branching. EM-2 and EM-3: the EM estimate under the pairwise and three-way log-linear models (EM from the uniform law, stopped when the estimate of the contrast changes by less than 10^{-5} or after 500 iterations; this is the maximum-likelihood estimate where the likelihood has an interior maximiser and an early-stopped iterate elsewhere, Table 24), on the fitted forms (optimized for the design-based score) and on forms optimized for the pairwise model, each against the same model fitted to branching records (the rule administered and padded to b questions). Last column: the pairwise EM estimate on fitted forms against the design-based mean under branching.

	Design-based	EM-2: forms / branching records		EM-3: forms / branching records		EM-2 on forms /
n	forms / branching	fitted forms	model forms	fitted forms	model forms	design-based branching
500	1.49 [1.04, 1.78]	1.00 [0.93, 1.18]	0.94 [0.84, 1.09]	1.46 [1.02, 1.77]	1.32 [1.00, 1.72]	0.96 [0.83, 1.18]
1,000	1.52 [1.03, 1.89]	1.04 [0.90, 1.17]	0.98 [0.88, 1.15]	1.46 [0.94, 1.76]	1.34 [1.00, 1.51]	0.96 [0.79, 1.21]
2,000	1.40 [1.05, 1.73]	1.03 [0.92, 1.28]	1.00 [0.85, 1.26]	1.30 [1.06, 1.63]	1.34 [1.00, 1.69]	0.91 [0.84, 1.25]
4,000	1.51 [1.03, 2.03]	1.07 [0.93, 1.74]	1.01 [0.93, 1.62]	1.35 [1.01, 1.80]	1.27 [1.00, 1.56]	0.97 [0.89, 1.64]
8,000	1.48 [1.04, 1.80]	1.04 [0.86, 2.25]	1.00 [0.83, 2.30]	1.26 [1.04, 1.58]	1.27 [1.00, 1.47]	0.93 [0.82, 1.92]
16,000	1.55 [1.03, 1.91]	1.04 [0.94, 3.36]	1.03 [0.81, 3.37]	1.33 [1.02, 1.67]	1.29 [1.00, 1.54]	1.09 [0.89, 2.88]
32,000	1.54 [1.05, 1.80]	1.08 [0.74, 4.97]	1.05 [0.59, 4.95]	1.50 [1.02, 1.93]	1.43 [1.00, 1.93]	1.10 [0.91, 4.51]

Table 23. A cross-fitted model-assisted estimator on the fitted forms that uses only the audit's own records (12 BRFSS trees at caps 4 and 5; 200 and 100 replications per tree and size at $b = 4$ and $b = 5$). Entries are medians over trees [minimum, maximum]. MSE ratios are relative to the design-based form score with tables fitted on the 20,000 reference records; coverage is that of the nominal 95% normal interval from the sample variance of the scores; with 100 to 200 replications the Monte Carlo standard error of |bias|/SD is 0.07 to 0.10, and the two estimators are computed on the same records.

	MSE ratio	coverage		|bias|/SD	
n	cross-fitted / reference	cross-fitted	reference	cross-fitted	reference
1,000	1.01 [0.95, 1.04]	0.952 [0.920, 0.980]	0.960 [0.920, 0.990]	0.04 [0.01, 0.14]	0.05 [0.00, 0.14]
4,000	1.01 [0.97, 1.05]	0.960 [0.870, 0.980]	0.962 [0.890, 0.980]	0.07 [0.01, 0.24]	0.07 [0.01, 0.25]
16,000	1.00 [0.98, 1.04]	0.952 [0.925, 0.980]	0.952 [0.925, 0.980]	0.08 [0.03, 0.21]	0.08 [0.02, 0.21]

to within a few percent (median ratio 1.00–1.01 over the three sizes, at most 1.05 for a single tree), a bias within Monte Carlo error (|bias|/SD at most 0.24, against 0.25 for the reference-table score on the same records), and coverage between 0.870 and 0.980 over trees and sizes (0.890–0.990 for the reference-table score; the low values are for the same tree and size). A design-unbiased analysis therefore needs no external reference. It does not capture the small-sample advantage of the pairwise EM estimate, as expected: for given form probabilities its variance cannot fall below that of the oracle tables, and the advantage of the model-based estimate is bought with the model's bias.

The pairwise working model when it is wrong. The constants $W^{(2)}$ of Table 19 are evaluated at laws that satisfy the pairwise model. The target laws do not, so we also describe the pairwise maximum-likelihood estimator at the target law P itself, by the theory of misspecified likelihood with data missing by design (White, 1982; Golden et al., 2019). Under a design q over interviews, a pseudo-true law is a maximizer P_q^* in the model of the expected observed-data log-likelihood $\sum_\iota q_\iota \sum_c P(c) \log P^*(c)$ over the transcript-and-label cells c of ι; for a tree the cell is a function of the complete record, so this is the exact likelihood although the questions are chosen adaptively. At an interior maximizer the estimate of the contrast converges to $\Delta_q^* = \mathbb{E}_{P_q^*}[t]$, and $\sqrt{n}(\hat{\Delta} - \Delta_q^*)$ is asymptotically normal with the sandwich variance $\nabla^\mathsf{T} A^+ B A^+ \nabla$, where, for the sufficient statistics χ of the model (the main effects and pairwise products of the items and the label), $\nabla = \mathrm{Cov}_{P^*}(t, \chi)$, $A = \mathrm{Cov}_{P^*}(\chi) - \sum_\iota q_\iota \sum_c P(c) \mathrm{Cov}_{P^*}(\chi \mid c)$ is the expected negative Hessian and $B = \sum_\iota q_\iota \sum_c P(c)\, s_c s_c^\mathsf{T}$ with $s_c = \mathbb{E}_{P^*}[\chi \mid c] - \mathbb{E}_{P^*}[\chi]$ the covariance of the score. The asymptotic mean squared error for the true contrast at n records is $(\Delta_q^* - \Delta)^2 + \nabla^\mathsf{T} A^+ B A^+ \nabla / n$. Five designs are compared at the BRFSS diabetes trees of caps four to six that depend on more than b items (17 runs): the optimal design-based forms (F); the forms (MF) and the branching design (MB) that are optimal for the pairwise model at the pairwise fit of P; the rule's own tree (T); and a pair-covering comparator (PC), which mixes MF with a 10% uniform distribution over all b-subsets of the rule's items.

Identification. The pairwise model is determined by the joint laws of the pairs of variables, so under fixed forms its parameters are identified from incomplete records when every pair of items is observed together with positive probability; this is the classical requirement on split-questionnaire and booklet designs (Raghunathan & Grizzle, 1995; Rhemtulla & Little, 2012). The condition is sufficient, not necessary, and for trees, where a pair is observed only after particular earlier answers, we use it as a diagnostic. A tree observes two items together only on profiles whose padded path contains both, and forms optimized for a design-based score have no reason to cover all pairs. For such designs the expected log-likelihood can

be flat or increase towards the boundary of the model, and maximum likelihood need not be defined; the table records for every design how many pairs it leaves uncovered. An interior point below is a stationary point of the expected log-likelihood in the interior of the model; starts are counted as reaching the same one when their limits of the contrast agree to six decimals.

Computation. Even under pair coverage the expected log-likelihood is not concave in the parameters when data are missing. We maximize it by a quasi-Newton method followed by Newton steps from 6 starts (the uniform law, the pairwise fit of P and random parameters), stop when the norm of the expected score is below 10^{-9}, and call a start interior if it stops there with all cell probabilities above 10^{-10} and parameters below 25 in absolute value (largest score norm among the interior points: 10^{-10}). The table reports the interior maximizer with the largest objective and the range over interior maximizers; the equation $Ax = \nabla$ is solvable at each of them (relative residual at most 10^{-11}). Four findings are stable over the 17 runs.

Table 24. The pairwise working model as an estimator of the true accuracy contrast at the target law, which does not satisfy the model (BRFSS diabetes trees that depend on more than b items). Designs: F, the optimal design-based forms; MF, forms optimal for the pairwise model at its fit; PC, the pair-covering comparator (MF mixed with a 10% uniform distribution over all forms of the rule's items); T, the rule's own tree; MB, the branching design optimal for the pairwise model. Uncovered: number of pairs of the rule's items that the design never observes together (zero for MF, PC and MB in every run). The expected log-likelihood is maximised from 6 starts. Bias: pseudo-true contrast minus true contrast at the interior maximiser with the largest objective, in percent of the standard deviation of t. Variance: sandwich variance times n relative to $\mathrm{Var}_P(t)$; a range is shown when the starts reach several interior maximisers. A dash: no start reached an interior stationary point. †: an interior maximiser exists but a start with a larger objective went to the boundary. MSE ratio: (bias^2 + variance$/n$) of PC over MB.

				uncovered		bias (% of SD of t)			sandwich variance					MSE, PC / MB		
b	target	seed	K	F	T	MF	PC	MB	F	MF	PC	T	MB	1k	4k	16k
4	random half	0	6	2	2	0.8	0.5	0.3	0.97 [0.97, 1.03]	0.95	0.89	–	1.57	0.58	0.60	0.70
4	random half	1	5	0	0	0.6	0.6	0.2	0.89	0.87	0.87	0.91 [0.87, 0.91]	0.81	1.11	1.21	1.59
4	random half	2	6	2	3	0.4	0.4	0.1	1.18	0.90	0.91	–	0.81	1.14	1.19	1.37
4	other sex	1	7	5	6	-0.3	-0.3	-0.2	–	0.93	0.94	1.00†	0.82	1.16	1.17	1.24
4	other sex	2	6	2	3	-0.3	-0.3	-0.3	0.94 [0.91, 0.94]	0.89	0.90	0.98	0.81	1.11	1.10	1.06
5	random half	0	6	0	0	-0.0	0.0	0.2	0.82	0.81	0.82	0.83	0.80 [0.80, 0.81]	1.02	1.00	0.96
5	random half	1	7	1	1	1.1	1.1	0.7	–	0.81	0.82	–	0.79 [0.79, 0.84]	1.12	1.32	1.74
5	random half	2	8	2	5	-0.0	0.0	-0.3	0.92†	0.90	0.90	–	0.86 [0.86, 1.96]	1.03	0.99	0.86
5	other sex	0	7	0	3	-0.0	-0.0	0.2	0.82	0.81	0.82	–	0.78	1.05	1.04	1.00
5	other sex	1	8	4	5	0.2	0.2	–	–	0.80	0.81	–	–	–	–	–
5	other sex	2	7	1	0	-0.1	-0.1	-0.5	0.93	0.80	0.80	0.80 [0.80, 0.86]	0.77	1.01	0.92	0.69
6	random half	0	10	5	5	–	0.6	0.3	–	–	0.84	–	0.73	1.17	1.24	1.46
6	random half	1	9	4	1	1.5	1.4	0.5	–	0.76 [0.76, 0.76]	0.76	–	0.74 [0.74, 0.82]†	1.25	1.87	3.52
6	random half	2	9	3	1	0.4	0.4	0.5	0.81†	0.80 [0.80, 0.80]	0.80	–	0.75	1.06	1.05	0.99
6	other sex	0	10	6	5	-0.1	-0.2	-0.1	–	0.81	0.82	–	0.78 [0.78, 0.78]	1.05	1.06	1.12
6	other sex	1	10	4	5	0.4	0.4	-0.1	–	0.76†	0.77	–	0.78 [0.74, 0.78]	1.01	1.08	1.33
6	other sex	2	10	7	3	0.1	-0.4	-0.7	–	0.80 [0.80, 0.82]	0.78	–	0.79 [0.79, 0.99]	0.95	0.84	0.62

First, identification follows pair coverage. MF, PC and MB observe every pair of the rule's items together in every run; F does in 3 runs and T in 3. Over the designs and runs that cover all pairs, all 6 starts reach one interior maximizer in 42 of 57 cases and none is found in 2; over those that leave a pair uncovered, the counts are 1 and 20 of 28. Coverage with a small probability is not always enough in practice: MF, whose smallest pair probability can be below 1%, has one maximizer from all starts in 12 runs and MB in 9, whereas PC, with its 10% floor, has one in all 17. Second, on PC the bias is at most 1.4% of the standard deviation of t, or 0.64 accuracy points, the sandwich variance is between 0.76 and 0.94 of $\mathrm{Var}_P(t)$, below the variance of the design-based mean on the tree, and the floor costs little: the variance of PC is 0.94–1.02 times that of MF where both are defined. Third, where PC and MB both have an interior maximizer (16 runs), the ratio of their sandwich variances is 0.57–1.15 (median 1.04), MB, optimal for the fitted law, has the larger variance at the true law in 3 runs, and with the bias included the mean squared error of PC relative to MB is a median 1.06, 1.07 and 1.09 at 1,000, 4,000 and 16,000 records (ranges 0.58–1.25 and 0.62–3.52 at the two ends). Fourth, the variance computed from the model's information matrix is within a factor 1.01 of the sandwich variance on PC, but up to 2.08 times too small on other designs, which contributes to the under-coverage of model-based intervals in Table 18 beyond the effect of bias. The design-based ratio at the same laws is 1.18–1.98 (median 1.52). The asymptotic bias agrees with the simulated audits: for the trees of caps four and five the calculation puts the bias of PC at a median 0.47 standard deviations at 32,000 records, and the EM estimate on the model forms of the audits, which have a 5% floor, has 0.42.

F.9. When Asking Changes the Answer

Let P describe latent complete records and let K_T be the distribution of the recorded transcript given a latent record under form T. A frozen score then estimates $\mathbb{E}_{P,q,K} H_T(O_T)$, which can differ from Δ; a branching policy has its own sequential kernel. We compute the resulting biases exactly on the screening trees of Appendix F.3. The answer at query position $j = 1, \dots, b$ flips independently with probability $\epsilon(j-1)/(b-1)$, so the first answer is exact and the last flips with probability ϵ; labels are unchanged. Branching routes on the recorded answers; fixed forms ask their coordinates in increasing index order and use the fitted design's scores. Summing over all 2^b flip patterns gives the exact mean and second moment of each design at the target law. This is an illustrative position model, not an estimated response-error process.

Table 25. Position effects on the screening trees: median absolute bias in units of the clean branching standard error at $n = 1000$, and the share of runs in which branching keeps the smaller RMSE at $n = 1000$ (the last answer flips with probability ϵ).

Data	ϵ	branching	fitted form	oracle form	branching RMSE smaller
BRFSS	0.01	0.15	0.21	0.22	24/24
BRFSS	0.03	0.45	0.64	0.65	24/24
BRFSS	0.05	0.75	1.08	1.09	24/24
BRFSS	0.1	1.54	2.17	2.20	24/24
Thyroid	0.01	0.02	0.24	0.23	24/24
Thyroid	0.03	0.06	0.71	0.70	24/24
Thyroid	0.05	0.11	1.18	1.17	24/24
Thyroid	0.1	0.21	2.37	2.34	23/24

F.10. Cost Regimes Under a Relaxable Cap

Suppose each feature costs one unit and each label c units, and the cap may be relaxed. A complete record costs $c + m$ and attains the full-information variance σ_P^2; a capped record costs at most $c + b$. For a screening tree, branching attains σ_P^2 at cost $c + b$ and therefore dominates complete records. Complete records are cheaper than the best fixed design exactly when $(c + m)/(c + b) < W_{\text{fixed}}/\sigma_P^2$, that is, when $c > (m - \zeta b)/(\zeta - 1)$ with $\zeta = W_{\text{fixed}}/\sigma_P^2$. For the BRFSS trees ($m = 12$) the median ζ is 1.49, 1.35, 1.58 at $b = 4, 5, 6$, so complete records undercut the best fixed design once a label costs more than about 12, 15, 4 feature questions, respectively, while branching is never undercut. Under an immutable cap the complete-record alternative is unavailable.

G. Proper Scores and Pairwise Targets

The score-design method applies to a bounded known record-level target $t(x, y)$, without requiring that it be a randomized classification gain. For a label observed separately, write

$$t(x, y) = a(x) + yc(x), \qquad a(x) = \frac{t(x, 1) + t(x, -1)}{2}, \quad c(x) = \frac{t(x, 1) - t(x, -1)}{2}.$$

On a finite domain, identification under arbitrary joint populations is equivalent to $a, c \in \mathcal{H}_{\mathcal{A}}$. On the full binary cube with cap b, this is $\deg(a), \deg(c) \le b$. Indeed the observation span on (x, y) is $\mathcal{H}_{\mathcal{A}} \otimes \operatorname{span}\{1, y\}$: a table may have separate entries for the two labels. Its orthogonal complement gives indistinguishable populations whenever the target lies outside that span. In particular the common identification criterion for fixed forms and branching still applies.

For $z = (y + 1)/2$ and $d = p_1 - p_0$, the improvement in Brier loss is

$$t_{\text{Br}}(x, y) = (p_0 - z)^2 - (p_1 - z)^2 = d(1 - p_0 - p_1) + yd.$$

Thus degree at most r for each probability function implies degree at most $2r$ for this target's feature-only component and at most r for its label component. The bound can be strict because terms cancel. The actual degree, or a certified approximation residual, determines the necessary cap. Doubling is attainable: take $p_0 = 1/2$ and $p_1 = 1/2 + \gamma(x_1x_2 + x_3x_4)$, with $0 < \gamma \le 1/4$. The accuracy contrast has degree two, whereas the Brier feature-only component is $-2\gamma^2(1 + x_1x_2x_3x_4)$ and has degree four. For a common depth-b tree whose leaf values determine both probabilities, the complete loss difference is still computable along that same path.

For log loss, clip each probability to a fixed interval $[\eta, 1-\eta]$, $0 < \eta < 1/2$, before defining the target. Then

$$a_{\log} = \tfrac{1}{2} \log \frac{p_1(1-p_1)}{p_0(1-p_0)},$$
$$c_{\log} = \tfrac{1}{2} \log \frac{p_1(1-p_0)}{p_0(1-p_1)}.$$

These formulas estimate improvement of the clipped rules. A logarithm of an ensemble average can introduce interactions of every order, even when each probability has a low-degree representation. A common tree partition again avoids that issue because the transformed values are constant on leaves.

The full-form program in Section E.5 replaces its right-hand side by $\widehat{t}_y(S)$ for each label. Its pointwise identity, population-specific variance objective, and invariance of unbiasedness to calibration error remain unchanged. The symmetric-label minimax formula $V(d, \mathcal{A})$ and its sharp ratio were proved for $Yd(X)$; this extension of the estimator does not assert that the identical formula is the global minimax coefficient for every proper score. The path simulation does preserve any bounded target's conditional mean and multiplies its conditional second moment by at most the number of padded path sets.

AUC needs a pair design. Let $X^+ \sim P(X \mid Y = 1)$ and $X^- \sim P(X \mid Y = -1)$ be independent. For a frozen score s, define $k_s(x^+, x^-) = \mathbf{1}\{s(x^+) > s(x^-)\} + \frac{1}{2}\mathbf{1}\{s(x^+) = s(x^-)\}$. The AUC difference is $\mathbb{E}[k_{s_1} - k_{s_0}]$ under the product of these two conditional laws. Disjoint labeled case–control pairs give iid pair targets; one can design unbiased scores on pair transcripts, retaining a separate cap on each member. This differs from a single-record target: using all case–control pairs introduces U-statistic dependence, and the sharp single-record coefficient ratio cannot be asserted for that design. Class prevalence, label-screening cost and the availability of both classes must also enter an acquisition comparison. We establish the record-level Brier/log-loss extension here and leave pair-design optimality unclaimed.

H. Reproducibility and Research Methods

The anonymous supplement contains the code of the solver and of every experiment, per-task results with certificates, logs, and the generators of all tables and figures. Source data are public and obtained separately; no individual survey records are redistributed.

Checks. The solver reproduces, on every task that overlaps with an independent dense implementation, the fixed constants, the explained-energy ends of the bracket and the certified branching constants; every interview it generates is checked to partition the profile cube (for $m \le 15$). Exact rational checks cover the hybrid-budget inequalities of Proposition C.9, small conic programs are solved independently, and the product counterexample of Appendix C.6 has a rational certificate checked on all 256 profiles. The potential inequality (D.3) is checked exhaustively on the pointer tree with two pointer levels and on random states for three and four levels (Appendix D.1); the label-dependent programs are certified by the same primal and dual bounds as the others; the model version of the solver reproduces the design-based constants when the model is saturated, and the inequalities of Theorem E.5(a) and (c) and the information identity and closed forms of (b) are checked by enumeration on random laws, trees and directions, as are the closed forms of $L_k(b)$ and the coefficient identities behind the bound $U_3(b)$ (for $b \le 13$, by a second, independently written set of scripts); and the generator of the tables asserts that runs computed by different scripts agree on their common quantities. These finite checks support the implementations and derivations; they do not replace the proofs.

Research methods. A generative language model participated in formulating research questions, deriving and checking proof arguments, implementing analyses, and drafting the manuscript; the authors remain responsible for all content.