



Sleep Patterns of College Students at a Public University

LeAnne M. Forquer PhD , Adrian E. Camden BS , Krista M. Gabriau BS & C. Merle Johnson PhD

To cite this article: LeAnne M. Forquer PhD , Adrian E. Camden BS , Krista M. Gabriau BS & C. Merle Johnson PhD (2008) Sleep Patterns of College Students at a Public University, Journal of American College Health, 56:5, 563-565, DOI: [10.3200/JACH.56.5.563-565](https://doi.org/10.3200/JACH.56.5.563-565)

To link to this article: <http://dx.doi.org/10.3200/JACH.56.5.563-565>



Published online: 06 Aug 2010.



Submit your article to this journal [↗](#)



Article views: 1881



View related articles [↗](#)



Citing articles: 39 View citing articles [↗](#)

Clinical and Program Note

Sleep Patterns of College Students at a Public University

LeAnne M. Forquer, PhD; Adrian E. Camden, BS; Krista M. Gabriau, BS;
C. Merle Johnson, PhD

Abstract. **Objective:** The authors' purpose in this study was to determine the sleep patterns of college students to identify problem areas and potential solutions. **Participants:** A total of 313 students returned completed surveys. **Methods:** A sleep survey was e-mailed to a random sample of students at a North Central university. Questions included individual sleep patterns, problems, and possible influencing factors. **Results:** Most students reported later bedtimes and rise times on weekends than they did on weekdays. More than 33% of the students took longer than 30 minutes to fall asleep, and 43% woke more than once nightly. More than 33% reported being tired during the day. The authors found no differences between freshmen, sophomores, juniors, seniors, and graduate students for time to fall asleep, number of night wakings, or total time slept each night. **Conclusions:** Many students have sleep problems that may interfere with daily performance, such as driving and academics. Circadian rhythm management, sleep hygiene, and white noise could ameliorate sleep difficulties.

Keywords: college students, night wakings, sleep, sleep aids, sleep deprivation

Adolescents and young adults, including college students, appear to be one of the most sleep-deprived groups in the United States.¹⁻³ Individuals in this age group require about 9 hours of sleep each night; however, most receive only 7 to 8 hours.¹ This sleep deprivation can have detrimental effects on performance, including driving⁴ and academics.⁵ According to Carskadon,¹ 55% of sleep-related accidents involve individuals younger than 25 years. In a survey by the American College Health Association⁵ involving students from 33 universities across the United States, researchers examined the top impediments to academic performance. Both men (23%) and women (25%) rated sleep difficulties as the third most common impedi-

ment, after stress and illness such as colds, flu, or sore throats. In this study we examined college students' sleep patterns to identify problem areas and potential solutions.

METHODS

Participants

Graduate and undergraduate students from a North Central university participated in an e-mail survey. We randomly selected these students from the approximately 20,000 currently enrolled. The Dean of Students approved the release of their university-issued e-mail addresses by the Registrar's office. The university's institutional review board approved this project.

Measures

To identify students' potential problem areas and factors influencing these problems, we conducted a sleep survey that included questions from the Pittsburgh Sleep Quality Index⁶ and the Sleep Hygiene Test.⁷

Procedure

We attached the survey to an e-mail that explained the study's purpose. We sent this e-mail to the university-issued e-mail addresses of students randomly selected to participate. All information was confidential; only a number identified participants. Students who decided to participate were instructed to complete the attached survey and return it via e-mail or campus mail. We sent all participants a reminder 2 weeks later with another copy of the survey. We conducted the study during the beginning of the spring semester, when sleep deprivation should be lowest.

RESULTS

We e-mailed surveys to 2,024 students; 44 of the surveys could not be delivered and 241 were completed. We e-mailed a second survey 2 weeks later, and this time 43

*At the time of the study, all authors were with the Department of Psychology at Central Michigan University in Mount Pleasant.
Copyright © 2008 Heldref Publications*

could not be delivered and 72 more were completed. A total of 313 students returned completed surveys, although 23 more surveys were returned without attachments. The sample of participating students was 62% female, 90% Caucasian, 93% unmarried, and 87% undergraduate; their mean age was 21.4 years ($SD = 4.3$). Most participants (90%) had roommates, including spouses (7%). Demographics were representative of the university.

The survey asked the students about their typical sleep patterns (see Table 1). When asked about their typical sleeping situation, most participants reported sleeping with roommates, although in separate beds. Bedtimes on weekends were more than 1 hour later and rise times were more than 2 hours later; thus, most participants slept more on weekend nights ($M = 8.6$ hours, $SD = 1.5$) than they did on

weekdays ($M = 7.2$ hours, $SD = 1.2$). Furthermore, women averaged 8 hours of sleep compared with 7.7 for men ($F[1, 312] = 5.27, p < .01$). Participants took an average of 25 minutes to fall asleep ($SD = 20$). An analysis of variance showed that women averaged 27.3 minutes to fall asleep, compared with 21.2 for men ($F[1, 312] = 6.93, p < .01$). We observed no sex differences in the number of night wakings ($M = 1.7, SD = 1$), nor any class differences (freshmen through graduate students) in the number of night wakings, time to fall asleep, or hours of nightly sleep. The most common reasons for night wakings included hearing noise from others (41%), going to the bathroom (40%), and being worried about something (33%).

When asked what a typical night's sleep consisted of, most participants answered that they slept all night (26%), had 1 waking (26%), or had 2–3 wakings (21%). The participants were also asked about sleep aids; half reported using sounds such as fans or music. Less common sleep aids included medication (6%). When asked how they felt during waking hours, 58% of participants reported being tired in the morning but okay once they got going. However, more than 33% reported that they started out energetic and then got tired or were tired all day.

COMMENT

College students reported later bedtimes and wake times on weekends than on weekdays. Of the participants, 33% took more than 30 minutes to fall asleep and 43% woke more than once a night. These data suggest sleep difficulties consistent with research of the National Sleep Foundation,⁸ which reported that more than 40% of Americans have difficulty falling asleep or have night wakings.

Our results support American College Health Association⁵ survey findings on college students' sleep difficulties, including long sleep latencies, short sleep time, and frequent night waking. Sleep problems may be worse than these self-reported responses, as students may give socially desirable answers such as not noting sexual encounters or alcohol abuse before bedtime. Thus, this survey may be limited by underreporting.

Improving sleep may enhance academic performance. Possible strategies include circadian rhythm management, sleep hygiene, and white noise.^{9,10} The circadian rhythm is the 24-hour day–night cycle that influences quantity and quality of sleep.⁹ The more stable and consistent this circadian rhythm, the better a person sleeps. This implies that individuals should go to bed and wake at the same time every day, including on weekends.¹⁰ Clearly, students in this sample disrupted circadian rhythms when weekday sleep is compared with weekend sleep. Improvements in sleep hygiene—including limiting naps to less than 1 hour, using beds only for sleeping (no reading, TV, or homework), and making sure the bedroom is comfortable^{9,10}—also promote sleep. White noise, which is continuous sound covering the entire range of human hearing from 20 to 20,000 Hertz (or approximations to white noise such as fans or humidifiers), also could improve sleep.^{9,10} College students need

TABLE 1. Participants' Sleep Patterns

Pattern	Value
Typical sleeping situation (%)	
Roommate in different room	41
Share a room	31
Share a bed	16
Alone	12
Mean bedtime (AM)	
Weekday	12:24
Weekend	1:54
Mean rise time (AM)	
Weekday	8:12
Weekend	10:30
Why do you awaken during the night? (Check all that apply)	
Noise from others	41
Need to go to bathroom	40
Worried about something	33
Bad dreams	20
Pain	9
Muscle spasms	7
Do you typically use sound to help you sleep? (%)	
Yes	47
No	53
If yes, what type of sound do you use? (%)	
Fan	55
Music	34
Humidifier	22
Television	8
Sound machine	3
Do you use any other aids to help you sleep? (%)	
Yes	10
No	90
If yes, what aids do you use? (%)	
Medication	60
Reading	13
Relaxation exercises	13
Earplugs	7
Alcohol	3

Note. The standard deviations for mean bedtimes and mean rise times are as follows: bedtime, $SD = 1.4$ and $SD = 1.6$, respectively, weekday and weekend; rise time, $SD = 1.4$ and $SD = 2.0$, respectively, weekday and weekend.

to address sleep problems with better sleep management, which may improve academic performance and driving.

NOTE

For comments and further information, address correspondence to Dr LeAnne M. Forquer, Division of Counselor Education and Psychology, Delta State University, Box 3142, Cleveland, MS, USA 38733. (e-mail: lforquer@deltastate.edu).

REFERENCES

1. Carskadon MA. *Adolescent Sleep Patterns: Biological, Social, and Psychological Influences*. New York: Cambridge University; 2002.
2. Pilcher JJ, Walters AS. How sleep deprivation affects psychological variables related to college students' cognitive performance. *J Am Coll Health*. 1997;46:121–126.
3. Tsai L, Li S. Sleep patterns in college students: gender and grade differences. *J Psychosom Res*. 2004;56:231–237.
4. Subramanian R. *Motor Vehicle Traffic Crashes as a Leading Cause of Death in the US, 2002*. <http://nhtsa.gov/people/crash/LCOD/Index.htm>. Accessed April 27, 2005.
5. American College Health Association. The American College Health Association–National College Health Assessment (ACHA–NCHA), spring 2003 reference group report. *J Am Coll Health*. 2005;53:199–210.
6. Buysse DJ, Reynolds CF, Monk TH, et al. The Pittsburgh Sleep Quality Index: a new instrument for psychiatric practice and research. *Psychiatry Res*. 1989;28:193–213.
7. *Sleep Hygiene Test–Abridged*. http://www.discoveryhealth.queendom.com/sleep_hygiene_abridged_access.html. Accessed September 22, 2004.
8. National Sleep Foundation. *The Basics of Sleep*. http://www.harvestmoonstudio.com/assets/SLEEP_HTML/pub_newsmaker.htm. Accessed September 28, 2005.
9. Breus M. *Sleep Dos and Don'ts: Sleep Hygiene Solutions for Better Sleep*. <http://my.webmd.com/content/Article/62/71839.htm>. Accessed June 2, 2005.
10. Dement WC. *How To Sleep Well*. <http://www.stanford.edu/~dement/howto.html>. Accessed June 2, 2005.

Call For Historical Photographs

The Executive Editors of the *Journal of American College Health* are currently seeking historical photos of college or university health centers and health workers for publication in JACH. JACH is the professional journal for the American College Health Association □ (www.acha.org). Founded in 1920, ACHA is the principal advocate and leadership organization for college and university health. JACH has been published since 1958, originally as *Student Medicine*.

Should the editors choose the photograph for publication, it will be published either inside the issue or on the cover, per the editors' discretion.

If you are interested in submitting, please visit <http://www.heldref.org/jach.php> for further details on submitting a photograph via the journal's online system. You may also contact the managing editor via e-mail at jach@heldref.org.

