1. The probability of a leap year selected at random contain 53 Sunday is:								
(a) 53/366 (b) 1/7 (c) 2/7 (d) 53/365								
2. A bag contains 3 red and 2 blue marbles. A marble is drawn at								
random. The probability of drawing a black ball is:								
(a) 3/5 (b) 2/5 (c) 0/5 (d) 1/5								
3. The probability that it will rain tomorrow is 0.85. What is the								
probability that it will not rain tomorrow								
(a) 0.25 (b) 0.145 (c) 3/20 (d) none of these								
4. What is the probability that a number selected from the numbers	5							
(1, 2, 3,,15) is a multiple of 4?								
(a) 1/5 (b) 4/5 (c) 2/15 (d) 1/3								
5. What are the total outcomes when we throw three coins?								
(a) 4 (b) 5 (c) 8 (d) 7								
6. The probability that a prime number selected at random from th	e							
numbers (1,2,3,35) is :								
(a) $12/35$ (b) $11/35$ (c) $13/35$ (d) none of these								
7. The sum of the probability of an event and non event is :								
(a) 2 (b) 1 (c) 0 (d) none of these.								
8. The following probabilities are given; choose the correct answe	r							
for that which is not possible.								
(a) $0.15$ (b) $2/7$ (c) $7/5$ (d) none of these.								
9. If three coins are tossed simultaneously, than the probability of								
getting at least two heads, is:								
(a) $1/4$ (b) $3/8$ (c) $\frac{1}{2}$ (d) $1/8$								
10. A letter is chosen at random from the letters of the word								
ASSASSINATION. The probability that the letter chosen has:								
(a) 6/13 (b) 7/13 (c) 1 (d) none of these								
11. A dice is thrown. Find the probability of getting an even number.								
(A) 2/3 (B) 1 (C) 5/6 (D) 1/2								
12. Two coins are thrown at the same time. Find the probability of								
getting both heads.								
(A) 3/4 (B) 1/4 (C) 1/2 (D) 0								
12 Two dies are thrown simultaneously. The probability of gotting	_							

13. Two dice are thrown simultaneously. The probability of getting a sum of 9 is:

(A) 1/10	(B) 3/10	(C) 1/9	(D) 4/	'9		
	rds are numb orime number.	ered from 1 to 1	00. Find the	probability of		
(A) 3/4	(B) 27/50	(C) 1/4	(D)	29/100		
of drawing				f the probability n the number of		
		contains 12 defe		One hulb ic		
taken out	at random fro	m this box. The				
non-defec (A) 143/15	tive bulb is: 50 (B) 14	47/150 (C)	) 1/25	(D) 1/50		
mixed tho	roughly. One o	numbers 2 to 10 card is drawn fronumber on card (C) 3/10	om this box r	andomly, then square.		
18. What (A) 1/7	17-3	lity of getting 53 (C) 2/7	3 Mondays in (D) 7			
probability	of getting a l	n a well shuffle king of red suit. (C) 7/52 (D		cards. Find the		
20. A game of chance consists of spinning an arrow which is equally likely to come to rest pointing to one of the number 1,2,312 ,then the probability that it will point to an odd number is:  (A) 1/6 (B) 1/12 (C) 7/12 (D) 5/12						
21. A game consists of tossing a one rupee coin 3 times and noting its outcome each time. Aryan wins if all the tosses give the same result i.e. three heads or three tails and loses otherwise. Then the probability that Aryan will lose the game.  (A) $3/4$ (B) $1/2$ (C) 1 (D) $1/4$						

22. Riya and Kajal are friends. Probability that both will have the same birthday is the same birthday is:								
(A) 364/365	(B) 31/365	(C) 1/365	(D) 1/133225					
23. A number $x$ is chosen at random from the numbers -2, -1, 0, 1, 2. Then the probability that $x^2 < 2$ is? (A) $1/5$ (B) $2/5$ (C) $3/5$ (D) $4/5$								
24. A jar contains 24 marbles. Some are red and others are white. If a marble is drawn at random from the jar, the probability that it is red is $2/3$ , then the number of white marbles in the jar is:  (A) 10 (B) 6 (C) 8 (D) 7								
25. A number is selected at random from first 50 natural numbers. Then the probability that it is a multiple of 3 and 4 is: (A) $7/50$ (B) $4/25$ (C) $1/25$ (D) $2/25$								
26. Consider a dice with the property that that probability of a face with n dots showing up is proportional to n. The probability of face showing 4 dots is?								
a) $\frac{1}{7}$	b) $\frac{5}{42}$	c) $\frac{1}{21}$	$\frac{4}{21}$					
		n 5 one day match	es are 50, 70, 82,					
	e standard devia b) 25.49	c) 25.29	d) 25.69					
28. Find median and mode of the messages received on 9 consecutive days 15, 11, 9, 5, 18, 4, 18, 13, 17.								
a) 13, 15	b) 13, 18		d) 13, 16					
29. A coin is tossed up 4 times. The probability that tails turn up in 3 cases is								
a) $\frac{1}{2}$ 30. X is a varia	b) $\frac{1}{3}$ te between 0 and	c) <sup>1</sup> / <sub>4</sub> d 3. The value of E c) 27						
31. The random variables X and Y have variances 0.2 and 0.5 respectively. Let Z= 5X-2Y. The variance of Z is?								

a) 3	b) 4	c) 5	d) 7	
32.Out of topic probability?		lues, which	one is not possib	le in
a) P(x) = 1 c) P(x) = 0.5	b)∑x I	P(x) = 3 () = -0.5		
	2 and E(z) = 4, b) 6			fficient data
34.The cov	ariance of two	independent	t random variable	is
a) 1	b) 0	c) - 1	d) Und	efined
<b>35.If Σ P(x</b> ) a) 0	) = <b>k</b> <sup>2</sup> – <b>8 then,</b> b) 1	the value of c) 3	1.00	ıfficient data
	<b>0.5 and x = 4, t</b> b) 0.5	t <b>hen E(x) = ?</b> c) 4	d) 2	
37.ln a disc is always?	rete probability	y distributio	ո, the sum of all բ	orobabilities
a) 0	b) Infinite	c) 1	d) Unde	efined
38.If the provided variance.	obability of hit	ting the targ	et is 0.4, find me	an and
a) 0.4, 0.24	b) 0.6, 0	0.24	c) 0.4, 0.16	d) 0.6, 0.16
	% and if 10 bor	nbs are dro	ped from a place oped, find mean a 4, 0.16	
a) 2		c) 8	d) 1 r standard norma	I distribution?

						d variance and varian		
a) E(X) b) E(X2)			(2)	c) $E(X2) - (E(X))2$			d) (E(X))2	
43.I	Mean of	a random	variable	X is giv	en by			
a) E	(X)	b) E(X2	2)	c) E(X2)	- (E(X))	2	d) (E(X))2	
44.N a) 0		a constant		c) a/2	•	d) 1		
<b>45.Variance of a constant 'a' is</b> . <b>a) 0</b> b) a c) a/2 d) 1								
46.F	ind the r	mean and	variance	of X?				
	X	0	1	2	3	4		
	f(x)	1/9	2/9	3/9	2/9	1/9		

c) 2, 2/3

d) 3, 2/3

الممسوحة ضوئيا بـ CamScanner

47. Find the expectation of a random variable X?

b) 3, 4/3

	X	0	1	2	3		
	f(x)	1/6	2/6	2/6	1/6		
C	).5		b) 1.5			c) 2.5	d) 3.5

48. In a Binomial Distribution, if p, q and n are probability of success, failure and number of trials respectively then variance is given by

\_\_\_\_\_

a) 2, 4/3

- a) np
- b) npq
- c) np2q

d) npq2

49. If 'X' is a random variable, taking values 'x', probability of success and failure being 'p' and 'q' respectively and 'n' trials being conducted, then what is the probability that 'X' takes values 'x'? Use **Binomial Distribution.** 

- a) P(X = x) = nCx px qx
- b) P(X = x) = nCx px q(n-x)
- c) P(X = x) = xCn qx p(n-x)
- d) P(x = x) = xCn pn qx

50. If 'p', 'q' and 'n' are probability pf success, failure and number of trials respectively in a Binomial Distribution, what is its Standard **Deviation?** 

- a)  $\sqrt{np}$
- b) $\sqrt{pq}$  c) (np)2
- d)  $\sqrt{npq}$