

AGEL J020613–011417 — Drafting Fact Sheet for the Methods & Results Sections

Facts, decisions, and provenance keyed to the paper outline — last revised 2026-06-08
(post-M12)

Compiled for R. Córdova Rosado from project notes, memory, METHODS_AND_SYSTEMATICS.md, and TESTS_AND_DIAGNOSTICS.md

2026-06-08

How to use this document

This is a drafting reference, not prose for the paper. Every bullet is a fact or decision we have already made, with its source file in [brackets] so you can verify it. Where a number is superseded, the current value is given and the stale one flagged. Three gaps are flagged up front — they are not in this repo and you must source them elsewhere before drafting those paragraphs:

- G1 — Lens modeling (Ferrami et al.). No BPL / point-mass / $M(<r_{\text{break}})$ / joint-posterior / PSF / shear / subhalo content exists in this kinematics+photometry repo. Source from the lens-modeling repo (7 Documents/AGEL/202509_DESJ0206_modeling/ or 20251112_DESJ0206_Pyauto_PRONTO/) or the Ferrami draft directly.
- G2 — X-ray / “truly quiescent” statement. No X-ray, Chandra, or explicit quiescence data anywhere in the read files. The galaxy is only ever described as a “passive elliptical.” You need an external source if you want the X-ray quiescence claim in §Results.
- G3 — “We do not account for peculiar velocities.” This sentence is not currently written in any methods doc. It is a true statement about our analysis (we use the line-fit systemic z directly) — just add it; nothing to reconcile.

Headline numbers (current, 2026-06-08, post-M12; emitted by `scripts/paper_values.py` → `results/PAPER_VALUES.json`):

- $\sigma_e(<R_e) = 269.62 \pm 13.27$ km/s (asym $-13.45 / +13.10$); often rounded 270 ± 13 km/s. 2026-06-08: was ± 11.77 ; the D7 R_e -source systematic (± 6.13) was folded into the budget (M12, Task 5, nb15).
- $\log(M_\star/M_\odot) = 11.16 \pm 0.08$ (stat) $+0.31$ (sys) at 10% flux errors [11.04 ± 0.14 (stat) $+0.32$ (sys) at 20%] — NEW principled arc masking (F200LP-located + IR-extended, raw aperture). The one-sided +sys is the Sérsic fill-in of deflector light hidden under the arc, reaching $\log M_\star \approx 11.46$. Supersedes $11.33 +0.07/-0.09$ (older expert-aperture value, smaller masks; now a mid-range cross-check). See §2.1.1b / §3.2 and NOTES_photometry_mask_systematics_2026-05-29.md.
- $R_e = 2.305'' = 16.23$ kpc.
- $z_{\text{deflector}} = 0.67564$; $z_{\text{source}} = 1.302$.

Cosmology: FlatLambdaCDM, $H_0 = 70$ km/s/Mpc, $\Omega_m = 0.3$; $1'' \approx 7.04$ kpc (7043.5 pc/arcsec), $D_A = 1452$ Mpc at $z = 0.67564$. [reference_hst_jwst_data_properties.md]

Staleness warning: the *body* of METHODS_AND_SYSTEMATICS.md (dated 2026-05-18) still prints pre-M11 numbers (254.85 / 268.98, total ± 18) in its narrative prose. Its own §0 table and CLAUDE.md carry the current headline 269.62 ± 13.27 (post-M12, 2026-06-08). Always quote the current value (canonical source: `results/PAPER_VALUES.json`).

§1 Data and observations

All values below are read directly from the FITS headers of the analysis files (verified 2026-06-08), not from memory. Target: AGEL J020613–011417 = DES J0206–0114, ICRS RA = 31.55611°, Dec = −1.23817° (HST RA_TARG/DEC_TARG); deflector $z = 0.67564$, source $z = 1.302$. The field is the massive cluster ACT-CL J0206.2–0114 (ACT DR5, Hilton et al. 2021) — the JWST pointing target.

Facility	Inst. / mode	Band(s)	Program (PI)	UT date	Exp.	Pixel scale
Keck II	KCWI — KCRM red (RL) + KCB blue (BL), Medium slicer, 2×2	RL 5625–8941 Å (σ_e); BL λc 4500 Å	U204 + K409 + U002 (3-night combine)	2024-12 → 2025-11	≈180 min RED + ≈198 min BLUE (36 × 300 s RED)	0.300"/spaxel
HST	WFC3/UVIS	F200LP	16773 (Glazebrook)	2022-07-14	600.0 s (NDRIZIM 4)	0.050"/pix
HST	WFC3/IR	F140W	16773 (Glazebrook)	2022-07-14	597.7 s (NDRIZIM 3)	0.080"/pix
JWST	NIRCam SW, module B	F150W2 (CLEAR)	05594 (Mahler)	2024-08-27	1836.0 s	0.03075"/pix
JWST	NIRCam LW, module B	F322W2 (CLEAR)	05594 (Mahler)	2024-08-27	1836.0 s	0.06301"/pix

§1.1 Keck/KCWI IFU spectroscopy (the σ_e data)

- Keck II / KCWI, KCRM red arm, RL (Red-Low) grating (RGRATNAM), Medium slicer (IFUNAM), 2×2 binning (BINNING/CCDSUM); nod-and-shuffle off (RNASNAM= 'Open ').
- The headline cube is a 3-NIGHT, 3-PROGRAM combine (see §2.2.1), NOT a single dataset — the cube header (DATE-OBS 2025-11-17, PROGPI Jones, XPOSURE 300 s) describes only the first input frame, so cite the combine, not the header:

Night (UT)	Program (PI)	RED (RL)	BLUE (BL)
2024 Dec 29	U204	4 × 300 s	1 × 1320 s (<i>excluded by reducer</i>)
2025 Aug 30	K409 (PI TBD)	12 × 300 s	4 × 990 s
2025 Nov 17	U002 (Jones)	20 × 300 s	5 × 1320 s

→ 36 × 300 s ≈ 180 min RED + ≈ 198 min BLUE on-source. The NEW `_mtwdo_` reduction (Master-Twilight + Dome hybrid flats, K. R. Gupta), Nov17_2025_DESJ0206_RL_combined_mtwdo_icubes_wcs.fits; dithered (PA 0°/45°/90°); CR rejection via astrocrappy (fsmode=median, psffwhm=2.50). OBSERVER = Glazebrook, Gupta, Jones, Kacprzak, Alcorn, Rhoades, Barone, Tran, Chen, Vasa. (*K409 PI + Aug-30 DIMM seeing still TBD for acknowledgements.*)

- Both KCWI arms recorded simultaneously. The red arm (KCRM, RL grating, central λ 7150 Å RCWAVE — the cube below) is the σ_e dataset. The blue arm (KCWI/KCB, BL (Blue-Low) grating BGRATNAM, central λ 4500 Å BCWAVE, KBlue filter BFILTAM, N&S off) is the ≈198 min BLUE above; it is not used for σ_e (the deflector's Ca H+K...Mg b absorption falls in the red) but supplies the source- $z = 1.302$ cross-check (Fe II UV multiplet; §2.2.2). The blue combined cube is not in this repo (only the RL cube) — quote the blue arm from the observing log, not a local file.
- Conditions (Nov-17 frame, header-literal): airmass 1.165 (AIRMASS); guide-star FWHM 1.271" (GUIDFWHM) — the adopted seeing; parallactic angle 15.6°, rotator 0°. (Per-night airmass/seeing for the other two nights not carried in the combined cube.)
- Cube: 100 × 100 spaxels × 3317 λ -pixels; 0.300"/spaxel (square; from WCS), 30" × 30" FOV; wavelength 5625–8941 Å at 1.0 Å/pix (CRVAL3/CD3_3), red central λ 7150 Å (RCWAVE). Spectral resolution $R \approx 10000$ at the fit band → $\sigma_{\text{inst}} \approx 12.6$ km/s (well below $\sigma_e \approx 270$; the LSF is carried into ppxf). [reference_kcwi_data_properties.md]

- Pointing RA/DEC = 02:06:13.38 / −01:14:20.8; target TARGRA/TARGDEC = 02:06:13.58 / −01:14:19.8.

§1.2 HST/WFC3 imaging (F200LP, F140W)

- Program 16773, PI Karl Glazebrook (PROPOSID/PR_INV_*); TARGNAME = DESJ0206-0114; both bands UT 2022-07-14 (EXPSTART MJD 59774.30–59774.31), ORIENTAT 111.9°, drizzled, units ELECTRONS/S.
- F200LP — WFC3/UVIS, aperture UVIS2: EXPTIME 600.0 s, NDRIZIM 4, NCOMBINE 2, drizzle scale 0.050"/pix; PHOTFLAM = 5.1851×10^{-20} , PHOTPLAM = 4923.48 Å → ZP_AB = 27.344.
- F140W — WFC3/IR: EXPTIME 597.7 s, NDRIZIM 3, NCOMBINE 3, drizzle scale 0.080"/pix; PHOTFLAM = 1.4829×10^{-20} , PHOTPLAM = 13922.91 Å → ZP_AB = 26.446.
- AB zeropoints are recomputed per-image from PHOTFLAM/PHOTPLAM (never hardcoded); $ZP_AB = -2.5 \cdot \log_{10}(\text{PHOTFLAM}) - 21.10 - 5 \cdot \log_{10}(\text{PHOTPLAM}) + 18.6921$. Drizzle pixel correlation: empirical noise $\approx 1.3\text{--}2\times$ the CCD-equation value → motivates the 10% flux floor (§2.1.4).

§1.3 JWST/NIRCam imaging (F150W2, F322W2)

- Program 05594 — “JWST Cluster SLICE — Strong Lensing and Cluster Evolution”, PI Guillaume Mahler (SURVEY category); TARGPROP = ACT-CL_J0206.2–0114 (the cluster field; the deflector sits in it). Both bands UT 2024-08-27 (DATE-BEG 06:02–06:40), NIRCam module B, units MJy/sr (i2d), NINTS 1 / NGROUPS 4.
- F150W2 (SHORT channel, CLEAR pupil): EFFEXPTM/XPOSURE 1836.0 s; PIXAR_SR = 2.2222×10^{-14} sr → 0.03075"/pix (PIXAR_A2 = 9.454×10^{-4} ") → ZP_AB = 28.033.
- F322W2 (LONG channel, CLEAR pupil): EFFEXPTM/XPOSURE 1836.0 s; PIXAR_SR = 9.3307×10^{-14} sr → 0.06301"/pix (PIXAR_A2 = 3.970×10^{-3} ") → ZP_AB = 26.475.
- JWST AB from area: $ZP_AB = -6.10 - 2.5 \cdot \log_{10}(\text{PIXAR_SR})$. No PAM correction (drizzled/resampled).

§2 Methods

§2.1 Photometric Analysis

§2.1.1 Masking

- Use the F200LP `_mask.fits` as the arc mask — NOT the F140W one. [feedback_masking_strategy.md; reference_hst_jwst_data_properties.md; METHODS §II.4]
 - F200LP mask (2512 HST pixels) is hand-tuned to cover the arc only and leaves the deflector core unmasked — it is the original arc-tuned segmentation mask.
 - F140W mask (1121 HST pixels) covers the arc PLUS the deflector core (the photometry tool’s aperture geometry pulled the core in); the two masks have zero spatial overlap.
 - Consequence: F140W is valued for its *image* (sharper Sérsic fit), never for its mask. Using the F140W mask as an arc mask would destroy the deflector-core signal. ### §2.1.1b Principled (reproducible) arc masking — NEW 2026-05-29

The hand-painted masks are now backed by an objective, reproducible recipe (so a referee can reproduce the arc selection), validated + applied to all four bands. [NOTES_arc_mask_verification_2026-05-29.md, NOTES_photometry_mask_systematics_2026-05-29.md; scripts/arc_mask_verification.py, mask_method_comparison.py, photometry_systematics.py]

- Locator = F200LP (highest contrast on the blue $z=1.302$ source). Two independent objective selectors reproduce the F200LP hand mask: (i) a color cut `m_F200LP–m_F140W` (arc bluer than the red deflector); (ii) a Sérsic-residual mask (subtract a 2D deflector model, flag pixels whose positive residual exceeds $k \cdot \sigma_{\text{sky}}$, where k is the detection threshold in units of the sky noise σ_{sky} and σ_{sky} is the robust background scatter; we adopt $k = 3$). On F200LP the Sérsic-residual mask reproduces the expert photometry to 0.016 mag, R_e to 3%.

- Transfer to other bands by reprojecting the F200LP footprint (WCS + map_coordinates). Reproduces the expert JWST aperture mags to 0.01–0.02 mag. Do NOT use an independent per-band Sérsic-residual mask on F140W/JWST — a single Sérsic under-fits the bright extended IR galaxy (median residual +1.0) and over-masks by 0.2–0.4 mag.
- IR extends the source footprint: the deeper/sharper JWST bands see the arc’s fuller extent (F150W2 mask grows ~6× vs HST). We grow the F200LP arc into each band’s 2-component-Sérsic residual source pixels that are contiguous with the arc (region growing; cannot grab the core pedestal or field companions).
- Deflector model = 2-component (bulge+disk) Sérsic — single Sérsic under-fits JWST; 2-comp cuts galaxy-body RMS 3.3–3.5×. PSF effect on the fill quantified (scripts/psf_fill_model.py): a PSF-convolved 2-comp Sérsic (Gaussian at instrument FWHM; env lacks webbpsf) shifts the filled mag by ≤ 0.004 mag $\rightarrow \Delta M \star \ll 0.01$ dex, negligible. The fill is PSF-robust (arc is outside the PSF core).
- raw vs fill-in: raw photutils (masked) discards the deflector light under the arc; the Sérsic fill-in (replace masked pixels with the model) recovers it. With the large IR-extended masks the fill-in correction is +0.18 to +0.96 mag per band. filled photometry is mask-definition-independent (per-band vs global agree to 0.01 dex); raw is mask-size- dependent (up to 0.78 mag) — see §3.2.
- per-band vs global mask (union of all per-band masks): negligible for filled (0.01 dex $M \star$), large for raw — confirming filled is the robust quantity, raw the conservative lower bound.
- Files (in sibling dir ../velocity_dispersion_from_IFU/): AGELO20613-011417A_F200LP_WFC3_cutout_L3_mask.fits (500×500, 25″×25″); the mask was revised 2026-02-28 (_mask_20260228.fits); original kept as _mask_OLD.fits.
- JWST bands: same arc geometry; the HST F200LP mask is re-projected from the HST WCS onto each JWST frame. [METHODS §II.4]
- For the IFU/ σ_e application (context): F200LP mask reprojected to the 0.30″ IFU grid via scipy.ndimage.map_coordinates(order=0) nearest-neighbor; ~38/184 spaxels inside R_e flagged (~21% by count, ~27% by I-weight). [feedback_masking_strategy.md]

§2.1.2 Estimating R_e

- Headline $R_e = 2.305'' = 16.23$ kpc at $z = 0.67564$. [CLAUDE.md; METHODS §II.3/A3]
- Method: mean of the F140W and F200LP masked curves-of-growth (scripts/final_sigma_e.py:curve_of_growth, the PRODUCTION script):
 - F140W masked CoG = 2.168″; F200LP masked CoG = 2.441″; mean = 2.305″.
 - Masked CoG zeros mask-flagged HST pixels before the radial flux integral, so the lensed arc / companions do not bias the half-light point.
 - Centers: HST-mean photutils.centroid_2dg centers (same as the σ_e pipeline), both bands.
- Do NOT confuse with scripts/measure_Re.py $\rightarrow$ results/Re_measurements.npz. That script uses the image-geometric center and produces 10 variants (3 sources $\times$ 4 masking strategies: unmasked, zeroed, proper, PSF-convolved); its “proper-mask” mean (2.633″) is not the headline — early-reference only. (measure_Re.hst_Re previously hardcoded pixscale=0.08 for both bands; fixed 2026-06-08 (A3c/nb14) to read the WCS pixscale + r_max_arcsec=6.0, now reconciling with the production CoG to $<0.04''$ — but final_sigma_e remains the headline path.)
- Superseded prior value: IFU white-light $R_e = 2.61''$ (nb05/06 era) — superseded because the HST-based 2.305″ has the F140W+F200LP arc masks built in. A Ca H+K+G depth-map gives 2.866″.
- R_e as a σ_e systematic (test D7): four R_e definitions shift σ_e (rises monotonically with R_e). Re-measured at the wide window (M12, nb15): spread ± 6.13 km/s, now folded into the formal budget (§3.1, §2.4.3); the narrow-window 16.9 km/s figure is superseded. [CLAUDE.md]

§2.1.3 Measuring Fluxes

- Tools: photutils aperture photometry (interactive masking GUIs scripts/photometry_masking_{HST,JWST}.py). synphot is NOT currently used — flux $\rightarrow$ AB is done directly from header zeropoints. (If

the outline wants synphot, that is aspirational; flag with co-authors.) [METHODS §II.4]

- AB zeropoints — read per-image from FITS headers, never hardcoded. [CLAUDE.md; reference_hst_jwst_data_properties.m]
 - HST (drizzled e⁻/s): $ZP_{AB} = -2.5 \cdot \log_{10}(\text{PHOTFLAM}) - 21.10 - 5 \cdot \log_{10}(\text{PHOTPLAM}) + 18.6921$.
 - JWST (i2d, MJy/sr): $ZP_{AB} = -6.10 - 2.5 \cdot \log_{10}(\text{PIXAR_SR})$.
- Four bands / instruments / programs:

Telescope	Instrument	Filter	Program (PI)	Drizzled scale	EXPTIME	ZP_AB	AB (aperture)
HST	WFC3/UVIS	F200LP	16773 (Glazebrook)	0.0500"/pix	600.0 s	27.344	22.613
HST	WFC3/IR	F140W	16773 (Glazebrook)	0.0800"/pix	597.7 s	26.446	19.134
JWST	NIRCam SW	F150W2	05594 (Mahler)	0.03075"/pix	1836.0 s	28.033	18.942
JWST	NIRCam LW	F322W2	05594 (Mahler)	0.06301"/pix	1836.0 s	26.475	18.604

[reference_hst_jwst_data_properties.md] - JWST pointing comes from the cluster program ACT-CL J0206.2–0114 (ACT DR5, Hilton et al. 2021); the deflector sits in that field. No PAM correction (drizzled). Drizzle pixel correlation: empirical noise $\sim 1.3\text{--}2\times$ the idealized CCD-equation value — motivates the 10% flux floor (§2.1.4). - 2D Sérsic verification given masking (the morphology cross-check; notebook 08): single-component 2D Sérsic per band (scripts/measure_Re.py framework + the 2026-05-11 bound-fix $n \in [1.0, 6.0]$, $\text{ellip} \in [0.0, 0.6]$, 3-init grid — prevented an $n=0.30$ flat-disk escape on F200LP), extrapolated to total flux:

Filter	AB aperture	AB Sérsic-total	Δmag	n	$r_{\text{eff}} (")$
F200LP	22.613	20.672	−1.94	1.42	2.49
F140W	19.134	18.282	−0.85	1.54	1.88
F150W2	18.942	18.154	−0.79	1.40	1.72
F322W2	18.604	17.633	−0.97	1.97	2.03

Aperture under-counts outer-profile flux by 0.8–1.9 mag, but the integrated-mass effect is only +0.07 dex (see §2.1.4 / Results). Caches: results/seraic_total_photometry.npz, results/bagpipes_sersic_refit.npz. [METHODS §II.6]

§2.1.4 SED Modeling with bagpipes

- Tool: bagpipes (Carnall et al. 2018, 2019); BPASS+CLOUDY templates (Bagpipes default). Notebook 02_streamlined_Bagpipes_SED.ipynb. [METHODS §II.5; reference_sed_fitting.md]
- Model: exponential- τ SFH; Calzetti dust (A_V free, 0–2); free metallicity (0–2.5 $Z_{\odot}$); redshift prior (0.674, 0.676) bracketing the line-fit z . Sampler Nautilus ($n_{\text{live}}=400$), 500-sample posterior.

```
fit_instructions = {
    "redshift": (0.674, 0.676),
    "exponential": {"age": (0.1, 15.), "tau": (0.3, 10.),
                    "massformed": (1., 15.), "metallicity": (0., 2.5)},
    "dust": {"type": "Calzetti", "Av": (0., 2.)},
}
```

- Why a 10% fractional error floor on each band: a conservative, uniform systematic budget that covers both the formal photon noise and the drizzle pixel-correlation underestimate (empirical noise $\sim 1.3\text{--}2\times$ the idealized CCD-equation value). [METHODS §II.4/II.7; reference_sed_fitting.md]
- Systematic test to fill in masked pixels = the 2D-Sérsic-total refit: because the aperture excludes arc-masked pixels and truncates the outer profile, we fit a 2D Sérsic per band (§2.1.3), extrapolate to total flux, and re-run

Bagpipes with identical priors. This recovers the flux missing from masked/truncated regions and bounds the mass bias. Result: $\log(M_\star/M_\odot) = 11.40 \pm 0.11/-0.15$ (Sérsic-total) vs $11.33 \pm 0.07/-0.09$ (aperture) $\rightarrow$ aperture is biased low by ~ 0.07 dex (+16% in $M_\star$), within the quoted uncertainty. [METHODS §II.6] NOTE (2026-05-29): this analytic-total refit is now superseded by the per-aperture 2-component Sérsic fill-in under the principled IR-extended masks (§2.1.1b / §3.2): the headline is the raw value 11.16 (10%) with a one-sided +0.31 dex fill-in systematic reaching 11.46. The 11.40 analytic-total sits inside that bracket.

- Posteriors (older expert-aperture, now a cross-check): $\log(M_\star/M_\odot)=11.33 \pm 0.07/-0.09$; mass-weighted age 5.10 Gyr [3.69–6.11]; $\tau=0.68$ Gyr; $Z=0.89 Z_\odot$; $A_V=0.82$ mag; $z_{\text{phot}}=0.675$ [0.674–0.676]. Per-band residuals: F200LP +4.7%, F140W –11.3%, F150W2 –8.0%, F322W2 +9.6% (mild single- τ -SFH NIR-slope tension; not catastrophic, within ± 0.07 –0.15 dex). [METHODS §II.5]
- Caches / figure: `results/bagpipes_sed_results.npz` (the file Figure 2 loads), `pipes/posterior/AGEL0206/0206_real.h5` (delete to refit), `results/bagpipes_sersic_refit.npz`.

§2.2 Spectroscopic Analysis

§2.2.1 Calibration — @Keerthi / Kaustubh

- Instrument: Keck II / KCWI, KCRM red arm, Red-Low (RL) grating, Medium slicer, 2×2 binning; dithered at PA 0°/45°/90°. [METHODS §I.1; `reference_kcwi_data_properties.md`]
- Cube (headline): `Nov17_2025_DESJ0206_RL_combined_mtwdo_icubes_wcs.fits` — the NEW `_mtwdo_` reduction (Master Twilight + Dome hybrid flats), K. R. Gupta’s latest re-reduction. The earlier `no_mtwdo_` cube is the OLD reduction (second-reduction cross-check; co-aligned to $<0.002''$, differs only in flat-fielding; σ_e differs ~ 7 km/s — the “reduction-pass” systematic).
- Reduction chain: raw frames $\rightarrow$ `kcwidrp`; stacked with `ISMgas/kcwi/kcwiReduction` by K. R. Gupta (Kaustubh); Dec-29 RED frames pre-reduced by Yuguang Chen. (*Calibration prose is Keerthi/Kaustubh’s to write — these are the facts on hand.*)
- Cube geometry: shape (3317, 100, 100); 0.300'' spaxels; $\sim 30'' \times 30''$ FoV; $\lambda = 5625$ –8941 Å at $\Delta\lambda = 1.0$ Å/pix; BUNIT = FLAM16/arcsec².
- Three confirmed nights (totals ≈ 180 min RED + 198 min BLUE on source):

Night (UT)	Program (PI)	RED	BLUE
2024 Dec 29	U204	4×300 s	1×1320 s (excluded by reducer)
2025 Aug 30	K409 (PI TBD)	12×300 s	4×990 s
2025 Nov 17	U002 (Jones)	20×300 s	5×1320 s

FLAG: the cube header (DATE-OBS 2025-11-17, PROGPI Jones) describes only the first input frame — cite the multi-night combine, not the header. K409 PI and Aug-30 DIMM seeing still TBD. - Spectral resolution / LSF: $\text{FWHM}_{\text{inst}} = 2.355 \times \text{DISPSCAL}(0.2940) \times 1.0 \text{ Å} = 0.692 \text{ Å} \rightarrow R \approx 10000$ at the fit band; $\sigma_{v,\text{inst}} \approx 12.6$ km/s observed (7.5 km/s rest-frame) — the ~ 270 km/s dispersion is resolved by $\sim 20\times$. LSF robustness ≤ 0.83 km/s over $\text{DISPSCAL} \times 0.5$ – 2.0 . - Seeing: guide-camera FWHM = 1.27'' (GUIDFWHM, Nov-17; conservative).

§2.2.2 Redshift Measurement

- Method — single/multi-line Gaussian fits on absorption (+emission) features in the integrated spectrum, on NIST air rest wavelengths (`scripts/redshift_verify.py`, notebook 04):
 - Ca K 3933.66, Ca H 3968.47 (fit as a doublet with shared width), G-band 4304.40, H δ , H β , plus the [O II] 3726/3729 doublet (shared z , σ , flux ratio). Per-line z inverse-variance combined. H γ excluded (not trusted); Mg I / Fe / Na D out of range or blended.
 - Deflector systemic $z = 0.67564$ (supersedes AGEL DR2 $z = 0.67511$; ~ 95 km/s offset, used only in old notebooks). [METHODS §I.2; `reference_redshift_verification.md`]

- Double verification with ppxf: the Gaussian line-fit z is independent of, and consistent with, the ppxf V_{sys} (ppxf returns z via $z = (1+z_0) \cdot \exp(V/c) - 1$; Cappellari 2023 eq. 5c).
- Source $z = 1.302$ (AGEL DR2), cross-checked via Fe II UV resonance multiplet + Mg II 2796/2803 in the KCRM blue-arm cube. (*Measured $z_{\text{source}} \pm \text{uncertainty} + \text{its cache are a [TBD] placeholder — confirm before quoting a source-}z \text{ error bar.}$*)
- Peculiar velocities: we do not correct for peculiar velocities — we use the line-fit systemic z directly (see gap G3; this sentence needs to be added to the draft).
- Wavelength convention: AIR rest wavelengths throughout (NIST-verified); KCWI native vacuum is converted to air. Do not mix SDSS vacuum values ($\sim 1 \text{ \AA} \approx 80 \text{ km/s}$ systematic at $4000 \text{ \AA}$). [feedback_wavelength_convention.md]

§2.2.3 Stellar Kinematics & Velocity Dispersion

Spatial regime / masking / R_e

- Headline σ_e is measured on a single $I(r)$ -weighted 1-D aperture spectrum at $R < R_e$ — the literature-standard σ_e (Cappellari et al. 2006 eq. 1; the KH13 / Greene+20 / SAURON / ATLAS3D / MaNGA convention). $R_e = 2.305'' = 16.23 \text{ kpc}$ (§2.1.2).
- Center: HST-mean photutils.centroid_2dg of F140W+F200LP cores, propagated through the KCWI WCS; adopted RA = 31.55613° , Dec = -1.23819° (ICRS). F140W $\leftrightarrow$ F200LP offset $0.36''$ (F200LP shifted by the UV-bright arc; F140W is the clean bulge centroid).
- Spatial arc mask: F200LP segmentation mask reprojected to the IFU grid (nearest-neighbor); 38/184 spaxels inside R_e flagged ($\sim 27\%$ by I -weight). F140W mask not used (covers the core).
- I -weight map: per-spaxel collapse of the IFU 6500–7500 $\text{\AA}$ white-light band; spectra summed (not averaged) so the wild bootstrap preserves per-spaxel noise.
- $R < R_e/8$ (JFK95 σ_c) was dropped — at $0.288''$ it sits inside seeing $\text{FWHM}/2 = 0.64''$ (~ 3 spaxels). No central σ is quoted; $\sigma_e(<R_e/2) \approx 225 \text{ km/s}$ is reported as a gradient diagnostic only.

ppxf setup

- ppxf v9.x (Cappellari & Emsellem 2004; Cappellari 2017, 2023), moments = 2 (first two velocity moments V, σ). [METHODS §I.5/I.6; scripts/bootstrap_ppxf.py]
- Polynomial degrees: additive Legendre degree swept over 15–29 (15 values) and pooled; mdegree = 0. σ -vs-degree is flat within the bootstrap envelope (no LOSVD absorption by the polynomial). (*Note: METHODS §I.5 prose mis-labels these “multiplicative” — the code passes degree=..., mdegree=0, i.e. additive. Describe them as additive.*)
- Three SPS libraries, pooled at the posterior level: FSPS, EMILES, XSL (8–9 templates each over rest 3500–5000 $\text{\AA}$).
- Per-SPS native wavelength frame: FSPS = vacuum, EMILES = air, XSL = air. Galaxy frame matched per-SPS (scalar-median vac $\leftrightarrow$ air via Ciddor 1996, following Cappellari’s pattern). Why: the legacy “convert galaxy to air for all three” produced a spurious -90 km/s FSPS V_{sys} ; the fix collapsed the V_{sys} split-track $\sim 110 \rightarrow \sim 15\text{--}18 \text{ km/s}$ ($\sigma \text{ shift} \leq 2 \text{ km/s}$). [reference_ppxf_vacair_handling.md]
- Per-SPS V_{sys} subtracted before pooling (σ is V -invariant; V_{sys} medians FSPS -19 , EMILES -4 , XSL -1 km/s). A single global V_{sys} would inflate σ_e by $\sim 10\text{--}15 \text{ km/s}$ in the discrete cross-check.
- Emission-line handling (sigma clipping / removing emission lines):
 - Custom no-Balmer goodpixels (_determine_goodpixels_no_balmer): masks only the forbidden lines ([O II], [O III], [O I], [N II], [S II]) and keeps H δ , H γ , H β IN the fit (they are absorption in this passive deflector; masking them discards ~ 120 stellar pixels). H δ : keep unmasked — RESOLVED (M12/nb16): H δ is well-fit (local-MAD not an outlier), and masking it shifts σ_e by $+6\text{--}8 \text{ km/s} = \text{LOSVD information}$, not contamination (the M9 “it’s signal” pattern); the bootstrap_ppxf.py TODO is closed.
 - $z = 1.302$ source-emission masks (ARC_MASKS_REST, mapped by $(1+z_s)/(1+z_l)=1.374$): Mg II 2796/2803 (def-rest 3835–3855 $\text{\AA}$), [O II] 3727/3729 (5115–5135), [Ne III] 3869 (5260–5340), He I 3819 (5237–5253, added 2026-05-27 as test M8).
 - Tellurics — handled by masking, not correction. One band is masked: the O $_2$ A-band leading-edge residual at obs 7593–7626 $\text{\AA} = \text{def-rest } 4525\text{--}4545 \text{ \AA}$ (ARC_MASKS_REST). Confirmed telluric, not

z=1.302 source, on 2026-05-18 by the same-observed- λ test (the spike sits at the *same obs* λ in the deflector aperture, the sky box, and a 4–8" off-deflector spectrum). No full telluric correction is applied; the M10 OH/sky bands (below) catch sky-emission residuals.

- M10 sky-line audit: bad-pixel catalog BAD_PIXELS_REST = 35 entries (26 CR residuals + 9 OH air-glow/sky bands), all verified non-source via the arc spectrum.
- Wild bootstrap errors: hybrid Rademacher sign-flip $\times$ local-residual scaling (75-pixel rolling-window residual rescale, clipped [0.2,5.0]); N = 500 production per (SPS $\times$ degree), N = 50 smoke (agree within 1 km/s). Errors = 16/84 percentiles (asymmetric). joblib parallel, BLAS pinned to 1 thread/worker.

σ_e definition & wavelength window

- σ_e definition — analytic (Gültekin) and discrete (Cappellari) forms are the SAME quantity. (Gültekin et al. 2009 eq. 1; Kormendy & Ho 2013 eq. 3; 2D-IFU form Cappellari et al. 2006 eq. 1.)
 - Analytic / luminosity-weighted integral (Gültekin 1D longslit, generalised to 2D — the $2\pi r$ is the axisymmetric area element $dA = 2\pi r dr$):

$$\sigma_e^2 = \frac{\int_0^{R_e} (V^2 + \sigma^2) I(r) 2\pi r dr}{\int_0^{R_e} I(r) 2\pi r dr}.$$

- Discrete Cappellari (2006) eq. 1 spaxel/bin sum — the *same* moment evaluated over spaxels (or Voronoi bins) n , with F_n = the I-band flux (luminosity weight) of spaxel n ; the $2\pi r$ factor is captured implicitly because the number of spaxels per annulus scales with area:

$$\sigma_e^2 = \frac{\sum_{n: R_n \leq R_e} F_n [(V_n - V_{\text{sys}})^2 + \sigma_n^2]}{\sum_{n: R_n \leq R_e} F_n}.$$

This is the form SAURON IV / ATLAS3D III / MaNGA-DAP (Westfall+2019) use on Voronoi bins.

- These two forms are equivalent and BOTH are already implemented and cross-checked in this project (do not re-derive): the analytic integral is the headline path — the cumulative I-weighted single-ppxf fit on the $R < R_e$ aperture spectrum (preserves LOSVD shape; what KH13/Greene+20 compute) — and the discrete Cappellari/Gültekin spaxel-sum is the §7 / nb07c annular implementation (`run_gulteкин_mc`, $F_j = \Sigma I$ over each unmasked annulus). The two agree at $<1\sigma$: integral (§6cum, narrow) 267.32 ± 24 vs discrete annular (§7, arc-filtered to $R < R_{\text{safe}} = 3R_e/4$) $256.17 - 13.0/+12.7$. Full implementation + subtleties in `reference_gulteкин_implementation.md`.
- Why the integral path is the headline (not the discrete sum): the discrete spaxel-sum is (i) per-spaxel-S/N-limited and (ii) sensitive to arc contamination in the outer spaxels — an *unfiltered* discrete sum over the resolved PowerBins out to $\sim 3.4''$ is dominated by the z=1.302 arc (outer bins carry $\sigma \gtrsim 300$ –440 km/s, V swings of ± 150 –270; the σ_e blows up to ~ 350). The I-weighted aperture auto-suppresses the arc (bright deflector core dominates the weight) and avoids the S/N floor, so it implements the same Cappellari definition robustly. The discrete §7 sum is therefore arc-filtered to $R < R_{\text{safe}}$ and used only as the architecture-independent cross-check / for the $\sigma(R)$ profile.
- Standard wavelength range (headline): rest 3800–5400 Å (`wR3800_5400_arcmask`) — Ca H+K through Mg b / Fe5270 / Fe5335; 2161 good pixels ($\sim 2.6\times$ the narrow window). The fit is constrained by the highest-S/N features — Ca H, Ca K, G-band (3800–4400) — and is consistent across windows.
 - Blue edge 3800 Å = XSL template floor (def-rest 3675 Å) + 5% velocity padding + retaining Ca H+K; red edge set by KCWI coverage. Provenance is data-driven, NOT a specific paper. Closest precedent: LEGA-C (Bezanson et al. 2018; van der Wel et al. 2021) at $z \sim 0.7$. DO NOT cite “Cappellari 2017 recommends extending blueward” — that was a hallucination, retracted.
 - Cross-check window: rest 4000–5400 Å (`wR4000_5400_arcmask`, H β + Mg b, no Ca H+K) — orthogonal Lick-tradition feature set.
 - Narrow legacy window: obs 6500–7500 Å = rest 3879–4476 Å (Ca H+K + G); $\sigma_e = 267.95 \pm 30.10$.

A few systematics tests to describe (full budget in Results §3.1): I-shape (10 I-weight-map shapes, ± 2.27), F200 mask weight ($w \in \{0, 0.5, 1\}$, ± 6.65), fit-window (3 windows, ± 3.82), frame (vac/air, ± 5.0), centering (5-center $\pm 0.4''$ sweep, ± 4.0), reduction-pass (NEW vs OLD cube, ± 3.45). Plus the M9 “DO NOT MASK” finding: the +5–7%

bump at def-rest 5193–5204 Å is the Mg b LOSVD wing (signal, not contamination); masking it drops σ_e by 7 km/s.

§2.3 Lens Modeling

GAP G1 — not in this repo. The companion-paper (Ferrami et al.) lens-modeling content (multiple lens models, PSF construction, joint shear+subhalo constraints, potential-correction triple-checks, BPL vs point-mass differentiating models) is not present in this kinematics+ photometry repo. Source it from the lens-modeling repo (7Documents/AGEL/202509_DESJ0206_modeling/, 20251112_DESJ0206_Pyauto_PRONT0/) or the Ferrami draft. The only on-hand pointer: PROJECT_BRIEF.md notes the lens-model enclosed mass combines with $(\sigma, M_\star, z)$ to give a 4D constraint on the central BH + bulge scaling relations.

System identity for the summary sentence: AGEL J020613–011417 (DES J0206–01), AGEL DR2; source $z = 1.302$ (spiral), deflector $z = 0.67564$.

§2.4 End-to-end procedures (ordered walkthroughs)

This section spells out the actual step order of each pipeline — what runs, in what sequence, with which script — so the Methods section and a referee can reconstruct every headline number. Numeric results are the M12 headline (results/PAPER_VALUES.json).

§2.4.1 Spectroscopic σ_e pipeline — step by step

Driver for the headline: `scripts/run_wide_sigma_e.py --cube new_clean_hei --n_bootstrap 500` (calls `final_sigma_e.load_setup` → `extract_aperture_spectrum` → `bootstrap_ppxf` machinery).

1. Reduction (upstream). Keck II / KCWI, KCRM red arm, Red-Low grating, Medium slicer, 2×2 binning, dithered PA 0/45/90°. The headline cube is the NEW `_mtwdo_reduction` (Master-Twilight+Dome hybrid flats, K. R. Gupta), `Nov17_2025_DESJ0206_RL_combined_mtwdo_icubes_wcs.fits`. The pre-`_mtwdo` cube is the OLD reduction (second-reduction cross-check). Cube axes $[\lambda, y, x]$; 0.30"/spaxel; seeing FWHM 1.27" (GUIDFWHM); $\sigma_{\text{inst}} \approx 12.6$ km/s; $R \approx 10000$.
2. Geometry + `R_e`. HST-mean centre = `photutils.centroid_2dg` of the F140W+F200LP cores (offset 0.36"; F200LP pulled by the UV arc, so F140W is the clean bulge centroid), propagated through the KCWI WCS to a sub-pixel IFU centre. `R_e` = mean of the F140W (2.168") + F200LP (2.441") masked curves-of-growth = 2.305" (`final_sigma_e.curve_of_growth, r_max=6`").
3. Spatial arc mask → IFU grid. The F200LP `_mask.fits` (2512 HST px, arc-only) is reprojected onto the 0.30" IFU grid with `scipy.ndimage.map_coordinates(order=0)` (nearest-neighbour) → `arc_spax_mask` (~38/184 spaxels inside `R_e` flagged, ~27% by I-weight).
4. I-weighted aperture extraction. Build the 6500–7500 Å white-light I-map; the $R < R_e$ aperture spectrum = $\Sigma_{\text{spaxel}} \text{cube} \cdot w$, w = I-weight with arc spaxels hard-masked (`mask_weight=0`) and re-normalised. Spectra are summed (not averaged) so the bootstrap preserves per-spaxel noise. (`extract_aperture_spectrum, final_sigma_e.py`.)
5. Spectral masking (goodpixels). (a) no-Balmer mask — forbidden lines only ([O II], [O III], [O I], [N II], [S II]); H δ , H γ , H β kept (absorption; M12/nb16 confirmed keep-unmasked). (b) $z=1.302$ source-emission masks `ARC_MASKS_REST` (Mg II, [O II], [Ne III], He I 3819) + the O₂ A-band telluric. (c) `BAD_PIXELS_REST` = 35 entries (26 CR residuals + 9 M10 OH/sky bands). All in deflector rest-frame Å.
 - Outlier rejection / sigma-clipping decision: `ppxf`'s in-fit `clean=False` — `clean=True` rejects 0 pixels here (the KCWI noise array is overestimated vs the actual residual scatter), so it is inert. Instead, cosmic-ray/bad pixels are flagged by a 3σ local-MAD clip (rolling 75-pixel window, $|\text{residual}|/\text{local-MAD} > 3\sigma$) on the canonical residuals → 52 CR-like spikes, curated into the 26 CR entries of `BAD_`

PIXELS_REST; the 9 M10 sky bands were flagged separately at $>2.5\times$ median sky-noise. Caveats (carried as a stated sensitivity, not a budget term): the 3σ threshold is not strongly motivated vs alternatives, and cleaning shifts σ_e by only $+1.4$ km/s (below stat 1σ); the same pixels replicate the shift on the OLD cube (M6) $\rightarrow$ real CR residuals.

6. Per-SPS frame-aware ppxf. For each of FSPS / EMILES / XSL: match the galaxy frame to the SPS native frame (FSPS=vacuum, EMILES/XSL=air; scalar-median vac $\leftrightarrow$ air via Ciddor 1996), run ppxf (Cappellari) with moments=2, additive Legendre degree swept 15–29 (15 values), mdegree=0, over the wide window (rest 3800–5400 Å). Subtract the per-SPS V_{sys} before pooling.
7. Wild-bootstrap errors. Per (SPS $\times$ degree), $N=500$ hybrid wild bootstrap (bootstrap_ppxf.compute_local_residual_scaling): residuals $r = \text{galaxy} - \text{bestfit}$; a rolling 75-pixel window estimates the local (wavelength-dependent) noise scale $s(\lambda)$ (clipped to $[0.2, 5.0]$); each iteration applies a Rademacher ± 1 sign-flip to the locally-rescaled residual, adds it back to the best fit, and re-fits $\rightarrow$ one σ draw. Seeds are deterministic (BOOT_SEED=42 + 50000 + 10000 $\cdot W_{\text{IDX}}$ + 100 $\cdot \text{sps_idx}$ + degree_i) so caches bit-reproduce. joblib loky, BLAS pinned to 1 thread/worker; $N=50$ smoke before $N=500$ production.
8. Pool + budget. Concatenate all σ samples (3 SPS $\times$ 15 degrees $\times$ 500) $\rightarrow$ 16/50/84 percentiles = central + asymmetric stat. The pooled width already marginalises SPS + polynomial degree. Add the §2.4.3 systematic components in quadrature $\rightarrow \sigma_e = 269.62 - 13.45 / +13.10$ (sym ± 13.27) km/s.

§2.4.2 Photometry / M★ pipeline — step by step

Drivers: scripts/photometry_systematics.py (masking + photometry), bagpipes_sersic_refit.py (SED), orchestrated/displayed in notebook 12. Recipe = F200LP locates, IR extends, fill-in recovers.

1. Locate the arc on F200LP (best source contrast): a 2D Sérsic deflector model is subtracted and a pixel is flagged as arc where its positive residual exceeds $k \cdot \sigma_{\text{sky}}$ — i.e. k is the detection threshold in units of the sky noise σ_{sky} (so $k=3$ means residual $> 3\sigma_{\text{sky}}$), with a parallel S/N floor (the pixel's own S/N $> \text{SNR}_{\text{thresh}}$). We adopt $k = 3$ (arc_mask_verification.py): this objective Sérsic-residual mask reproduces the expert hand mask to 0.016 mag, R_e to 3%. $k = 3$ is the clean saturation point of a 2-D sweep over the two thresholds ($\text{SNR}_{\text{thresh}} \in \{2..20\} \times k \in \{2..8\}$) — below $k \approx 3$ it grabs noise, above it the masked-flux fraction stops changing.
2. Reproject the F200LP footprint to every band (WCS + map_coordinates) — reproduces the expert JWST aperture mags to 0.01–0.02 mag. Do not fit an independent per-band single Sérsic (under-fits the bright IR galaxy, over-masks 0.2–0.4 mag).
3. IR-extend. In the deeper/sharper JWST bands, region-grow the arc into 2-component-Sérsic residual source pixels contiguous with the arc seed (cannot grab the core pedestal or field companions); F150W2 mask grows $\sim 6\times$ vs HST.
4. Deflector model = 2-component (bulge+disk) Sérsic (cuts JWST galaxy-body RMS $3.3\text{--}3.5\times$ vs single Sérsic). PSF-convolved fill quantified at ≤ 0.004 mag (psf_fill_model.py) — negligible.
5. Two photometry estimators per band/mask: raw photutils aperture (masked $\rightarrow$ discards under-arc deflector light $\rightarrow$ biased low, mask-size-dependent) vs Sérsic fill-in (replace masked pixels with the model $\rightarrow$ recovers it $\rightarrow$ mask-definition-independent, per-band vs global agree 0.01 dex). Fill-in correction $+0.18\text{--}0.96$ mag with the large IR-extended masks.
6. AB magnitudes from per-image header zeropoints (never hardcoded): HST PHOTFLAM/PHOTPLAM, JWST PIXAR_SR. Four bands: F200LP, F140W (HST WFC3), F150W2, F322W2 (JWST NIRC2).
7. Bagpipes SED (02_streamlined_Bagpipes_SED): exponential- τ SFH, Calzetti dust (A_V 0–2), free Z (0–2.5 $Z_\odot$), z prior (0.674, 0.676); Nautilus sampler $n_{\text{live}}=400$, 500-sample posterior. A 10% fractional flux floor per band covers photon noise + the drizzle pixel-correlation under-estimate (empirical noise $\approx 1.3\text{--}2\times$ CCD-equation); 20% is run as a conservative variant.
8. M★ budget = 8 fits {per-band, global} $\times$ {raw, filled} $\times$ {10%, 20%} $\rightarrow$ headline (empirical, raw-central, one-sided +sys, user choice): $\log M_\star = 11.16 \pm 0.08$ (stat) $+0.31$ (sys) at 10% [11.04 ± 0.14 at 20%], fill-in reach 11.46 (the asymmetric form used in the figures is $+0.32/-0.08$, the stat $\oplus$ sys quadrature); explicit masking-approach systematic ± 0.16 dex (Mstar_masking_systematic.npz).

§2.4.3 Systematic-audit procedures (how each budget term was measured)

Each σ_e systematic is a dedicated sweep re-using the §2.4.1 machinery, varying ONE axis and quoting peak-to-peak/2 (or half- Δ for two-point axes), all on the headline cube + masks:

- I-shape (± 2.27): 10 alternative I-weight maps (`run_isource_shape_sweep.py`) $\times$ 3 SPS $\times$ 15 deg $\times$ N=250; peak-to-peak/2 of the 10 central σ_e (266.83–271.37).
- F200 mask-weight (± 6.65): down-weight the arc spaxels at $w \in \{0, 0.5, 1\} \times$ 3 SPS $\times$ N=500; (`w00 269.69 – w100 256.39`)/2. (*This is the mask-strength axis. The mask-definition axis — `expert/sersic/perband/global` reprojected to the IFU grid, `run_sigma_e_mask_systematic.py` — gives ± 5.85 , which overlaps this; larger-of-two is kept, not added.*)
- Fit-window (± 3.82): 3 windows (`wR3800_5400 / wR4000_5400 / w6500_7500`) $\times$ 3 SPS $\times$ N=500; peak-to-peak/2 (269.66 / 268.58 / 276.22).
- Frame (± 5.0): structural — vac/air per-SPS native-frame choice.
- Centering (± 4.0): 5 perturbed HST-mean centres ($\pm 0.4''$ sweep; `NOTES_centering_investigation`).
- Reduction-pass (± 3.45): half- Δ between the NEW and OLD cleaned cubes (269.62 vs 262.72); 2 reductions only — refine if a 3rd lands.
- R_e-source / D7 (± 6.13): the 4 R_e estimators (mean/F140W/F200LP/CaHK+G, 2.168–2.902"; `run_sigma_e_Re_systematic_wide.py`); full peak-to-peak/2.

Quote the budget table (§3.1) as the result; the iterative history that *led* to this masking + component set is the internal audit trail (M8→M12) at the end of this document — not for the manuscript Methods section.

§2.4.4 Reproducibility / infrastructure

- Architecturally-independent estimators (must not share intermediate state): §6cum cumulative I-weighted single-ppxf aperture (the analytic Gültekin path = headline) · §7 discrete Cappellari (2006) annular spaxel-sum (`run_gultekin_mc`, arc-filtered to $R < R_{\text{safe}}$) · nb07e arc-spectrum subtraction. The three agree at $< 1\sigma$ (267.32 integral vs 256.17 discrete, narrow).
- Caching + seed discipline. Every expensive bootstrap writes a `.npz` keyed by its parameters (`results/<sweep>/<params>_N{N}.npz`); a cached result is never re-run. Seeds are deterministic (§2.4.1 step 7) so re-runs bit-reproduce; N=50 smoke before N=500 production; joblib with BLAS=1/worker to avoid over-subscription.
- Deterministic values registry (single source of truth). `scripts/paper_values.py` recomputes every headline number from the result caches (with per-entry provenance + formula) → emits `results/PAPER_VALUES.json`. The ApJL figures load that JSON (no hard-coded literals); the summary docs are validated by `paper_values.py --check <files>`, an anchored, precision-aware drift linter (parses only the canonical $\sigma_e = N \pm M / \log M \star = N$ statements; dated snapshots carry a `pv-skip-file` marker). Regenerate + verify: `python scripts/paper_values.py && python scripts/paper_values.py --check *.md`.

§3 Results

Figure 2 (left panel) — the integrated spectrum + SED

- Left: KCWI/KCRM Red-Low integrated, I(r)-convolved spectrum of the deflector elliptical, with the ppxf best fit overlaid (and inline residuals — Rodrigo’s standard nested-gridspec style). Loads from the wide-window fit + bootstrap pool.
- Right: HST+JWST four-band SED + Bagpipes posterior model. Loads `results/bagpipes_sed_results.npz`.
- Figure file: `AGEL0206_sigma_e_SED_final_wide.pdf` (per outline) / `results/AGEL0206_spectra_SED_fit.pdf`.

§3.1 Velocity Dispersion Measurement

- Headline: $\sigma_e(<R_e) = 269.62 \pm 13.27$ km/s (asym $-13.45/+13.10$), via the Gültekin (2009) luminosity-weighted formalism evaluated with the Cappellari ppxf implementation on the I(r)-weighted $R<R_e$ aperture spectrum, plus the wild-bootstrap error pool.
- Reproduce: `scripts/run_wide_sigma_e.py --cube new_clean_hei --n_bootstrap 500`. Caches: `results/run_wide_sigma_e/new_clean_hei/wR3800_5400_arcmask_{fsps,emiles,xsl}_T*_N500.npz`.

R_e measurement and checks (for the §3.1 paragraph): $R_e = 2.305'' = 16.23$ kpc, the F140W (2.168'') + F200LP (2.441'') masked-CoG mean; supersedes IFU-only 2.61''; masked CoG removes arc/companion bias.

Systematic error budget (post-M12, 2026-06-08; computed by `scripts/paper_values.py`):

Component	$\pm$ km/s	Why
stat (N=500)	4.6 ($-5.16/+4.13$)	wild-bootstrap pool over 3 SPS $\times$ 15 degrees; already marginalizes SPS (between-SPS ± 2.04) + degree spread
I-shape (10 shapes)	2.27	peak-to-peak/2 of 10 I-weight-map shapes (266.83–271.37)
F200 mask (3 weights)	6.65	(w00 269.69 – w100 256.39)/2; cleaned cube has more signal pixels $\rightarrow$ more arc-dilution leverage. (Arc-mask-definition cross-check ± 5.85 overlaps this, not added — M12/nb13)
frame (vac/air)	5.0	structural per-SPS native-frame choice
centering (HST WCS)	4.0	5-center $\pm 0.4''$ sweep; reduction-independent
fit-window (3 windows)	3.82	wR3800_5400 269.66 / wR4000_5400 268.58 / w6500_7500 276.22; dropped from carried ± 15 on the cleaned cube
reduction-pass	3.45	(269.62 – 262.72)/2, NEW vs OLD cleaned cube; only 2 reductions — refine if a 3rd lands
R_e -source (D7 wide)	6.13	NEW M12 (2026-06-08, nb15): peak-to-peak/2 over 4 R_e estimators (mean 2.305''/F140W 2.168''/F200LP 2.441''/CaHK+G 2.902'' $\rightarrow$ 269.62/267.44/272.44/279.69). User chose full spread; CaHK+G's +10 is the rising $\sigma(R)$ gradient. Light-family-only = ± 2.50
TOTAL (sym)	13.27	quadrature
TOTAL (asym)	$-13.45 / +13.10$	preserves stat skew

- No separate SPS line is folded in: the SPS-library spread collapsed from ~ 26 km/s at the narrow window (FSFS 253.0 < EMILES 267.5 < XSL 279.7) to ~ 4 km/s at the wide window (FSFS 271.3 / EMILES 267.2 / XSL 270.4; PAPER_VALUES.json:per_sps_p50) — a key argument for the wide window; it now sits inside the pooled stat width (between-SPS ± 2.04).
- Quadrature is deliberately conservative (components not strictly independent).
- R_e -source systematic IS now folded in (± 6.13 , M12 2026-06-08): re-measured at the wide window (nb15) and included in the budget above — the narrow-window ± 16.9 is superseded. The full 4-estimator spread was adopted (user decision); the CaHK+G 2.90'' estimator drives it (+10 km/s = rising $\sigma(R)$, see below), while the light- R_e family alone gives ± 2.50 .

Systematics paragraph order (per outline): first R_e and I(r) (I-shape ± 2.27 , R_e -source flagged), then the spectrum (mask ± 6.65 , frame ± 5.0 , window ± 3.82 , reduction ± 3.45 , centering ± 4.0).

Cross-checks (narrow window, architecturally independent, superseded for headline but valid):

- §6cum cumulative I-weighted aperture ppxf: 267.32 ± 24 ; $\sigma_e(<R_e/2) \approx 225.78$; $\sigma_e(<R_e/8) \approx 209.18$.
- §7 discrete Gültekin annular sum (equal-N inside $R_{\text{safe}}=3R_e/4$): $256.17 -13.0/+12.7$.

- §7b flat- σ extrapolation: $274.37 - 16.2/+17.4$.
- nb07e arc-spectrum subtraction matches §6cum bit-identically (<1 km/s).
- Narrow Ca H+K + G window (nb09d): 267.95 ± 30.10 — headline within ~ 2 km/s.

Headline evolution (mask-audit arc): 268.98 (clean) $\rightarrow 271.87$ (+He I, M8) $\rightarrow 269.62$ (+M10 sky audit) $\rightarrow \pm 11.77$ (M11 re-derivation) $\rightarrow \pm 13.27$ (M12, +R_e-source ± 6.13 , 2026-06-08). He I pushes σ up ~ 2.9 , the 9 sky bands pull it down ~ 2.2 (near-cancel); central value 269.62 stable across M10 $\rightarrow$ M12 (only the error grew).

Compare with Greene (2016/2020) and KH13 choices: $\sigma_e(<R_e)$ here uses the Cappellari et al. 2006 eq. 1 luminosity-weighted single-aperture definition, exactly as adopted by Kormendy & Ho 2013 (M $\bullet$ - σ eq. 3) and Greene et al. 2020 ARA&A (fig. 5). We dropped $R < R_e/8$ (JFK95 σ_c) because seeing under-resolves it. For a pure elliptical, galaxy $R_e \approx$ bulge R_e , so KH13’s “bulge R_e ” equals the measured total R_e — no bulge/disk decomposition needed.

σ_e physical-meaning discussion: the deflector is a dispersion-supported elliptical bulge; $\sigma(r)$ should *decrease* gently with radius — any rising $\sigma(r)$ in raw output is a continuum-dilution artifact from the $z=1.302$ arc, not a physical gradient. Resolved kinematics (nb11): 17 central spaxels at $S/N \geq 5$, $\sigma \in [144, 251]$ km/s (median 201), declining with R , no coherent rotation above noise.

§3.2 Stellar Mass Estimate

- $\log(M_\star/M_\odot) = 11.16 \pm 0.08$ (stat) $+0.31$ (sys) at 10% flux errors [$11.04 \pm 0.14 +0.32$ at 20%] — NEW headline from the principled F200LP-located + IR-extended arc masking with raw aperture photometry (scripts/photometry_systematics.py; results/photometry_systematics_Mstar.npz; fig results/figures/Mstar_budget.png). Empirical / conservative: raw masked photometry discards the deflector light under the (now IR-extended) arc, so the central is biased LOW and the systematic is one-sided UP.
- The systematic is intentionally uneven (could be higher): the Sérsic fill-in — which restores the deflector light under the masked arc and is mask-definition-independent (per-band vs global agree to 0.01 dex) — gives $\log(M_\star/M_\odot) = 11.46$ (10%) / 11.36 (20%), i.e. the $+0.31$ dex upper reach. The mask-definition term is negligible (± 0.003 – 0.034 dex).
- Flux-error choice matters at ~ 0.10 dex: 10% $\rightarrow$ 20% lowers the central ($11.16 \rightarrow 11.04$, looser likelihood lets the prior pull mass down) and widens stat ($\pm 0.08 \rightarrow \pm 0.14$). Both quoted.
- Explicit masking-approach systematic on $M_\star = \pm 0.16$ dex (10%: ± 0.160 , 20%: ± 0.162) — peak-to-peak/2 across all masking approaches {expert-aperture, per-band/global $\times$ raw/filled} (results/Mstar_masking_systematic.npz, fig Mstar_masking_systematic.png). Decomposition: under-arc light (raw $\leftrightarrow$ filled) ± 0.15 (dominant; the adopted one-sided $+0.31$), mask-definition (per-band $\leftrightarrow$ global) ± 0.004 (negligible — filled is mask-def-independent), mask-extent (expert $\leftrightarrow$ IR-extended) 0.18 . This ± 0.16 is the symmetric statement of the same effect quoted one-sided ($+0.31$) in the headline.
- Superseded values (cross-checks, smaller masks): older expert-aperture $\log(M_\star/M_\odot) = 11.33 +0.07/-0.09$ (sits mid-range); 2D-Sérsic-total refit $11.40 +0.11/-0.15$. Consistent with the new raw $\leftrightarrow$ filled bracket [$11.16, 11.46$].
- “Typical elliptical at $z \sim 0.7$ ”: $M_\star \approx 1.4 \times 10^{11} M_\odot$ (raw headline) to $\approx 2.9 \times 10^{11}$ (fill-in upper) at $z = 0.6756$, passive/quiescent SFH (mass-weighted age ~ 5 Gyr, low sSFR). Place on the $z \approx 0.5$ – 1.0 mass-size relation (van der Wel et al. 2014; Mowla et al. 2019) and the strong-lens-deflector analog sample (Sonnenfeld et al. 2015, SL2S V — best z -match). [notebook 10 verified citations]
- X-ray / quiescence: GAP G2 — no X-ray data in repo. Source externally if you want the “truly quiescent” claim; otherwise frame from the passive SED only.

§3.3 Mass Estimates from Lens Modeling

GAP G1 — see §2.3. The range of lens-model results, the joint-posterior mass + uncertainty, the BPL vs point-mass differentiating models, $M(<r_{\text{break}})$, the pedagogical-figure intuition, the small table reproduced from Ferrami et al., the comparison to local galaxies, and the “shape of the mass profile matters” argument are all in the lens-modeling repo / Ferrami draft, not here. The only in-

repo pointer is the 4D-constraint sentence in PROJECT_BRIEF.md. Pull these numbers from the companion paper before drafting §3.3.

Key citations (verified in notebook 10)

- ppxf: Cappellari & Emsellem (2004); Cappellari (2017, MNRAS 466, 798); Cappellari (2023).
- σ_e aperture definition: Cappellari et al. (2006, MNRAS 366, 1126) eq. 1; Gültekin et al. (2009) eq. 1; Kormendy & Ho (2013, ARA&A 51, 511) eq. 3 (scatter 0.29 dex); Greene, Strader & Ho (2020, ARA&A 58, 257) fig. 5 [$\triangle$ AUDIT FLAG — see below]; Saglia et al. (2016).
- z=0 IFS calibration: Cappellari et al. (2013, ATLAS3D XV & XX); Krajnović et al. (2013, XVII); Zhu et al. (2024, MaNGA DynPop III).
- Mass-size / analogs: van der Wel et al. (2014); Mowla et al. (2019); Ge et al. (2021); Sonnenfeld et al. (2015, SL2S V) [$\triangle$ AUDIT FLAG — see below].
- Vacuum $\leftrightarrow$ air: Ciddor (1996). Wide-window precedent: Bezanson et al. (2018); van der Wel et al. (2021).
- Source-z catalog: AGEL DR2 (Carleton et al., in prep). Cluster: Hilton et al. (2021, ACT DR5).

Citation audit + referee challenges (2026-06-08)

Citation audit (vs the user’s Zotero library; “not in library” $\neq$ fabricated — most are real foundational papers simply not yet added to Zotero, flagged so the .bib can be completed):

- Verified in-library, metadata exact: Cappellari (2017, MNRAS 466, 798) ✓; Kormendy & Ho (2013, ARA&A 51, 511) ✓; Greene, Strader & Ho (2020, ARA&A 58, 257) ✓ (*metadata only — see flag*); Hilton et al. (2021, ApJS 253, 3 = ACT SZ-cluster catalog) ✓; Cappellari (2023, MNRAS 526, 3273) ✓. Also in-library and relevant: McConnell & Ma (2013) — a massive-BH $M_\bullet-\sigma$ / $M_\bullet-M_{\text{bulge}}$ anchor.
- $\triangle$ FLAG 1 — Greene, Strader & Ho (2020, ARA&A 58, 257): bibliographic metadata is CORRECT, but this review is titled “Intermediate-Mass Black Holes.” It does provide $M_\bullet-\sigma$ / $M_\bullet-M_\star$ relations (legitimately citable as a *comparison* relation, as Fig. 4 uses it), but for a $\sim 10^9 M_\odot$ SMBH the *primary* anchors should be KH13 / McConnell & Ma 2013 / Saglia+2016, with Greene+20 as one of the overplotted comparison relations. ACTION: verify the “fig. 5” number (the scaling-relation figure in Greene+20 may not be fig. 5) and ensure the text frames it as a comparison, not the primary massive-BH relation. (*Not a fabrication — a framing/figure-number check.*)
- $\triangle$ FLAG 2 — Sonnenfeld “SL2S V” (2015): the library contains SL2S IV (Sonnenfeld et al. 2013, ApJ 777, 98), not paper V (2015, ApJ 800, 94). ACTION: decide which you mean and either correct to “IV (2013)” or add the 2015 V paper to Zotero. Do not let the 2013 record stand in for a 2015 citation.
- $\triangle$ FLAG 3 — Bezanson et al. (2018): no Bezanson-first-author 2018 item in the library. If you mean the LEGA-C velocity-dispersion paper (Bezanson et al. 2018, ApJ 858, 60), add it / verify.
- To add to Zotero before the .bib is complete (real papers, just absent; metadata NOT independently verified here — confirm on add): Cappellari & Emsellem (2004), Cappellari et al. (2006, SAURON IV), Gültekin et al. (2009), Saglia et al. (2016), Cappellari et al. (2013, ATLAS3D), Krajnović et al. (2013), Zhu et al. (2024, MaNGA DynPop III), van der Wel et al. (2014), Mowla et al. (2019), Ge et al. (2021), Ciddor (1996), van der Wel et al. (2021, LEGA-C DR3).

Referee challenges to the current doc (anticipate / address in drafting):

1. R_e is not converged (A3c, nb14). The headline $R_e = 2.305''$ is a *raw* curve-of-growth at $r_{\text{max}}=6''$; the CoG never flattens (no sky pedestal subtraction), so R_e climbs with r_{max} and sits at the top of a 2.1–2.5'' method family (sky-sub 2.14'', Sérsic 2.18''). Strongest soft spot. A referee will ask for a Sérsic / sky-subtracted / PSF-deconvolved R_e . Currently flagged, not fixed. Knock-on: R_e sets both the σ_e aperture and (weakly) $M_\star$.
2. R_e -source budget term uses the FULL 4-estimator spread (± 6.13). Including the CaHK+G 2.90'' estimator — which is larger than the light- R_e family and whose +10 km/s is the rising- $\sigma(R)$ gradient — arguably

conflates aperture-uncertainty with $\sigma(R)$ -physics. Defensible (conservative), but a referee could argue for the light-family $\pm 2.50 \rightarrow \pm 12.0$ total). Documented both ways.

3. Rising outer $\sigma(r)$ attributed to arc dilution. The claim “ $\sigma(r)$ should decrease; any rise is the $z=1.302$ arc” is the right physical call, but it is *also* what lets us drop the high- σ outer bins; make the arc-contamination evidence explicit (EW / continuum-dilution diagnostic, nb07e) so it does not read as motivated reasoning.
4. He I 3819 source-emission mask rests on a 3-pixel residual cluster — defensible (consistent across reductions, matches $z=1.302$), but small; a referee may probe it. The M8/M9/M10 audit trail covers this.
5. $\sigma_{\text{inst}} = 12.6$ km/s vs $\sigma_e \approx 270$: well-resolved, not a challenge — but state the LSF treatment.

Flagged open items (carry into drafting)

1. ~~R_e-source systematic not folded in~~ RESOLVED 2026-06-08 (M12, nb15): re-measured at the wide window (± 6.13) and folded into the budget $\rightarrow$ headline $\pm 11.77 \rightarrow \pm 13.27$.
2. Reduction-pass ± 3.45 rests on only 2 reductions — refine if a 3rd lands.
3. ~~H δ may still need targeted masking~~ RESOLVED 2026-06-08 (M12, nb16): decision = keep unmasked (H δ is well-fit, not a contaminant; masking it would discard LOSVD information). TODO in `bootstrap_ppxf.py` closed. 3b. NEW (A3c, nb14): raw CoG R_e is r_{max} -non-convergent; headline 2.305" is the top of a 2.1–2.5" family (R_e method systematic $\pm 0.08''$), not folded into $\sigma_e/M\star$ — flagged for a decision.
4. Asymmetric-error tables: older 4-corner/M10 tables print $-17.92/+17.65$ (pre-M11), M11 was $-11.98/+11.57$ (± 11.77); current (M12) is $-13.45/+13.10$ (± 13.27). Quote the M12 value (canonical: `results/PAPER_VALUES.json`).
5. No X-ray / quiescent data in repo (G2). 6. No lens-model content in repo (G1).
6. “We do not account for peculiar velocities” needs to be written in (G3 — true, just add it).
7. Source- z (1.302) error bar + its cache are a [TBD] placeholder.
8. K409 (Aug-30) PI and per-night DIMM seeing still TBD for acknowledgements.

Internal appendix — σ_e measurement audit trail (M8→M12) — NOT for the manuscript

This is the iterative history of how the final §2.4.3 masking + systematic-component set was reached. The Methods section should present the final procedure (§2.4) and the budget (§3.1), not this trail. Kept here as internal provenance / referee-rebuttal backup. Each step is also a TESTS row with its own cache.

- M8 — He I 3819 (added as a source-emission mask). A 3-pixel positive-residual cluster at def-rest 5244–5248 Å = source-rest 3818–3820 Å matches He I 3819.6 from the $z=1.302$ source; consistent across both reductions $\rightarrow$ astrophysical, not a CR. Masked. σ_e (NEW) 268.98 $\rightarrow$ 271.87.
- M9 — 5193–5204 Å bump (deliberately NOT masked). A +5–7% feature; checked against the arc spectrum (no source counterpart) and the noise spectrum (no sky counterpart) $\rightarrow$ it is the Mg b LOSVD wing (signal). Smoke-masking it drops σ_e by 7 km/s $\rightarrow$ DO NOT mask. (Same “it’s signal, not contamination” logic later applied to H δ in M12.)
- M10 — full sky-line audit. Flag every band with `cube noise_sky > 2.5 × median` across the fit window, and cross-check each against the arc spectrum to confirm no source counterpart $\rightarrow$ 9 OH/sky bands added to `BAD_PIXELS_REST` (26 $\rightarrow$ 35 entries). σ_e (NEW) 271.87 $\rightarrow$ 269.62.
- M11 — cube-matched systematic re-derivation. Re-ran the I-shape / mask-weight / fit-window sweeps on the NEW cube + M10 masks (replacing carried OLD-cube values). The fit-window term collapsed $\pm 15 \rightarrow \pm 3.82$ on the cleaned cube; total sys $\pm 17.16 \rightarrow \pm 10.81$ (total ± 11.77).
- M12 (2026-06-08) — R_e-source fold-in + two cross-checks. Folded the wide-window R_e-source systematic (± 6.13) into the budget $\rightarrow$ sys ± 12.43 , total ± 13.27 . Added the arc-mask-definition cross-check (± 5.85 ; overlaps the F200-mask term, not double-counted) and resolved the H δ open item (keep unmasked — well-fit, masking it discards LOSVD information). Central σ_e 269.62 unchanged across M10→M12; only the error grew.

Net effect of the trail: central σ_e settled at 269.62 km/s by M10 and has not moved since; the error bar evolved ± 17.78 (pre-M11) $\rightarrow \pm 11.77$ (M11) $\rightarrow \pm 13.27$ (M12). The mask set (He I + 35-entry sky/CR + no-Balmer) and the 7-component budget that the final §2.4 procedure uses are the M12 result.