

MODELLING FOR ENGINEERING AND HUMAN BEHAVIOUR 2018

Instituto Universitario de Matemática Multidisciplinar

**MATHEMATICAL
MODELLING IN
ENGINEERING
& HUMAN
BEHAVIOUR 2018**

*Valencia, Spain
July 16th-18th,
2018*



L. Jódar, J. C. Cortés and L. Acedo(Editors)

**Instituto Universitario de
Matemática Multidisciplinar**


im²

Instituto de Matemática Multidisciplinar



**UNIVERSITAT
POLITÈCNICA
DE VALÈNCIA**

Ciudad Politécnica de la Innovación



MODELLING FOR ENGINEERING, & HUMAN BEHAVIOUR 2018

Instituto Universitario de Matemática Multidisciplinar

Universitat Politècnica de València

Valencia 46022, SPAIN

Edited by

Lucas Jódar, Juan Carlos Cortés and Luis Acedo

Instituto Universitario de Matemática Multidisciplinar

Universitat Politècnica de València

I.S.B.N.: 978-84-09-07541-6

CONTENTS

1. **A model for making choices with fuzzy soft sets in an intertemporal framework**, by J. C. R. Alcantud, and M. J. Muñoz Pag: 1-5
2. **Methylphenidate and the Self-Regulation Therapy increase happiness and reduce depression: a dynamical mathematical model**, by S. Amigó, J. C. Micó, and A. Caselles Pag: 6-9
3. **A procedure to predict the short-term glucose level in a diabetic patient which captures the uncertainty of the data**, by C. Burgos, J. C. Cortés, D. Martínez-Rodríguez, J. I. Hidalgo, and R.J. Villanueva.....Pag: 10-15
4. **Assessing organizational risk in industry by evaluating interdependencies among human factors through the DEMATEL methodology**, by S. Carpitella, F. Carpitella, A. Certa, J. Benítez, and J. IzquierdoPag: 16-24
5. **Selection of an anti-torpedo decoy for the new frigate F-110 by using the GMUBO method**, by R. M. Carreño, J. Martínez, and J. BenitoPag: 25-30
6. **A modelling methodology based on General Systems Theory**, by A. Caselles .Pag: 31-34
7. **Dynamics of the general factor of personality as a consequence of alcohol consumption**, by S. Amigó, A. Caselles, J. C. Micó, M. T. Sanz, and D. SolerPag: 35-38
8. **An optimal eighth-order scheme for multiple roots applied to some real life problems**, by R. Behl, E. Martínez, F. Cevallos, and A. S. AlshomraniPag: 39-43
9. **Optimal Control of Plant Virus Propagation**, by B. Chen and M. Jackson Pag: 44-49
10. **On the inclusion of memory in Traub-type iterative methods for solving nonlinear equations**, by F. I. Chicharro, A. Cordero, N. Garrido, and J. R. Torregrosa .. Pag: 50-55
11. **Mean square analysis of non-autonomous second-order linear differential equations with randomness**, by J. Calatayud, J. C. Cortés, M. Jornet and L. Villafuerte Pag: 56-62
12. **A Statistical Model with a Lotka-Volterra Structure for Microbiota Data**, by I. Creus, A. Moya, and F. J. SantonjaPag: 63-68
13. **Some new Hermite matrix polynomials series expansions and their applications in hyperbolic matrix sine and cosine approximation**, by E. Defez, J. Ibáñez, J. Peinado, P. Alonso, J. M. Alonso, and J. Sastre Pag: 69-78

14. **A novel optimization technique for railway wheel rolling noise reduction**, by J. Gutiérrez, X. García, J. Martínez, E. Nadal, and F. D. Denia Pag: 79-84
15. **Improving the order of convergence of Traub-type derivative-free methods**, by F. I. Chicharro, A. Cordero, N. Garrido, and J. R. TorregrosaPag: 85-90
16. **Efficient decoupling technique applied to the numerical time integration of advanced interaction models for railway dynamics**, by J. Giner, J. Marínez, F. D. Denia, and L. Baeza Pag: 91-96
17. **Matrix-free block Newton method to compute the dominant λ -modes of a nuclear power reactor**, by A. Carreño, L. Bergamaschi, A. Martínez, A. Vidal, D. Ginestar, and G. VerdúPag: 97-103
18. **A new automatic gonad differentiation for salmon gender identification based on Echography image treatment**, by A. Sancho, L. Andrés, B. Baydal, and J. Real .. Pag: 104-109
19. **A New Earthwork Measurement System based on Stereoscopic Vision by Unmanned Aerial System flights**, by V. Espert, P. Moscoso, T. Real, M. Martínez, and J. RealPag: 110-115
20. **A New Forest Measurement and Monitoring System Based on Unmanned Aerial Vehicles Imaging**, by F. Ribes, V. Ramos, V. Espert, and J. RealPag: 116-121
21. **A new non-intrusive and real time monitoring technique for pavement execution based on Unmanned Aerial Vehicles flights**, by T. Real, P. Moscoso, V. Espert, A. Sancho, and J. Real Pag: 122-127
22. **A New Road Type Response Roughness Measurement System for existent defects localization and quantification**, by F. J. Veá, C. Masanet, M. Ballester, R. Redón, and J. RealPag: 128-133
23. **Application of an analytical solution based on beams on elastic foundation model for precast railway transition wedge design automatization**, by J. L. Pérez, M. Labrado, T. Real, A. Zorzona, and J. RealPag: 134-139
24. **Development of an innovative wheel damage detection system based on track vibration response on frequency domain**, by R. Auñon, B. Baydal, S. Nuñez, and J. RealPag: 140-145
25. **Mathematical characterization of liquefaction phenomena for structure foundation monitoring**, by P. Moscoso, R. Sancho, E. Colomer, and J. Real Pag: 146-151
26. **Neural Network application for concrete compression strength evolution prediction**, by T. Real, M. Labrado, B. Baydal, and J. Real Pag: 152-157
27. **Numerical simulation of lateral railway dynamic effects for a new stabilizer sleeper design**, by F. J. Fernández, T. Real, A. Zorzona, and J. RealPag: 158-162

28. **Operational costs optimization method in transport systems for open-pit mines**, by F. Halles, F. Ribes, C. Masanet, R. Redón, and J. RealPag: 163-168
29. **Structural Railway Bridge Health monitoring by means of data analysis**, by F. Ribes, C. Zamorano, P. Moscoso, and J. RealPag: 169-174
30. **A spatial model for mean house mortgage appraisal value in boroughs of the city of Valencia**, by M. A. López, N. Guadalajara, A. Iftimi, and A. Usai 175-180
31. **Third order root-finding methods based on a generalization of Gander's result**, by S. Busquier, J. M. Gutiérrez, and H. Ramos Pag: 181-184
32. **ASSESSMENT OF A GRAPHIC MODEL FOR SOLVING DELAY TIME MODEL INSPECTION CASES OF REPAIRABLE MACHINERY. PREDICTION OF RISK WHEN SELECTING INSPECTION PERIODS**, by F. Pascual, E. Larrodé, and V. Muerza Pag: 185-189
33. **A high order iterative scheme of fixed point for solving nonlinear Fredholm integral equations**, by M. A. Hernández, M. Ibáñez, E. Martínez, and S. Singh Pag: 190-194
34. **Some parametric families improving Newton's method**, by A. Cordero, S. Masallén, and J. R. Torregrosa Pag: 195-200
35. **Modeling consumer behavior in Spain**, by P. Merello, L. Jódar, G. Douklia, and E. de la PozaPag: 201-208
36. **Hamiltonian approach to human personality dynamics: an experiment with methylphenidate**, by J. C. Micó, S. Amigó, and A. CasellesPag: 209-212
37. **A Pattern Recognition Bayesian Model for the appearance of Pathologies in Automated Systems**, by M. Alacreu, N. Montes, E. García, and A. Falco ..Pag: 213-218
38. **A study of the seasonal forcing in SIRS models for Respiratory Syncytial Virus (RSV) using a constant period of temporary immunity**, by L. Acedo, J. A. Morano, and R. J. VillanuevaPag: 219-226
39. **Improving urban freight distribution through techniques of multicriteria decision making. An AHP-GIS approach**, by V. Muerza, C. Thaller, and E. LarrodéPag: 227-232
40. **Nonlinear transport through thin heterogeneous membranes**, by A. Muntean Pag: 233-236
41. **Application of the transfer matrix method for modelling Cardan mechanism of a real vehicle**, by P. Hubrý and T. Nhlík Pag: 237-242
42. **The RVT method to solve random non-autonomous second-order linear differential equations about singular-regular points**, by J. C. Cortés, A. Navarro, J. V. Romero, and M. D. Roselló Pag: 243-248

43. **On some properties of the PageRank versatility**, by F. Pedroche, R. Criado, E. García, and M. RomancePag: 249-254
44. **Network clustering strategies for setting degree predictors based on deep learning architectures**, by F. J. Pérez, E. Navarro, J. M. García, and J. Alberto ConejeroPag: 255-261
45. **Qualitative preserving stable difference methods for solving nonlocal biological dynamic problems**, by M. A. Piqueras, R. Company, and L. JódarPag: 262-267
46. **Probabilistic solution of a random model to study the effectiveness of anti-epileptic drugs**, by E. M. Sánchez-Orgaz, I. Barrachina, A. Navarro, and M. Ramos Pag: 268-273
47. **Weighted graphs to redefine the centrality measures**, by M. D. López, J. Rodrigo, C. Puente, and J. A. OlivasPag: 274-279
48. **Numerical solution to the random heat equation with zero Cauchy-type boundary conditions**, by J. C. Cortés, A. Navarro, J. V. Romero, and M. D. Roselló .. Pag: 280-285
49. **A Multistate Model for Non Muscle Invasive Bladder Carcinoma**, by C. Santamaría, B. García, and G. Rubio Pag: 286-291
50. **Birth rate and population pyramid: A stochastic dynamical model**, by J. C. Micó, D. Soler, M. T. Sanz, A. Caselles, and S. Amigó Pag: 292-297
51. **Application of the finite element method in the analysis of oscillations of rotating parts of machine mechanisms**, by P. Hubrý, and D. Smetanová Pag: 298-302
52. **Using Integer Linear Programming to minimize the cost of the thermal refurbishment of a faade: An application to building 1B of the Universitat Politècnica de València, Spain**, by D. Soler, A. Salandin, and M. Bevivino Pag: 303-308
53. **Modeling the Effects of the Immune System on Bone Fracture Healing**, by I. Trejo, H. Kojouharov, and B. Chen-CharpentierPag: 309-314
54. **Metamaterial Acoustics on the Einstein Cylinder**, by M. M. Tung Pag: 315-324
55. **Extrapolated Stabilized Explicit Runge-Kutta methods**, by J. Martín and A. Kleefeld Pag: 325-331
56. **Modelling and simulation of biological pest control in broccoli production**, by L. V. Vela-Arévalo, R. A. Ku-Carrilo, and S. E. Delgadillo-Alemán Pag: 332-337
57. **Preliminary study of fuel assembly vibrations in a nuclear reactor**, by A. Vidal, D. Ginestar, A. Carreño and G. VerdúPag: 338-343
58. **Evolution and prediction with uncertainty of the bladder cancer of a patient using a dynamic model**, by C. Burgos, N. García, D. Martínez, and R. J. Villanueva Pag: 344-348

59. **Dynamics of a family of Ermakov-Kalitlin type methods**, by A. Cordero, J. R. Torregrosa, and P. Vindel Pag: 349-353
60. **A Family of Optimal Fourth Order Methods for Multiple Roots of Non-linear Equations**, by F. Zafar, A. Cordero, and J. R. TorregrosaPag: 354-359
61. **Randomizing the von Bertalanffy growth model: Theoretical analysis and computing**, by J. Calatayud, J.-C. Cortés, and M. Jornet Pag: 360-365
62. **A Gauss-Legendre Product Quadrature for the Neutron Transport Equation**, by A. Bernal, S. Morató, R. Miró, and G. Verdú Pag: 366-371
63. **PGD path planning for dynamic obstacle robotic problems**, by L. Hilario, N. Montés, M. C. Mora, E. Nadal, A. Falcó, F. Chinesta and J. L. Duval Pag: 372-376

Some new Hermite matrix polynomials series expansions and their applications in hyperbolic matrix sine and cosine approximation ^{*}

E. Defez[★], J. Ibáñez[†], J. Peinado[§], P. Alonso[‡], J. M. Alonso[†], J. Sastre[‡]

[★] Instituto de Matemática Multidisciplinar.

[†] Instituto de Instrumentación para Imagen Molecular.

[§] Departamento de Sistemas Informáticos y Computación.

[‡] Grupo Interdisciplinar de Computación y Comunicaciones.

[‡] Instituto de Telecomunicaciones y Aplicaciones Multimedia.

Universitat Politècnica de València, Camino de Vera s/n, 46022, Valencia, España.

edefez@imm.upv.es, {jjibanez, jpeinado, palonso, jmalonso}@dsic.upv.es, jorsasma@iteam.upv.es

1 Introduction and notation

Hermite matrix polynomial $H_n(x, A)$ has the generating function, see [1]:

$$e^{xt\sqrt{2A}} = e^{t^2} \sum_{n \geq 0} \frac{H_n(x, A)}{n!} t^n, \quad (1)$$

from following expressions for the matrix hyperbolic sine and cosine are derived:

$$\left. \begin{aligned} \cosh(xt\sqrt{2A}) &= e^{t^2} \sum_{n \geq 0} \frac{H_{2n}(x, A)}{(2n)!} t^{2n} \\ \sinh(xt\sqrt{2A}) &= e^{t^2} \sum_{n \geq 0} \frac{H_{2n+1}(x, A)}{(2n+1)!} t^{2n+1} \end{aligned} \right\}, \quad x \in \mathbb{R}, |t| < \infty. \quad (2)$$

***Acknowledgements:** This work has been partially supported by Spanish Ministerio de Economía y Competitividad and European Regional Development Fund (ERDF) grants TIN2017-89314-P and by the Programa de Apoyo a la Investigación y Desarrollo 2018 of the Universitat Politècnica de València (PAID-06-18) grants SP20180016.

Recently we have shown the following formulas which are a generalization of formulas (2):

$$\left. \begin{aligned} \sum_{n \geq 0} \frac{H_{2n+1}(x, A)}{(2n)!} t^{2n} &= e^{-t^2} \left[H_1(x, A) \cosh \left(xt\sqrt{2A} \right) - 2t \sinh \left(xt\sqrt{2A} \right) \right], \\ \sum_{n \geq 0} \frac{H_{2n+2}(x, A)}{(2n+1)!} t^{2n+1} &= e^{-t^2} \left[H_1(x, A) \sinh \left(xt\sqrt{2A} \right) - 2t \cosh \left(xt\sqrt{2A} \right) \right], \\ \sum_{n \geq 0} \frac{H_{2n+3}(x, A)}{(2n+1)!} t^{2n+1} &= e^{-t^2} \left[(H_2(x, A) + 4t^2 I) \sinh \left(xt\sqrt{2A} \right) - 4t H_1(x, A) \cosh \left(xt\sqrt{2A} \right) \right]. \end{aligned} \right\} \quad (3)$$

We will use formulas (3) to obtain a new expansion of the hyperbolic matrix sine and cosine in Hermite matrix polynomials series.

Throughout this paper, we denote by $\mathbb{C}^{r \times r}$ the set of all the complex square matrices of size r . We denote by Θ and I , respectively, the zero and the identity matrix in $\mathbb{C}^{r \times r}$. If $A \in \mathbb{C}^{r \times r}$, we denote by $\sigma(A)$ the set of all the eigenvalues of A . For a real number x , $\lfloor x \rfloor$ denotes the lowest integer not less than x and $\lceil x \rceil$ denotes the highest integer not exceeding x .

We recall that for a positive stable matrix $A \in \mathbb{C}^{r \times r}$ the n -th Hermite matrix polynomial is defined in [1] by:

$$H_n(x, A) = n! \sum_{k=0}^{\lfloor \frac{n}{2} \rfloor} \frac{(-1)^k \left(\sqrt{2A} \right)^{n-2k}}{k!(n-2k)!} x^{n-2k}, \quad (4)$$

which satisfies the three-term matrix recurrence:

$$\left. \begin{aligned} H_m(x, A) &= x\sqrt{2A}H_{m-1}(x, A) - 2(m-1)H_{m-2}(x, A), \quad m \geq 1, \\ H_{-1}(x, A) &= \Theta, \quad H_0(x, A) = I. \end{aligned} \right\} \quad (5)$$

2 Some new Hermite matrix series expansions for the hyperbolic matrix cosine and sine

Let $A \in \mathbb{C}^{r \times r}$ be a positive stable matrix, then the matrix polynomial $H_1(x, A) = \sqrt{2A}x$ is invertible if $x \neq 0$. Substituting $\sinh \left(xt\sqrt{2A} \right)$ given in (2) into the first expression of (3) we obtain the following new rational expression for the hyperbolic matrix cosine in terms of Hermite matrix polynomials:

$$\begin{aligned} \cosh\left(xt\sqrt{2A}\right) &= e^{t^2} \left(\sum_{n \geq 0} \frac{H_{2n+1}(x, A)}{(2n)!} \left(1 + \frac{2t^2}{2n+1}\right) t^{2n} \right) [H_1(x, A)]^{-1}, \\ x \in \mathbb{R} \sim \{0\}, |t| < +\infty. \end{aligned} \quad (6)$$

Substituting $\sinh\left(xt\sqrt{2A}\right)$ given in (2) into the second expression of (3) and using the three-term matrix recurrence (5) we obtain the expression of $\cosh\left(xt\sqrt{2A}\right)$ given in (2).

On the other hand, replacing the expression of $\sin\left(xt\sqrt{2A}\right)$ given in (2) into the third expression of (3), we obtain another new rational expression for the hyperbolic matrix cosine in terms of Hermite matrix polynomials:

$$\begin{aligned} \cosh\left(xt\sqrt{2A}\right) &= \\ &= \frac{-e^{t^2}}{4} \left[\sum_{n \geq 0} \frac{H_{2n+3}(x, A)}{(2n+1)!} t^{2n} - (H_2(x, A) + 4t^2 I) \star \left(\sum_{n \geq 0} \frac{H_{2n+1}(x, A)}{(2n+1)!} t^{2n+1} \right) \right] [H_1(x, A)]^{-1}, \\ x \in \mathbb{R} \sim \{0\}, |t| < +\infty. \end{aligned} \quad (7)$$

Comparing (7) with (6), we observe that it always has a matrix product more when evaluating (7), the matrix product remarked by symbol “ \star ” in (7). Due to the importance of reducing the number of matrix products, see [2–4] for more details, we will focus mainly on the expansion (6).

From (4), it follows that, for $x \neq 0$:

$$\begin{aligned} H_{2n+1}(x, A) [H_1(x, A)]^{-1} &= \frac{(2n+1)!}{x} \sum_{k=0}^n \frac{(-1)^k x^{2(n-k)+1} (2A)^{n-k}}{k!(2(n-k)+1)!} \\ &= \tilde{H}_{2n+1}(x, A), \end{aligned} \quad (8)$$

where

$$\tilde{H}_n(x, A) = n! \sum_{k=0}^{\lfloor \frac{n}{2} \rfloor} \frac{(-1)^k (\sqrt{2A})^{n-2k-1}}{k!(n-2k)!} x^{n-2k}, \quad (9)$$

so the right side of (8) is still defined in the case where the matrix A is singular. In this way, we can re-write the relation (6) in terms of the matrix polynomial $\tilde{H}_{2n+1}(x, A)$:

$$\cosh\left(xt\sqrt{2A}\right) = e^{t^2} \left(\sum_{n \geq 0} \frac{\tilde{H}_{2n+1}(x, A)}{(2n)!} \left(1 + \frac{2t^2}{2n+1}\right) t^{2n} \right), \quad (10)$$

$x \in \mathbb{R}, |t| < +\infty.$

Replacing the matrix A by matrix $A^2/2$ in (10) we can avoid the square roots of matrices, and taking $x = \lambda, \lambda \neq 0, t = 1/\lambda$, we finally obtain

$$\cosh(A) = e^{\frac{1}{\lambda^2}} \left(\sum_{n \geq 0} \frac{\tilde{H}_{2n+1}\left(\lambda, \frac{1}{2}A^2\right)}{(2n)!\lambda^{2n+1}} \left(1 + \frac{2}{(2n+1)\lambda^2}\right) \right), \quad 0 < \lambda < +\infty. \quad (11)$$

3 Numerical approximations

Truncating the given series (11) until order m , we obtain the approximation $CH_m(\lambda, A) \approx \cosh(A)$ defined by

$$CH_m(\lambda, A) = e^{\frac{1}{\lambda^2}} \left(\sum_{n=0}^m \frac{\tilde{H}_{2n+1}\left(\lambda, \frac{1}{2}A^2\right)}{(2n)!\lambda^{2n+1}} \left(1 + \frac{2}{(2n+1)\lambda^2}\right) \right), \quad 0 < \lambda < +\infty. \quad (12)$$

Working analogously to the proof of the formula (3.6) of [5] one gets, for $x \neq 0$ the following bound:

$$\left\| \tilde{H}_{2n+1}\left(x, \frac{1}{2}A^2\right) \right\|_2 \leq (2n+1)! \frac{e \sinh\left(|x| \|A^2\|_2^{1/2}\right)}{|x| \|A^2\|_2^{1/2}}. \quad (13)$$

Then we can obtain the following expression for the approximation error:

$$\begin{aligned} \|\cosh(A) - CH_m(\lambda, A)\|_2 &\leq e^{\frac{1}{\lambda^2}} \sum_{n \geq m+1} \frac{\left\| \tilde{H}_{2n+1}\left(\lambda, \frac{1}{2}A^2\right) \right\|_2}{(2n)!\lambda^{2n+1}} \left(1 + \frac{2}{(2n+1)\lambda^2}\right) \\ &\leq \frac{e^{1+\frac{1}{\lambda^2}} \sinh\left(\lambda \|A^2\|_2^{1/2}\right)}{\lambda^2 \|A^2\|_2^{1/2}} \sum_{n \geq m+1} \frac{2n+1}{\lambda^{2n}} \left(1 + \frac{2}{(2n+1)\lambda^2}\right). \end{aligned} \quad (14)$$

Taking $\lambda > 1$ it follows that $\frac{2}{(2n+1)\lambda^2} < 1$, and one gets

$$\begin{aligned} \sum_{n \geq m+1} \frac{2n+1}{\lambda^{2n}} \left(1 + \frac{2}{(2n+1)\lambda^2}\right) &\leq 2 \sum_{n \geq m+1} \frac{2n+1}{\lambda^{2n}} \\ &= \frac{4 + (4m+6)(\lambda^2 - 1)}{\lambda^{2m} (\lambda^2 - 1)^2}, \end{aligned}$$

m	z_m	λ_m
2	0.0020000000061361199	909.39256098888882
4	0.079956209874370632	99.997970988888895
6	0.34561400005673254	39.999499988888893
9	1.1120032200657	17.997896988889799
12	2.2373014291079998	11.882978988901458
16	4.1086396680000004	7.9999999964157498

Table 1: Values of z_m and λ_m for $\cosh(A)$.

	$m_1 = 2$	$m_2 = 4$	$m_3 = 6$	$m_4 = 9$	$m_5 = 12$	$m_6 = 16$
\bar{m}_k	1	2	3	5	7	11
\tilde{m}_k	1	2	4	10	13	17
$f_{m_k}(\max)$	0	0	$1.9 \cdot 10^{-17}$	$6.0 \cdot 10^{-19}$	$1.4 \cdot 10^{-26}$	$1.3 \cdot 10^{-35}$

Table 2: Values \bar{m}_k , \tilde{m}_k , and f_{\max} .

thus from (14) we finally obtain:

$$\|\cosh(A) - CH_m(\lambda, A)\|_2 \leq \frac{e^{1+\frac{1}{\lambda^2}} \sinh\left(\lambda \|A^2\|_2^{1/2}\right) (4 + (4m+6)(\lambda^2 - 1))}{\|A^2\|_2^{1/2} \lambda^{2m+2} (\lambda^2 - 1)^2}. \quad (15)$$

From this expression (15) we derived the optimal values $(\lambda_m; z_m)$ such that

$$z_m = \max \left\{ z = \|A^2\|_2; \frac{e^{1+\frac{1}{\lambda^2}} \sinh\left(\lambda z^{1/2}\right) (4 + (4m+6)(\lambda^2 - 1))}{z^{1/2} \lambda^{2m+2} (\lambda^2 - 1)^2} < u \right\}$$

where u is the unit roundoff in IEEE double precision arithmetic, $u = 2^{-53}$. The optimal values of m , z and λ have been obtained with MATLAB. The results are given in the Table 1.

If $\cosh(A)$ is calculated from the Taylor series, then the absolute forward error of the Hermite approximation of $\cosh(A)$, denoted by E_f , can be computed as

$$E_f = \|\cosh(A) - P_{m_k}(B)\| = \left\| \sum_{i \geq \bar{m}_k} f_{m_k, i} B^i \right\| \cong \left\| \sum_{i \geq \tilde{m}_k} f_{m_k, i} B^i \right\|,$$

where the values of \bar{m}_k and \tilde{m}_k for each $m_k \in \{2, 4, 6, 9, 12, 16\}$ appear in the Table 2.

Scaling factor s and the order of Hermite approximation m_k are obtained by the following:

Theorem 3.1 ([6]) Let $h_l(x) = \sum_{i \geq l} p_i x^i$ be a power series with radius of convergence w , $\tilde{h}_l(x) = \sum_{i \geq l} |p_i| x^i$, $B \in \mathbb{C}^{n \times n}$ with $\rho(B) < w$, $l \in \mathbb{N}$ and $t \in \mathbb{N}$ with $1 \leq t \leq l$. If t_0 is the multiple of t such that $l \leq t_0 \leq l + t - 1$ and

$$\beta_t = \max\{d_j^{1/j} : j = t, l, l+1, \dots, t_0-1, t_0+1, t_0+2, \dots, l+t-1\},$$

where d_j is an upper bound for $\|B^j\|$, $d_j \geq \|B^j\|$, then

$$\|h_l(B)\| \leq \tilde{h}_l(\beta_t).$$

We have empirically verified that by neglecting the coefficients whose absolute value is lower than u , the efficiency results are far superior to the state-of-the-art algorithms, with also excellent accuracy.

4 Numerical experiments

The MATLAB's implementation `coshmtayher` is a modification of the MATLAB's code `coshher` given in [5], replacing the original Hermite approximation `coshher` by the new Hermite matrix polynomial obtained from (11). In this section, we compare the new MATLAB function developed in this paper, `coshmtayher`, with the functions `coshher` and `funmcosh`:

- `coshmtayher`. New code based on the new developments of Hermites matrix polynomials (11).
- `coshher`. Code based on the Hermite series for the hyperbolic matrix cosine [5].
- `funmcosh`. MATLAB function `funm` for compute matrix functions, i. e. the hyperbolic matrix cosine.

The tests have been develop using MATLAB (R2017b), runing on an Apple Macintosh iMac 27" (iMac retina 5K 27" late 2015) with a quadcore INTEL i7-6700K 4 Ghz processor and 16 Gb of RAM.

The following sets of matrices have been used:

- a) One hundred diagonalizable matrices of size 128×128 . Table 3 show the percentage of cases in which the relative errors of `coshmtayher` (new Hermite code) are lower, greater or equal than the relative errors of `coshher` (Hermite code) and `funmcosh` (funm code). Table 4 shows the matrix products of each method. Graphics with the Normwise relative errors, see [7, p. 253] and Performance Profile, see [7, p. 254], are given in Figure 1.
- b) One hundred non diagonalizables matrices of size 128×128 with multiple eigenvalues randomly generated. Table 5 shows the percentage of cases in

which the relative errors of `coshmtayher` are lower, greater or equal than the relative errors of `coshher` and `funmcosh`. Table 6 shows the matrix products of each method. Graphics of the Normwise relative errors and the Performance Profile are given in Figure 2.

- c) Ten matrices from the Eigtool MATLAB [8] package with size 128×128 , and thirty matrices from the function `matrix` of the Matrix Computation Toolbox [9] with dimensions lower or equal than 128. These matrices have been chosen because they have more varied and significant characteristics. Table 7 shows the percentage of cases in which the relative errors of `coshmtayher` are lower, greater or equal than the relative errors of `coshher` and `funmcosh`. Table 8 shows the matrix products of each method. Graphics of the Normwise relative errors and the Performance Profile are given Figure 3.

$E(\text{coshmtayher}) < E(\text{coshher})$	47.50%
$E(\text{coshmtayher}) > E(\text{coshher})$	50.00%
$E(\text{coshmtayher}) = E(\text{coshher})$	3.00%
$E(\text{coshmtayher}) < E(\text{funmcosh})$	100.00%
$E(\text{coshmtayher}) > E(\text{funmcosh})$	0.00%
$E(\text{coshmtayher}) = E(\text{funmcosh})$	0.00%

Table 3: Comparative between the methods

<i>cosmtayher</i>	<i>coshher</i>	<i>funmcosh</i>
671	973	1500

Table 4: Matrix products

$E(\text{coshmtayher}) < E(\text{coshher})$	52.50%
$E(\text{coshmtayher}) > E(\text{coshher})$	47.00%
$E(\text{coshmtayher}) = E(\text{coshher})$	1.00%
$E(\text{coshmtayher}) < E(\text{funmcosh})$	100.00%
$E(\text{coshmtayher}) > E(\text{funmcosh})$	0.00%
$E(\text{coshmtayher}) = E(\text{funmcosh})$	0.00%

Table 5: Comparative between the methods




Figure 1: Diagonalizable matrices

<i>cosmtayher</i>	<i>coshher</i>	<i>funmcosh</i>
685	989	1500

Table 6: Matrix products




Figure 2: Non diagonalizable matrices

$E(\text{coshmtayher}) < E(\text{coshher})$	57.50%
$E(\text{coshmtayher}) > E(\text{coshher})$	30.00%
$E(\text{coshmtayher}) = E(\text{coshher})$	12.50%
$E(\text{coshmtayher}) < E(\text{funmcosh})$	97.50%
$E(\text{coshmtayher}) > E(\text{funmcosh})$	2.50%
$E(\text{coshmtayher}) = E(\text{funmcosh})$	0.00%

Table 7: Comparative between the methods

<i>cosmtayher</i>	<i>coshher</i>	<i>funmcosh</i>
191	315	600

Table 8: Matrix products




Figure 3: Matrices from the Eigtool and the Matrix Computation Toolbox packages

5 Conclusions

The more accurate are the implementations based on the Hermite series: the initial MATLAB implementation (*coshher*) and the proposed MATLAB implementation based on (11) (*coshmtayher*). Also, the new implementation (*coshmtayher*) have considerably lower computational costs than the other functions.

References

- [1] J. Jódar, R. Company, Hermite matrix polynomials and second order matrix differential equations, *Approximation Theory and its Applications* 12 (2) (1996) 20–30.
- [2] J. Sastre, J. Ibáñez, E. Defez, P. Ruiz, New scaling-squaring Taylor algorithms for computing the matrix exponential, *SIAM Journal on Scientific Computing* 37 (1) (2015) A439–A455.
- [3] P. Alonso, J. Peinado, J. Ibáñez, J. Sastre, E. Defez, Computing matrix trigonometric functions with gpus through matlab, *The Journal of Supercomputing* (2018) 1–14.

- [4] J. Sastre, Efficient evaluation of matrix polynomials, *Linear Algebra and its Applications* 539 (2018) 229–250.
- [5] E. Defez, J. Sastre, J. Ibáñez, J. Peinado, Solving engineering models using hyperbolic matrix functions, *Applied Mathematical Modelling* 40 (4) (2016) 2837–2844.
- [6] J. Sastre, J. Ibáñez, P. Ruiz, E. Defez, Efficient computation of the matrix cosine, *Applied Mathematics and Computation* 219 (14) (2013) 7575–7585.
- [7] N. J. Higham, *Functions of Matrices: Theory and Computation*, SIAM, Philadelphia, PA, USA, 2008.
- [8] T. Wright, Eigtool, version 2.1, URL: web.comlab.ox.ac.uk/pseudospectra/eigtool.
- [9] N. J. Higham, *The test matrix toolbox for MATLAB (Version 3.0)*, University of Manchester Manchester, 1995.