

# Stochastic Calculus

Based on lecture notes by Prof. Xunyu Zhou

Michaelmas Term, 2015

## Contents

<b>0</b>	<b>Conventions and abbreviations</b>	<b>2</b>
<b>1</b>	<b>Preliminaries</b>	<b>2</b>
1.1	Probability Space . . . . .	2
1.2	Convergence and Uniform Integrability . . . . .	4
1.3	Stochastic Processes and Filtration . . . . .	6
1.4	Martingales . . . . .	7
<b>2</b>	<b>Brownian Motion</b>	<b>9</b>
2.1	Definition . . . . .	9
2.2	Markov property and finite-dimensional distributions . . . . .	10
2.3	The reflection principle and martingale property . . . . .	12
2.4	Quadratic variation process . . . . .	13
<b>3</b>	<b>Itô's calculus</b>	<b>15</b>
3.1	Stochastic integrals for BM . . . . .	15
3.2	Stochastic integrals for adapted processes . . . . .	16
3.3	Itô's integration for semimartingales . . . . .	17
3.4	Itô's formula . . . . .	18
3.5	Stochastic exponentials and Girsanov theorem . . . . .	20
<b>4</b>	<b>Stochastic Differential Equations</b>	<b>23</b>
4.1	Introduction . . . . .	23
4.2	Several examples . . . . .	24
4.2.1	Linear SDEs . . . . .	24
4.2.2	Geometric BM . . . . .	24
4.2.3	Cameron-Martin's formula . . . . .	24
4.3	Existence and uniqueness . . . . .	25

## 0 Conventions and abbreviations

Vectors are column vectors.

a.s.	almost surely
BM	Brownian motion
CDF	cumulative distribution function
r.v.	random variable / random vector
s.t.	such that
PDF	probability density function
SDE	stochastic differential equation
w.r.t.	with respect to

## 1 Preliminaries

### 1.1 Probability Space

**Definition 1.1.1.** Let a set  $\Omega$  be non-empty and  $\mathcal{F} \subseteq 2^\Omega$ , called a class, be non-empty. We call  $\mathcal{F}$

(i) a  $\pi$ -system if  $A, B \in \mathcal{F} \implies A \cap B \in \mathcal{F}$ .

(ii) a  $\lambda$ -system if

$$\begin{cases} \Omega \in \mathcal{F} \\ A, B \in \mathcal{F} \text{ and } A \subseteq B \implies B \setminus A \in \mathcal{F} \\ A_i \in \mathcal{F}, A_i \uparrow A, i = 1, 2, \dots^1 \implies A \in \mathcal{F} \end{cases}$$

(iii) a  $\sigma$ -field (or  $\sigma$ -algebra) if

$$\begin{cases} \Omega \in \mathcal{F} \\ A, B \in \mathcal{F} \implies B \setminus A \in \mathcal{F} \\ A_i \in \mathcal{F}, i = 1, 2, \dots \implies \bigcup_{i=1}^{\infty} A_i \in \mathcal{F} \end{cases}$$

*Remark.*  $\mathcal{F}$  is a  $\sigma$ -field if and only if  $\mathcal{F}$  is both a  $\pi$ -system and a  $\lambda$ -system.

For any class  $\mathcal{A} \subseteq 2^\Omega$ ,  $\sigma(\mathcal{A})$  is the smallest  $\sigma$ -field containing  $\mathcal{A}$ , which is called the  $\sigma$ -field generated by  $\mathcal{A}$ .

**Lemma 1.1.2** (Monotone Class Theorem). Let  $\mathcal{A} \subseteq \mathcal{F} \subseteq 2^\Omega$ .

If  $\mathcal{A}$  is a  $\pi$ -system and  $\mathcal{F}$  is a  $\lambda$ -system, then  $\sigma(\mathcal{A}) \subseteq \mathcal{F}$ .

**Definition 1.1.3.** Let  $\{\mathcal{F}_\alpha\}$  be a (possibly uncountable) family of  $\sigma$ -fields on  $\Omega$ . Then we define

$$\begin{aligned} \bigvee_{\alpha} \mathcal{F}_\alpha &:= \sigma \left( \bigcup_{\alpha} \mathcal{F}_\alpha \right) && \text{the smallest } \sigma\text{-field containing all } \mathcal{F}_\alpha; \\ \bigwedge_{\alpha} \mathcal{F}_\alpha &:= \bigcap_{\alpha} \mathcal{F}_\alpha && \text{the largest } \sigma\text{-field contained in } \mathcal{F}_\alpha. \end{aligned}$$

**Definition 1.1.4.** Let  $\Omega$  be non-empty and let  $\mathcal{F}$  be a  $\sigma$ -field on  $\Omega$ .

Then we call  $(\Omega, \mathcal{F})$  a measurable space.

A map  $\mathbf{P} : \mathcal{F} \rightarrow [0, 1]$  is called a probability measure on  $(\Omega, \mathcal{F})$  if

(i)  $\mathbf{P}(\Omega) = 1, \mathbf{P}(\emptyset) = 0$ ;

(ii)  $\forall A_i \in \mathcal{F}$  with  $A_i \cap A_j = \emptyset \ \forall j \neq i \implies \mathbf{P}(\bigcup_{i=1}^{\infty} A_i) = \sum_{i=1}^{\infty} \mathbf{P}(A_i)$ .

Then we call  $(\Omega, \mathcal{F}, \mathbf{P})$  a probability space.

We call  $\Omega$  a sample space,  $A \in \mathcal{F}$  an event and  $\omega \in \Omega$  a sample or outcome.

---

<sup>1</sup>  $A_i \uparrow A$  means  $A_1 \subseteq A_2 \subseteq \dots$  and  $A = \bigcup_{i=1}^{\infty} A_i$ .

**Definition 1.1.5** (Independence). *Two events  $A, B$  are called independent if  $\mathbf{P}(A \cap B) = \mathbf{P}(A)\mathbf{P}(B)$ . An event  $A$  is independent of a  $\sigma$ -field  $\mathcal{F}$  if  $A$  is independent of  $B \forall B \in \mathcal{F}$ . Two  $\sigma$ -fields  $\mathcal{F}, \mathcal{G}$  are independent if all  $A \in \mathcal{F}$  are independent of  $\mathcal{G}$ .*

If an event  $A$  is s.t.  $\mathbf{P}(A) = 1$ , we may denote it as “ $A$  holds,  $\mathbf{P}$ -a.s.” or just a.s. (almost surely) if the probability measure in question is clear.

$A$  is called  $\mathbf{P}$ -null if  $\mathbf{P}(A) = 0$ .

**Definition 1.1.6.**  $(\Omega, \mathcal{F}, \mathbf{P})$  is called complete if for any  $\mathbf{P}$ -null set  $A \in \mathcal{F}$ , we have  $B \in \mathcal{F}$  whenever  $B \subseteq A$ .

**Definition 1.1.7.** Given  $(\Omega, \mathcal{F}, \mathbf{P})$ .

Let  $X : \Omega \rightarrow \mathbb{R}^d$  be measurable, i.e. for every Borel measurable set  $B$  of  $\mathbb{R}^d$ ,

$$X^{-1}(B) := \{\omega : X(\omega) \in B\} \in \mathcal{F}.$$

Then  $X$  is called a random variable (or random vector, if  $d > 1$ ).

If  $X$  is integrable, denoted as  $X \in \mathcal{L}^1(\Omega, \mathcal{F}, \mathbf{P})$ , i.e.  $\int_{\Omega} |X(\omega)| d\mathbf{P}(\omega) < \infty$ , then its mean (or expectation) is

$$E[X] := \int_{\Omega} X(\omega) d\mathbf{P}(\omega).$$

In general, for  $p \geq 1$ ,  $X$  is called  $p$ -th integrable, denoted as  $X \in \mathcal{L}^p(\Omega, \mathcal{F}, \mathbf{P})$  if  $\int_{\Omega} |X(\omega)|^p d\mathbf{P}(\omega) < \infty$ .

**Definition 1.1.8.** If  $X, Y \in \mathcal{L}^2(\Omega, \mathcal{F}, \mathbf{P})$  then  $\text{Cov}(X, Y) := \mathbf{E}[(X - \mathbf{E}[X])(Y - \mathbf{E}[Y])^T]$ .

We can then define the variance as  $\text{Var}(X) = \text{Cov}(X, X)$ .

**Definition 1.1.9.** Let  $X \equiv (X_1, X_2, \dots, X_d)^T$  be a r.v..

We define the cumulative distribution function (CDF)

$$F(x) \equiv F(x_1, x_2, \dots, x_d) := \mathbf{P}(\{\omega : X_i(\omega) \leq x_i, i = 1, 2, \dots, d\}).$$

**Definition 1.1.10.** If there exists an  $f \in \mathcal{L}^1(\mathbb{R}^d)$  s.t.

$$F(x) = \int_{-\infty}^{x_d} \cdots \int_{-\infty}^{x_1} f(y_1, \dots, y_d) dy_1 \dots dy_d,$$

then  $f$  is called the probability density function (PDF) of  $F$ .

If  $X$  is a one-dimensional r.v., then

$$\begin{aligned} \mathbf{E}[X] &= \int_{\mathbb{R}} x dF(x) = \int_{-\infty}^{\infty} x f(x) dx && \text{if } f \text{ exists} \\ \mathbf{E}[g(X)] &= \int_{\mathbb{R}} g(x) dF(x) = \int_{-\infty}^{\infty} g(x) f(x) dx && \text{if } f \text{ exists and } g : \mathbb{R} \rightarrow \mathbb{R} \text{ measurable} \end{aligned}$$

*Example 1.* If

$$f(x) = [(2\pi)^d |\Sigma|]^{-1/2} \exp \left[ -\frac{1}{2} (x - \mu)^T \Sigma^{-1} (x - \mu) \right], \quad x \in \mathbb{R}^d$$

where  $\mu \in \mathbb{R}^d$ ,  $\Sigma \in \mathbb{R}^{d \times d}$  and  $\Sigma^T = \Sigma > 0^2$ , then we say that  $X$  is a normal r.v., denoted  $X \sim N(\mu, \Sigma)$ .  $X$  is also called *Gaussian*.

**Definition 1.1.11.** Let  $X$  be an integrable r.v. on  $(\Omega, \mathcal{F}, \mathbf{P})$  and  $\mathcal{G} \subseteq \mathcal{F}$ .

The conditional expectation of  $\mathbf{E}[X|\mathcal{G}]$  of  $X$  given  $\mathcal{G}$  is a r.v. (unique up to a.s.) satisfying

(i)  $\mathbf{E}[X|\mathcal{G}]$  is measurable w.r.t.  $\mathcal{G}$  (also called a  $\mathcal{G}$ -r.v.);

---

<sup>2</sup>  $\Sigma > 0$  means that  $\Sigma$  is positive definite, i.e.  $x^T \Sigma x > 0 \forall x \in \mathbb{R}^d \setminus \{0\}$ .

(ii)  $\mathbf{E}[X; A] := \mathbf{E}[X \mathbb{1}_A] = \mathbf{E}[\mathbf{E}[X|\mathcal{G}]; A] \quad \forall A \in \mathcal{G}$ .

*Remark.* (i) If  $X$  is  $\mathcal{F}$ -measurable and  $Y$  is  $\mathcal{G}$ -measurable then

$$\mathbf{E}[XY|\mathcal{G}] = Y\mathbf{E}[X|\mathcal{G}], \text{ a.s.,}$$

so long as both sides are well defined (cf. integrability requirement for a conditional r.v.).

(ii) Let  $\mathcal{G}_1 \subseteq \mathcal{G}_2 \subseteq \mathcal{F}$ , then

$$\mathbf{E}[\mathbf{E}[X|\mathcal{G}_1]|\mathcal{G}_2] = \mathbf{E}[X|\mathcal{G}_1] = \mathbf{E}[\mathbf{E}[X|\mathcal{G}_2]|\mathcal{G}_1], \text{ a.s..}$$

(iii) Jensen's inequality. Suppose  $\varphi : \mathbb{R}^d \rightarrow \mathbb{R}$  is a convex function<sup>3</sup>, then

$$\varphi(\mathbf{E}[X|\mathcal{G}]) \leq \mathbf{E}[\varphi(X)|\mathcal{G}].$$

A  $\sigma$ -field  $\sigma(X)$  generated by a r.v.  $X$  is the smallest  $\sigma$ -field contained in  $\mathcal{F}$  w.r.t. which  $X$  is measurable.

Similiarly, we can define  $\sigma(X_\alpha : \alpha \in A)$ <sup>4</sup>, for a collection of random variables.

We define  $\mathbf{E}[X|Y] := \mathbf{E}[X|\sigma(Y)]$ .

$X$  is independent of  $\mathcal{G}$  if  $\sigma(X)$  is independent of  $\mathcal{G}$ , which is equivalent to

$$\mathbf{E}[f(X)|\mathcal{G}] = \mathbf{E}[f(X)] \text{ for all bounded Borel-measurable functions } f.$$

Random variables  $X_1, \dots, X_n$  are mutually independent if  $\sigma(X_1), \dots, \sigma(X_n)$  are mutually independent, which is equivalent to

$$\mathbf{E}\left[\prod_{i=1}^n f_i(X_i)\right] = \prod_{i=1}^n \mathbf{E}[f_i(X_i)] \text{ for all bounded Borel-measurable functions } f_1, \dots, f_n.$$

In particular, if  $X, Y$  are independent, then  $\mathbf{E}[XY] = \mathbf{E}[X]\mathbf{E}[Y]$ .

## 1.2 Convergence and Uniform Integrability

**Definition 1.2.1.** Let  $X_n, n = 1, 2, 3, \dots$  and  $X : (\Omega, \mathcal{F}, \mathbf{P}) \rightarrow \mathbb{R}^d$  be r.v.s. We say  $X_n$  converges to  $X$

- almost surely (a.s.) if  $\lim_{n \rightarrow \infty} |X_n - X| = 0$  a.s.,
- in probability if  $\forall \varepsilon > 0, \lim_{n \rightarrow \infty} P(|X_n - X| > \varepsilon) = 0$ ,
- in  $\mathcal{L}^1$  if  $\lim_{n \rightarrow \infty} \mathbf{E}|X_n - X| = 0$ .

*Remark.*

a.s. convergence  $\implies$  convergence in probability

$\mathcal{L}^1$  convergence  $\implies$  convergence in probability

convergence in probability  $\implies$  there exists a subsequence that converges a.s.

**Lemma 1.2.2** (Fatou's Lemma). If  $X_n \geq 0$ <sup>5</sup>, then

$$\mathbf{E}\left[\liminf_{n \rightarrow \infty} X_n\right] \leq \liminf_{n \rightarrow \infty} \mathbf{E}[X_n].$$

*Remark.* 1. Fatou's lemma holds when  $X_n \geq Y$  and  $\mathbf{E}[Y] > -\infty$  (consider that  $X_n - Y \geq 0$ ).

<sup>3</sup>A function  $f$  is convex if  $f(tx_1 + (1-t)x_2) \leq tf(x_1) + (1-t)f(x_2) \quad \forall x_1, x_2 \in \mathbb{R}, \forall t \in [0, 1]$

<sup>4</sup> $\sigma(X_\alpha : \alpha \in A) = \sigma(\{\omega \in \Omega : X_\alpha(\omega) \in B\} : \alpha \in A, B \in \mathcal{B})$ .

<sup>5</sup>a.s. is implied here and in all other inequalities involving r.v.s.

2. When  $X_n \leq 0$ , applying the lemma to  $-X_n$  gives us

$$\mathbf{E} \left[ \limsup_{n \rightarrow \infty} X_n \right] \geq \limsup_{n \rightarrow \infty} \mathbf{E}[X_n].$$

**Lemma 1.2.3** (Monotone Convergence Theorem). *If  $X_n \uparrow X$  and  $\mathbf{E}[X_1] > -\infty$ , then*

$$\mathbf{E}[X] = \mathbf{E} \left[ \lim_{n \rightarrow \infty} X_n \right] = \lim_{n \rightarrow \infty} \mathbf{E}[X_n].$$

**Lemma 1.2.4** (Dominated Convergence Theorem). *If  $X_n \rightarrow X$  a.s. and  $|X_n| \leq Y$  with  $\mathbf{E}[Y] < \infty$ , then*

$$\mathbf{E}[X] = \mathbf{E} \left[ \lim_{n \rightarrow \infty} X_n \right] = \lim_{n \rightarrow \infty} \mathbf{E}[X_n].$$

$$\text{If } X \in \mathcal{L}^1(\Omega, \mathcal{F}, \mathbf{P}), \text{ then } \lim_{N \rightarrow \infty} \int_{\{|X| \geq N\}} |X| d\mathbf{P} = 0. \quad (1.2.1)$$

$$\text{If } X \in \mathcal{L}^1(\Omega, \mathcal{F}, \mathbf{P}), \text{ then } \int_A |X| d\mathbf{P} \rightarrow 0 \text{ as } \mu(A) \rightarrow \infty. \quad (1.2.2)$$

**Definition 1.2.5.** *Let  $\Gamma \subseteq \mathcal{L}^1(\Omega, \mathcal{F}, \mathbf{P})$  be given.*

*$\Gamma$  is said to be uniformly integrable (UI) if*

$$\lim_{N \rightarrow \infty} \sup_{X \in \Gamma} \int_{\{|X| \geq N\}} |X| d\mathbf{P} = 0.$$

*Remark.* 1. Any finite family of integrable r.v.s is UI. (Note that because the family is finite, the supremum becomes a maximum.)

2. Let  $\Gamma \subseteq \mathcal{L}^1(\Omega, \mathcal{F}, \mathbf{P})$ , and suppose there is  $Y \in \mathcal{L}^1(\Omega, \mathcal{F}, \mathbf{P})$ , s.t.  $|X| \leq Y \forall X \in \Gamma$ , then  $\Gamma$  is UI by (1.2.1).

3. Let  $\Gamma \subseteq \mathcal{L}^1(\Omega, \mathcal{F}, \mathbf{P})$  and suppose  $\exists p > 1$  s.t.  $\sup_{X \in \Gamma} \mathbf{E}|X|^p < \infty$ , then  $\Gamma$  is UI.

**Theorem 1.2.6.**  *$\Gamma \subseteq \mathcal{L}^1(\Omega, \mathcal{F}, \mathbf{P})$  is UI if and only if*

$$1. \sup_{X \in \Gamma} \mathbf{E}|X| < \infty,$$

$$2. \forall \varepsilon > 0, \exists \delta > 0 \text{ s.t. } \int_A |X| d\mathbf{P} < \varepsilon, \forall X \in \Gamma, \text{ whenever } A \in \mathcal{F}^6 \text{ and } \mathbf{P}(A) < \delta.$$

---

<sup>6</sup> $A$  here is like a more general form of  $\{|X| \geq N\}$  in the definition of UI.

**Proof. Necessity**

For any  $A \in \mathcal{F}$  and  $N > 0$ ,

$$\begin{aligned} \int_A |X| d\mathbf{P} &= \int_{A \cap \{|X| < N\}} |X| d\mathbf{P} + \int_{A \cap \{|X| \geq N\}} |X| d\mathbf{P} \\ &\leq N\mathbf{P}(A \cap \{|X| < N\}) + \int_{A \cap \{|X| \geq N\}} |X| d\mathbf{P} \\ &\leq N\mathbf{P}(A) + \underbrace{\int_{A \cap \{|X| \geq N\}} |X| d\mathbf{P}}_{\text{c.f. UI definition}}. \end{aligned}$$

Let  $A = \Omega$ . Then, for  $N$  sufficiently large,

$$\begin{aligned} \mathbf{E}[X] &\leq N\mathbf{P}(\Omega) + \int_{\{|X| \geq N\}} |X| d\mathbf{P} \\ &\leq N + 1. \end{aligned}$$

For any  $\varepsilon > 0$ , choose  $N > 0$  s.t.

$$\sup_{X \in \Gamma} \int_{\{|X| \geq N\}} |X| d\mathbf{P} \leq \frac{\varepsilon}{2}.$$

Let  $\delta = \varepsilon/(2N)$ . Then, so long as  $P(A) \leq \delta$ , we have

$$\int_{|A|} |X| d\mathbf{P} \leq N \times \delta + \frac{\varepsilon}{2} = N \times \frac{\varepsilon}{2N} + \frac{\varepsilon}{2} = \varepsilon, \quad \forall X \in \Gamma.$$

**Sufficiency** Refer to exercise sheet 2, question 3. □

**Theorem 1.2.7.** Let  $\{X_n\} \in \mathcal{L}^1(\Omega, \mathcal{F}, \mathbf{P})$ . Then  $\mathbf{E}|X_n - X| \rightarrow 0$  as  $n \rightarrow \infty$  for some  $X$  if and only if  $\{X_n\}$  is UI and  $X_n \rightarrow X$  in probability as  $n \rightarrow \infty$ .

Let  $\{A_n\} \subseteq \mathcal{F}$ . Define

$$\{A_n \text{ i.o.}\} \equiv \{A_n \text{ occurs infinitely often}\} \equiv \limsup_{n \rightarrow \infty} A_n := \bigcap_{n=1}^{\infty} \bigcup_{k=n}^{\infty} A_k.$$

**Lemma 1.2.8** (Borel-Cantelli Lemma). 1. If  $\sum_{n=1}^{\infty} \mathbf{P}(A_n) < \infty$ , then  $\mathbf{P}(\{A_n \text{ i.o.}\}) = 0$ ,

2. If  $\sum_{n=1}^{\infty} \mathbf{P}(A_n) = \infty$  and  $A_1, A_2, \dots$  are independent, then  $\mathbf{P}(\{A_n \text{ i.o.}\}) = 1$ .

### 1.3 Stochastic Processes and Filtration

Let  $T$  denote either  $\mathbb{Z}^+ := \{0, 1, 2, \dots\}$  or  $\mathbb{R}^+ = [0, \infty)$ .

**Definition 1.3.1.** A family  $X \equiv (X_t)_{t \in T}$  of r.v.s from  $(\Omega, \mathcal{F}, \mathbf{P})$  to  $\mathbb{R}^d$  is called a stochastic process.

$(X_t)_{t \in T}$  is also a mapping from  $T \times \Omega \rightarrow \mathbb{R}^d$ , which is sometimes also called a *random function*.

For any  $\omega \in \Omega$ , the function  $t \rightarrow X_t(\omega)$  is called a *sample path*. The process is called *continuous* (or càdlàg / RCLL<sup>7</sup>) if  $T = [0, \infty)$  and the sample is continuous (resp. càdlàg / RCLL) for almost all  $\omega \in \Omega$ .

Let  $X = (X_t)_{t \geq 0}$  be a stochastic process and  $0 \leq t_1 < t_2 < \dots < t_n$  be an arbitrary partition. The joint distribution of r.v.s  $(X_{t_1}, X_{t_2}, \dots, X_{t_n})$ :

$$\mu_{t_1, \dots, t_n}(dx_1, \dots, dx_n) = \mathbf{P}(X_{t_1} \in dx_1, \dots, X_{t_n} \in dx_n)$$

is called a *finite dimensional distribution* of  $(X_t)_{t \geq 0}$ .

---

<sup>7</sup>Right continuous with left limits.

**Definition 1.3.2.** Two processes  $X_t$  and  $Y_t$  are (stochastically) equivalent if  $X_t = Y_t$  a.s.,  $\forall t \in T$ .

For a measurable space  $(\Omega, \mathcal{F})$ , we introduce a montone family of  $\sigma$ -fields  $\mathcal{F}_t \subseteq \mathcal{F}$ ,  $\forall t \in T$ , satisfying

$$\mathcal{F}_{t_1} \subseteq \mathcal{F}_{t_2} \quad \forall 0 \leq t_1 \leq t_2, \quad t_1, t_2 \in T.$$

Such a family is called a *filtration*, and  $(\Omega, \mathcal{F}, \mathbf{P}; (\mathcal{F}_t)_{t \in T})$  is called a *filtered probability space*.

Set

$$\begin{aligned} \mathcal{F}_{t+} &:= \bigcap_{s>t} \mathcal{F}_s, \quad \forall t \geq 0, \\ \mathcal{F}_{t-} &:= \bigcup_{s<t} \mathcal{F}_s, \quad \forall t \geq 0. \end{aligned}$$

If  $\mathcal{F}_{t+} = \mathcal{F}_t$  (resp.  $\mathcal{F}_{t-} = \mathcal{F}_t$ ), then we say that the filtration is right (resp. left) continuous.

If  $(X_t)_{t \in T}$  is a process, then  $\mathcal{F}_t := \sigma\{X_s, s \in T, s \leq t\}$  is called the filtration generated by  $(X_t)_{t \in T}$ .

If  $(X_t)_{t \geq 0}$  is càdlàg, then it generates a right continuous filtration.

**Definition 1.3.3.** We say that  $(\Omega, \mathcal{F}, \mathbf{P})$  satisfies the usual conditions if it is complete,  $\mathcal{F}_0$  contains all the  $\mathbf{P}$ -null sets in  $\mathcal{F}$ , and  $(\mathcal{F}_t)_{t \geq 0}$  is right-continuous.

*Remark.* In future, we always assume a filtered probability space satisfies the usual conditions.

**Definition 1.3.4.** Let  $(\Omega, \mathcal{F}, \mathbf{P}; (\mathcal{F}_t)_{t \geq 0})$  be a filtered probability space, and  $(X_t)_{t \geq 0}$  a process.

1.  $(X_t)_{t \geq 0}$  is called measurable if the map  $(t, \omega) \mapsto X_t(\omega)$  is  $(\mathcal{B}(\mathbb{R}^+) \times \mathcal{F})/\mathcal{B}(\mathbb{R}^d)$ -measurable.
2.  $(X_t)_{t \geq 0}$  is called  $\mathcal{F}_t$ -adapted if  $\forall t \geq 0$ , the map  $\omega \mapsto X_t(\omega)$  is  $\mathcal{F}_t/\mathcal{B}(\mathbb{R}^d)$ -measurable.

## 1.4 Martingales

Let  $\{X_n : n \in \mathbb{Z}\}$  be a discrete time process on a filtered probability space  $(\Omega, \mathcal{F}, \mathbf{P}; (\mathcal{F}_n)_{n \in \mathbb{Z}^+})$ .

If  $X_n$  is  $\mathcal{F}_{n-1}$  measurable  $\forall n \geq 1$  and  $X_0$  is  $\mathcal{F}_0$  measurable, then we say that  $\{X_n\}$  is *predictable* or *previsible*.

On  $(\Omega, \mathcal{F}, \mathbf{P}; (\mathcal{F}_n)_{n \in \mathbb{Z}^+})$ , a measurable function  $\tau : \Omega \rightarrow \mathbb{Z}^+ \cup \{+\infty\}$  is called a *stopping time* (or *random time*) w.r.t. the filtration  $\{\mathcal{F}_n\}$  if  $\{\omega : \tau(\omega) = n\} \in \mathcal{F}_n \quad \forall n \in \mathbb{Z}^+$ .

Given  $\tau$ , the  $\sigma$ -field

$$\mathcal{F}_\tau := \{A \in \mathcal{F} : A \cap \{\tau \leq n\} \in \mathcal{F}_n \quad \forall n \in \mathbb{Z}^+\}$$

represents the information available up to the random time  $\tau$ .

**Definition 1.4.1.** Let  $\{X_n\}$  be adapted to  $(\mathcal{F}_n)$  on  $(\Omega, \mathcal{F}, \mathbf{P}; (\mathcal{F}_n)_{n \in \mathbb{Z}^+})$ . Assume that each  $X_n \in \mathcal{L}^1(\Omega, \mathcal{F}, \mathbf{P})$ .

1.  $\{X_n\}$  is called a martingale if  $\mathbf{E}[X_{n+1}|\mathcal{F}_n] = X_n \quad \forall n \in \mathbb{Z}^+$ .
2.  $\{X_n\}$  is called a supermartingale (resp. submartingale) if  $\mathbf{E}[X_{n+1}|\mathcal{F}_n] \leq X_n$  (resp.  $\mathbf{E}[X_{n+1}|\mathcal{F}_n] \geq X_n$ ).

**Theorem 1.4.2** (Doob's optional sampling theorem / optional stopping theorem). Let  $\{X_n\}$  be a martingale (resp. supermartingale) and  $\sigma, \tau$  be two bounded stopping times with  $\sigma \leq \tau$ <sup>8</sup>. Then  $\mathbf{E}[X_\tau|\mathcal{F}_\sigma] = X_\sigma$  a.s. (resp.  $\mathbf{E}[X_\tau|\mathcal{F}_\sigma] \leq X_\sigma$  a.s.).

**Theorem 1.4.3** (Kolmogorov's inequality). Let  $\{X_n\}$  be a martingale and  $X_n \in \mathcal{L}^2(\Omega, \mathcal{F}, \mathbf{P}) \quad \forall n \in \mathbb{Z}^+$ . Then

$$P\left(\max_{k \leq n} |X_k| \geq \lambda\right) \leq \frac{\mathbf{E}[|X_n|^2]}{\lambda^2}.$$

---

<sup>8</sup>Recall that this is a.s. because  $\sigma$  and  $\tau$  are random.

*Remark.* Compare this with Markov's inequality,

$$P\left(\max_{k \leq n} |X_k| \geq \lambda\right) \leq \frac{\mathbf{E}[\max_{k \leq n} |X_k|^2]}{\lambda^2}.$$

Note that Kolmogorov's inequality is "sharper", because we have been able to take advantage of  $X_n$  being a martingale.

**Theorem 1.4.4** (Convergence Theorem). *1. Let  $\{X_n\}$  be a supermartingale. If  $\sup_n \mathbf{E}|X_n| < \infty$ , then  $\exists X_\infty \in \mathcal{L}^1(\Omega, \mathcal{F}, \mathbf{P})$  s.t.  $X_n \rightarrow X_\infty$  a.s. as  $n \rightarrow \infty$ . Moreover, if  $\{X_n\}$  is non-negative, then  $\mathbf{E}[X_\infty | \mathcal{F}_n] \leq X_n$  a.s.  $\forall n$ .*  
*2. Let  $\{X_n\}$  be a martingale and also be UI. Then  $\exists X_\infty \in \mathcal{L}^1(\Omega, \mathcal{F}, \mathbf{P})$  s.t.  $X_n \rightarrow X_\infty$  a.s. and in  $\mathcal{L}^1$ . Moreover,  $X_n = \mathbf{E}[X_\infty | \mathcal{F}_n]$ .*

*Remark.* Applying Fatou's lemma (note that we have  $X_n$  non-negative), we have

$$\mathbf{E}[X_\infty | \mathcal{F}_n] = \mathbf{E}\left[\lim_{m \rightarrow \infty} X_m | \mathcal{F}_n\right] \leq \lim_{m \rightarrow \infty} \mathbf{E}[X_m | \mathcal{F}_n] = \lim_{m \rightarrow \infty} X_n = X_n.$$

Note that  $\mathbf{E}[X_m | \mathcal{F}_n] \leq X_n$  is only true for  $m > n$ , but we're taking the limit as  $m \rightarrow \infty$  so this isn't a problem.

Now we consider the continuous time case. Let  $(\Omega, \mathcal{F}, \mathbf{P}; (\mathcal{F}_t)_{t \geq 0})$  be a filtered probability space. A map  $\tau : \Omega \rightarrow [0, \infty]$  is an  $(\mathcal{F}_t)$ -stopping time if

$$\{\omega : \tau(\omega) \leq t\} \in \mathcal{F}_t, \quad \forall t \geq 0.$$

Define

$$\mathcal{F}_\tau := \{A \in \mathcal{F} : \{\tau \leq t\} \cap A \in \mathcal{F}_t, \quad \forall t \geq 0\}.$$

**Definition 1.4.5.** A real-valued  $(\mathcal{F}_t)_{t \geq 0}$ -adapted process  $(X_t)_{t \geq 0}$  is called an  $(\mathcal{F}_t)$ -martingale (resp. supermartingale, submartingale) if  $\forall t \geq 0, X_t \in \mathcal{L}^1(\Omega, \mathcal{F}, \mathbf{P})$  with  $\mathbf{E}[X_t | \mathcal{F}_s] = X_s$  (resp.  $\leq, \geq$ ) a.s.,  $\forall s \leq t$ .

**Theorem 1.4.6.** Let  $p \geq 1$  and  $(X_t)_{t \geq 0}$  be a right continuous  $(\mathcal{F}_t)$ -martingale with  $\mathbf{E}|X_t|^p < \infty \quad \forall t \geq 0$ . Then for any  $T > 0$ :

1. (Doob's martingale inequality)  $P(\sup_{0 \leq t \leq T} |X_t| \geq \lambda) \leq \mathbf{E}|X_T|^p / \lambda^p \quad \forall \lambda > 0$ .
2. For  $p > 1$ ,  $\mathbf{E}[\sup_{0 \leq t \leq T} |X_t|^p] \leq (p/[p-1])^p \mathbf{E}|X_T|^p$ .

**Theorem 1.4.7** (Doob's optional sampling theorem / optional stopping theorem). Let  $(X_t)_{t \geq 0}$  be a right continuous martingale (resp. supermartingale, submartingale) on  $(\Omega, \mathcal{F}, \mathbf{P}; (\mathcal{F}_t)_{t \geq 0})$ .

1. If  $\sigma \leq \tau$  are two bounded stopping times, then

$$\mathbf{E}[X_\tau | \mathcal{F}_\sigma] = X_\sigma \text{ a.s. (resp. } \leq, \geq \text{).} \quad (1.4.1)$$

2. If  $(X_t)_{t \geq 0}$  are UI and  $\sigma \leq \tau \leq \infty$ , then (1.4.1) holds.

*Remark.* Suppose you're playing a fair game, and you have some strategy for when you stop playing ( $\tau$ ). The optional sampling theorem tells us that you cannot devise a winning strategy (or a losing strategy, for that matter).

Let  $(X_t)_{t \geq 0}$  be a process and  $\tau$  a stopping time. Then


$$X_{t \wedge \tau}(\omega) := \begin{cases} X_t(\omega) & \text{if } t < \tau(\omega) \\ X_{\tau(\omega)}(\omega) & \text{if } t \geq \tau(\omega) \end{cases}.$$

$X_{t \wedge \tau}(\omega)$  is called a stopped process, refer to figure 1.

**Definition 1.4.8.** An  $(\mathcal{F}_t)$ -adapted process  $(X_t)_{t \geq 0}$  on a filtered space  $(\Omega, \mathcal{F}, \mathbf{P}; (\mathcal{F}_t)_{t \geq 0})$  is called a local martingale if there is an increasing sequence  $\{\tau_n\}$  of finite stopping times  $\tau_n \uparrow \infty$  as  $n \rightarrow \infty$  a.s. and s.t.  $(X_{t \wedge \tau_n})_{t \geq 0}$  is a martingale for each  $n$ .



FIGURE 1: Stopped process.



## 2 Brownian Motion

### 2.1 Definition

**Definition 2.1.1.** A process  $B = (B_t)_{t \geq 0}$  on  $(\Omega, \mathcal{F}, \mathbf{P})$ <sup>9</sup> taking values in  $\mathbb{R}^d$  is called Brownian motion (BM) if

1. (Independent Increments)<sup>10</sup> For any  $0 \leq t_0 < t_1 < \dots < t_n$ , the r.v.s  $B_{t_0}, B_{t_1} - B_{t_0}, \dots, B_{t_n} - B_{t_{n-1}}$  are mutually independent.
2. (Normality) For any  $t > s \geq 0$ , the r.v.  $B_t - B_s \sim N(0, (t - s)I)$ , that is,  $B_t - B_s$  has pdf

$$p(t - s, x) = \frac{1}{(2\pi[t - s])^{d/2}} \exp\left(-\frac{|x|^2}{2(t - s)}\right), \quad x \in \mathbb{R}^d.$$

3. (Continuity)  $(B_t)_{t \geq 0}$  is continuous a.s..

If  $\mathbf{P}(B_0 = 0) = 1$ , then  $(B_t)_{t \geq 0}$  is called *standard Brownian motion*. We typically assume that this is what we are working with.

*Example 2.* Let  $(B_t)_{t \geq 0}$  be a BM in  $\mathbb{R}$ . Then

$$\mathbf{E}|B_t - B_s|^p = C_p |t - s|^{p/2} \quad \forall t > s \geq 0, \quad p \geq 0,$$

where

$$C_p := \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} |x|^p e^{-|x|^2/2} dx.$$

Indeed,

$$\begin{aligned} \mathbf{E}|B_t - B_s|^p &= \frac{1}{\sqrt{2\pi(t-s)}} \int_{-\infty}^{\infty} |x|^p \exp\left(-\frac{|x|^2}{2(t-s)}\right) dx \\ &= \frac{1}{\sqrt{2\pi(t-s)}} \int_{-\infty}^{\infty} (t-s)^p |y|^p e^{-|y|^2/2} \sqrt{t-s} dy \\ &= C_p (t-s)^{p/2}, \end{aligned}$$

<sup>9</sup>Note that is *not* a filtered probability space.

<sup>10</sup>This property is very important.

where we made the change of variables  $y = x/\sqrt{t-s}$ .

## Properties of BM

1.  $(B_t)_{t \geq 0}$  is nowhere differentiable, a.s..
2.  $\lim_{t \rightarrow \infty} B_t/t = 0$ , a.s..
3.  $\text{Cov}(B_t, B_s) = s \wedge t$ .
4.  $\tilde{B}_t := B_{t+s} - B_s$ , for any given  $s \geq 0$ , is also a standard Brownian motion. This property is also true for random  $s \geq 0$ .

*Remark.* For point 2, consider

$$\frac{B_n}{n} = \frac{(B_n - B_{n-1}) + \cdots + (B_1 - B_0)}{n} \rightarrow \mathbf{E}[N(0, 1)] = 0 \text{ a.s..}$$

The result holds due to the strong law of large numbers and the normality property of BM (in particular,  $B_i - B_{i-1} \sim N(0, 1)$ ).

This is not a proof! It just provides some intuition.

For point 3, suppose  $t \geq s$ . Then

$$\begin{aligned} \text{Cov}(B_t, B_s) &= \text{Cov}([B_t - B_s] + B_s, B_s) \\ &= \text{Cov}(B_t - B_s, B_s) + \text{Cov}(B_s, B_s) \\ &= 0 + \text{Var}(B_s) \\ &= s, \end{aligned}$$

Where we have used independent increments and normality of BM. The result as given above follows from symmetry.

*Remark* (Scaling). 1.  $M_t := \lambda B_{t/\lambda^2} \forall \lambda \neq 0$  is also a standard BM. In particular,  $-B_t$  is a standard BM.

2. Let  $d = 1$ . Define

$$W_t = \begin{cases} tB_{1/t} & t > 0 \\ 0 & t = 0 \end{cases}.$$

Then  $W_t$  is a BM.

It is easy to check that  $\text{Var}(W_t) = t$ .

For continuity, we need to check that  $\lim_{t \rightarrow 0} tB_{1/t} = 0$ . Then

$$\lim_{t \rightarrow 0} tB_{1/t} = \lim_{t \rightarrow 0} \frac{B_{1/t}}{1/t} = 0 \text{ a.s..}$$

The final equality follows from the second property of BM described above.

## 2.2 Markov property and finite-dimensional distributions

Recall:

$$p(t, x) = \frac{1}{(2\pi t)^{d/2}} e^{-|x|^2/2t}, \quad t > 0, \quad x \in \mathbb{R}^d.$$

We also have that

$$p(t+s, x-y) = p(t, x-z)p(s, z-y) \quad \forall t, s > 0, x, y, z \in \mathbb{R}^d.$$

Refer to exercise sheet 4, problem 2.

**Lemma 2.2.1.** *If  $t > s > 0$ , then the joint distribution  $(B_t, B_s)$  is given by*

$$P(B_s \in dx, B_t \in dy) = p(s, x)p(t-s, y-x) \, dx dy.$$

*Proof.* Since  $B_s$  and  $B_t - B_s$  are independent,  $(B_s, B_t - B_s)$  has joint density function  $p(s, x_1)p(t - s, x_2)$ . Hence, for all bounded Borel measurable functions  $f$ :

$$\begin{aligned} E[f(B_s, B_t)] &= \mathbf{E}[f(B_s, B_t - B_s + B_s)] \\ &= \int \int f(x_1, x_2 + x_1) p(s, x_1) p(t - s, x_2) dx_1 dx_2 \\ &= \int \int f(x, y) p(s, x) p(t - s, y - x) dx dy. \end{aligned}$$

In the final equality, we performed the change of variables  $x_1 = x$ ,  $x_1 + x_2 = y$ . In this case, the Jacobian is 1, so  $dx_1 dx_2 = dx dy$ .

This proves the claim.  $\square$

Let  $\mathcal{F}_t^0 := \sigma\{B_s, s \leq t\}$ ,  $t \geq 0$ . This is the *natural filtration generated by the BM*.

$B_t - B_s$  is independent of  $\mathcal{F}_s^0 \forall t > s \geq 0$ , because of independent increments.

**Theorem 2.2.2** (Markov Property). *Let  $t > s \geq 0$  and  $f$  be a bounded Borel measurable function. Then*

$$\mathbf{E}[f(B_t) | \mathcal{F}_s^0] = \mathbf{E}[f(B_t) | B_s].$$

*This is sometimes called the “history independent” or “memoryless” property.*

*Proof.* It suffices to prove for  $f \in C_b^\infty(\mathbb{R}^d)$ , the set of infinitely differentiable (“smooth”), bounded functions.<sup>11</sup>

For such  $f$ , we have the Taylor expansion

$$f(x + y) = \lim_{N \rightarrow \infty} \sum_{n=1}^N f_n(x) g_n(y) \text{ for some functions } f_n, g_n.$$

Then

$$\begin{aligned} \mathbf{E}[f(B_t) | \mathcal{F}_s^0] &= \mathbf{E}[f(B_t - B_s + B_s) | \mathcal{F}_s^0] \\ &= \mathbf{E} \left[ \lim_{N \rightarrow \infty} \sum_{n=1}^N f_n(B_s) g_n(B_t - B_s) | \mathcal{F}_s^0 \right] \\ &= \lim_{N \rightarrow \infty} \sum_{n=1}^N \mathbf{E}[f_n(B_s) g_n(B_t - B_s) | \mathcal{F}_s^0] \\ &= \lim_{N \rightarrow \infty} \sum_{n=1}^N \mathbf{E}[g_n(B_t - B_s) | \mathcal{F}_s^0] f_n(B_s) \\ &= \lim_{N \rightarrow \infty} \sum_{n=1}^N \mathbf{E}[g_n(B_t - B_s)] f_n(B_s) \end{aligned}$$

where we move the limit out of the expectation by the dominated convergence theorem, we move the  $f_n(B_s)$  term out because  $B_s$  is  $\mathcal{F}_s^0$  measurable and we remove the condition on  $\mathcal{F}_s^0$  because of independent increments.

If we repeat the above derivation, but with  $\mathcal{F}_s^0$  replaced by  $B_s$ , we get the same result.  $\square$

We now consider a more general version of lemma 2.2.1.

**Theorem 2.2.3** (Finite-Dimensional Distributions). *For any  $0 < t_1 < t_2 < \dots < t_n$ , the  $\mathbb{R}^{nd}$  valued r.v.  $(B_{t_1}, B_{t_2}, \dots, B_{t_n})$  has a joint density function*


$$p(t_1, x_1) p(t_2 - t_1, x_2 - x_1) \dots p(t_n - t_{n-1}, x_n - x_{n-1}).$$

<sup>11</sup>Any Borel measurable function can be approximated by functions in this space, cf. pp. 27 - 28 in Stochastic Differential Equations, Øksendal (2000).

## 2.3 The reflection principle and martingale property

Let  $B = (B_t)_{t \geq 0}$  be a standard BM on  $(\Omega, \mathcal{F}, \mathbf{P}; \mathcal{F}_t^0)$  in  $\mathbb{R}$ . Let  $b > 0$ ,  $b > a$  and  $\tau_b = \inf\{t \geq 0 : B_t = b\}$  (a stopping time).

FIGURE 2: The reflection principle.



Consider

$$\mathbf{P}\left(\max_{0 \leq s \leq t} B_s \geq b, B_t \leq a\right) = \mathbf{P}\left(\max_{0 \leq s \leq t} B_s \geq b, B_t \geq b + (b - a)\right) = \mathbf{P}(B_t \geq 2b - a) = \frac{1}{\sqrt{2\pi t}} \int_{2b-a}^{\infty} e^{-x^2/2t} dx.$$

The first equality is clear from figure 2, the second follows because the second event implies the first, and the third equality holds because  $B_t \sim N(0, t)$ .

Now we consider the case where  $\max_{0 \leq s \leq t} B_s \leq b$ ,

$$\mathbf{P}\left(\max_{0 \leq s \leq t} B_s \leq b, B_t \leq a\right) = \mathbf{P}(B_t \leq a) - \mathbf{P}(B_t \leq a - 2b) = \mathbf{P}(a - 2b \leq B_t \leq a) = \int_{a-2b}^a \frac{1}{\sqrt{2\pi t}} e^{-x^2/2t} dx.$$

Let  $M_t := \max_{0 \leq s \leq t} B_s$ . This is called the *running maximum* process.

We now consider the joint distribution of  $M_t$  and  $B_t$ , which are highly correlated with each other.

**Theorem 2.3.1.** *The joint distribution of  $(M_t, B_t)$  is*

$$P(M_t \in db, B_t \in da) = \frac{2(2b - a)}{\sqrt{2\pi t^3}} \exp\left(-\frac{(2b - a)^2}{2t}\right) dadb,$$

over the region  $\{b - a : b \geq a, b > 0\}$ .

Now,

$$\begin{aligned} P(M_t \geq c) &= \frac{2}{\sqrt{2\pi t^3}} \int_c^\infty \int_{-\infty}^b (2b - a) \exp\left(-\frac{(2b - a)^2}{2t}\right) dadb \\ &= \frac{2}{\sqrt{2\pi t^3}} \int_c^\infty \int_b^\infty x \exp\left(-\frac{x^2}{2t}\right) dx db \\ &= \frac{2}{\sqrt{2\pi}} \int_{c/\sqrt{t}}^\infty \exp\left(-\frac{x^2}{2}\right) dx \\ &= 2[1 - N(c/\sqrt{t})] \\ &= 2\mathbf{P}(B_t \geq c), \end{aligned}$$

where  $N(x) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^x e^{-y^2/2} dy$ .

In the above, we first used the change of variable  $x := 2b - a$ , followed by  $x := b/\sqrt{t}$ .

A martingale  $(M_t)_{t \geq 0}$  is called *square integrable* if  $\mathbf{E}[M_t^2] < \infty \forall t \geq 0$ .

We define the class  $\mathcal{M}_2^c := \{\text{all continuous, square integrable martingales}\}$ .

**Theorem 2.3.2.** 1.  $B_t \equiv (B_t^1 \ B_t^2 \ \dots \ B_t^d)$  is a continuous, square integrable martingale.

2.  $M_t := B_t^i B_t^j - \delta_{ij} t$  is a continuous martingale, where  $\delta_{ij}$  is Kronecker's delta.

*Proof.* 1. We only show that  $B_t$  is a martingale:

$$\begin{aligned} \mathbf{E}[B_t | \mathcal{F}_s^0] &= \mathbf{E}[B_t - B_s + B_s | \mathcal{F}_s^0] \\ &= \mathbf{E}[B_t - B_s | \mathcal{F}_s^0] + \mathbf{E}[B_s | \mathcal{F}_s^0] \\ &= \mathbf{E}[B_t - B_s] + B_s \\ &= 0 + B_s \\ &= B_s. \end{aligned}$$

2. We only need to show for BM in  $\mathbb{R}$  that

$$\mathbf{E}[B_t^2 - t | \mathcal{F}_s^0] = B_s^2 - s.$$

Rearranging, we see that

$$\begin{aligned} \mathbf{E}[B_t^2 - B_s^2 | \mathcal{F}_s^0] &= \mathbf{E}[(B_t - B_s)^2 | \mathcal{F}_s^0] + \mathbf{E}[2B_s(B_t - B_s) | \mathcal{F}_s^0] \\ &= \mathbf{E}[(B_t - B_s)^2 | \mathcal{F}_s^0] + 2B_s \underbrace{\mathbf{E}[B_t - B_s | \mathcal{F}_s^0]}_{=0} \\ &= \text{Var}(B_t - B_s) + \mathbf{E}[B_t - B_s]^2 \\ &= t - s + 0^2 \\ &= t - s. \end{aligned}$$

□

## 2.4 Quadratic variation process

**Lemma 2.4.1.** Let  $\mathcal{D} := \{0 = t_0 < t_1 < \dots < t_n = t\}$  be a finite partition of  $[0, t]$ , and let  $V_D := \sum_{l=1}^n |B_{t_l} - B_{t_{l-1}}|^2$ , the quadratic variation of  $B$  over  $D$ . Then  $\mathbf{E}[V_D] = t$ ,  $\text{Var}(V_D) = 2 \sum_{l=1}^n (t_l - t_{l-1})^2$ .

*Proof.* The expectation is

$$\begin{aligned} \mathbf{E}[V_D] &= \sum_{l=1}^n \mathbf{E}[|B_{t_l} - B_{t_{l-1}}|^2] \\ &= \sum_{l=1}^n |t_l - t_{l-1}| \\ &= t, \end{aligned}$$

and the variance is

$$\begin{aligned}
\text{Var}(V_D) &= \mathbf{E} \left[ (V_D - \mathbf{E}V_D)^2 \right] \\
&= \mathbf{E} \left[ \left( \sum_{l=1}^n |B_{t_l} - B_{t_{l-1}}|^2 - |t_l - t_{l-1}| \right)^2 \right] \\
&= \sum_{k,l=1}^n \mathbf{E} \left[ (|B_{t_k} - B_{t_{k-1}}|^2 - |t_k - t_{k-1}|) (|B_{t_l} - B_{t_{l-1}}|^2 - |t_l - t_{l-1}|) \right] \\
&= \sum_{k=1}^n \mathbf{E} \left[ (|B_{t_k} - B_{t_{k-1}}|^2 - |t_k - t_{k-1}|)^2 \right] + \\
&\quad \sum_{k \neq l} \mathbf{E} \left[ (|B_{t_k} - B_{t_{k-1}}|^2 - |t_k - t_{k-1}|) (|B_{t_l} - B_{t_{l-1}}|^2 - |t_l - t_{l-1}|) \right] \\
&= \sum_{k=1}^n \mathbf{E} \left[ (B_{t_k} - B_{t_{k-1}})^4 - 2(t_k - t_{k-1})(B_{t_k} - B_{t_{k-1}}) + (t_k - t_{k-1})^2 \right] \\
&= \sum_{k=1}^n \{ 3(t_k - t_{k-1})^2 - 2(t_k - t_{k-1})^2 + (t_k - t_{k-1})^2 \} \\
&= 2 \sum_{l=1}^n (t_k - t_{k-1})^2.
\end{aligned}$$

□

**Theorem 2.4.2.** We have  $\lim_{m(D) \rightarrow 0} \sum_{l=1}^n |B_{t_l} - B_{t_{l-1}}|^2 = t$  in  $\mathcal{L}^2$  and in probability, for any  $t \geq 0$ , where  $\mathcal{D}$  runs over all finite partitions of  $[0, t]$  and  $m(D) := \max_l |t_l - t_{l-1}|$ .

*Proof.* We only need to show convergence in  $\mathcal{L}^2$  as this implies convergence in  $\mathcal{L}^1$  which in turn implies convergence in probability.

$$\begin{aligned}
\mathbf{E} \left| \sum_{l=1}^n |B_{t_l} - B_{t_{l-1}}|^2 - t \right|^2 &= \mathbf{E} |V_D - \mathbf{E}V_D|^2 \\
&= \text{Var}(V_D) \\
&= 2 \sum_{l=1}^n (t_l - t_{l-1})^2 \\
&\leq 2m(D) \sum_{l=1}^n (t_l - t_{l-1}) \\
&= 2t \cdot m(D) \rightarrow 0 \text{ as } m(D) \rightarrow 0.
\end{aligned}$$

□

Let  $M = (M_t)_{t \geq 0} \in \mathcal{M}_c^2$ . As with BM, the limit

$$\langle M \rangle_t := \lim_{m(D) \rightarrow 0} \sum_{l=1}^n |M_{t_l} - M_{t_{l-1}}|^2$$

exists in  $\mathcal{L}^2$  and hence also in probability.

$\{\langle M \rangle_t\}_{t \geq 0}$  is called the *quadratic variation process* of  $(M_t)_{t \geq 0}$ , or simply the *bracket process*. It is an adapted, continuous, increasing process with  $\langle M \rangle_0 = 0$ .

**Doob-Meyer decomposition** If  $M = (M_t)_{t \geq 0}$  is a continuous martingale and  $A = (A_t)_{t \geq 0}$  is adapted, continuous and increasing, then  $X := M + A$  is a submartingale.

The reverse is the Doob-Meyer decomposition. In particular, the decomposition is unique.

**Theorem 2.4.3.** Let  $M \in \mathcal{M}_c^2$  be given. Then  $\langle M \rangle_t$  is the unique, continuous, adapted and increasing process initially zero s.t.  $M_t^2 - \langle M \rangle_t$  is a martingale.

*Proof.* By Jensen's inequality,  $\mathbf{E}[M_t^2 | \mathcal{F}_s] \geq \mathbf{E}[M_t | \mathcal{F}_s]^2 = M_s^2$ , so  $M_t^2$  is a submartingale. Hence, by the Doob-Meyer decomposition,

$$M_t^2 = N_t + A_t,$$

where  $N_t$  is a martingale and  $A_t$  is an increasing process. This decomposition is unique.

On the other hand,  $M_t^2 = (M_t^2 - \langle M \rangle_t) + \langle M \rangle_t$ , and since the decomposition above is unique and  $\langle M \rangle_t$  is increasing,  $\langle M \rangle_t$  is the unique process described in the theorem statement.  $\square$

**Theorem 2.4.4.** Let  $M, N \in \mathcal{M}_c^2$  and let

$$\langle M, N \rangle_t := \frac{1}{4}(\langle M + N \rangle_t - \langle M - N \rangle_t),$$

called the bracket process of  $M, N$ . Then  $\langle M, N \rangle_t$  is the unique, adapted, continuous process having finite variation almost surely and with initial zero, s.t.  $M_t N_t - \langle M, N \rangle_t$  is a martingale.

Moreover,  $\sum_{m(\mathcal{D}) \rightarrow 0} (M_{t_l} - M_{t_{l-1}})(N_{t_l} - N_{t_{l-1}}) = \langle M, N \rangle_t$  in probability where  $\mathcal{D} = \{0 = t_0 < t_1 < \dots < t_n = t\}$  and  $m(\mathcal{D}) = \max_l |t_l - t_{l-1}|$ .

### 3 Itô's calculus

$B = (B_t)_{t \geq 0}$  is a standard BM on  $(\Omega, \mathcal{F}, \mathbf{P})$  and  $\mathcal{F}_t^0 = \sigma\{B_s, 0 \leq s \leq t\}$  is the natural filtration.


Our objective is to define  $\int_0^t F_s dB_s \forall t \geq 0$  for some process  $(F_t)_{t \geq 0}$ .

#### 3.1 Stochastic integrals for BM

By convention, we shall call a process *adapted* if it is measurable and  $\mathcal{F}_t^0$ -adapted.

An adapted process  $F = (F_t)_{t \geq 0}$  is called a *simple process* if  $F_t(\omega) = f_0(\omega) \mathbb{1}_{\{0\}}(t) + \sum_{i=0}^{\infty} \mathbb{1}_{(t_i, t_{i+1}]}(t) f_i(\omega)$ , where  $0 = t_0 < t_1 < \dots$  so that for any finite  $T \geq 0$ , there are finitely many  $t_i \in [0, T]$ , each  $f_i$  is a  $\mathcal{F}_{t_i}^0$ -measurable r.v.,  $f_0$  is  $\mathcal{F}_0^0$ -measurable, and  $F$  is uniformly bounded.

FIGURE 3: A simple process.



Let  $\mathcal{L}_0$  denote the space of all simple processes.

Let  $F \in \mathcal{L}_0$ , then define

$$\int_0^t F_s dB_s \equiv I(F)_t := \sum_{i=0}^{\infty} f_i(B_{t_{i+1} \wedge t} - B_{t_i \wedge t}).$$

Note that this sum is finite, since we stop at  $t$ .

Clearly,  $I(F)$  is continuous in  $t$  and  $\mathcal{F}_t^0$ -adapted ( $f_i$  is  $\mathcal{F}_{t_i}$ -measurable, therefore  $\mathcal{F}_t$ -measurable, since  $\mathcal{F}_{t_i} \subseteq \mathcal{F}_t$  if  $t_i \leq t$ ).

**Lemma 3.1.1.**  $(I(F))_{t \geq 0}$  is a martingale, i.e.


$$\mathbf{E}[I(F)_t - I(F)_s | \mathcal{F}_s^0] = 0, \quad \forall t \geq s \geq 0.$$

*Proof.* Assume  $t_j < s \leq t_{j+1}$ ,  $t_k < t \leq t_{k+1}$  for some  $j, k$ .

Then  $j \leq k$ . Consider  $j+1 < k$ , in which case

$$I(F)_t - I(F)_s = \sum_{i=j+1}^{k-1} f_i(B_{t_{i+1}} - B_{t_i}) + f_j(B_{t_{j+1}} - B_s) + f_k(B_t - B_{t_k}).$$

FIGURE 4



Consider the expectation of the terms in the sum. By the tower property, we have

$$\mathbf{E}[f_i(B_{t_{i+1}} - B_{t_i}) | \mathcal{F}_s^0] = \mathbf{E}\{\mathbf{E}[f_i(B_{t_{i+1}} - B_{t_i}) | \mathcal{F}_{t_i}^0] | \mathcal{F}_s^0\} = \mathbf{E}\{f_i \mathbf{E}[(B_{t_{i+1}} - B_{t_i}) | \mathcal{F}_{t_i}^0] | \mathcal{F}_s^0\} = 0, \quad \forall j+1 \leq i \leq k-1.$$

Similarly, the terms involving  $f_j$  and  $f_k$  have conditional expectation zero also. Hence the expectation of the difference is zero and so the given process is a martingale.  $\square$

**Lemma 3.1.2.**  $I(F) \in \mathcal{M}_c^2$  and  $\langle I(F) \rangle_t = \int_0^t F_s^2 ds$ .

*Proof.* To prove  $I(F)_t^2 - \int_0^t F_s^2 ds$  is a martingale, use the same idea as in the proof of lemma 3.1.1.  $\square$

**Lemma 3.1.3** (Itô's isometry).  $F \mapsto I(F)$  is linear and  $\mathbf{E}[I(F)_t^2] = \mathbf{E}\left[\int_0^t F_s^2 ds\right]$ .

*Proof.*  $I(F)_t^2 - \int_0^t F_s^2 ds$  is a martingale, hence  $\mathbf{E}[I(F)_t^2 - \int_0^t F_s^2 ds] = 0$ . Then by linearity of expectation and rearranging, we have the result.  $\square$

### 3.2 Stochastic integrals for adapted processes

Now, for  $T > 0$  define

$$\mathcal{L}_{\mathcal{F}}^2(0, T; \mathbb{R}) = \left\{ F = (F_t)_{t \geq 0} : F \text{ is adapted and } \mathbf{E}\left[\int_0^T F_t^2 dt\right] < \infty \right\} \supseteq \mathcal{L}_0.$$

In fact,  $\mathcal{L}_0$  is dense in  $\mathcal{L}_{\mathcal{F}}^2$ .

$\forall F \in \mathcal{L}_{\mathcal{F}}^2(0, T; \mathbb{R})$ ,  $\exists F_n \in \mathcal{L}_0$  s.t.  $\mathbf{E}\left[\int_0^T |F_{n,t} - F_t|^2 dt\right] \rightarrow 0$  as  $n \rightarrow \infty$ .

$$\mathbf{E}|I(F_n)_T - I(F_m)_T|^2 = \mathbf{E}|I(F_n - F_m)_T|^2 = \mathbf{E} \int_0^T |F_{n,s} - F_{m,s}|^2 ds \rightarrow 0 \text{ as } n, m \rightarrow \infty.$$

$$I(F_n) \rightarrow I(F).$$

We define  $I(F) := \lim_{n \rightarrow \infty} I(F_n)$  as the Itô integral against  $B$ , denoted as  $I(F)_t = \int_0^t F_s dB_s$ .



**Theorem 3.2.1.** If  $F = (F_t)_{t \geq 0} \in \mathcal{L}_{\mathcal{F}}^2(0, T; \mathbb{R})$ , then both  $\int_0^t F_s dB_s$  and  $\left(\int_0^t F_s dB_s\right)^2 - \int_0^t F_s^2 ds$  are martingales.

Moreover,  $\mathbf{E}[\int_0^t F_s dB_s]^2 = \mathbf{E} \int_0^t F_s^2 ds \forall t \geq 0$ , and  $\mathbf{E} \left[ \left( \int_s^t F_u dB_u \right)^2 \middle| \mathcal{F}_s \right] = \mathbf{E}[\int_s^t F_u^2 du | \mathcal{F}_s] \forall t \geq s > 0$ .

Finally,  $\langle \int_0^t F_s dB_s, \int_0^t G_s dB_s \rangle_t = \int_0^t F_s G_s ds, \forall F, G \in \mathcal{L}_{\mathcal{F}}^2(0, T; \mathbb{R})$ .

We now define  $\int_0^t F_s dM_s$  where  $M \in \mathcal{M}_c^2$ .

Suppose  $F \in \mathcal{L}_0$ :  $F_t = f_0 \mathbb{1}_{\{0\}}(t) + \sum_{i=0}^{\infty} f_i \mathbb{1}_{(t_i, t_{i+1}]}(t)$ .

Define  $I^M(F) := \sum_{i=0}^{\infty} f_i (M_{t \wedge t_{i+1}} - M_{t \wedge t_i})$ .

Then

1.  $I^M(F) \in \mathcal{M}_c^2$ ;
2.  $\langle I^M(F) \rangle_t = \int_0^t F_s^2 d\langle M \rangle_s$ ;
3. (Itô's isometry)  $E(\int_0^t F_s dM_s)^2 = \mathbf{E}[\int_0^t F_s^2 d\langle M \rangle_s]$ .

$$\mathcal{L}_F^{2,M}(0, T; \mathbb{R}) := \left\{ F = (F_t)_{t \geq 0} : F \text{ is adapted, and } \mathbf{E} \int_0^t F_s^2 d\langle M \rangle_s < \infty \right\}.$$

One can define  $\int_0^t F_s dM_s$  for  $F \in \mathcal{L}_{\mathcal{F}}^{2,M}(0, T; \mathbb{R})$  via a limiting argument.

$\int_0^t F_s dM_s$  is a continuous martingale,  $\mathbf{E}(\int_0^t F_s dM_s)^2 = \mathbf{E}[\int_0^t F_s^2 d\langle M \rangle_s]$ .

$\langle \int_0^t F_s dM_s \rangle = \int_0^t F_s^2 d\langle M \rangle_s$ .

$\langle \int_0^t F_s dM_s, \int_0^t G_s dN_s \rangle = \int_0^t F_s G_s d\langle M, N \rangle_s$ .

### 3.3 Itô's integration for semimartingales

Let  $M = (M_t)_{t \geq 0}$  be a continuous local martingale with  $M_0 = 0$ . Then there exists  $\{\tau_n\}$  with  $\tau_n \uparrow \infty$  as  $n \rightarrow \infty$  a.s. and  $(M_n)_{t \geq 0} := (M_{t \wedge \tau_n})_{t \geq 0}$  is a continuous martingale.

Define  $\langle M \rangle_t := \langle M_n \rangle_t \forall t \leq \tau_n$ , which is an adapted, continuous and increasing process, with  $\langle M \rangle_0 = 0$  and  $M_t^2 - \langle M \rangle_t$  is a local martingale.

Now we define  $\int_0^t F_s dM_s$  for  $F$  which is left-continuous, adapted and  $\int_0^T F_s^2 d\langle M \rangle_s < \infty$  a.s.  $\forall T > 0$ .

$\int_0^t F_s dM_s := \int_0^t F_s d(M_n)_s$  if  $t \leq \tilde{\tau}_n$  where  $\tilde{\tau}_n \uparrow \infty$ .

Both  $\int_0^t F_s dM_s$  and  $(\int_0^t F_s dM_s)^2 - \int_0^t F_s^2 d\langle M \rangle_s$  are local martingales.

**Definition 3.3.1.** An adapted continuous process  $X = (X_t)_{t \geq 0}$  is called a semimartingale if  $X_t = M_t + V_t$  where  $(M_t)_{t \geq 0}$  is a continuous local martingale and  $(V_t)_{t \geq 0}$  is a process with finite variation in any finite interval.

$\int_0^t F_s dX_s := \int_0^t F_s dM_s + \int_0^t F_s dV_s$ .

$\int_0^t F_s dX_s = \lim_{m(\mathcal{D}) \rightarrow 0} \sum_l (F_{l-1}(X_{t_l} - X_{t_{l-1}}))$  in probability.

If  $X, Y$  are semimartingales, then

$$\langle X, Y \rangle_t := \lim_{m(\mathcal{D}) \rightarrow 0} \sum_{l=1}^n (X_{t_l} - X_{t_{l-1}})(Y_{t_l} - Y_{t_{l-1}}) \text{ in probability}$$

where  $\mathcal{D} = \{0 = t_0 < t_1 < \dots < t_n = t\}$ .

## Properties

1. (Bilinearity)  $\langle \alpha X + Y, \beta U + V \rangle = \alpha \beta \langle X, U \rangle + \alpha \langle X, V \rangle + \beta \langle Y, U \rangle + \langle Y, V \rangle$ .
2.  $\langle X, Y \rangle = 0$  if  $X$  is continuous with finite variation.
3. If  $X_t = M_t + V_t$ ,  $Y_t = N_t + U_t$ , where  $M, N$  are martingales,  $V, U$  have finite variation and so  $X, Y$  are semimartingales, then  $\langle X, Y \rangle_t = \langle M, N \rangle_t$ .

## 3.4 Itô's formula

Let  $(X_t)_{t \geq 0}$  be a continuous semimartingale. For a partition  $0 = t_0 < t_1 < \dots < t_n = t$ ,

$$\begin{aligned} X_{t_j}^2 - X_{t_{j-1}}^2 &= (X_{t_j} - X_{t_{j-1}})^2 + 2X_{t_{j-1}}(X_{t_j} - X_{t_{j-1}}) \\ \implies X_t^2 - X_0^2 &= 2 \sum_{j=1}^n X_{t_{j-1}}(X_{t_j} - X_{t_{j-1}}) + \sum_{j=1}^n (X_{t_j} - X_{t_{j-1}})^2. \end{aligned}$$

Letting  $m(\mathcal{D}) \rightarrow 0$ ,

$$X_t^2 - X_0^2 = 2 \int_0^t X_s dX_s + \langle X \rangle_t. \quad (3.4.1)$$

Compare this with the formula for integration by parts:

$$\int_0^t x_s dy_s = x_s y_s \Big|_0^t - \int_0^t y_s dx_s.$$

Setting  $y_s := x_s$  and rearranging, we get

$$x_t^2 - x_0^2 = 2 \int_0^t x_s dx_s.$$

Note that in the stochastic version there is a quadratic variation term, but this is absent in the deterministic version.

This is the simplest case of Itô's formula.

**Lemma 3.4.1** (Integration by parts). *Let  $X, Y$  be two continuous semimartingales. Then*

$$X_t Y_t - X_0 Y_0 = \int_0^t X_s dY_s + \int_0^t Y_s dX_s + \langle X, Y \rangle_t.$$

**Theorem 3.4.2** (Itô's formula). *Let  $(X_t)_{t \geq 0} = (X_t^1, \dots, X_t^d)_{t \geq 0}$  be a continuous, semimartingale in  $\mathbb{R}^d$  with  $X_t^i = M_t^i + A_t^i$  where  $M_t^1, \dots, M_t^d$  are continuous local martingales and  $A_t^1, \dots, A_t^d$  are continuous, adapted processes with finite variation.*

*Let  $f \in C^2(\mathbb{R}^d; \mathbb{R})$ <sup>12</sup>. Then*

$$f(X_t) - f(X_0) = \sum_{i=1}^d \int_0^t \frac{\partial f(X_s)}{\partial x_i} dX_s^i + \frac{1}{2} \sum_{i,j=1}^d \int_0^t \frac{\partial^2 f(X_s)}{\partial x_i \partial x_j} d\langle M^i, M^j \rangle_s.$$
<sup>13</sup>

*Proof.* We prove in the case  $d = 1$  and  $X_t = M_t \in \mathcal{M}_c^2$ .

We need to show

$$f(M_t) - f(M_0) = \int_0^t f'(M_s) dM_s + \frac{1}{2} \int_0^t f''(M_s) d\langle M \rangle_s. \quad (3.4.2)$$

This is true for  $f(x) = x^2$  (this is just (3.4.1)).

<sup>12</sup>  $f : \mathbb{R}^d \rightarrow \mathbb{R}$  and  $f \in C^2$ .

<sup>13</sup>  $\partial x_i$  is differentiating with respect to the  $i$ -th parameter of  $f$ .

Now suppose (3.4.2) is true for  $f(x) = x^n$ :

$$M_t^n - M_0^n = n \int_0^t M_s^{n-1} dM_s + \frac{1}{2}n(n-1) \int_0^t M_s^{n-2} d\langle M \rangle_s.$$

Applying integration by parts (lemma 3.4.1) to  $M^n$  and  $M$ , we have

$$\begin{aligned} M_t^{n+1} - M_0^{n+1} &= \int_0^t M_s^n dM_s + \int_0^t M_s dM_s^n + \langle M^n, M \rangle_t \\ &= \int_0^t M_s^n dM_s + \int_0^t M_s (nM_s^{n-1} + \frac{1}{2}n(n-1)M_s^{n-2} d\langle M \rangle_s) \\ &\quad + \left\langle M_0^n + n \int_0^t M_s^{n-1} dM_s + \frac{1}{2}n(n-1) \int_0^t M_s^{n-2} d\langle M \rangle_s, \int_0^t dM_s \right\rangle_t \\ &= \int_0^t M_s^n dM_s + n \int_0^t M_s^n dM_s + \frac{1}{2}n(n-1) \int_0^t M_s^{n-1} d\langle M \rangle_s \\ &\quad + n \int_0^t M_s^{n-1} d\langle M \rangle_s \\ &= (n+1) \int_0^t M_s^n dM_s + \frac{1}{2}n(n+1) \int_0^t M_s^{n-1} d\langle M \rangle_s \end{aligned}$$

□

**Corollary 3.4.3** (Itô's formula for diffusion processes). *Let  $(X_t)_{t \geq 0} = (X_t^1, \dots, X_t^d)_{t \geq 0}$  be given as*

$$X_t^i = X_0^i + \int_0^t b_s^i ds + \sum_{j=1}^n \int_0^t \sigma_s^{i,j} dB_s^j, \quad i = 1, 2, \dots, d$$

where  $(B_t)_{t \geq 0} = (B_t^1, \dots, B_t^n)_{t \geq 0}$  is a standard BM in  $\mathbb{R}^n$ , and  $f \in C^{1,2}([0, \infty] \times \mathbb{R}^d; \mathbb{R})$ . Then

$$f(t, X_t) - f(0, X_0) = \int_0^t \frac{\partial f(s, X_s)}{\partial s} ds + \sum_{i=1}^d \int_0^t \frac{\partial f(s, X_s)}{\partial x_i} dX_s^i + \frac{1}{2} \sum_{i,j=1}^d \int_0^t a_s^{i,j} \frac{\partial^2 f(s, X_s)}{\partial x_i \partial x_j} ds,$$

where  $a_s^{i,j} = \sum_{k=1}^n \sigma_s^{ik} \sigma_s^{kj}$ ,  $i, j = 1, 2, \dots, d$ .

*Example 3.* Let  $B = (B^1, \dots, B_n)$  be a standard BM. Then for any  $f \in C^2(\mathbb{R}^n; \mathbb{R})$ :

$$\begin{aligned} f(B_t) - f(B_0) &= \sum_{i=1}^n \int_0^t \frac{\partial f(B_s)}{\partial x_i} dB_s^i + \frac{1}{2} \sum_{i=1}^n \int_0^t \frac{\partial^2 f(B_s)}{\partial x_i^2} ds \\ &= \int_0^t \nabla f(B_s) \cdot dB_s + \frac{1}{2} \int_0^t \Delta f(B_s) ds. \end{aligned}$$

Hence,  $f(B_t) - f(B_0) - \frac{1}{2} \int_0^t \Delta f(B_s) ds$  is a local martingale.

*Example 4.* We will show that  $\mathbf{E}|\int_0^t \sigma_s dB_s|^2 = \mathbf{E} \int_0^t \sigma_s^2 ds$  for bounded  $\sigma$ .

Let  $M_t := \int_0^t \sigma_s dB_s$ .

$f(x) = x^2$ ,  $f'(x) = 2x$ ,  $f''(x) = 2$ ,  $\langle M \rangle_t = \int_0^t \sigma_s^2 ds$ .

$$\begin{aligned} f(M_t) - f(M_0) &= 2 \int_0^t M_s dM_s + \frac{1}{2} \cdot 2 \int_0^t \sigma_s^2 ds \\ \implies \left| \int_0^t \sigma_s dB_s \right|^2 &= 2 \int_0^t M_s dM_s + \int_0^t \sigma_s^2 ds \\ \implies \mathbf{E} \left| \int_0^t \sigma_s dB_s \right|^2 &= \mathbf{E} \int_0^t \sigma_s^2 ds. \end{aligned}$$

### 3.5 Stochastic exponentials and Girsanov theorem

Consider a deterministic ODE,

$$\begin{cases} dz_t &= z_t dx_t \\ z_0 &= 1 \end{cases} \implies z_t = e^{x_t}.$$

What about the stochastic case?

$$\begin{cases} dZ_t &= Z_t dX_t \\ Z_0 &= 1 \end{cases}, \quad (3.5.1)$$

where  $X_t$  is a continuous semimartingale.

The real meaning of (3.5.1) is

$$Z_t = 1 + \int_0^t Z_s dX_s. \quad (3.5.2)$$

Try  $Z_t = \exp(X_t + V_t)$ , where  $(V_t)_{t \geq 0}$ , a finite variation process, is to be determined.

Applying Itô's formula,

$$\begin{aligned} Z_t &= Z_0 + \int_0^t \exp(X_s + V_s) d(X_s + V_s) + \frac{1}{2} \int_0^t \exp(X_s + V_s) d\langle X \rangle_s^{14} \\ &= 1 + \int_0^t Z_s d(X_s + V_s) + \frac{1}{2} \int_0^t Z_s d\langle X \rangle_s. \end{aligned} \quad (3.5.3)$$

This suggests we should set  $V_t := -\frac{1}{2}\langle X \rangle_t$ . Compare (3.5.2) with (3.5.3).

**Lemma 3.5.1.** *Let  $X_t$  be a semimartingale with  $X_0 = 0$ . Then*

$$\mathcal{E}(X) := \exp(X_t - \frac{1}{2}\langle X \rangle_t), \quad (3.5.4)$$

*called the stochastic exponential of  $X$ , is the solution to (3.5.2).*

**Lemma 3.5.2.** *Any continuous, non-negative local martingale is a supermartingale.*

*Proof.* Refer to exercise sheet 7, question 1. □

**Corollary 3.5.3.** *Let  $M = (M_t)_{t \geq 0}$  be a continuous local martingale with  $M_0 = 0$ . Then  $\mathcal{E}(M)$  is a supermartingale. In particular,*

$$\mathbf{E}[\exp(M_t - \frac{1}{2}\langle M \rangle_t)] \leq 1 \quad \forall t \geq 0.$$

*Moreover,  $\mathcal{E}(M)$  is a martingale up to time  $T > 0$  if and only if*

$$\mathbf{E}[\exp(M_T - \frac{1}{2}\langle M \rangle_T)] = 1. \quad (3.5.5)$$

**Theorem 3.5.4** (Novikov's condition). *Let  $M = (M_t)_{t \geq 0}$  be a continuous local martingale with  $M_0 = 0$ . If  $\mathbf{E} \exp(\frac{1}{2}\langle M \rangle_T) < \infty$ , then  $\mathcal{E}(M)$  is a martingale up to time  $T$ .*

*Proof.* Refer to Prof. Qian's lecture notes, theorem 4.6.9. □

*Example 5.* Let  $B = (B_t)_{t \geq 0}$  be a standard BM and  $F = (F_t)_{t \geq 0} \in \mathcal{L}_{\mathcal{F}}^2(0, T; \mathbb{R})$ . Assume

$$\mathbf{E} \exp\left(\frac{1}{2} \int_0^T F_t^2 dt\right) < \infty,$$

then  $X_t = \mathcal{E}\left[\int_0^t F_s dB_s\right] \equiv \exp\left(\int_0^t F_s dB_s - \frac{1}{2} \int_0^t F_s^2 ds\right)$  is a martingale up to  $T$ .

**Theorem 3.5.5** (Lévy's characterisation). *A process  $M = (M_t)_{t \geq 0}$  with  $M_0 = 0$  is a standard BM if and only if it is a continuous local martingale with  $\langle M \rangle_t = t$ .*

<sup>14</sup>  $d\langle X + V \rangle_s = d\langle X \rangle_s$  because  $V$  has finite variation.

*Proof.* We prove the backwards direction.

$\forall T > 0, \lambda \in \mathbb{R}$ :

$$\mathbf{E} \exp(\tfrac{1}{2} \langle \lambda M \rangle_T) = \mathbf{E} \exp(\tfrac{1}{2} \lambda^2 \langle M \rangle_T) = \mathbf{E} \exp(\tfrac{1}{2} \lambda^2 T) < \infty.$$

Hence,  $\mathcal{E}(\lambda M) = \exp(\lambda M_t - \tfrac{1}{2} \lambda^2 t)$  is a martingale up to  $T$ , by Novikov's condition.

$$\begin{aligned} \mathbf{E}[e^{\lambda M_t - \frac{1}{2} \lambda^2 t} | \mathcal{F}_s] &= e^{\lambda M_s - \frac{1}{2} \lambda^2 s} \quad \forall 0 \leq s \leq t \leq T \\ \iff \mathbf{E}[e^{\lambda(M_t - M_s)} | \mathcal{F}_s] &= e^{\frac{1}{2} \lambda^2 (t-s)} \\ \iff \mathbf{E}[e^{\lambda(M_t - M_s)}] &= e^{\frac{1}{2} \lambda^2 (t-s)}. \end{aligned}$$

So,  $(M_t)_{t \geq 0}$  has independent increments.

Note that the last line is just the expression for the MGF of a  $N(0, t-s)$  r.v., so  $M_t - M_s \sim N(0, t-s)$ .  $\square$

Suppose we have a filtered probability space  $(\Omega, \mathcal{F}, \mathbf{P}; (\mathcal{F}_t)_{t \geq 0})$  and  $T > 0$ .

Let  $Q$  be another probability measure s.t.

$$\int_{\Omega} X(\omega) d\mathbf{Q}(\omega) = \int_{\Omega} X(\omega) \xi(\omega) d\mathbf{P}(\omega) \quad \forall \text{ bounded } \mathcal{F}_T\text{-measurable r.v.s } X \quad (3.5.6)$$

where  $0 < \xi \in \mathcal{L}^1(\Omega, \mathcal{F}_T, \mathbf{P})$ .

We write this as

$$\left. \frac{d\mathbf{Q}}{d\mathbf{P}} \right|_{\mathcal{F}_t} = \xi.$$

We can rewrite (3.5.6) as

$$\mathbf{E}^{\mathbf{Q}}[X] = \mathbf{E}^{\mathbf{P}}[X\xi] \implies \mathbf{E}^{\mathbf{P}}[\xi] = 1,$$

where the implication follows by setting  $X \equiv 1$ .

Now, let  $X$  be  $\mathcal{F}_t$ -measurable,  $\forall t \leq T$ . Then

$$\mathbf{E}^{\mathbf{Q}}[X] = \mathbf{E}^{\mathbf{P}}[X\xi] = \mathbf{E}^{\mathbf{P}}[\mathbf{E}^{\mathbf{P}}[X\xi | \mathcal{F}_t]] = \mathbf{E}^{\mathbf{P}}[X \mathbf{E}^{\mathbf{P}}[\xi | \mathcal{F}_t]].$$

So,  $\forall t \leq T, \left. \frac{d\mathbf{Q}}{d\mathbf{P}} \right|_{\mathcal{F}_t} = \mathbf{E}^{\mathbf{P}}[\xi | \mathcal{F}_t] > 0$ <sup>15</sup>.

Conversely, let  $Z = (Z_t)_{t \geq 0}$  be a continuous, positive martingale up to  $T$  with  $Z_0 = 1$ , on  $(\Omega, \mathcal{F}, \mathbf{P}; (\mathcal{F}_t)_{t \geq 0})$ .

Define a measure on  $(\Omega, \mathcal{F}_T)$  by

$$\left. \frac{d\mathbf{Q}}{d\mathbf{P}} \right|_{\mathcal{F}_t} = Z_t.$$

For any  $\mathcal{F}_t$ -measurable r.v.  $X$ ,

$$\mathbf{E}^{\mathbf{Q}}[X] = \mathbf{E}^{\mathbf{P}}[\mathbf{E}^{\mathbf{P}}[Z_T X | \mathcal{F}_t]] = \mathbf{E}^{\mathbf{P}}[X \mathbf{E}^{\mathbf{P}}[Z_T | \mathcal{F}_t]] = \mathbf{E}^{\mathbf{P}}[X Z_t].$$

**Theorem 3.5.6** (Girsanov's theorem). *Let  $(M_t)_{t \geq 0}$  be a continuous local martingale and  $(Z_t)_{t \geq 0}$  a continuous, positive martingale up to  $T$  on  $(\Omega, \mathcal{F}, \mathbf{P}; (\mathcal{F}_t)_{t \geq 0})$  with  $Z_0 = 1$ . Then*

$$X_t := M_t - \int_0^t \frac{1}{Z_s} d\langle M, Z \rangle_s$$

*is a continuous local martingale on  $(\Omega, \mathcal{F}, \mathbf{Q}; (\mathcal{F}_t)_{t \geq 0})$  up to  $T$ .*

<sup>15</sup>This expectation is a process in  $E$ , in fact, is a martingale up to  $T$ . Put  $Y_t := \mathbf{E}[\xi | \mathcal{F}_t]$ . Then

$$\mathbf{E}[Y_t | \mathcal{F}_s] = \mathbf{E}[\mathbf{E}[\xi | \mathcal{F}_t] | \mathcal{F}_s] = \mathbf{E}[\xi | \mathcal{F}_s] = Y_s, \quad \forall s \leq t \leq T.$$

*Proof.* We prove for the case when  $Z$  and  $X$  are bounded and  $M \in \mathcal{M}_c^2$ . Using integration by parts,

$$\begin{aligned} Z_t X_t &= Z_0 X_0 + \int_0^t Z_s dX_s + \int_0^t X_s dZ_s + \langle Z, X \rangle_t \\ &= Z_0 X_0 + \int_0^t Z_s dM_s - \underbrace{\int_0^t Z_s \cdot \frac{1}{Z_s} d\langle M, Z \rangle_s}_{=\langle M, Z \rangle_t - \langle M, Z \rangle_0} + \int_0^t X_s dZ_s + \underbrace{\langle Z, X \rangle_t}_{=\langle M, Z \rangle_t} \\ &= Z_0 X_0 + \int_0^t Z_s dM_s + \int_0^t X_s dZ_s + \langle M, Z \rangle_0, \end{aligned}$$

which is a  $\mathbf{P}$ -martingale.

Now,  $\forall A \in \mathcal{F}_s, s \leq t$ ,

$$\begin{aligned} \mathbf{E}^{\mathbf{Q}}[\mathbb{1}_A(X_t - X_s)] &= \mathbf{E}^{\mathbf{Q}}[\mathbb{1}_A X_t] - \mathbf{E}^{\mathbf{Q}}[\mathbb{1}_A X_s] \\ &= \mathbf{E}^{\mathbf{P}}[\mathbb{1}_A X_t Z_t] - \mathbf{E}^{\mathbf{P}}[\mathbb{1}_A X_s Z_s] \\ &= \mathbf{E}^{\mathbf{P}}[\mathbb{1}_A(X_t Z_t - X_s Z_s)] \\ &= 0. \end{aligned}$$

□

Since  $Z_t > 0$ , applying Itô's lemma to  $\log Z_t$

$$\log Z_t - \log Z_0 = \int_0^t \frac{1}{Z_s} dZ_s - \frac{1}{2} \int_0^t \frac{1}{Z_s^2} d\langle Z \rangle_s.$$

Hence,  $Z_t = \mathcal{E}(N)_t$  where  $N_t := \int_0^t \frac{1}{Z_s} dZ_s$  is a continuous local martingale.

$$Z_t = 1 + \int_0^t Z_s dN_s.$$

$$\langle M, Z \rangle_t = \left\langle \int_0^t dM_s, \int_0^t Z_s dN_s \right\rangle_t = \int_0^t Z_s d\langle M, N \rangle_s.$$

**Corollary 3.5.7.** *Let  $N_t$  be a continuous martingale on  $(\Omega, \mathcal{F}, \mathbf{P}; (\mathcal{F}_t)_{t \geq 0})$ ,  $N_0 = 0$ , s.t.  $\mathcal{E}(N)_t$  is a continuous martingale up to  $T$ .*

*Define a probability measure  $\mathbf{Q}$  on  $(\Omega, \mathcal{F}_T)$  by*

$$\left. \frac{d\mathbf{Q}}{d\mathbf{P}} \right|_{\mathcal{F}_t} = \mathcal{E}(N)_t, \forall t \leq T.$$

*If  $(M_t)_{t \geq 0}$  is a continuous local martingale under  $P$ , then  $X_t := M_t - \langle N, M \rangle_t$  is a continuous local martingale under  $\mathbf{Q}$  up to  $T$ .*

*Example 6.* Consider the stock price

$$dS_t = S_t(r_t dt + dW_t) = S_t d\tilde{W}_t.$$

Let  $N_t := -\int_0^t r_s dW_s$ .

$\mathbf{E}[\exp(\frac{1}{2}\langle N \rangle_t)] = \mathbf{E}[\exp(\frac{1}{2}\int_0^t r_s^2 ds)] < \infty$ , then  $\mathcal{E}(N)_t = \exp(-\int_0^t r_s dW_s + \frac{1}{2}\int_0^t r_s^2 ds)$  is a martingale up to  $T$ , by Novikov's condition.

Under  $\frac{d\mathbf{Q}}{d\mathbf{P}}|_{\mathcal{F}_t} = \mathcal{E}(N)_t$ ,  $\tilde{W}_t := W_t - \langle N, W \rangle_t = W_t + \int_0^t r_s ds$ <sup>16</sup> is a continuous local martingale under  $\mathbf{Q}$ .

However,  $\langle \tilde{W} \rangle_t = \langle W \rangle_t = t \implies \tilde{W}$  is a BM under  $\mathbf{Q}$ .

In this case,  $dS_t = S_t d\tilde{W}_t \implies S_t$  is a martingale under  $\mathbf{Q}$ .

So, e.g.  $\mathbf{E}^{\mathbf{Q}}[S_T] = \mathbf{E}^{\mathbf{Q}}[S_0]$ , where  $S_T$  is the future price and  $S_0$  is the current price.  $\mathbf{Q}$  is called the “risk-neutral measure” and  $\mathbf{P}$  is called the “physical measure”.

---

<sup>16</sup>  $\langle N, W \rangle_t = -\left\langle \int_0^t r_s dW_s, \int_0^t dW_s \right\rangle_t = -\int_0^t r_s ds$ .

$M_t := \int_0^t F_s dB_s$  is an  $\mathcal{F}_t^0$ -martingale if  $F \in \mathcal{L}_{\mathcal{F}}^2(0, T; \mathbb{R})$ .

**Theorem 3.5.8** (Martingale representation theorem). *Let  $M = (M_t)_{t \geq 0}$  be a square-integrable martingale on  $(\Omega, \mathcal{F}, \mathbf{P}; (\mathcal{F}_t^0)_{t \geq 0})$ . Then there exists  $F = (F_t)_{t \geq 0} \in \mathcal{L}_{\mathcal{F}}^2(0, T; \mathbb{R})$  s.t.  $M_t = M_0 + \int_0^t F_s dB_s$ .*

## 4 Stochastic Differential Equations

### 4.1 Introduction

Consider

$$dX_t^j = f_0^j(t, X_t)dt + \sum_{i=1}^n f_i^j(t, X_t)dB_t^i, \quad j = 1, 2, \dots, N \quad (4.1.1)$$

where  $B_t = (B_t^1, \dots, B_t^n)_{t \geq 0}$  is a standard BM in  $\mathbb{R}^n$  on  $(\Omega, \mathcal{F}, \mathbf{P}; (\mathcal{F}_t)_{t \geq 0})$ , and

$$f_i^j : [0, \infty) \times \mathbb{R}^N \rightarrow \mathbb{R}, \quad i = 1, 2, \dots, n, \quad j = 1, 2, \dots, N$$

are Borel measurable functions.

Note that (4.1.1) should be interpreted as

$$X_t^j = X_0^j + \int_0^t f_0^j(s, X_s)ds + \sum_{i=1}^n \int_0^t f_i^j(s, X_s)dB_s^i. \quad (4.1.2)$$

**Definition 4.1.1.** 1. An adapted, continuous,  $\mathbb{R}^N$ -valued process  $X = (X_t)_{t \geq 0} = (X_t^1, \dots, X_t^N)_{t \geq 0}$  on  $(\Omega, \mathcal{F}, \mathbf{P}; (\mathcal{F}_t)_{t \geq 0})$  is a weak solution of (4.1.1) if there is a standard BM  $W = (W_t)_{t \geq 0}$  in  $\mathbb{R}^n$ , adapted to  $(\mathcal{F}_t)_{t \geq 0}$ , s.t. (4.1.2) holds with  $B$  replaced by  $W$ .

2. Given a standard BM  $B = (B_t)_{t \geq 0}$  in  $\mathbb{R}^n$  on  $(\Omega, \mathcal{F}, \mathbf{P})$  with filtration  $(\mathcal{F}_t^0)_{t \geq 0}$ , an adapted continuous process  $X = (X_t)_{t \geq 0}$  on  $(\Omega, \mathcal{F}, \mathbf{P}; (\mathcal{F}_t^0)_{t \geq 0})$  is a strong solution of (4.1.1) if (4.1.2) holds.

**Definition 4.1.2.** 1. We say that (4.1.1) has pathwise uniqueness if whenever  $(X, B)$  and  $(\tilde{X}, B)$  are two weak solutions defined on the same space  $(\Omega, \mathcal{F}, \mathbf{P}; (\mathcal{F}_t)_{t \geq 0})$ , and  $X_0 = \tilde{X}_0$ , then  $X = \tilde{X}$  a.s..

2. We say that (4.1.1) has uniqueness in law if whenever  $(X, B)$  and  $(\tilde{X}, \tilde{B})$  are two weak solutions (possibly defined on different probability spaces) and  $X_0$  and  $\tilde{X}_0$  have the same distribution, then  $X$  and  $\tilde{X}$  have all the same finite-dimensional distributions.

**Theorem 4.1.3** (Yamada-Watanabe). *Pathwise uniqueness implies uniqueness in law.*

*Example 7* (Hiroshi Tanaka). Consider a one-dimensional SDE

$$X_t = \int_0^t \text{sgn}(X_s)dB_s, \quad 0 \leq t < \infty$$

where  $\text{sgn}(x) = \begin{cases} 1 & \text{if } x \geq 0 \\ -1 & \text{if } x < 0 \end{cases}$

Then

1. Uniqueness in law holds:  $X_t$  is a continuous martingale and  $\langle X \rangle_t = \int_0^t 1ds = t \implies X_t$  is BM.
2. If  $(X, B)$  is a weak solution:

$$-X_t = \int_0^t -\text{sgn}(X_s)dB_s = \int_0^t \text{sgn}(-X_s)dB_s \implies (-X, B) \text{ is also a weak solution.}$$

Hence pathwise uniqueness does not hold.

3. Let  $W_t$  be any standard BM on  $(\Omega, \mathcal{F}, \mathbf{P})$  and then let  $B_t := \int_0^t \text{sgn}(W_s)dW_s$ .  $(B_t)_{t \geq 0}$  is BM and  $\int_0^t \text{sgn}(W_s)dB_s = \int_0^t \underbrace{\text{sgn}(W_s)^2}_{=1}dW_s = W_t$ .

Hence  $(W_t, B_t)$  is a weak solution.

4. There is no strong solution (see Roger & Williams 1990, p. 151).

<sup>17</sup>The stochastic differential equation (SDE) is defined by the  $f$ s. The underlying probability space and the BM are *not* part of the definition of the SDE.

## 4.2 Several examples

### 4.2.1 Linear SDEs

Consider

$$dX_t = \beta X_t dt + \sigma dB_t, \quad (4.2.1)$$

where  $X_t$  and  $B_t$  are  $N \times 1$  column vectors,  $\beta$  is an  $N \times N$  matrix and  $\sigma$  is an  $N \times n$  matrix.

Using integration by parts,

$$\begin{aligned} e^{-\beta t} X_t - e^{-\beta 0} X_0 &= \int_0^t e^{-\beta s} dX_s + \int_0^t d(e^{-\beta s}) X_s^{18} \\ &= \int_0^t e^{-\beta s} dX_s - \int_0^t e^{-\beta s} \beta X_s ds \\ &= \int_0^t e^{-\beta s} (dX_s - \beta X_s ds) \\ &= \int_0^t e^{-\beta s} \sigma dB_s \\ \Rightarrow X_t &= e^{\beta t} X_0 + \int_0^t e^{\beta(t-s)} \sigma dB_s. \end{aligned}$$

### Ornstein-Uhlenbeck process

$$dX_t = dB_t - (AX_t)dt.$$

### 4.2.2 Geometric BM

The Black-Scholes model is

$$dS_t = S_t(\mu_t dt + \sigma_t dB_t).$$

Regarding the parenthesised term as “ $X_t$ ”, we see that we have a stochastic exponential:

$$\begin{aligned} S_t &= \mathcal{E}(X)_t \\ &= \mathcal{E}\left(\int_0^t \mu_s ds + \int_0^t \sigma_s dB_s\right) \\ &= S_0 \exp\left(\int_0^t \mu_s ds + \int_0^t \sigma_s dB_s - \frac{1}{2} \int_0^t \sigma_s^2 ds\right) \\ &= S_0 \exp\left(\int_0^t [\mu_s - \frac{1}{2} \sigma_s^2] ds + \int_0^t \sigma_s dB_s\right). \end{aligned}$$

If  $\mu_t \equiv \mu$  and  $\sigma_t \equiv \sigma$  then

$$S_t = S_0 \exp(\sigma B_t + [\mu - \frac{1}{2} \sigma^2]t),$$

hence  $S_t$  follows a lognormal distribution, because  $B_t \sim N(0, t)$  and the remaining terms are constants.

### 4.2.3 Cameron-Martin's formula

Consider the SDE

$$dX_t = dB_t + b(t, X_t)dt \quad (4.2.2)$$

where  $b$  is bounded and measurable on  $[0, T] \times \mathbb{R}$ .

Let  $(W_t)_{t \geq 0}$  be a standard BM on  $(\Omega, \mathcal{F}, \mathbf{P}; (\mathcal{F}_t)_{t \geq 0})$ .

Define

$$\left. \frac{d\mathbf{Q}}{d\mathbf{P}} \right|_{\mathcal{F}_t} = \mathcal{E}(N)_t, \quad t \geq 0,$$

---

<sup>18</sup>Recall that the matrix exponential is defined as  $e^{\beta t} := \sum_{k=0}^{\infty} \frac{t^k}{k!} \beta^k$  and so we have  $\frac{d}{dt} e^{\beta t} = \beta e^{\beta t} = e^{\beta t} \beta$ .



where  $N_t := \int_0^t b(s, W_s) dW_s$  is a  $\mathbf{P}$ -martingale with  $\langle N \rangle_t = \int_0^t b^2(s, X_s) ds$ . By Novikov's condition,

$$\mathcal{E}(N)_t \equiv \exp \left( \int_0^t b(s, W_s) dW_s - \frac{1}{2} \int_0^t b^2(s, W_s) ds \right)$$

is a  $\mathbf{P}$ -martingale.

By corollary 3.5.7,

$$B_t := W_t - \langle W, N \rangle_t = W_t - \int_0^t b(s, W_s) ds$$

is a continuous local  $\mathbf{Q}$ -martingale. Moreover,  $\langle B \rangle_t = \langle W \rangle_t = t$ .

Hence,  $(B_t)_{t \geq 0}$  is a standard BM under  $\mathbf{Q}$ .

However,  $dW_t = dB_t + b(t, W_t)dt \implies (W, B)$  is a weak solution of (4.2.2) on  $(\Omega, \mathcal{F}, \mathbf{Q})$ .

**Theorem 4.2.1** (Cameron-Martin's formula). *Let  $b(t, x) = (b^1(t, x), \dots, b^n(t, x))$  be bounded measurable functions on  $[0, T] \times \mathbb{R}^n$  and  $W_t = (W_t^1, \dots, W_t^n)$  be a standard BM on  $(\Omega, \mathcal{F}, \mathbf{P}; (\mathcal{F}_t)_{t \geq 0})$ . Define probability measure  $\mathbf{Q}$  by*

$$\left. \frac{d\mathbf{Q}}{d\mathbf{P}} \right|_{\mathcal{F}_t} := \exp \left( \sum_{k=1}^n \int_0^t b^k(s, W_s) dW_s^k - \frac{1}{2} \sum_{k=1}^n \int_0^t |b^k(s, W_s)|^2 ds \right), \quad 0 \leq t \leq T.$$

Then  $(W_t)_{t \geq 0}$  under  $\mathbf{Q}$  is a solution to

$$dX_t^j = dB_t^j + b^j(t, X_t)dt$$

for some BM  $B_t = (B_t^1, \dots, B_t^n)$  under  $\mathbf{Q}$ .

### 4.3 Existence and uniqueness

Consider the SDE

$$dX_t^j = f_0^j(t, X_t)dt + \sum_{i=1}^n f_i^j(t, X_t)dB_t^i, \quad j = 1, 2, \dots, N \quad (4.3.1)$$

**Theorem 4.3.1.** *Suppose that  $f_i^j$   $i = 1, 2, \dots, n, j = 1, 2, \dots, N$  satisfy the Lipschitz condition*

$$|f_i^j(t, x) - f_i^j(t, y)| \leq c|x - y| \quad \forall x, y \in \mathbb{R}^N, \quad t \geq 0$$

for some constant  $c > 0$  and the linear growth condition

$$|f_i^j(t, x)| \leq c(1 + |x|) \quad \forall x \in \mathbb{R}^N, \quad t \geq 0.$$

Then for any  $\eta \in \mathcal{L}^2(\Omega, \mathcal{F}, \mathbf{P})$  and a standard BM  $B_t$  in  $\mathbb{R}^n$ , there is a strong solution  $(X_t)_{t \geq 0}$  of (4.3.1) with  $X_0 = \eta$ . Moreover, the pathwise uniqueness holds, which implies uniqueness in law by theorem 4.1.3 (Yamada-Watanabe).

Consider example 7 (Hiroshi Tanaka). There is no strong solution and no pathwise uniqueness. Note that sgn doesn't satisfy the Lipschitz condition.

*Example 8* (Itô and Watanabe, 1978). The SDE

$$dX_t = 3X_t^{2/3}dB_t + 3X_t^{1/3}dt$$

has infinitely many strong solutions of the form

$$X_t^{(\theta)} := \begin{cases} 0 & \text{if } 0 \leq t < \tau_\theta \\ B_t^3 & \text{if } \tau_\theta \leq t < \infty \end{cases}$$

where  $0 \leq \theta < \infty$  and  $\tau_\theta := \inf\{t \geq \theta : B_t = 0\}$ .

If  $t < \tau_\theta$ , then  $X_t^{(\theta)} = 0$  and so clearly it satisfies the SDE.

When  $t \geq \tau_\theta$ , we see that

$$dB_t^3 = 3B_t^2dB_t + \frac{1}{2} \cdot 3 \cdot 2B_tdt = 3(B_t^3)^{2/3} + 3(B_t^3)^{1/3}dt.$$

by applying Itô's formula.