



# Involvement of the ubiquitin-proteasome system in the expression of extracellular matrix genes in retinal pigment epithelial cells

J. Emanuel Ramos de Carvalho<sup>a,\*</sup>, Milan T. Verwoert<sup>a</sup>, Ilse M.C. Vogels<sup>a</sup>, Eric A. Reits<sup>b</sup>, Cornelis J.F. Van Noorden<sup>a,b</sup>, Ingeborg Klaassen<sup>a</sup>, Reinier O. Schlingemann<sup>a</sup>

<sup>a</sup> Ocular Angiogenesis Group, Departments of Ophthalmology and Medical Biology, Academic Medical Center, University of Amsterdam, Amsterdam, The Netherlands

<sup>b</sup> Department of Medical Biology, Academic Medical Center, University of Amsterdam, Amsterdam, The Netherlands

## ARTICLE INFO

### Keywords:

RPE  
Fibrosis  
Retina  
TGFβ  
CTGF  
PPARγ  
Proteasome  
Epoxomicin

## ABSTRACT

Emerging evidence suggests that dysfunction of the ubiquitin-proteasome system is involved in the pathogenesis of numerous senile degenerative diseases including retinal disorders. The aim of this study was to assess whether there is a link between proteasome regulation and retinal pigment epithelium (RPE)-mediated expression of extracellular matrix genes. For this purpose, human retinal pigment epithelial cells (ARPE-19) were treated with different concentrations of transforming growth factor-β (TGFβ), connective tissue growth factor (CTGF), interferon-γ (IFNγ) and the irreversible proteasome inhibitor epoxomicin. First, cytotoxicity and proliferation assays were carried out. The expression of proteasome-related genes and proteins was assessed and proteasome activity was determined. Then, expression of fibrosis-associated factors fibronectin (FN), fibronectin EDA domain (FN EDA), metalloproteinase-2 (MMP-2), tissue inhibitor of metalloproteinases-1 (TIMP-1) and peroxisome proliferator-associated receptor-γ (PPARγ) was assessed. The proteasome inhibitor epoxomicin strongly arrested cell cycle progression and down-regulated TGFβ gene expression, which in turn was shown to induce expression of pro-fibrogenic genes in ARPE-19 cells. Furthermore, epoxomicin induced a directional shift in the balance between MMP-2 and TIMP-1 and was associated with down-regulation of transcription of extracellular matrix genes FN and FN-EDA and up-regulation of the anti-fibrogenic factor PPARγ. In addition, both CTGF and TGFβ were shown to affect expression of proteasome-associated mRNA and protein levels. Our results suggest a link between proteasome activity and pro-fibrogenic mechanisms in the RPE, which could imply a role for proteasome-modulating agents in the treatment of retinal disorders characterized by RPE-mediated fibrogenic responses.

## 1. Introduction

Age-related macular degeneration (AMD) is a progressive disease of the central retina-choroid tissue complex and one of the leading causes of blindness worldwide [1]. The retinal pigment epithelium (RPE), a polarized monolayer of epithelial cells that separates the neural retina from the vascularized choroid, has been implied to play an important role in the pathogenesis of the disease. Early AMD is characterized by focal drusen deposits in the macula, mostly located between the basal lamina of the RPE and the inner collagenous layer of Bruch's membrane [2]. Drusen contain carbohydrates, zinc and nearly 150 proteins

including vitronectin, apolipoproteins E and B, clusterin, connective tissue growth factor (CTGF) and complement system components [3,4]. Advanced AMD is divided into nonexudative or dry AMD which affects 8% of patients and is characterized by macular RPE atrophy and ensuing photoreceptor degeneration, and exudative or neovascular AMD (nAMD) which affects 5% of patients and is characterized by the development of choroidal neovascularization (CNV) [5]. CNV may ultimately lead to the development of a fibrous plaque or disciform scar that leads to secondary atrophy of the neurosensory retina and irreversible and untreatable loss of macular visual function [6–10]. The advent of anti-vascular endothelial growth factor (VEGF) therapy has

**Abbreviations:** AMD, age-related macular degeneration; ARPE-19, human retinal pigment epithelial cells; CNV, choroidal neovascularization; CTGF, connective tissue growth factor; EMT, epithelial-mesenchymal transition; ECM, extracellular matrix; FN, fibronectin; FN EDA, fibronectin EDA domain; IFNγ, interferon-γ; MMP-2, matrix metalloproteinase-2; nAMD, neovascular age-related macular degeneration; PPARγ, peroxisome proliferator-associated receptor-γ; RPE, retinal pigment epithelium; TIMP-1, tissue inhibitor of metalloproteinases-1; TGFβ, transforming growth factor-β; UPS, ubiquitin-proteasome system

\* Correspondence to: Ocular Angiogenesis Group, Departments of Ophthalmology and Medical Biology, Academic Medical Center, PO Box 22660, 1100 DD Amsterdam, The Netherlands.

E-mail addresses: [j.e.decarvalho@amc.uva.nl](mailto:j.e.decarvalho@amc.uva.nl) (J.E. Ramos de Carvalho), [milantv@hotmail.com](mailto:milantv@hotmail.com) (M.T. Verwoert), [i.m.vogels@amc.uva.nl](mailto:i.m.vogels@amc.uva.nl) (I.M.C. Vogels), [e.a.reits@amc.uva.nl](mailto:e.a.reits@amc.uva.nl) (E.A. Reits), [c.j.vannoorden@amc.uva.nl](mailto:c.j.vannoorden@amc.uva.nl) (C.J.F. Van Noorden), [i.klaassen@amc.uva.nl](mailto:i.klaassen@amc.uva.nl) (I. Klaassen), [r.o.schlingemann@amc.uva.nl](mailto:r.o.schlingemann@amc.uva.nl) (R.O. Schlingemann).

<https://doi.org/10.1016/j.bbrep.2018.01.005>

Received 28 November 2016; Received in revised form 8 January 2018; Accepted 9 January 2018

Available online 28 January 2018

2405-5808/ © 2018 The Authors. Published by Elsevier B.V. This is an open access article under the CC BY-NC-ND license

(<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

greatly improved the prognosis of nAMD patients, stabilizing or even improving visual function [11–13]. Subretinal fibrosis, however, is a common ensuing process of CNV membrane formation, occurring in approximately half of anti-VEGF treated eyes [7,10,14].

Fibrosis may be considered as a deregulated wound healing response to tissue damage [10,15,16]. Angiogenesis occurs in this process as an initial trigger for fibrin deposition, tissue repair, oxygen supply and recruitment of inflammatory cells to the wound [10,17]. In AMD, angiogenesis occurs in the subretinal or sub-RPE space, leading to exudation, hemorrhage and eventually fibrosis. During this process, various types of cells such as RPE cells, glial cells, fibroblasts, myofibroblast-like cells and macrophages infiltrate and/or proliferate, secreting pro-angiogenic and pro-fibrogenic factors that interact with inflammatory cytokines and growth factors. Prevention of visual loss in AMD may therefore depend on the development of successful therapeutic regimens that can halt subretinal fibrosis and preserve the RPE.

The fibrogenic response is stimulated by inflammatory-derived cytokines and growth factors, including transforming growth factor- $\beta$  (TGF $\beta$ ) [18], an ubiquitously expressed growth factor belonging to the large superfamily of activins/bone morphogenetic proteins [19] and connective tissue growth factor (CTGF), a member of the connective tissue growth factor/cysteine-rich 61/nephroblastoma overexpressed (CCN) family of extracellular matrix (ECM) proteins, also known as CCN2 [20,21]. The expression of CTGF is regulated by TGF $\beta$  [22–24] and, likewise, CTGF has been shown to be an important mediator of TGF $\beta$  signaling and its effects in different cell types [24–32]. We and others have shown that both TGF $\beta$  and CTGF are major players in the fibrogenic response in the retina [4,16,21,33–40].

The ubiquitin-proteasome system (UPS), a multi-catalytic cytoplasmic and nuclear protein complex present in all eukaryotic cells, is responsible for non-lysosomal proteolysis and thus maintenance of a normal protein homeostasis in cells [41]. Mounting evidence suggests that UPS dysfunction is a major pathogenic mechanism in senile degenerative disorders [42], including AMD and other ophthalmic conditions [43–53]. Proteasomes diffuse rapidly in the cytoplasm and nucleus where they encounter intracellular proteins that are appropriately tagged or misfolded. Proteins are tagged by ubiquitination processes and as such recognized by the 19S regulatory particle of the proteasome [54]. Ubiquitin has been shown to be uniformly expressed in the RPE-Bruch's membrane complex of patients afflicted with AMD [53]. The 19S regulatory particle, combined with the 20S catalytic core, forms the standard proteasome. Within the proteasome core, specialized catalytic subunits are responsible for the cleavage of the carboxyl termini of proteins. There are 3 catalytic subunits in the standard proteasome:  $\beta$ 1 for acidic amino acids,  $\beta$ 2 for basic amino acids, and  $\beta$ 5 for hydrophobic amino acids. The standard proteasome may in some instances undergo a change in configuration into the immunoproteasome. This is achieved upon replacement of the constitutive subunits in the standard proteasome by inducible subunits,  $\beta$ 1i,  $\beta$ 2i, and  $\beta$ 5i [55,56]. Although uninjured RPE contains a baseline level of immunoproteasome subunits [52], cellular stress, such as retinal injury by cytotoxic T-lymphocytes [47], optic nerve trauma [57], aging mechanisms [44,48], complement overactivation [52], chronic oxidative stress [58] and exposure to pro-inflammatory cytokines [52,59] may increase the number of active immunoproteasome subunits. Therefore, the ratio between the nascent ( $\beta$ 1,  $\beta$ 2, and  $\beta$ 5) and inducible subunits ( $\beta$ 1i,  $\beta$ 2i, and  $\beta$ 5i) may be used as a marker of cellular stress [47,48,58].

The aim of the present study was to characterize the involvement of the proteasome pathway in TGF $\beta$  and CTGF-mediated expression of ECM genes in RPE cells. Likewise, potential anti-fibrogenic effects of the selective proteasome inhibitor epoxomicin were assessed in ARPE-19 cell cultures.

## 2. Materials and methods

### 2.1. Culture, maintenance and treatment of ARPE-19 cells

Experiments were conducted using ARPE-19 cells, a human RPE cell line that has structural and functional properties that are characteristic of RPE cells in vivo. Monolayers of cells cultured on transwell filters reached a transepithelial resistance of 30–40  $\Omega$  cm<sup>2</sup> after 3 weeks of culture and expressed CRALBP, as detected by RT-PCR. Cells were cultured at 37 °C in 5% CO<sub>2</sub> in gelatin-coated T75 cell culture flasks (Corning, Lowell, MA, USA) in Dulbecco Modified Eagle Medium (DMEM; Gibco Life Technologies, Carlsbad, CA, USA), low glucose, pyruvate in the presence of 1% penicillin/streptomycin and 10% fetal calf serum. Cell growth was monitored and medium was changed twice a week. For passaging of cells, TrypLE Express (Invitrogen, Carlsbad, CA, USA) was used and cell suspensions were diluted 3-fold. For experiments, cells were cultured in 6-well plates. Upon confluence, cells were washed once with phosphate-buffered saline (PBS), serum starved for 24 h and then treated with various concentrations of the selective and irreversible proteasome inhibitor epoxomicin (Sigma-Aldrich, St. Louis, MO, USA), rhCTGF (ProSpec-Tany TechnoGene, Rehovot, Israel), rhTGF $\beta$ 1 (ProSpec) and interferon- $\gamma$  (IFN $\gamma$ ) (PBL Biomedical, Piscataway, NJ, USA). All experiments were performed in triplicate and repeated at least twice.

### 2.2. Protein extraction

Cells were harvested using TrypLE Express (Invitrogen), collected in Eppendorf tubes and centrifuged for 10 min at 400g. Supernatant was removed and the pellet was suspended in TSDG buffer (10 mM Tris, pH 7.5, 25 mM KCl, 10 mM NaCl, 1.1 mM MgCl<sub>2</sub>, 0.1 mM EDTA, and 8% glycerol), 5 mM ATP and 1x protease inhibitor (Roche Applied Science, Penzberg, Germany).

Cells were lysed with 3 cycles of freezing in liquid nitrogen and thawed at room temp. After centrifugation (15 min, 10,000g), the protein concentration was determined using a Bradford protein assay (Serva, Heidelberg, Germany). All experiments were performed in triplicate and repeated at least twice.

### 2.3. Cell cycle and cell viability assays

To assess the viability of healthy ARPE-19 cells and to assess the toxic effects of different stimulants used throughout assays, the PrestoBlue cytotoxicity assay (Invitrogen) was performed according to the manufacturer's instructions. The assays were carried out in 96-well plates (roughly 10,000–25,000 cells per well). After cells were conditioned and washed, PrestoBlue reagent was added to each well. The plates were subsequently incubated at 37 °C for the recommended time period (20–30 min). After incubation, the solution containing PrestoBlue reagent from the wells of the assay plates was transferred to new wells in a 96-well plate, and absorbance was read on a plate reader (Bio-Rad, Hercules, CA, USA) with the excitation/emission wavelengths set at 570/600 nm.

To evaluate the effects of different stimulants on cell proliferation, the Click-iT EdU Alexa Fluor 488 imaging kit (Invitrogen) was applied according to the protocol provided by the manufacturer. Briefly, ARPE-19 cells at 30–50% confluence were treated with EdU (10  $\mu$ M). EdU was added 2 h prior to the addition of CTGF and TGF $\beta$  (both 24 h incubation) and epoxomicin (16 h incubation). Subsequently, cells were fixed, permeabilized, and click-labeled. As a negative control, untreated cells were used. Following incubation, fluorescence readout was determined using a FACS LSRII (Becton Dickinson, Breda, The Netherlands) to determine percentages of EdU-proliferative cells in S and M phase and EdU-negative quiescent cells in G0 and G1 phase. The experiment was performed in triplicate and repeated twice (N = 2).

**Table 1**  
Primer details.

Gene	GenBank	Forward primer	Reverse primer	Size (bp)	T <sub>m</sub> (°C)
PSME1	NM_006263	CAGCCCCATGTGGGTGATTATC	GCTTCTCGAAGTTCTTCAGGATGAT	139	82
PSMA7	NM_002792	CCTGGAAGGCCAATGCCATAG	TTTGCCACCTGACTGAACCACTTC	149	82
PSMB5	NM_002797	CCATGATCTGTGGCTGGGATAAG	GGTCATAGGAATAGCCCCGATC	144	83
PSMB8	NM_004159	CTGGAGGCGTTGTCAATATGTACC	GCAGCAGGTCACTGACATCTGTAC	81	76
VEGFA	NM_003375	GGCAGAAGGAGGAGGGCAGAAT	CACCAGGGTCTCGATTGGATGG	91	80
FN1	NM_002026	TGGGACCGTCAGGAGAGAAATG	CAGGAGCAAAATGGCACCAGAT	167	82
FN EDA	XM_005246414	GCAGTGACCAACATTGATCGC	ACCCTGTACCTGGAAACTTGCC	110	80
MMP2	NM_004530	GGAATGCCATCCCCGATAACC	CCAGCTTCAGGTAATAGGCACCCCT	93	83
TIMP1	NM_003254	ACTTCCACAGGTCCCAACACCG	AGGGAAACACTGTGCATTCTCTAC	180	84
PPARG	NM_138712	CCTCGGAAAGCCTTTTGGTGAC	AAACCTGGGCGGTCTCCACT	135	79

## 2.4. Proteasome activity labeling

Proteasome subunits were labeled in the lysate with 0.5  $\mu$ M activity-based probe BODIPY-epoxomicin for 1 h at 37 °C (BodipyFl-Ahx3L3VS, MV121, provided by H. Overkleeft, Institute of Chemistry, Leiden, The Netherlands) [60] and sample buffer (350 mM Tris-HCl pH 6.8, 10% SDS, 30% glycerol, 6%  $\beta$ -mercaptoethanol, 0.02% bromophenol blue), added to 20  $\mu$ g protein lysate. Samples were boiled for 5 min and loaded on a 12.5% SDS-PAGE gel. As a positive control, maximal proteasome activity was determined after treatment with IFN $\gamma$  (50 U). After running the proteins on the gel, fluorescence imaging was performed on a Trio Typhoon (GE Medical Systems, Little Chalfont, UK) using the 580 bandpass filter to detect the probe directly on the gel. Proteasome total activity values were normalized according to the total proteasome content in cells as indicated by the levels of the  $\alpha$ 7 subunit of the 20S proteasome (1:1000; MCP72; Enzo Life Sciences, Zandhoven, Belgium). The experiment was performed in triplicate and repeated three times (N = 3).

## 2.5. Western blot analysis

Proteins were isolated using a 1% Triton X-100 cell lysis buffer (10 mM Hepes, 150 mM NaCl, 10% glycerol, 1.5 mM MgCl<sub>2</sub>, 1 mM EGTA, 1% Triton X-100 and 1x Complete Protease Inhibitors; Roche Biochemicals, Almere, The Netherlands). All samples were run on SDS-PAGE under denaturing conditions. Briefly, 20  $\mu$ g of protein was loaded on a 12.5% SDS-PAGE gel, and after electrophoresis transferred to nitrocellulose membranes using a Trans-Blot Turbo Transfer System (Bio-Rad). Membranes were incubated overnight or longer at 4 °C with a monoclonal antibody against the  $\alpha$ 7 subunit of the 20S proteasome (1:1000; BML-PW8110-0025; Enzo Life Sciences) and the following polyclonal antibodies: anti- $\beta$ 5 subunit (1:1000; BML-PW8895-0100; Enzo Life Sciences) and anti- $\beta$ 5i subunit (1:1000; ab3329; Abcam, Cambridge, UK). Anti  $\beta$ -actin (1:10,000; a5441; Sigma-Aldrich, St. Louis, MI, USA) was determined for the loading control. Intensity of bands was quantified by densitometric analysis using Odyssey (LI-COR Biosciences, Lincoln, NE, USA). Quantification was performed with Image studio Lite 4.0 (LI-COR). Values were normalized using  $\beta$ -actin (1:10,000; ab8227; Abcam). The experiment was performed in triplicate and repeated three times (N = 3).

## 2.6. RNA isolation and mRNA quantification

Total RNA was isolated from ARPE-19 cell cultures using TRIzol reagent (Invitrogen) according to the manufacturer's instructions. ARPE-19 cells were stimulated with TGF $\beta$  (5 ng, 30 ng or 50 ng for 24 h), CTGF (50 ng or 200 ng for 24 h), epoxomicin (50 nM, 100 nM, 250 nM, 500 nM for 16 h) and IFN $\gamma$  (50 U for 72 h) in 6-well plates. In addition, in order to assess whether epoxomicin treatment was able to counteract TGF $\beta$ -mediated responses, TGF $\beta$ -stimulated ARPE-19 cells were treated with 500 nM epoxomicin. Total RNA (1  $\mu$ g) was treated

with DNase I (amplification grade; Life Technologies) and reverse transcribed into first strand cDNA using a Maxima® First Strand cDNA Synthesis Kit (Thermo Scientific, Roskilde, Denmark). Real-time qPCR was performed using a CFX96 system (Bio-Rad) as described previously [61]. Primer details are given in Table 1. Ct-values were converted to absolute amounts with the formula  $2^{-Ct}$  and taken relative to the absolute amounts of control samples, that were set to 1. The experiment was performed in triplicate and repeated three times (N = 3).

## 2.7. Enzyme-linked immunosorbent assay (ELISA)

ARPE-19 cell samples treated with epoxomicin (50 nM, 100 nM, 250 nM, 500 nM for 16 h) and supernatant was collected. Concentrations of activated TGF $\beta$ 2 were determined by Quantikine ELISA assays according to the manufacturer's protocol (R&D Systems, Minneapolis MS, USA). The experiment was performed in triplicate and repeated twice (N = 2).

## 2.8. Statistical analysis

Data are presented as fold change, with a fold change of 1.0 meaning the same level as control samples. Asterisks (\*) indicate a significant change relative to the control samples. Differences between experimental conditions were calculated with one-way or two-way ANOVA with P < 0.05 indicating a statistical difference. Statistical analysis of data was performed using IBM SPSS 20 (SPSS, Chicago, IL, USA).

## 3. Results

### 3.1. TGF $\beta$ and epoxomicin arrest cell cycle progression

CTGF (50 and 200 ng for 24 h), TGF $\beta$  (5 and 50 ng for 24 h) and epoxomicin (50 and 500 nM for 16 h) did not induce any significant cytotoxicity in ARPE-19 cells (data not shown). TGF $\beta$  significantly reduced cell proliferation (35–37% cell cycle arrest in G0/G1 phase corresponding to a 20% increase when compared to control), whereas CTGF did not show an anti-proliferative effect (Fig. 1). On the other hand, epoxomicin, a cell-permeable potent and selective irreversible proteasome inhibitor [62], strongly arrested cell cycle progression (Fig. 1). These results suggest that epoxomicin has significant anti-proliferative effects in ARPE-19 cells.

### 3.2. CTGF and TGF $\beta$ affect proteasome subunit mRNA levels

As expected, treatment with high doses of epoxomicin significantly decreased mRNA levels of PA28 $\alpha$  and  $\beta$ 5i (data not shown), whereas IFN $\gamma$  increased mRNA levels of  $\alpha$ 7 and  $\beta$ 5 (Fig. 2). CTGF marginally increased mRNA expression of subunit  $\beta$ 5 (Fig. 2). Upon TGF $\beta$  stimulation, mRNA levels of the immunoproteasome subunit  $\beta$ 5i and proteasome regulatory subunit PA28 $\alpha$  were decreased.




Fig. 1. Proliferation of ARPE-19 cells in the presence of TGFβ, CTGF, TGFβ and CTGF or epoxomicin expressed as percentage of cells in the S and M phase versus cells in the G0 and G1 phase after flow cytometric analysis of the percentage of cells that had incorporated EdU. \*Significant difference from control of percentage of cells in G0 and G1 phase. \*\*Significant difference from control of percentage of cells in the S and M phase. The experiment was performed in triplicate and repeated twice (N = 2).




Fig. 2. Proteasome mRNA levels induced by IFNγ, TGFβ or CTGF. mRNA levels of PA28α, β5, β5i, and α7 subunits of the proteasome in ARPE-19 cells, after stimulation with IFNγ, TGFβ and low and high concentrations of CTGF. IFNγ upregulates mRNA expression of α7 and β5 subunits whereas TGFβ downregulates mRNA expression of PA28α and β5i subunits and CTGF upregulates mRNA expression of β5 subunit. Values represent mRNA expression levels (mean ± SD) relative to untreated control cells. \*, Significant change (P < 0.05); \*\*\*, significant change (P < 0.001). The experiment was performed in triplicate (N = 3).

### 3.3. CTGF increases the β5i/β5 ratio

To assess whether the induced changes in expression of proteasome-associated genes was associated with increased protein expression levels of the respective subunits, we performed western blotting on ARPE-19 cells treated with CTGF (50 ng), TGFβ (50 ng), a combination of CTGF and TGFβ, and IFNγ as positive control (Fig. 3). The ratio of β5i and β5 (β5i/β5) was taken as a marker of immunoproteasome activation. As expected, IFNγ induced maximal immunoproteasome activation, as indicated by a 28-fold change in the β5i/β5 ratio (data not shown). Protein levels of β5i were slightly higher in the presence of CTGF (Fig. 3A,B) which translated in a 19% increased β5i/β5 ratio (Fig. 3C), whereas TGFβ stimulation down-regulated the expression level of the proteasome β5i subunit by 33% (Fig. 3A,B), whereas the β5i/β5 ratio was not significantly affected (Fig. 3C).

These results indicate that, to a limited extent, expression of immunoproteasome β5i subunit protein is up-regulated by CTGF and down-regulated by TGFβ.

### 3.4. CTGF upregulates proteasome activity in ARPE-19 cells

To assess whether proteasome gene expression and protein level changes induced by CTGF and TGFβ affect proteolytic activity of the proteasome, the activity of individual subunits after treatment of cells with CTGF, TGFβ or IFNγ was determined (Fig. 4). IFNγ, as expected, induced a substantial increase (4–5-fold change, P = 0.049) in activity of all subunits. CTGF (50 ng) significantly up-regulated the activity of the β1/β5i complex (2.6-fold change, P = 0.005) and β5/β1i complex (1.5-fold change, P = 0.026). TGFβ (30 ng) did not affect proteasomal activity, whereas a combination of CTGF (50 ng) and 6 h later TGFβ (5 ng) did not induce changes in the activity of the β1/β5i and β5/β1i complexes.

These results demonstrate that CTGF upregulates the activity of specific proteasome subunits, probably mediated by a change in the configuration of the standard proteasome into the immunoproteasome and up-regulation in mRNA and protein levels of proteasome β5 and β5i subunits.

### 3.5. TGFβ up-regulates mRNA levels of ECM-associated genes

In order to characterize the effects of TGFβ and CTGF on the transcription of ECM-associated genes, we assessed the mRNA levels of CTGF, TGFβ1 and TGFβ2, VEGF, fibronectin (FN), fibronectin EDA domain (FN EDA), metalloproteinase-2 (MMP-2), tissue inhibitor of metalloproteinases-1 (TIMP-1) and peroxisome proliferator-associated receptor-γ (PPARγ) upon stimulation with CTGF, TGFβ, and CTGF followed by TGFβ (Fig. 5).

TGFβ up-regulated mRNA levels of CTGF and VEGF (Fig. 5A). The same effect was observed when ARPE-19 cells were treated with CTGF followed by TGFβ (data not shown). This effect was dependent on the concentration of TGFβ, which implies that TGFβ may be the main mediator of this response in ARPE-19 cells.


With respect to fibrosis-related genes, TGFβ, but not CTGF, up-regulated mRNA levels of FN EDA, FN and MMP-2 (Fig. 5B). Transcript levels of the anti-fibrogenic factor PPARγ were down-regulated in the presence of TGFβ (Fig. 5B). Again, these effects were dependent on the concentration of TGFβ, irrespective of simultaneous treatment with different concentrations of CTGF (data not shown). These results confirm the role of TGFβ as a major pro-fibrogenic mediator in RPE cells.

### 3.6. Proteasome inhibition by epoxomicin down-regulates expression of ECM-associated genes

In order to test the effects of proteasome modulation on mRNA expression of CTGF, TGFβ1, TGFβ2, VEGF, FN, FN EDA, TIMP-1, MMP-2 and the anti-fibrogenic protein PPARγ, we assessed the effects of different concentrations of epoxomicin and TGFβ plus epoxomicin (Fig. 6).

Treatment of ARPE-19 cells with epoxomicin resulted in decreased mRNA levels of TGFβ1 and TGFβ2 (Fig. 6A). Mean levels of activated TGFβ2 protein were strongly reduced (to undetected levels when compared to untreated cells) after treatment of ARPE-19 cells with 50–500 nM epoxomicin (data not shown). At low concentrations (50 and 100 nM), epoxomicin down-regulated VEGF mRNA expression (Fig. 6A). Furthermore, treatment with epoxomicin resulted in decreased mRNA levels of FN, FN EDA, TIMP-1 and a corresponding increase in MMP-2 mRNA levels (Fig. 6B). Down-regulation of mRNA expression of TGFβ1, TGFβ2, FN EDA and VEGF with 500 nM epoxomicin was also observed in TGFβ-treated ARPE-19 cells. In addition, epoxomicin, alone or in the presence of TGFβ, down-regulated the expression of the anti-fibrogenic mediator PPARγ (Fig. 6B). These results





**Fig. 3.** Proteasome-specific subunit protein expression and  $\beta 5i:\beta 5$  ratios upon IFN $\gamma$ , CTGF, TGF $\beta$  or CTGF and TGF $\beta$  stimulation of ARPE-19 cells. (A) Western blot showing protein levels of  $\beta 5i$ ,  $\alpha 7$  and actin (loading control) in cells that had been incubated with IFN $\gamma$ , CTGF or TGF $\beta$ . (B) Protein levels of  $\beta 5$  and  $\beta 5i$  subunits were assessed by western blot with actin as loading control and  $\alpha 7$  subunit as proteasome content control. (C) Quantitative data of the average ratio of  $\beta 5i$  and  $\beta 5$  relative to control samples after incubation in the presence or absence of CTGF, TGF $\beta$  or CTGF and TGF $\beta$ . Data are expressed as the mean  $\pm$  SD. \*, Significant change ( $P < 0.05$ ); \*\*, significant change ( $P < 0.01$ ). The experiment was performed in triplicate ( $N = 3$ ).

suggest that epoxomicin, even upon TGF $\beta$  secretion, counteracts the pro-fibrogenic transcription effects of TGF $\beta$ .


#### 4. Discussion

This study attributes a role to the proteasome pathway in modulation of part of the fibrogenic response of RPE cells which is a multifactorial response dependent on activation and suppression of a myriad of growth factors and cytokines. For the purpose of this study, we selected TGF $\beta$  and CTGF as both are regarded as important mediators of pathological fibrosis in the eye and other organs [4,20–26,28,29,31,33,35–40,63–74]. Likewise, emerging evidence suggests a link between the fibrogenic response, proteasome modulation and TGF $\beta$  signaling in multiple systemic conditions [75–85].


Routine passaging of ARPE-19 cells was used as an *in vitro* wound response model to study the fibrogenic response in the retina. Epithelial cells (such as the RPE in the retina) are considered to be the major mediators of fibrogenic responses to tissue injury [10,71]. The RPE is a highly polarized monolayer of epithelial pigmented cells between the choroid and the neurosensory retina that plays a crucial role in the maintenance of visual function [86]. RPE cells proliferate and undergo epithelial-mesenchymal transition (EMT) when dissociated into single cells [87,88], whereas sheets of RPE cells in culture preserve their

morphology for a longer period of time [89,90]. Disorganization and extensive damage to the RPE such as during subretinal CNV membrane formation, is a prerequisite for development of subretinal fibrosis [16]. Accordingly, subretinal fibrosis is frequently reported in late stages of nAMD [7,14]. Fibrosis in other organs such as lung, kidney, liver, skin and heart follows pathogenic pathways similar to subretinal fibrosis development in nAMD [10,91]. In all these tissues, an intact epithelium is considered protective against fibrosis development [92]. Although the ARPE-19 cell line was deemed appropriate for the purpose of this study, confirmation of the attained results in an *in vivo* model is warranted. Our study is focused on the fibrogenic response of RPE cells, however, we acknowledge that RPE cells are only one of many other cell types involved in RPE-mediated fibrosis.

Our results demonstrate that CTGF is associated with activation of the proteasome as demonstrated by the increased proteolytic activity of specific proteasome complexes, namely  $\beta 1/\beta 5i$  and  $\beta 5/\beta 1i$ . The proteasome activity probe assay is unable to discriminate between the activities of the various proteasome subunits. Nevertheless, it is likely that the observed changes in proteasome activity stem from  $\beta 5$  and  $\beta 5i$  increased proteolytic activity since these are known to be the rate-limiting subunits at the level of the RPE [49,51]. Accordingly, these changes in proteasome activity are accompanied by a slight but significant up-regulation in  $\beta 5$  and  $\beta 5i$  protein expression and  $\beta 5$  mRNA




**Fig. 4.** Increased specific proteasome activity upon CTGF stimulation of ARPE-19 cells. (A) After treatment with IFN $\gamma$  (50 U), TGF $\beta$  (50 ng), CTGF (50 ng) or CTGF (50 ng) followed 6 h later by TGF $\beta$  (5 ng), ARPE-19 cells were harvested and proteasomes were labeled with a Bodipy-Ep activity probe. Proteasome activities were assessed by western blotting. Quantitative data of the proteasome activity are presented for proteasome subunit  $\beta 2$  (B),  $\beta 1/\beta 5i$  complex (C), and  $\beta 5/\beta 1i$  complex (D). \*, Significant change ( $P < 0.05$ ); \*\*, significant change ( $P < 0.01$ ). The experiment was performed in triplicate ( $N = 3$ ).



**Fig. 5.** TGF $\beta$  upregulates mRNA expression of CTGF, VEGF and pro-fibrogenic genes and downregulates mRNA expression of the anti-fibrogenic factor PPAR $\gamma$ . After stimulation with TGF $\beta$ , CTGF and IFN $\gamma$ , mRNA levels of (A) CTGF, TGF $\beta$ 1, TGF $\beta$ 2, VEGF and (B) FN, FN EDA, MMP-2, TIMP-1 and PPAR $\gamma$  were assessed in ARPE-19 cells. Values represent mRNA expression levels (mean  $\pm$  SD) relative to untreated control cells. \*, Significant change ( $P < 0.05$ ); \*\*, significant change ( $P < 0.01$ ); \*\*\*, significant change ( $P < 0.001$ ). The experiment was performed in triplicate ( $N = 3$ ).

levels. Conversely, TGF $\beta$  was associated with a down-regulation of  $\beta$ 5i and proteasome regulatory subunit PA28 $\alpha$  mRNA levels. Association of PA28 $\alpha$  with the 20S catalytic core has been shown to increase proteasome activity [93–95]. Studies have shown that expression of PA28 $\alpha$  tends to decline in aged retina [44]. Unlike the effects of CTGF, proteasome activity assays in the presence of TGF $\beta$  demonstrated no changes in the proteolytic activity of specific proteasome subunit complexes. Recent evidence has suggested that the immunoproteasome, besides its role in immune surveillance, may be considered as a rescue mechanism in response to cellular stress [44,47,48,52, 57–59,93,96–98]. Correspondingly, immunoproteasome activation has been demonstrated in the RPE of a mouse model of age-related atrophic degeneration of the RPE [52] as well as in the retina of AMD human donors [44].

There is converging evidence for TGF $\beta$  as an important pro-fibrogenic factor in the RPE. RPE cells from CNV membranes are strongly immunoreactive for TGF $\beta$  [99] and the RPE has been shown to be an intraocular secretion site of TGF $\beta$  [100]. Elevated mRNA levels of TGF $\beta$  (TGF $\beta$ 2 isoform) have been demonstrated in the RPE-choroid complex and retina of AMD patients [101] and in the vitreous of patients with proliferative vitreoretinopathy and proliferative diabetic retinopathy [70,102]. TGF $\beta$  can induce EMT of RPE cells in suspension, but fails to do so when RPE cells have well-established cell-cell contacts [88]. Earlier studies have confirmed that TGF $\beta$  is an inducer of a number of growth factors such as CTGF, platelet-derived growth factor, fibroblast growth factors, and VEGF, as well as TGF $\beta$  itself [103,104]. Likewise, in the presence of a TGF $\beta$  signaling inhibitor (A-83-01), RPE cells were more tolerant to continuous wound response triggers (such as routine passaging of cell cultures) and retained the capacity to acquire a pigmented epithelial morphology [72]. In addition, inhibition of TGF $\beta$  signaling did not prevent RPE differentiation or productive RPE-mediated wound repair [72]. Semaphorin 3A, a TGF $\beta$  inhibitor, suppressed laser-induced CNV formation in mice by inhibition of the Smad2/3



**Fig. 6.** Epoxomicin (Ep) downregulates mRNA expression of TGF $\beta$ 1, TGF $\beta$ 2, VEGF, FN, FN EDA, TIMP-1 and upregulates mRNA expression of CTGF and PPAR $\gamma$ . In the presence of TGF $\beta$ , epoxomicin downregulates mRNA expression of TGF $\beta$ 1, TGF $\beta$ 2, VEGF and FN EDA and expression of PPAR $\gamma$  is up-regulated. After treatment with increasing concentrations of epoxomicin in untreated and TGF $\beta$ -treated ARPE-19 cells, mRNA levels of (A) CTGF, TGF $\beta$ 1, TGF $\beta$ 2, VEGF, (B) FN, FN EDA, MMP-2, TIMP-1 and PPAR $\gamma$  were assessed. Values represent mRNA expression levels (mean  $\pm$  SD) relative to untreated control cells. \*, Significant change ( $P < 0.05$ ); \*\*, significant change ( $P < 0.01$ ); \*\*\*, significant change ( $P < 0.001$ ). The experiment was performed in triplicate ( $N = 3$ ).

signaling pathway [105]. These and our results suggest that TGF $\beta$  may be regarded as a fibrogenic marker in disrupted RPE cells and that targeting of TGF $\beta$ -mediated effects may improve wound repair mechanisms. Our results demonstrate that inhibition of the proteasome by epoxomicin is associated with down-regulation of the expression of TGF $\beta$  (both isoforms TGF $\beta$ 1 and TGF $\beta$ 2) and a complete blockage of TGF $\beta$ 2 activity. These two isoforms were studied because *in vivo* roles and expression of the different TGF $\beta$  isoforms may not be uniform, although *in vitro* experiments often elicit similar responses [106].


CTGF expression is regulated by several signaling mechanisms including pathways of TGF $\beta$ /Smad [107]. TGF $\beta$  is a major inducer of CTGF [20,28] whereas CTGF has been shown to synergistically enhance the effects of TGF $\beta$ . Our results attribute an independent role for CTGF and TGF $\beta$  regarding modulation of the proteasome and fibrogenic response in RPE cells. When given consecutively, the effects of CTGF and TGF $\beta$  on proteasome expression and activity remained unchanged. Similarly, the pro-fibrogenic effects of TGF $\beta$  were not synergized by CTGF treatment. In other cell types, such as in hepatocytes, CTGF has been demonstrated to affect TGF $\beta$  signaling by facilitating binding of TGF $\beta$  to its receptor, down-regulation of the negative feedback loop via Smad7 and inhibition of receptor binding and signaling of the physiological TGF $\beta$  antagonist BMP-7 [28]. Likewise, CTGF is considered to be a downstream mediator of certain effects attributed to TGF $\beta$  such as cell proliferation, migration, adhesion, ECM production and EMT [20]. In the eye, CTGF has been shown to accumulate in basal deposits and in Bruch's membrane of early AMD specimens [4]. In accordance with our results, CTGF secretion has been shown to be linked to up-regulation of the expression of the ECM components FN, laminin and MMP-2 in ARPE-19 cells by mechanisms involving activation of ERK and p38 MAPK signaling pathways [4]. Furthermore, vitreous levels of CTGF correlated strongly with degree of fibrosis in vitreoretinal disorders

such as proliferative vitreoretinopathy, proliferative diabetic retinopathy and macular hole [33]. Although expression of CTGF may occur independently of TGF $\beta$  in other retinal cell types [32], our results suggest CTGF expression may be regulated by TGF $\beta$ -mediated pathways in RPE cells. In the presence of TGF $\beta$ , epoxomicin suppressed the strong up-regulation induced by TGF $\beta$  on CTGF mRNA levels. Since the pro-fibrogenic effects of TGF $\beta$  were more significant than those of CTGF, targeting of the TGF $\beta$  pathway instead of CTGF may have a more substantial anti-fibrogenic effect in subretinal fibrosis. Potential anti-fibrogenic effects were demonstrated after simultaneous treatment of RPE cell cultures with anti-VEGF (bevacizumab) and a CTGF inhibitor, but not when the CTGF inhibitor was administered alone [63]. Targeting the TGF $\beta$  pathway, however, could be less attractive due to concomitant inhibition of anti-inflammatory properties attributed to TGF $\beta$  alongside other important cellular effects [21].

Epoxomicin is one of the most selective inhibitors of the proteasome. Indeed, proteasomal subunits are the only cellular proteins covalently modified by the biotinylated derivatives of epoxomicin with no other proteolytic enzymes inhibited along this process [108]. Treatment of RPE cells with epoxomicin down-regulated the expression of the pro-fibrogenic ECM mRNA levels of FN and FN-EDA. In the presence of TGF $\beta$ , epoxomicin still down-regulated mRNA expression of FN EDA, VEGF, TGF $\beta$ 1 and TGF $\beta$ 2. FN, a glycoprotein that mediates cellular adhesion and migration of RPE cells, is one of the components of the ECM that is expressed in early phases of fibrosis [109]. FN is composed of two cross-linked subunits. Alternative splicing of the FN gene transcript results in several variants. One isoform (FN-EDA) has an extra domain in cellular FN [110]. Expression of FN-EDA is significantly increased in specific stages of embryonic development, during wound healing processes in the adult and in several fibrogenic diseases [69].

Furthermore, epoxomicin led to a shift in the balance between MMP-2 and TIMP-1, as mRNA levels of MMP-2 were up-regulated and TIMP-1 levels were down-regulated. As demonstrated in other cell lines [111], TGF $\beta$  increased MMP-2 levels without affecting mRNA levels of TIMP-1. RPE cells are known to express MMP-2 and TIMP-1 [112,113]. MMPs are involved in a number of normal and physiological responses such as degradation of the basal lamina, remodeling of ECM, connective tissue turnover, angiogenesis and wound repair mechanisms [111]. TIMPs are the natural inhibitors of the functional effects of MMPs [114]. TIMPs have been shown to suppress angiogenesis and promote fibrosis by inhibiting the degradation and processing of ECM proteins [115]. TIMP-1, in particular, has a pivotal role in the fibrogenic response [115]. In vitreous samples of patients afflicted with proliferative diabetic retinopathy, TIMP-1 and MMP-2 were shown to be involved in angiogenesis, with TIMP-1 possibly acting as a natural anti-angiogenic factor [38]. Moreover, the balance between MMPs and TIMPs may be important for the integrity of ECM components, including, amongst others, collagens, vitronectin, fibronectin, laminin, elastin, and proteoglycans, and as such it may be regarded as indicative of the initiation and progression of the fibrogenic response [16,116]. The observed changes in the MMP/TIMP-1 balance by epoxomicin may result in improved breakdown of ECM components and subsequent attenuation of RPE cells' migration and fibrogenic responses. On the other hand, these effects on the MMP/TIMP ratio might be transitory, as evidenced in a previous study in which TGF $\beta$  inhibition had a late inhibitory effect on MMP-2 mRNA levels [75].

The mechanism(s) by which proteasome inhibition protects against fibrosis remain unknown. Our results suggest that PPAR $\gamma$  modulation, alongside inhibition of TGF $\beta$  expression, may explain the anti-fibrogenic properties of epoxomicin. PPAR $\gamma$ , a member of the nuclear receptor superfamily, is a ligand-activated transcription factor known to be involved in various distinct physiological processes including fat cell differentiation, glucose homeostasis, lipid metabolism, aging and inflammatory and immune responses [117,118]. Furthermore, PPAR $\gamma$  possesses important anti-angiogenic and anti-fibrogenic properties and is involved in the oxidative stress response [119]. In the retina, PPAR $\gamma$



**Fig. 7.** A role for proteasome inhibition in the modulation of fibrogenic mechanisms mediated by RPE cells. TGF $\beta$  activates multiple pathways, including the Smad, Rho-like GTPase, PI3K/AKT and MAPK pathways, resulting in the transcription of several pro-fibrogenic genes such as CTGF, FN, FN EDA, VEGF and down-regulation of PPAR $\gamma$  transcription. These effects contribute to epithelial-mesenchymal transition processes in RPE cells and initiation of fibrosis. Proteasome inhibition halts cell cycle progression and downregulates transcription of FN, FN EDA, TGF $\beta$ , VEGF and TIMP-1 whilst transcription of the anti-fibrogenic factor PPAR $\gamma$  is up-regulated, also upon exposure to TGF $\beta$ . Effects of proteasome inhibition are depicted as dotted lines. Abbreviations: connective tissue growth factor (CTGF); fibronectin (FN); fibronectin EDA (FN EDA); MAP kinase pathway (MAPK); phosphatidylinositol-3-kinase pathway (PI3K/AKT); retinal pigment epithelium cells (RPE); Rho-like GTPase pathway (RhoA/ROCK); tissue inhibitor metalloproteinase-1 (TIMP-1); transforming growth factor  $\beta$  (TGF $\beta$ ); transforming growth factor receptor (TGF $\beta$ R); vascular endothelial growth factor (VEGF).

has been shown to be involved in multiple molecular processes, including VEGF-induced choroidal angiogenesis response [120], photo-receptor renewal process [121], retinal neuroprotection [122] and protection from oxidative stress [123,124]. Degradation of PPAR $\gamma$  has been reported to occur via the proteasome [125]. Inhibition of TGF $\beta$  signaling by PPAR $\gamma$  has been attributed to restriction of Smad 2,3 binding to TGF $\beta$ -responsive promoters [76]. After phosphorylation, Smad2,3 forms a complex with other Smad proteins, which in turn facilitate translocation to the nucleus. In the nucleus, coactivators or repressors (such as PPAR $\gamma$ ) regulate the binding of the Smad complex with DNA [82]. Our results demonstrate inhibition of the proteasome was able to counteract TGF $\beta$ -mediated down-regulation of PPAR $\gamma$ , an effect also demonstrated in other cell types [125]. Proteasome inhibitors may also impair late TGF $\beta$ -mediated responses by up-regulation of transcriptional corepressors such as Ski novel gene N and cellular Ski [82].

Proteasome inhibition as a means to suppress pathological fibrogenic and proliferative responses has been proposed in various experimental studies. Proteasome inhibitors have been found to inhibit proliferation and induce apoptosis in renal interstitial fibroblasts [85], prevent development of experimental dermal fibrosis [79], attenuate diabetic nephropathy [78] and prevent hepatic fibrosis [126]. In Fig. 7 we propose a RPE-mediated fibrosis model and the signaling pathways affected by proteasome inhibition in the RPE.

## 5. Conclusion

In this study, we highlight specific fibrogenic and proliferative responses of RPE cells to proteasomal inhibition and propose mechanisms by which proteasomal inhibition may regulate TGF $\beta$  expression and signaling. Accordingly, the anti-fibrogenic properties of proteasome inhibitors may have a therapeutic role in RPE-mediated fibrosis. Further *in vivo* studies are required to elucidate the clinical value of these findings.



## Funding

This study was supported by grants of the Landelijke Stichting voor Blinden en Slechtienden (LSBS), Stichting Blindenpenning and MaculaFonds all having contributed through UitZicht (Grant UitZicht 2011-15) and Stichting Oogfonds. The publishing of this article was supported by Edward en Marianne Blaauw Fonds voor Oogheelkunde (Edward and Marianne Blaauw Fund for Ophthalmology). The funding organizations had no participation in the design or conduct of this study, data collection, management, analysis, interpretation, preparation, review or approval of this manuscript.

## Appendix A. Transparency document

Transparency document associated with this article can be found in the online version at <http://dx.doi.org/10.1016/j.bbrep.2018.01.005>

## References

- [1] C.A. Augood, J.R. Vingerling, P.T.V.M. De Jong, U. Chakravarthy, J. Seland, G. Soubrane, L. Tomazzoli, F. Topouzis, G. Bentham, M. Rahu, J. Vioque, I.S. Young, A.E. Fletcher, Prevalence of age-related maculopathy in older Europeans - the European Eye Study (EUREYE), *Arch. Ophthalmol.* 124 (2006) 529–535.
- [2] R.F. Spaide, C.A. Curcio, Drusen characterization with multimodal imaging, *Retina* 30 (2010) 1441–1454.
- [3] J.W. Crabb, M. Miyagi, X.R. Gu, K. Shadrach, K.A. West, H. Sakaguchi, M. Kamei, A. Hasan, L. Yan, M.E. Rayborn, R.G. Salomon, J.G. Hollyfield, Drusen proteome analysis: an approach to the etiology of age-related macular degeneration, *Proc. Natl. Acad. Sci. USA* 99 (2002) 14682–14687.
- [4] N. Nagai, A. Klimava, W.H. Lee, K. Izumi-Nagai, J.T. Handa, CTGF is increased in basal deposits and regulates matrix production through the ERK (p42/p44(mapk)) MAPK and the p38 MAPK signaling pathways, *Investig. Ophthalmol. Vis. Sci.* 50 (2009) 1903–1910.
- [5] R. Klein, C.F. Chou, B.E.K. Klein, X.Z. Zhang, S.M. Meuer, J.B. Saaddine, Prevalence of age-related macular degeneration in the US population, *Arch. Ophthalmol.* 129 (2011) 75–80.
- [6] D. Kent, C. Sheridan, Choroidal neovascularization: a wound healing perspective, *Mol. Vis.* 9 (2003) 747–755.
- [7] S.B. Bloch, H. Lund-Andersen, B. Sander, M. Larsen, Subfoveal fibrosis in eyes with neovascular age-related macular degeneration treated with intravitreal ranibizumab, *Am. J. Ophthalmol.* 156 (2013) 116–124.
- [8] W.R. Green, C. Enger, Age-related macular degeneration histopathologic studies - the 1992 Zimmerman, Lorenz, E Lecture, *Ophthalmology* 100 (1993) 1519–1535.
- [9] L.S. Lim, P. Mitchell, J.M. Seddon, F.G. Holz, T.Y. Wong, Age-related macular degeneration, *Lancet* 379 (2012) 1728–1738.
- [10] R.O. Schlingemann, Role of growth factors and the wound healing response in age-related macular degeneration, *Graef. Arch. Clin. Exp.* 242 (2004) 91–101.
- [11] F.G. Holz, S. Schmitz-Valckenberg, M. Fleckenstein, Recent developments in the treatment of age-related macular degeneration, *J. Clin. Invest.* 124 (2014) 1430–1438.
- [12] A.N. Witmer, H.G. Blaauwgeers, H.A. Weich, K. Alitalo, G.F.J.M. Vrensen, R.O. Schlingemann, Expression patterns of VEGF receptors in healthy human and monkey retina: involvement of VEGFs in physiology? *Investig. Ophthalmol. Vis. Sci.* 43 (2002) U654–U654.
- [13] A.N. Witmer, G.F.J.M. Vrensen, C.J.F. Van Noorden, R.O. Schlingemann, Vascular endothelial growth factors and angiogenesis in eye disease, *Prog. Retinal Eye Res.* 22 (2003) 1–29.
- [14] E. Daniel, C.A. Toth, J.E. Grunwald, G.J. Jaffe, D.F. Martin, S.L. Fine, J.Y. Huang, G.S. Ying, S.A. Hagstrom, K. Winter, M.G. Maguire, Risk of scar in the comparison of age-related macular degeneration treatments trials, *Ophthalmology* 121 (2014) 656–666.
- [15] T.A. Wynn, Common and unique mechanisms regulate fibrosis in various fibroproliferative diseases, *J. Clin. Invest.* 117 (2007) 524–529.
- [16] K. Ishikawa, R. Kannan, D.R. Hinton, Molecular mechanisms of subretinal fibrosis in age-related macular degeneration, *Exp. Eye Res.* (2015).
- [17] N.S. Greaves, K.J. Ashcroft, M. Baguneid, A. Bayat, Current understanding of molecular and cellular mechanisms in fibroplasia and angiogenesis during acute wound healing, *J. Dermatol. Sci.* 72 (2013) 206–217.
- [18] F. Verrecchia, A. Mauviel, Transforming growth factor-beta signaling through the Smad pathway: role in extracellular matrix gene expression and regulation, *J. Invest. Dermatol.* 118 (2002) 211–215.
- [19] Y.G. Shi, J. Massague, Mechanisms of TGF-beta signaling from cell membrane to the nucleus, *Cell* 113 (2003) 685–700.
- [20] A. Leask, D.J. Abraham, The role of connective tissue growth factor, a multi-functional matricellular protein, in fibroblast biology, *Biochem. Cell Biol.* 81 (2003) 355–363.
- [21] I. Klaassen, R.J. van Geest, E.J. Kuiper, C.J.F. van Noorden, R.O. Schlingemann, The role of CTGF in diabetic retinopathy, *Exp. Eye Res.* 133 (2015) 37–48.
- [22] S.A. Black, P.C. Trackman, Transforming growth factor-beta 1 (TGF beta 1) stimulates connective tissue growth factor (CCN2/CTGF) expression in human gingival fibroblasts through a RhoA-independent, Rac1/Cdc42-dependent mechanism - statins with forskolin block TGF beta 1-induced CCN2/CTGF expression, *J. Biol. Chem.* 283 (2008) 10835–10847.
- [23] C. Wunsche, A. Koch, R. Goldschmeding, S. Schwalm, D.M. zu Heringdorf, A. Huwiler, J. Pfeilschifter, Transforming growth factor beta 2 (TGF-beta(2))-induced connective tissue growth factor (CTGF) expression requires sphingosine 1-phosphate receptor 5 (S1P(5)) in human mesangial cells, *BBA-Mol. Cell Biol. Lipid* 1851 (2015) 519–526.
- [24] Y. Liu, H. Liu, C. Meyer, J. Li, S. Nadalin, A. Konigsrainer, H.L. Weng, S. Dooley, P. ten Dijke, Transforming growth factor-beta (TGF-beta)-mediated connective tissue growth factor (CTGF) expression in hepatic stellate cells requires stat3 signaling activation, *J. Biol. Chem.* 288 (2013) 30708–30719.
- [25] T. Nagaraja, L. Chen, A. Balasubramanian, J.E. Groopman, K. Ghoshal, S.T. Jacob, A. Leask, D.R. Brigstock, A.R. Anand, R.K. Ganju, Activation of the connective tissue growth factor (CTGF)-transforming growth factor beta 1 (TGF-beta 1) axis in hepatitis C virus-expressing hepatocytes, *PLoS One* 7 (2012) e46526.
- [26] M.M. Chen, A. Lam, J.A. Abraham, G.F. Schreiner, A.H. Joly, CTGF expression is induced by TGF-beta in cardiac fibroblasts and cardiac myocytes: a potential role in heart fibrosis, *J. Mol. Cell. Cardiol.* 32 (2000) 1805–1819.
- [27] C. Parada, J.Y. Li, J. Iwata, A. Suzuki, Y. Chai, CTGF mediates smad-dependent transforming growth factor beta signaling to regulate mesenchymal cell proliferation during palate development, *Mol. Cell. Biol.* 33 (2013) 3482–3493.
- [28] O.A. Gressner, A.M. Gressner, Connective tissue growth factor: a fibrogenic master switch in fibrotic liver diseases, *Liver Int.* 28 (2008) 1065–1079.
- [29] J.A. Arnott, E. Nuglozeh, M. Rico, T.A. Owen, F.F. Safadi, S. Popoff, Connective tissue growth factor (CTGF) is a downstream mediator of transforming growth factor-beta 1 (TGF-beta 1) in osteoblasts, *Mol. Biol. Cell* 15 (2004) 417a–417a.
- [30] S.Y. Ren, A. Babelova, K. Moreth, C.Y. Xin, W. Eberhardt, A. Doller, H. Pavenstadt, L. Schaefer, J. Pfeilschifter, A. Huwiler, Transforming growth factor-beta 2 upregulates sphingosine kinase-1 activity, which in turn attenuates the fibrotic response to TGF-beta 2 by impeding CTGF expression, *Kidney Int.* 76 (2009) 857–867.
- [31] S. Wu, M. Duncan, G. Grotendorst, E. Bancalari, Connective tissue growth factor (CTGF) mediates transforming growth factor-beta (TGF-beta) induced inhibition of lung morphogenesis in mouse embryo, *Pediatr. Res.* 53 (2003) 29a–29a.
- [32] R.J. Van Geest, I. Klaassen, I.M.C. Vogels, C.J.F. Van Noorden, R.O. Schlingemann, Differential TGF-beta signaling in retinal vascular cells: a role in diabetic retinopathy? *Investig. Ophthalmol. Vis. Sci.* 51 (2010) 1857–1865.
- [33] E.J. Kuiper, M.D. de Smet, J.C. van Meurs, H.S. Tan, M.W.T. Tanck, N. Oliver, F.A. van Nieuwenhoven, R. Goldschmeding, R.O. Schlingemann, Association of connective tissue growth factor with fibrosis in vitreoretinal disorders in the human eye, *Arch. Ophthalmol.* 124 (2006) 1457–1462.
- [34] E.J. Kuiper, M.D. de Smet, J.C. van Meurs, S. Tan, F.A. van Nieuwenhoven, N. Oliver, L. van Wieten, R. Goldschmeding, R.O. Schlingemann, Connective Tissue Growth Factor in vitreous correlates with fibrosis in vitreoretinal disorders in the human eye, *Investig. Ophthalmol. Vis. Sci.* 45 (2004) U375–U375.
- [35] E.J. Kuiper, F.A. Van Nieuwenhoven, M.D. de Smet, J.C. van Meurs, M.W. Tanck, N. Oliver, I. Klaassen, C.J.F. Van Noorden, R. Goldschmeding, R.O. Schlingemann, The angio-fibrotic switch of VEGF and CTGF in proliferative diabetic retinopathy, *PLoS One* 3 (2008).
- [36] E.J. Kuiper, R. van Zijderveld, P. Roestenberg, K.M. Lyons, R. Goldschmeding, I. Klaassen, C.J.F. Van Noorden, R.O. Schlingemann, Connective tissue growth factor is necessary for retinal capillary basal lamina thickening in diabetic mice, *J. Histochem. Cytochem.* 56 (2008) 785–792.
- [37] E.J. Kuiper, A.N. Witmer, I. Klaassen, N. Oliver, R. Goldschmeding, R.O. Schlingemann, Differential expression of connective tissue growth factor in microglia and pericytes in the human diabetic retina, *Br. J. Ophthalmol.* 88 (2004) 1082–1087.
- [38] R.J. Van Geest, I. Klaassen, S.Y. Lesnik-Oberstein, H.S. Tan, M. Mura, R. Goldschmeding, C.J.F. Van Noorden, R.O. Schlingemann, Vitreous TIMP-1 levels associate with neovascularization and TGF-beta 2 levels but not with fibrosis in the clinical course of proliferative diabetic retinopathy, *J. Cell Commun. Signal.* 7 (2013) 1–9.
- [39] R.J. Van Geest, J.W. Leeuwis, A. Dendooven, F. Pfister, K. Bosch, K.A. Hoebe, I.M.C. Vogels, D.M. Van der Giezen, N. Dietrich, H.P. Hammes, R. Goldschmeding, I. Klaassen, C.J.F. Van Noorden, R.O. Schlingemann, Connective tissue growth factor is involved in structural retinal vascular changes in long-term experimental diabetes, *J. Histochem. Cytochem.* 62 (2014) 109–118.
- [40] R.J. Van Geest, S.Y. Lesnik-Oberstein, H.S. Tan, M. Mura, R. Goldschmeding, C.J.F. Van Noorden, I. Klaassen, R.O. Schlingemann, A shift in the balance of vascular endothelial growth factor and connective tissue growth factor by bevacizumab causes the angiofibrotic switch in proliferative diabetic retinopathy, *Br. J. Ophthalmol.* 96 (2012) 587–590.
- [41] O. Coux, K. Tanaka, A.L. Goldberg, Structure and functions of the 20S and 26S proteasomes, *Annu. Rev. Biochem.* 65 (1996) 801–847.
- [42] F. Carvalhal Marques, Y. Volovik, E. Cohen, The roles of cellular and organismal aging in the development of late-onset maladies, *Annu. Rev. Pathol.* 10 (2015) 1–23.
- [43] Q.X. Ding, S. Martin, E. Dimayuga, A.J. Bruce-Keller, J.N. Keller, LMP2 knock-out mice have reduced proteasome activities and increased levels of oxidatively damaged proteins, *Antioxid. Redox Signal.* 8 (2006) 130–135.
- [44] C.M. Ethen, S.A. Hussong, C. Reilly, X. Feng, T.W. Olsen, D.A. Ferrington, Transformation of the proteasome with age-related macular degeneration, *FEBS Lett.* 581 (2007) 885–890.
- [45] A.F. Fernandes, W.M. Guo, X.Y. Zhang, M. Gallagher, M. Ivan, A. Taylor,



- P. Pereira, F. Shang, Proteasome-dependent regulation of signal transduction in retinal pigment epithelial cells, *Exp. Eye Res.* 83 (2006) 1472–1481.
- [46] A.F. Fernandes, J.L. Zhou, X.Y. Zhang, Q.N. Bian, J. Sparrow, A. Taylor, P. Pereira, F. Shang, Oxidative inactivation of the proteasome in retinal pigment epithelial cells - a potential link between oxidative stress and up-regulation of interleukin-8, *J. Biol. Chem.* 283 (2008) 20745–20753.
- [47] D.A. Ferrington, S.A. Hussong, H. Roehrich, R.J. Kapphahn, S.M. Kavanaugh, N.D. Heuss, D.S. Gregerson, Immunoproteasome responds to injury in the retina and brain, *J. Neurochem.* 106 (2008) 158–169.
- [48] S.A. Hussong, H. Roehrich, R.J. Kapphahn, M. Maldonado, M.T. Pardue, D.A. Ferrington, A novel role for the immunoproteasome in retinal function, *Investig. Ophthalmol. Vis. Sci.* 52 (2011) 714–723.
- [49] R.J. Kapphahn, E.J. Bigelow, D.A. Ferrington, Age-dependent inhibition of proteasome chymotrypsin-like activity in the retina, *Exp. Eye Res.* 84 (2007) 646–654.
- [50] Y. Li, Y.S. Wang, X.F. Shen, Y.N. Hui, J. Han, W. Zhao, J. Zhu, Alterations of activity and intracellular distribution of the 20S proteasome in ageing retinal pigment epithelial cells, *Exp. Gerontol.* 43 (2008) 1114–1122.
- [51] J.L. Louie, R.J. Kapphahn, D.A. Ferrington, Proteasome function and protein oxidation in the aged retina, *Exp. Eye Res.* 75 (2002) 271–284.
- [52] J.E.R. de Carvalho, I. Klaassen, I.M.C. Vogels, S. Schipper-Krom, C.J.F. Van Noorden, E. Reits, T.G.M.F. Gorgels, A.A.B. Bergen, R.O. Schlingemann, Complement factor C3a alters proteasome function in human RPE cells and in an animal model of age-related RPE degeneration, *Investig. Ophthalmol. Vis. Sci.* 54 (2013) 6489–6501.
- [53] J. Viiri, M. Amadio, N. Marchesi, J.M. Hyttinen, N. Kivinen, R. Sironen, K. Rilla, S. Akhtar, A. Provenzano, V.G. D'Agostino, S. Govoni, A. Pascale, H. Agostini, G. Petrovski, A. Salminen, K. Kaarniranta, Autophagy activation clears ELAVL1/HuR-mediated accumulation of SQSTM1/p62 during proteasomal inhibition in human retinal pigment epithelial cells, *PLoS One* 8 (2013) e69563.
- [54] E.A.J. Reits, A.M. Benham, B. Plougastel, J. Neefjes, J. Trowsdale, Dynamics of proteasome distribution in living cells, *EMBO J.* 16 (1997) 6087–6094.
- [55] B. Dahlmann, T. Ruppert, L. Kuehn, S. Merforth, P.M. Klotzel, Different proteasome subtypes in a single tissue exhibit different enzymatic properties, *J. Mol. Biol.* 303 (2000) 643–653.
- [56] N. Klare, M. Seeger, K. Janek, P.R. Jungblut, B. Dahlmann, Intermediate-type 20S proteasomes in HeLa cells: "asymmetric" subunit composition, diversity and adaptation, *J. Mol. Biol.* 373 (2007) 1–10.
- [57] N.J. Schuld, S.A. Hussong, R.J. Kapphahn, U. Lehmann, H. Roehrich, A.A. Rageh, N.D. Heuss, W. Bratten, D.S. Gregerson, D.A. Ferrington, Immunoproteasome deficiency protects in the retina after optic nerve crush, *PLoS One* 10 (2015) e0126768.
- [58] S.A. Hussong, R.J. Kapphahn, S.L. Phillips, M. Maldonado, D.A. Ferrington, Immunoproteasome deficiency alters retinal proteasome's response to stress, *J. Neurochem.* 113 (2010) 1481–1490.
- [59] D.S. Gregerson, K.L. Lew, S.W. McPherson, N.D. Heuss, D.A. Ferrington, RPE cells resist bystander killing by CTLs, but are highly susceptible to antigen-dependent CTL killing, *Investig. Ophthalmol. Vis. Sci.* 47 (2006) 5385–5394.
- [60] M. Verdoes, B.I. Florea, V. Menendez-Benito, C.J. Maynard, M.D. Witte, W.A. Van der Linden, A.M.C.H. Van den Nieuwendijk, T. Hofmann, C.R. Berkens, F.W.B. van Leeuwen, T.A. Groothuis, M.A. Leeuwenburgh, H. Ovaa, J.J. Neefjes, D.V. Filippov, G.A. Van der Marel, N.P. Dantuma, H.S. Overkleeft, A fluorescent broad-spectrum proteasome inhibitor for labeling proteasomes in vitro and in vivo, *Chem. Biol.* 13 (2006) 1217–1226.
- [61] I. Klaassen, J.M. Hughes, I.M.C. Vogels, C.G. Schalkwijk, C.J.F. Van Noorden, R.O. Schlingemann, Altered expression of genes related to blood-retina barrier disruption in streptozotocin-induced diabetes, *Exp. Eye Res.* 89 (2009) 4–15.
- [62] L.H. Meng, R. Mohan, B.H.B. Kwok, M. Elofsson, N. Sin, C.M. Crews, Epoxomicin, a potent and selective proteasome inhibitor, exhibits in vivo antiinflammatory activity, *Proc. Nat. Acad. Sci. USA* 96 (1999) 10403–10408.
- [63] A. Bagheri, Z.S. Soheili, H. Ahmadiani, S. Samiei, N. Sheibani, S.D. Astanah, M.R. Kanavi, A. Mohammadiani, Simultaneous application of bevacizumab and anti-CTGF antibody effectively suppresses proangiogenic and profibrotic factors in human RPE cells, *Mol. Vis.* 21 (2015) 378–390.
- [64] C.L. Chen, C.M. Liang, Y.H. Chen, M.C. Tai, D.W. Lu, J.T. Chen, Bevacizumab modulates epithelial-to-mesenchymal transition in the retinal pigment epithelial cells via connective tissue growth factor up-regulation, *Acta Ophthalmol.* 90 (2012) e389–e398.
- [65] M.R. Duncan, K.S. Frazier, S. Abramson, S. Williams, H. Klapper, X.F. Huang, G.R. Grotendorst, Connective tissue growth factor mediates transforming growth factor beta-induced collagen synthesis: down-regulation by cAMP, *FASEB J.* 13 (1999) 1774–1786.
- [66] C.M. Guo, Y.S. Wang, D. Hu, Q.H. Han, J.B. Wang, X. Hou, Y.N. Hui, Modulation of migration and Ca<sup>2+</sup> signaling in retinal pigment epithelium cells by recombinant human CTGF, *Curr. Eye Res.* 34 (2009) 852–862.
- [67] S. He, M.L. Jin, V. Worpel, D.R. Hinton, A role for connective tissue growth factor in the pathogenesis of choroidal neovascularization, *Arch. Ophthalmol.* 121 (2003) 1283–1288.
- [68] D.R. Hinton, S. He, M.L. Jin, E. Barron, S.J. Ryan, Novel growth factors involved in the pathogenesis of proliferative vitreoretinopathy, *Eye* 16 (2002) 422–428.
- [69] R. Khankan, N. Oliver, S.K. He, S.J. Ryan, D.R. Hinton, Regulation of fibronectin-EDA through CTGF domain-specific interactions with TGF beta 2 and its receptor TGF beta RII, *Investig. Ophthalmol. Vis. Sci.* 52 (2011) 5068–5078.
- [70] T. Kita, Y. Hata, K. Kano, M. Miura, S. Nakao, Y. Noda, H. Shimokawa, T. Ishibashi, Transforming growth factor-beta 2 and connective tissue growth factor in proliferative vitreoretinal diseases - possible involvement of hyalocytes and therapeutic potential of Rho kinase inhibitor, *Diabetes* 56 (2007) 231–238.
- [71] A. Leask, D.J. Abraham, TGF-beta signaling and the fibrotic response, *FASEB J.* 18 (2004) 816–827.
- [72] M.J. Radeke, C.M. Radeke, Y.H. Shih, J. Hu, D. Bok, L.V. Johnson, P.J. Coffey, Restoration of mesenchymal retinal pigmented epithelial cells by TGF beta pathway inhibitors: implications for age-related macular degeneration, *Genome Med.* 7 (2015).
- [73] P. Roestenberg, F.A. van Nieuwenhoven, L. Wieten, P. Boer, T. Diekmann, A.M. Tiller, W.M. Wiersinga, N. Oliver, W. Usinger, S. Weitz, R.O. Schlingemann, R. Goldschmeding, Connective tissue growth factor is increased in plasma of type 1 diabetic patients with nephropathy, *Diabetes Care* 27 (2004) 1164–1170.
- [74] D. Watanabe, H. Takagi, K. Suzuma, H. Oh, H. Ohashi, Y. Honda, Expression of connective tissue growth factor and its potential role in choroidal neovascularization, *Retina* 25 (2005) 911–918.
- [75] N. Awasthi, S.T. Wang-Su, B.J. Wagner, Downregulation of MMP-2 and -9 by proteasome inhibition: a possible mechanism to decrease LEC migration and prevent posterior capsular opacification, *Investig. Ophthalmol. Vis. Sci.* 49 (2008) 1998–2003.
- [76] S. Fineschi, W. Reith, P.A. Guerne, J.M. Dayer, C. Chizzolini, Proteasome blockade exerts an antifibrotic activity by coordinately down-regulating type I collagen and tissue inhibitor of metalloproteinase-1 and up-regulating metalloproteinase-1 production in human dermal fibroblasts, *FASEB J.* 20 (2006) 562–564.
- [77] C. Gruendler, Y. Lin, J. Farley, T.W. Wang, Proteasomal degradation of Smad1 induced by bone morphogenetic proteins, *J. Biol. Chem.* 276 (2001) 46533–46543.
- [78] W. Huang, C. Yang, Q.L. Nan, C.L. Gao, H. Feng, F. Gou, G. Chen, Z.H. Zhang, P.J. Yan, J. Peng, Y. Xu, The proteasome inhibitor, MG132, attenuates diabetic nephropathy by inhibiting SnoN degradation in vivo and in vitro, *Biomed. Res. Int.* (2014).
- [79] S.S. Koca, M. Ozgen, F. Dagli, M. Tuzcu, I.H. Ozerkan, K. Sahin, A. Isik, Proteasome inhibition prevents development of experimental dermal fibrosis, *Inflammation* 35 (2012) 810–817.
- [80] G.M. Mutlu, G.R.S. Budinger, M.H. Wu, A.P. Lam, A. Zirk, S. Rivera, D. Urich, S.E. Chiarella, L.H.T. Go, A.K. Ghosh, M. Selman, A. Pardo, J. Varga, D.W. Kamp, N.S. Chandel, J.I. Sznajder, M. Jain, Proteasomal inhibition after injury prevents fibrosis by modulating TGF-beta(1) signalling, *Thorax* 67 (2012) 139–146.
- [81] T. Sakairi, K. Hiromura, S. Takahashi, H. Hamatani, S. Takeuchi, M. Tomioka, A. Maeshima, T. Kuroiwa, Y. Nojima, Effects of proteasome inhibitors on rat renal fibrosis in vitro and in vivo, *Nephrology* 16 (2011) 76–86.
- [82] C.H. Weiss, G.R. Budinger, G.M. Mutlu, M. Jain, Proteasomal regulation of pulmonary fibrosis, *Proc. Am. Thorac. Soc.* 7 (2010) 77–83.
- [83] F. Zhang, M. Laiho, On and off: proteasome and TGF-beta signaling, *Exp. Cell Res.* 291 (2003) 275–281.
- [84] F. Zhang, M. Monkkonen, S. Roth, M. Laiho, Proteasomal activity modulates TGF-beta signaling in a gene-specific manner, *FEBS Lett.* 527 (2002) 58–62.
- [85] B.B. Zhu, Y.M. Jin, L. Han, H. Chen, F. Zhong, W.M. Wang, N. Chen, Proteasome inhibitor YM150 promotes proliferation and induces apoptosis in renal interstitial fibroblasts, *Pharmacol. Rep.* 65 (2013) 1357–1365.
- [86] O. Strauss, The retinal pigment epithelium in visual function, *Physiol. Rev.* 85 (2005) 845–881.
- [87] S. Grisanti, C. Guidry, Transdifferentiation of retinal-pigment epithelial-cells from epithelial to mesenchymal phenotype, *Investig. Ophthalmol. Vis. Sci.* 36 (1995) 391–405.
- [88] S. Tamiya, L.H. Liu, H.J. Kaplan, Epithelial-mesenchymal transition and proliferation of retinal pigment epithelial cells initiated upon loss of cell-cell contact, *Investig. Ophthalmol. Vis. Sci.* 51 (2010) 2755–2763.
- [89] L.V. Delpriori, B.M. Glaser, H.A. Quigley, M.E. Dorman, W.R. Green, Morphology of pig retinal-pigment epithelium maintained in organ-culture, *Arch. Ophthalmol.* 106 (1988) 1286–1290.
- [90] M. Kamei, J.M. Lewis, A. Hayashi, K. Sakagami, M. Ohji, Y. Tano, A new wound healing model of retinal pigment epithelial cells in sheet culture, *Curr. Eye Res.* 15 (1996) 714–718.
- [91] M. Friedlander, Fibrosis and diseases of the eye, *J. Clin. Invest.* 117 (2007) 576–586.
- [92] S.L. Friedman, D. Sheppard, J.S. Duffield, S. Violette, Therapy for fibrotic diseases: nearing the starting line, *Sci. Transl. Med.* 5 (2013).
- [93] D.A. Ferrington, A.D. Husom, L.V. Thompson, Altered proteasome structure, function, and oxidation in aged muscle, *FASEB J.* 19 (2005) 664.
- [94] J. Reidlinger, A.M. Pike, P.J. Savory, R.Z. Murray, A.J. Rivett, Catalytic properties of 26 S and 20 S proteasomes and radiolabeling of MB1, LMP7, and C7 subunits associated with trypsin-like and chymotrypsin-like activities, *J. Biol. Chem.* 272 (1997) 24899–24905.
- [95] W. Dubiel, K. Pratt, K. Ferrell, M. Rechsteiner, Purification of An 11-S regulator of the multicatalytic protease, *J. Biol. Chem.* 267 (1992) 22369–22377.
- [96] D.A. Ferrington, D.S. Gregerson, Immunoproteasomes: structure, function, and antigen presentation, *Prog. Mol. Biol. Transl. Sci.* 109 (2012) 75–112.
- [97] A.L. Goldberg, P. Cascio, T. Saric, K.L. Rock, The importance of the proteasome and subsequent proteolytic steps in the generation of antigenic peptides, *Mo. Immunol.* 39 (2002) 147–164.
- [98] A.D. Husom, E.A. Peters, E.A. Kolling, N.A. Fugere, L.V. Thompson, D.A. Ferrington, Altered proteasome function and subunit composition in aged muscle, *Arch. Biochem. Biophys.* 421 (2004) 67–76.
- [99] R. Amin, J.E. Puklin, R.N. Frank, Growth-factor localization in choroidal neovascular membranes of age-related macular degeneration, *Investig. Ophthalmol. Vis. Sci.* 35 (1994) 3178–3188.
- [100] H. Tanihara, M. Yoshida, M. Matsumoto, N. Yoshimura, Identification of

- transforming growth-factor-beta expressed in cultured human retinal-pigment epithelial-cells, *Investig. Ophthalmol. Vis. Sci.* 34 (1993) 413–419.
- [101] A.M. Newman, N.B. Gallo, L.S. Hancox, N.J. Miller, C.M. Radeke, M.A. Maloney, J.B. Cooper, G.S. Hageman, D.H. Anderson, L.V. Johnson, M.J. Radeke, Systems-level analysis of age-related macular degeneration reveals global biomarkers and phenotype-specific functional networks, *Genome Med.* 4 (2012).
- [102] T.B. Connor, A.B. Roberts, M.B. Sporn, D. Danielpour, L.L. Dart, R.G. Michels, S. Debustros, C. Enger, H. Kato, M. Lansing, H. Hayashi, B.M. Glaser, Correlation of fibrosis and transforming growth factor-beta type-2 levels in the eye, *J. Clin. Investig.* 83 (1989) 1661–1666.
- [103] E. Vanobberghenschilling, N.S. Roche, K.C. Flanders, M.B. Sporn, A.B. Roberts, Transforming growth factor-beta-1 positively regulates its own expression in normal and transformed-cells, *J. Biol. Chem.* 263 (1988) 7741–7746.
- [104] A. Holmes, D.J. Abraham, S. Sa, S.W. Xu, C.M. Black, A. Leask, CTGF and SMADs, maintenance of scleroderma phenotype is independent of SMAD signaling, *J. Biol. Chem.* 276 (2001) 10594–10601.
- [105] Y.J. Bai, S.T. Liang, W.Z. Yu, M. Zhao, L.Z. Huang, M.W. Zhao, X.X. Li, Semaphorin 3A blocks the formation of pathologic choroidal neovascularization induced by transforming growth factor beta, *Mol. Vis.* 20 (2014) 1258–1270.
- [106] S. Saika, TGF beta pathobiology in the eye, *Lab. Investig.* 86 (2006) 106–115.
- [107] S.K. Leivonen, L. Hakkinen, D. Liu, V.M. Kahari, Smad3 and extracellular signal-regulated kinase 1/2 coordinately mediate transforming growth factor-beta-induced expression of connective tissue growth factor in human fibroblasts, *J. Investig. Dermatol.* 124 (2005) 1162–1169.
- [108] A.F. Kisselev, A.L. Goldberg, Proteasome inhibitors: from research tools to drug candidates, *Chem. Biol.* 8 (2001) 739–758.
- [109] P.A. Campochiaro, B.M. Glaser, Platelet-derived growth-factor is chemotactic for human retinal-pigment epithelial-cells, *Arch. Ophthalmol.* 103 (1985) 576–579.
- [110] T. Vartio, L. Laitinen, O. Narvanen, M. Cutolo, L.E. Thornell, L. Zardi, I. Virtanen, Differential expression of the Ed sequence-containing form of cellular fibronectin in embryonic and adult human-tissues, *J. Cell Sci.* 88 (1987) 419–430.
- [111] W. Eichler, U. Friedrichs, A. Thies, C. Tratz, P. Wiedemann, Modulation of matrix metalloproteinase and TIMP-1 expression by cytokines in human RPE cells, *Investig. Ophthalmol. Vis. Sci.* 43 (2002) 2767–2773.
- [112] E.K. An, X.N. Lu, J. Flippin, J.M. Devaney, B. Halligan, E. Hoffman, K. Csaky, Y. Hathout, Secreted proteome profiling in human RPE cell cultures derived from donors with age related macular degeneration and age matched healthy donors (vol 5, pg 2599, 2006), *J. Proteome Res.* 6 (2007) 1615–1615.
- [113] J.P. Alexander, J.M.B. Bradley, J.D. Gabourel, T.S. Acott, Expression of matrix metalloproteinases and inhibitor by human retinal-pigment epithelium, *Investig. Ophthalmol. Vis. Sci.* 31 (1990) 2520–2528.
- [114] K. Brew, D. Dinakarpanian, H. Nagase, Tissue inhibitors of metalloproteinases: evolution, structure and function, *Biochim. Biophys. Acta* 1477 (2000) 267–283.
- [115] J.P. Iredale, Tissue inhibitors of metalloproteinases in liver fibrosis, *Int. J. Biochem. Cell Biol.* 29 (1997) 43–54.
- [116] M.A. Moses, J. Sudhalter, R. Langer, Identification of an inhibitor of neovascularization from cartilage, *Science* 248 (1990) 1408–1410.
- [117] E.D. Rosen, B.M. Spiegelman, PPARgamma: a nuclear regulator of metabolism, differentiation, and cell growth, *J. Biol. Chem.* 276 (2001) 37731–37734.
- [118] P. Tontonoz, E. Hu, R.A. Graves, A.I. Budavari, B.M. Spiegelman, mPPAR gamma 2: tissue-specific regulator of an adipocyte enhancer, *Genes Dev.* 8 (1994) 1224–1234.
- [119] S. Zhang, H. Gu, N. Hu, Role of peroxisome proliferator-activated receptor gamma in ocular diseases, *J. Ophthalmol.* 2015 (2015) 275435.
- [120] T. Murata, S. He, M. Hangai, T. Ishibashi, X.P. Xi, S. Kim, W.A. Hsueh, S.J. Ryan, R.E. Law, D.R. Hinton, Peroxisome proliferator-activated receptor-gamma ligands inhibit choroidal neovascularization, *Investig. Ophthalmol. Vis. Sci.* 41 (2000) 2309–2317.
- [121] A.V. Ershov, N.G. Bazan, Photoreceptor phagocytosis selectively activates PPARgamma expression in retinal pigment epithelial cells, *J. Neurosci. Res.* 60 (2000) 328–337.
- [122] J. Zhu, J. Zhang, M. Ji, H. Gu, Y. Xu, C. Chen, N. Hu, The role of peroxisome proliferator-activated receptor and effects of its agonist, pioglitazone, on a rat model of optic nerve crush: PPARgamma in retinal neuroprotection, *PLoS One* 8 (2013) e68935.
- [123] S. Qin, A.P. McLaughlin, G.W. De Vries, Protection of RPE cells from oxidative injury by 15-deoxy-delta12,14-prostaglandin J2 by augmenting GSH and activating MAPK, *Investig. Ophthalmol. Vis. Sci.* 47 (2006) 5098–5105.
- [124] S.F. Qin, Y.M. Lu, G.A. Rodrigues, Resveratrol protects RPE cells from sodium iodate by modulating PPAR alpha and PPAR delta, *Exp. Eye Res.* 118 (2014) 100–108.
- [125] J.E. Milam, V.G. Keshamouni, S.H. Phan, B. Hu, S.R. Gangireddy, C.M. Hogaboam, T.J. Standiford, V.J. Thannickal, R.C. Reddy, PPAR-gamma agonists inhibit profibrotic phenotypes in human lung fibroblasts and bleomycin-induced pulmonary fibrosis, *Am. J. Physiol. Lung Cell Mol. Physiol.* 294 (2008) L891–L901.
- [126] H. Chen, Y.W. He, W.Q. Liu, J.H. Zhang, Rosiglitazone prevents murine hepatic fibrosis induced by *Schistosoma japonicum*, *World J. Gastroenterol.* 14 (2008) 2905–2911.