



# Function of lipid binding proteins of parasitic helminths: still a long road

Jose F. Lombardo<sup>1</sup> · Jorge L. Pórfido<sup>2</sup> · Martín S. Sisti<sup>1</sup> · A. Nahili Giorello<sup>1</sup> · Santiago Rodríguez<sup>1</sup> · Betina Córscico<sup>1</sup> · Gisela R. Franchini<sup>1,3</sup>

Received: 2 August 2021 / Accepted: 7 February 2022

© The Author(s), under exclusive licence to Springer-Verlag GmbH Germany, part of Springer Nature 2022

## Abstract

Infections with parasitic helminths cause severe debilitating and sometimes lethal diseases in humans and domestic animals on a global scale. Unable to synthesize de novo their own fatty acids and sterols, helminth parasites (nematodes, trematodes, cestodes) rely on their hosts for their supply. These organisms produce and secrete a wide range of lipid binding proteins that are, in most cases, structurally different from the ones found in their hosts, placing them as possible novel therapeutic targets. In this sense, a lot of effort has been made towards the structure determination of these proteins, but their precise function is still unknown. In this review, we aim to present the current knowledge on the functions of LBPs present in parasitic helminths as well as novel members of this highly heterogeneous group of proteins.

**Keywords** Lipid binding proteins · FABPs · Nematodes · Platyhelminthes · FAR · NPAs

## Introduction

Knowing the biochemistry of parasitic helminths in detail may reveal differential enzymes or metabolic pathways that could represent an Achilles' heel for these organisms. Those proteins or enzymes could be targets for novel therapeutic strategies as well as molecular markers to be used in diagnostics. However, it is a really hard task to perform studies on the biochemistry of parasitic helminths, and many times it is almost impossible. This difficulty is based on the fact that these experiments may require keeping the complete life cycle (or at least part of it) under laboratory conditions, which can be complex and expensive, including welfare

protocols and keeping animals as alternative hosts. Although this has been achieved in some cases, e.g., for *Echinococcus multilocularis* (Spiliotis and Brehm 2009) and the hookworm *Necator americanus* (Jian et al. 2003), this is still not possible for many parasitic species. Particularly, those parasites affecting wild fauna where the collection of live samples depends on roadkill findings according to different country regulations. However, this is not the case for plant parasites where different biological approaches have been accomplished, and there is enough information to look into the possible biological function of different types of proteins (see below). In this regard, an impressive advance on helminthology has been achieved based on the massive data retrieved by whole genomic sequencing and genome-wide expression profiling methods at the level of the transcriptome and proteomes. All this data provides highly valuable information platforms to seek for alternative biochemical pathways, variations in certain canonical routes, different expression patterns, specific proteins, etc.

In particular, parasitic helminths present a restricted lipid metabolism; this means that many biochemical pathways are dampened or completely absent in different members of this group. The parasitic helminths considered in this review include species of the phylum Platyhelminth, commonly known as “flatworms” and phylum Nematoda or “roundworms”. Particularly, neither flatworms nor roundworms

---

Guest Editor: Anja Taubert

---

✉ Gisela R. Franchini  
gfranchini@conicet.gov.ar

<sup>1</sup> Instituto de Investigaciones Bioquímicas de La Plata (INIBIOLP), Facultad de Ciencias Médicas, Universidad Nacional de La Plata (UNLP)-Consejo Nacional de Investigaciones Científicas Y Técnicas (CONICET), La Plata, Argentina

<sup>2</sup> Institut Pasteur de Montevideo, Montevideo, Uruguay

<sup>3</sup> Departamento de Ciencias Biológicas, Facultad de Ciencias Exactas, UNLP, La Plata, Argentina

can synthesize fatty acid *de novo* (Barrett 1981). In some parasitic nematodes, the enzymes are present but with really low activity so no significant product is obtained (Barrett 2009). Nevertheless, incorporated fatty acids are involved in the synthesis of complex lipids such as triacylglycerol and phospholipids, suggesting their importance in maintaining different cellular structures. Although a partial mevalonate pathway is present for the synthesis of dolichol, quinones' side chains, and isoprenoids for protein prenylation (Hiepe et al. 2006), the complete *de novo* synthesis pathway of cholesterol is absent in both phyla (Barrett 1981; Frayha and Smyth 1983). In this sense, the acquisition and transport of several lipid molecules are crucial to these organisms, and the proteins and receptors involved in lipid transport and exchange may provide potential targets for chemo- and immunotherapy.

Lipid binding proteins (LBPs) of parasitic helminths have been studied for decades and are considered relevant because they are usually found in the excretion/secretion (E/S) products of these organisms. Most of them are structurally different from those of their host and as such they are usually targets of a strong immune response (Kennedy et al. 1990; Tomlinson et al. 1989). Hence, a lot of effort has been dedicated to study the structure of these proteins to improve novel drug design and the setting up of more specific diagnostic methods. Besides their possible utility, the precise biological functions of LBPs are still unknown. They have been proposed to participate in the acquisition and distribution of lipids within tissues of the parasite and/or modulation of the host's local tissue environment and its innate and acquired immune systems.

Addressing the function of these proteins is not a straightforward task, since they are not catalytic enzymes but carrier proteins. Hence, one of the few ways to assess LBPs' biological activity is to evaluate their binding capacity. A key step towards understanding the functions of these proteins might be to describe the ligands that are bound in their

natural environment, as well as studies performed on tissues from parasites or whole organisms. The most informative experiments used to unravel protein functions involve reverse genetics approaches. These would imply the knock-down (or knock out) expression of the protein of interest within the parasite. Unfortunately, this technique has been achieved successfully only in a few examples of parasitic species (Britton et al. 2016; Dutta et al. 2015; Guidi et al. 2015; Kimber et al. 2007; Lilley et al. 2012; MacDonald et al. 2014; Pérez et al. 2019). In this scenario, appropriate model organisms are needed to shed light on the functions. In this review, we highlight the improvements in the assessment of the biological functions of LBPs from parasitic helminths as well as the description of novel members of this diverse group.

## A highly heterogeneous group

Altogether, parasitic helminths produce and secrete a broad range of different LBPs. At least five different types of LBPs have been described and they are distributed in different taxa as is summarized in Table 1. Below we describe in brief each type of LBP trying to address the most significant improvements on their biological functions.

### Hydrophobic ligand binding proteins (HLBPs)

HLBPs is a family of proteins that are specific to cestodes in phylum Platyhelminthes. This unusual family is composed of intracellular and extracellular members that bind a variety of fatty acids, retinoids, and some sterols as reviewed in Alvite and Esteves (2012). In their native state, HLBPs are described as lipoprotein particles that could reach a size of about 230 kDa. This complex structure consists of multiple small protein subunits of approximately 8 kDa (although this

**Table 1** Distribution of LBPs across groups of parasitic helminths

Type of LBP	Phylum <i>Platyhelminthes</i>		Phylum <i>Nematoda</i>				
	<i>Cestodes</i>	<i>Trematodes</i>	<i>Clade I</i>	<i>Clade II</i>	<i>Clade III</i>	<i>Clade IV</i>	<i>Clade V</i>
HLBP	✓						
FABP	✓	✓	✓	✓	✓	✓	✓
nemFABP				*	✓	✓	✓
FAR				✓	✓	✓	✓
NPA				✓	✓	✓	✓
Dorylipophorin			✓				

Bibliographic references for each group are cited in each section. \*A search for nemFABPs has been done in clade II using Blast tool from WormBase Parasite (<https://parasite.wormbase.org/Tools/Blast?db=core>), and no hit was found with the presence of leader signal peptide. Nevertheless, there is only one species with a determined genome so we cannot rule out the presence of nemFABPs in the clade

might change within species) interacting with lipids (Lee et al. 2007; Silva-Álvarez et al. 2015).

One of the most studied examples is antigen B (EgAgB) from *Echinococcus granulosus* sensu lato (s. l.). EgAgB is one of the major molecules synthesized in large amounts by the cyst (metacestode larval stage), and it is present in most tissues of the parasite as well as in protoscoleces (infective stage for definitive host) (González et al. 1996; Oriol et al. 1971; Sanchez et al. 1991, 1993). From the molecular point of view, it conforms to a lipoprotein particle with a protein component consisting of 8 kDa subunits that are rich in amphipathic alpha-helices (Fernández et al. 1996; Lightowlers et al. 1989). For the case of *E. granulosus* s. l., these subunits are encoded by a multigene family, dubbed *EgAgB8/1–EgAgB8/5* (Arend et al. 2004; Chemale et al. 2001; Frosch et al. 1994; Haag et al. 2004; Kamenetzky et al. 2005; Muzulin et al. 2008; Zhang et al. 2010). The lipid moiety found to be interacting with native EgAgB showed that not only fatty acids and triacylglycerols are bound but also a wide range of other hydrophobic compounds like cholesterol esters, cholesterol, and phospholipids (Obal et al. 2012).

In vitro functional analysis of recombinant lipid-free rEgAgB8/2 and rEgAgB8/3 showed a selective capacity to bind lipids, showing affinity at least for 16- and 18-C fatty acids, but not for cholesterol, indicating that these components of the natural EgAgB lipoprotein particles would not interact directly with cholesterol (Silva-Álvarez et al. 2016). Additionally, the capacity of these subunits to transfer lipids to membranes was also assayed showing that different subunits of EgAgB8 are able to deliver their cargo to phospholipid membranes, supporting the hypothesis that EgAgB is involved in lipid transport between parasite and host tissues (Silva-Álvarez et al. 2015). However, the capacity of EgAgB particles to transfer fatty acids to the parasite or to the host's cells remains to be formally demonstrated. In this regard, assays performed with a related macromolecule found in *Taenia solium* metacestode strongly support this hypothesis (Lee et al. 2007).

Remarkably, EgAgB and particularly its predominant EgAgB8/1 apolipoprotein are potential ligands for monocyte and macrophage receptors as described in Silva-Álvarez et al. (2016). These receptors may also be involved in plasma lipoprotein recognition and induce an anti-inflammatory phenotype in macrophages upon recognition of EgAgB (Silva-Álvarez et al. 2016, 2018). Moreover, it has been proposed to be the most specific *Echinococcus* antigen for serodiagnosis (Siracusano et al. 2008).

Molecules of this type have also been described in other cestodes including *Moniezia expansa*, *Hymenolepis diminuta*, *Taenia crassiceps*, *Taenia solium*, and *Taenia hydatigena* (Barrett et al. 1997; Jia et al. 2011; Saghir et al. 2001; Sako et al. 2000; Zarlenga et al. 1994). A primary sequence comparison of HLBP found in these species


revealed that they are significantly diverse but present well-conserved molecular characteristics like two alpha-helical domains that are presumed to be involved in protein-membrane interactions (Lee et al. 2007). Notably, HLBP from *M. expansa* and *H. diminuta* do not show an N-terminal hydrophobic leader sequence indicating that they might exert their function intracellularly.

## Fatty acid binding proteins (FABPs)

FABPs are small intracellular proteins (around 15 kDa) that reversibly bind fatty acids and other hydrophobic ligands. Members of this family have been found throughout the animal kingdom, both in invertebrates and vertebrates, but no counterparts have been found in plants or fungi (Haunerland and Spener 2004). Ancestral intracellular FABP genes are supposed to have arisen after the separation of animals from fungi and plants (Schaap et al. 2002).

The first FABP from invertebrates was found in *Schistosoma gregaria* (Haunerland and Chisholm 1990), almost 20 years after the first vertebrate FABP was described. After this finding, many proteins of the family have been described in nematodes, insects, mollusks, and flatworms (Esteves and Ehrlich 2006). The first flatworm FABP to be described was Sm14 from *Schistosoma mansoni* (Moser et al. 1991), followed by FABPs of *S. japonicum*, *Fasciola hepatica*, *Fasciola gigantica*, and *E. granulosus* s.l. (Becker et al. 1994; Esteves et al. 1993; Estuningsih et al. 1997; Rodríguez-Pérez et al. 1992). As in vertebrate's FABPs, the sequence identity between different FABPs from invertebrates varies significantly, but they still share their overall 3D protein structure. It consists of a  $\beta$ -barrel, composed of ten antiparallel  $\beta$ -strands ( $\beta$ A– $\beta$ J), capped with two  $\alpha$ -helices ( $\alpha$ I and  $\alpha$ II). The barrel is solvent accessible, and the helices have been proposed as a gate that regulates the entrance or exit of ligands see Fig. 1 (Haunerland and Spener 2004; Scapin et al. 1992; Storch and Corsico 2008; Storch and McDermott 2009; Zimmerman and Veerkamp 2002).

FABPs from vertebrates are grouped into four subfamilies based on their primary sequence analysis (Marcelino et al. 2006) and each of these characteristically binds certain types of ligands. When compared to mammalian FABPs, proteins from parasitic helminths show higher similarity to those that belong to subfamily IV (Esteves and Ehrlich 2006; Marcelino et al. 2006; Smathers and Petersen 2011). Group IV is the largest subfamily and includes FABP3, FABP4, FABP5, FABP7, FABP8, FABP9, and FABP12 (Liu et al. 2008) from vertebrates and binds a variety of ligands including long-chain fatty acids, eicosanoids, and retinoids. In particular, there is a high conservation of a triad of residues that play a central role in the mechanism of binding of fatty acids to those FABPs. In relation to the genetic structure, it is worth mentioning that the position of the introns in



**Fig. 1** X-ray structure of EgFABP1 (PDB: 108 V) from *Echinococcus granulosus*. Copurifying ligand, palmitic acid, is depicted in green

invertebrate FABPs seems to be substantially conserved, although their number and length vary considerably (Esteves and Ehrlich 2006).

Recently, data mining in genomes from the parasitic plathelminthes, *Echinococcus multilocularis* and *E. granulosus s.l.*, revealed that this family of proteins is far more complex than previously reported in cestodes. Six genes with different coding sequences for FABPs were found in each organism, with the sole exception of EmFABP1.1 and EmFABP1.2, in *E. multilocularis*, that are identical to one another (Pórfido et al. 2020). All of these variants have been cloned and produced recombinantly presenting structural features (inferred from spectroscopic data as well as in silico prediction) that resemble those of previously reported FABPs from vertebrates and invertebrates (Bélgamo et al. 2020; Pórfido et al. 2020). Notably, for EmFABP4, the most striking feature is that the primary sequence is considerably longer (176 amino acids) than that for a regular FABP (around 130 amino acids). In this case, a typical FABP fold is predicted, but no specific structure is assigned to the C terminus of the protein (Pórfido et al. 2020). Surprisingly, EmFABP4 was purified

as a dimer, and this state is stable in solution (Bélgamo et al. 2020). However, whether it exists as a dimer in its biological environment remains to be elucidated. It is noticeable that a large subfamily of FABPs, including some with C-terminal extensions, has also been reported in *Fasciola* spp. (Bélgamo et al. 2020; Morphew et al. 2016). Interestingly, these longer variants are different from the so-called nemFABPs (see below) since the unusual extensions in flatworms' FABPs are exclusively on the C-terminal sequence.

Published transcriptomic information on *E. multilocularis* and *E. granulosus s.l.* suggest that all *fabp* genes are transcribed in different stages of these organisms (Huang et al. 2016; Tsai et al. 2013; Zheng et al. 2013a, b). Moreover, according to the different sets of data, *emfabp1.1* and *emfabp1.2* are the most highly expressed FABP genes in *E. multilocularis* (Pórfido et al. 2020).

It is worth mentioning that specific inhibitors of mammalian FABP4 have been developed and employed as effective therapeutic agents of diseases such as diabetes and atherosclerosis (Barf et al. 2009; Furuhashi et al. 2007; Wang et al. 2016). Based on the degree of conservation between FABPs from helminths to group IV from vertebrate FABPs, members of this family from *E. multilocularis* and *E. granulosus* are being considered as druggables (Bélgamo et al. 2020), and this could also be extended to FABPs from other helminth species as well.

As expected, FABPs are also present in both parasitic and free-living nematodes. Particularly, in the free-living species, *Caenorhabditis elegans*, there are nine isoforms of FABPs (dubbed *lbp-1* to *lbp-9*) with tissue specific expression patterns (Plenefisch et al. 2000). As previously said, it is difficult to perform reverse genetic experiments on most parasitic helminths, but in the case of nematodes, the use of *C. elegans* as a model to study protein function is an excellent option. In this sense, knockdown experiments on *lbp-5* of *C. elegans* have shown strong physiological disruptions (Xu et al. 2011). *lbp-5* knockdown causes a decrease in expression of  $\beta$ -oxidation genes, by affecting NHR-49 signaling (homolog of mammalian NHR-4 and functional homolog of PPAR- $\alpha$ ), which contributes to fat storage increase. An analysis of the localization of LBP-5 in cells from *C. elegans* revealed that it is found both in the nucleus and the cytoplasm, suggesting a possible role in transporting fatty acids into the nucleus regulating the expression of genes involved in fat metabolism (Xu et al. 2011).

### The particular case of “nemFABPs”

All the FABPs described so far in nematodes form two distant groups suggesting at least two different phylogenetic origins (Plenefisch et al. 2000; Zheng et al. 2013a, b). One of these groups includes sequences related with muscle-type intracellular FABPs from vertebrates as is the case of




LBP-5 mentioned above (Plenefisch et al. 2000). The other group includes some FABPs that present particular characteristics that are found almost exclusively in the phylum Nematoda; hence, they were dubbed “nemFABPs.” One of the most striking differences found in “nemFABPs” is the presence of a cleavable secretion signal leader peptide, and hence they are secreted from the cell, presumably by traditional secretory pathways (Mei et al. 1997). Until now, only FABPs genes from *Trichinella spiralis* are found to be the exception to the two groups mentioned before. This is an interesting finding, since *T. spiralis* is a member of Clade I of nematodes which is quite an underrepresented group and may show interesting features (Blaxter and Koutsovoulos 2015).

As in vertebrate cytosolic FABPs, “nemFABPs” present the characteristic beta barrel fold with an alpha-helical cap. The first of these to be described from nematodes were Asp-18 from the large intestinal roundworm, *Ascaris suum*, and Bm-FAB-1 from the agent of lymphatic filariasis, *Brugia malayi* (Michalski et al. 2002). These proteins are not only produced by human and animal parasites (Zheng et al. 2013a, b) but, as already mentioned, are also found in free-living forms like *C. elegans*. These proteins were found to be gender specific as well as developmentally regulated and have been proposed to be associated with reproduction since they are found in the perivitelline fluid of nematode eggs harboring the embryos (Li et al. 2004; Mei et al. 1997; Michalski et al. 2002; Michalski and Weil 1999; Plenefisch et al. 2000). One of these, Ce-LBP-1, appears only to be synthesized and secreted by developing embryos within the egg (Plenefisch et al. 2000).

Notably, their primary sequences are typically 10–19 residues longer than other FABPs, and it was proposed that some of these additional residues are accommodated in extended loops (Mei et al. 1997). The structure of Asp-18 has been solved at the atomic level by X-ray crystallography and NMR confirming the presence of two extended loops, one immediately adjacent to the ligand portal and another distantly located on the opposite side of the molecule see Fig. 2. Additionally, the triad of amino acid side chains that tether bound fatty acids is differently arranged and composed compared to what is found in mammalian FABPs (Ibáñez-Shimabukuro et al. 2019). A preliminary analysis of their ligand preferences in a biological context such as *E. coli* showed that Asp-18 binds exclusively fatty acids, particularly 18 carbons vaccenic acid with the highest affinity (Ibáñez-Shimabukuro et al. 2019).

The fact that some nemFABPs are found in eggs, more precisely in the perivitelline fluid allows us to hypothesize that they might be involved in the uptake of lipids from perivitelline fluid. Unfortunately, no further functional experiments have been performed other than in vitro assays.



**Fig. 2** X-ray structure of Asp-18 (PDB: 6I8X) from *Ascaris suum* in complex with vaccenate (depicted in light blue). Extended loops are shown in dashed circles

### Fatty acid and retinol binding proteins (FAR)

The FAR family is a novel class of proteins, also exclusively found in nematodes. They have an approximate size of 20 kDa and are able to bind both fatty acids and retinol. Genes encoding FAR proteins have been described in many nematode species, both free-living and parasitic forms (Garofalo et al. 2002; Kennedy et al. 1997) and each species produces several isoforms. FARs are present in ES products, as is the case of Ac-FAR-1 from the intestinal hookworm *Ancylostoma caninum* (Basavaraju et al. 2003; Kennedy et al. 1997). The study of FAR proteins started in 1995 with Ov-FAR-1 (Tree et al. 1995), a protein secreted by *Onchocerca volvulus*, a causative agent of river blindness. Since then, the FAR family has expanded greatly; being described not only in filarial species but also in hookworms, plant parasites, and the free-living nematode *C. elegans*, as well as other *Caenorhabditis* species. Notably, none of these proteins are found in clade I of phylum Nematoda. The knowledge we have about them nowadays seems vast but also dispersed, with many studies done in separated groups of proteins within the family.

From the molecular point of view, FARs range from 17 to 30 kDa in size and have relatively conserved glycosylation sites, but glycosylation patterns vary from one protein to another (Nirmalan et al. 1999). They also have potential sites of phosphorylation for conserved type II casein kinase (Prior et al. 2001). They are rich in alpha-helices, show high stability, and do not have structural analogues in other animal groups. To date, the crystallographic structure of FAR

proteins has been resolved for two members of the family, Ce-FAR-7 from *C. elegans* (Jordanova et al. 2009) and Na-FAR-1 from *Necator americanus* (Rey-Burusco et al. 2015) (see Fig. 3). Ce-FAR-7 presents two discrete cavities in which it could locate different types of ligands; however, based on its sequence identity, this protein might not be representative of parasite proteins. The overall structure is a flattened ellipsoid which seems to adopt different degrees of expansion depending on whether ligands are absent or bound to the central cavity. Recent studies using in silico techniques compared Na-FAR-1 with classical  $\beta$ -barrel FABPs. Particularly, the use of extended molecular dynamics simulations and principal component analysis allowed to identify different conformations adopted by each system during the simulations. Na-FAR-1 encompasses a complex highly flexible internal ligand-binding domain that allows reversible switching between distinct states in the protein. Besides, the ligand inside the cavity experiences large conformational changes between bent and stretched conformations. These changes in the ligand conformation follow changes in the cavity size dictated by the transient protein conformation. On the contrary, protein–ligand complex in  $\beta$ -barrel FABPs fluctuates around a unique conformation (Barletta et al. 2019). These results are in agreement with the observations that FAR proteins bind a broad range of hydrophobic ligands while FABPs bind almost uniquely fatty acids.


Na-FAR-1 and possibly other FARs may have a wider repertoire for hydrophobic ligand binding; this is supported by the fact that a range of neutral and polar lipids co-purify with the bacterially expressed recombinant protein (Rey-Burusco et al. 2015). Many studies show that these proteins are able to bind fatty acids (mainly oleic acid) and retinol, but recent evidence shows phospholipids as other possible ligands (Rey-Burusco et al. 2015). As observed with FABPs,

FAR proteins have a well described and conserved structure but have a wide range of sequence similarity even between isoforms. For example, Na-FAR-1 has a sequence identity that ranges from 44 to 60% with FARs from other nematodes from the *Rhabditida* (*Ancylostoma*, *Ostertagia*, and *Caenorhabditis*) class, but this identity descends to 25–38% with other classes like *Spirurida* (*Onchocerca*, *Brugia*, *Acanthocheilonema*) and *Diplogasterida* (*Globodera* and *Meloidogyne*) (Rey Burusco, 2014). For *C. elegans*, Ce-FAR-8 and Ce-FAR-7 are so divergent (19–23%) that has even led to cataloging them as outliers, disregarding their importance.

In regard to expression levels, different isoforms are found through all developmental stages, some are highly expressed during infective stages other during reproductive or prereproductive stages. An important note is that no FAR proteins are highly expressed in eggs. Moreover, in most cases, more than one FAR isoform is found at the same stage and they show a different pattern of expression, as is clear for the case of *C. elegans* (Garofalo et al. 2002). This evidence suggests that FAR proteins might fulfill different roles through development rather than having one static biological function.

FARs are the major components of human, animal, and plant parasite secretions (Basavaraju et al. 2003; Kennedy et al. 1997). As an example, we have the case of Na-FAR-1, found in ES products thanks to immunological assays with antibodies against the recombinant protein (Rey-Burusco et al. 2015). A proteogenomic analysis of ES products revealed that FAR proteins are among the top 10 most abundant protein families in the ES products of adult *N. americanus* (Logan et al. 2020). These findings have made it possible to use FAR proteins as diagnostic tools for some parasitic diseases (Burbelo et al. 2009). This raises the hypothesis that they could play roles in the interaction with

**Fig. 3** X-ray structures of **A** Na-FAR-1 (PDB: 4XCP) from *Necator americanus* and **B** Ce-FAR-7 (PDB: 2W9Y) from *Caenorhabditis elegans*. For Na-FAR-1, the ligands (palmitic acid) are depicted in blue



the host and in pathogenesis by interfering with host defense signals (Bradley et al. 2001; Hewitson et al. 2009; Iberkleid et al. 2015). In fact, recent results demonstrated that injecting recombinant FAR proteins from entomopathogenic nematode *Steinernema carpocapsae* can reduce survival of host *Drosophila melanogaster* when exposed to bacterial infections, showing that these proteins can interfere with immune response. Although further studies are needed, the mechanism seems to indeed involve sequestering lipidic signals, thus disrupting the normal functioning of host defenses (Parks et al. 2021). Apart from this, it should be noted that Ace-FAR-1 from *Ancylostoma ceylanicum* has been shown to be potentially useful as a vaccine because it is capable of conferring immunity in challenge tests in laboratory animals (Fairfax et al. 2009). On the other hand, these parasitism-focused hypotheses fall short when we look at the eight FAR proteins described for *C. elegans*, where seven of them present a signal peptide with Ce-FAR-7 lacking this sequence. It is thus important to formulate new hypotheses including the possible function of these secreted proteins in free-living worms.

### FAR proteins in plant parasites

The first example we have is Gp-FAR-1 from *Globodera pallida*. From its molecular description, analyses of its binding specificity (encompassing linoleic and linolenic acid) and localization in the hypodermis of J2 stage (invasive stage of the parasite) suggested a crucial role in countering plant defense mechanisms. Lipxygenase-mediated peroxidation of linolenic acid is an early step in the octadecanoid signaling pathway, which leads to the synthesis of the systemic plant defense signal transducer jasmonic acid. A pronounced inhibition of lipxygenase-mediated breakdown of linoleic acid was observed in the presence of recombinant Gp-FAR-1 (Prior et al. 2001).

The study of the root knot nematode, *Meloidogyne javanica*, and its FAR proteins Mj-FAR-1 and Mj-FAR-2 was the first work to place FAR proteins from plant parasites as a unique node in a phylogenetic tree showing a clear evolutionary adaptation to plant parasitism. Mj-FAR-1 accumulates in the cuticle and is secreted by hypodermis, being found in the intercellular space between the parasite and the host (Iberkleid et al. 2013, 2015). These findings so far are aiming towards an immunomodulatory function, disrupting plant defense mechanisms, and facilitating parasitism. On the other hand, Mi-FAR-1 from *Meloidogyne incognita* is expressed in J2, J3, and J4. In this last stage, it might be playing a role in the pre-reproductive stage, presenting the possible relationship with development, egg-laying, and reproduction. This was also supported by the silencing of Mi-FAR-1 with RNAi that showed a decreased number of females as well as the fertility and production of eggs when

exposed to bacteria (Phani et al. 2017). This decrease in fertility was also seen in Ab-FAR-1 from *Aphelenchoides besseyi*, “white tip nematode”, where the retinol binding capabilities of FAR proteins could be related to the needs of retinol for the synthesis of collagen and egg development (Cheng et al. 2013; Wang et al. 2018).

In accordance with both hypotheses, FAR proteins in *Bursaphelenchus mucronatus* are among the genes with increased expression in the more virulent strains of this plant parasite. A silencing experiment showed a decrease of infective capacity and offspring, associating FARs with both development and reproduction functions as well as disturbing inter-intracellular defense signals in plants (Zhou et al. 2016). Similar experiments and results are presented with Pp-FAR-1 from *Pratylenchus penetrans* (Vieira et al. 2015), highly expressed during infection and showing a significant reduction in reproduction upon silencing, and Rs-FAR-1 from *Radopholus similis* (Zhang et al. 2015), as this protein regulates expression of allene oxide synthase and it is upregulated during early infection in *Arabidopsis thaliana*.

At last, work done in *Heterodera avenae* described two FAR proteins, Ha-FAR-1 and Ha-FAR-2, with different molecular structures (Le et al. 2016). While Ha-FAR-1 has the conserved phosphorylation site and no glycosylation site, Ha-FAR-2 has a glycosylation site and no phosphorylation site, also showing weaker binding capabilities. Expression of both is seen in hypodermis, suggesting secretion, but Ha-FAR-1 is greater than Ha-FAR-2, and although they are expressed through all stages, Ha-FAR1 is predominant in the J4 stage (prereproductive), and Ha-FAR2 is predominant in J2 (post-parasitic). Structurally, they show a 28% similarity, a difference also seen in their weight, 17 kDa for Ha-FAR-1, and 30kDa for Ha-FAR-2; and isoelectric point, 5.62 for Ha-FAR-1 and 9.02 for Ha-FAR-2, which might indicate that they work in different environments (Le et al. 2016).

### FAR proteins in *Caenorhabditis* genus

Recent studies done in *Caenorhabditis bovis*, a close relative of *C. elegans*, have rekindled the interest in FAR-8, an outlier of the FAR protein family (Garofalo et al. 2002), as a possible factor in the evolution towards parasitic behavior. *C. bovis* is unusual for a *Caenorhabditis* species, as it has been isolated several times from the outer auditory canals of Zebu cattle in Eastern Africa and Gyr cattle in South America, presenting a deeply different habitat (Cardona et al. 2010; Kiontke and Sudhaus 2006). *C. bovis* is believed to be the causative agent of bovine parasitic otitis. However, there is not enough information about the biology to classify it as a “true” parasite; *C. bovis* might instead be an opportunistic colonizer of niches created by other pathogens. The publication of the genome of *C. bovis* showed that the genome and gene set of *C. bovis*, compared to *C. elegans* is smaller



but at least three gene families have undergone an expansion and all of them have been related to parasitism (Stevens et al. 2020). The FAR family of genes in *C. bovis* has been almost completely lost, except for FAR-8 which is repeated two times. This suggests that there might be a relationship between the adaptation to a completely different niche and the conservation and expansion of this FAR protein. These findings indicate how poorly understood these proteins are and that molecular analysis alone provides little information of the possible biological functions and role of these proteins in the major plot that is the evolution in nematodes.


### Nematode polypeptide antigens/allergens (NPAs)

Nematodes present yet another well-described LBP group, the NPAs. These proteins are found in the pseudocoelomic fluid and connective tissue of nematodes, as well as in the E/S products where they induce a highly immunogenic response from the host (Kennedy 2011).

NPAs present a quite particular translation mechanism since they are produced as a large polypeptide that is post-translationally cleaved by proteases rendering globular units of 15 kDa (Kennedy 2011). These units are helix-rich and can present similar or divergent amino acid sequences, depending on the species (Kennedy et al. 1995; Kuang et al. 2009).

Only one of these proteins has its structure determined by nuclear magnetic resonance (NMR), ABA-1A from *Ascaris suum* (Meenan et al. 2011). This protein adopts a novel seven-helical fold comprising a long central helix that participates in two four-helical bundles. There are two discrete hydrophobic ligand-binding pockets, one in each of the N-terminal and C-terminal bundles (Fig. 4). It is important to note that this is the first structure of a unit of any tandemly repetitive polypeptide yet reported. To date, the structure and ligand-binding characteristics of these proteins have been demonstrated only for single units of the polypeptide, but not for two or more units in tandem from the same polypeptide array as these functions might change if interactions with other subunits are present.

From the functional point of view, NPAs bind small lipids such as fatty acids and retinol as has been demonstrated using in vitro fluorescence experiments (Moore et al. 1999). Additionally, it has been shown that NPAs, particularly ABA-1A, physically interact with membranes and they unload their lipid cargo by releasing it into the aqueous phase (McDermott et al. 2002). They have been described in several species of disease-causing nematodes, including *Ascaris lumbricoides*, *A. suum* and *B. malayi* of humans, and *Dictyocaulus viviparus*, *Ostertagia ostertagi*, *Haemonchus contortus*, and *Dirofilaria immitis* of domestic animals (Britton et al. 1995; Kuang et al. 2009; Poole et al. 1996; Selkirk et al. 1993; Solovyova et al. 2003; Xia et al. 2000).



**Fig. 4** NMR structure of ABA-1A (PDB: 2XV9) from *Ascaris suum*. Black arrows are showing the two described ligand binding sites

Notably, as with other LBPs, NPAs are also found in free-living species from the phylum. Indeed the *npa-1* gene from *C. elegans* has been involved in some morphological, and life span phenotypes obtained in gunshot RNAi experiments (Ceol and Horvitz 2004; Curran and Ruvkun 2007). Unfortunately, no further functional experiments have been performed on these extremely interesting proteins.

### Novel LBPs of parasitic helminths: a bottomless pit

It is well known that we are still in a lack of knowledge of novel species across the biodiversity worldwide and hence a lot of biomolecules are still to be described. A good example is the case of the dorylipophorin (GeneBank: MW014827.1) isolated from the pseudocoelomic fluid of *Diectophyme renale*, the giant kidney worm (Giorello et al. 2017). To date, *D. renale* is a clade I nematode that has no genomic, transcriptomic, or proteomic data, yet it is an important parasitic infection affecting domestic and wild fauna, not mentioning that it is a zoonosis (Eiras et al. 2021; Paras et al. 2018). The phylogenetically closest species for which genomic data is available is *Sobolophyme baturini* that belongs to the same subclass Dorylaimia but is a member of an entirely different family of nematode parasites. It is important to note that other members of clade I include the human parasites *Trichuris trichiura* and *Trichinella spiralis*.

Dorylipophorin has proven to selectively bind fatty acids from a highly heterogeneous environment (Giorello et al.



2017). N-terminal and internal peptide amino-acid sequences of this protein indicate a relationship with a cysteine- and histidine-rich protein of unknown function from *Trichinella spiralis* and *Trichuris muris* which are found to be the major component of E/S products (Radoslavov et al. 2010; Tritten et al. 2017). Recently, the structure of P43, the major secreted protein from *T. muris*, was resolved to the atomic level, and it was shown to bind the signaling molecule interleukin-13 (Bancroft et al. 2019). Since dorylipophorin and P43 present 50% sequence identity, these observations give room to hypothesize that dorylipophorin might be important in the attenuation of host's immune response. Notably, within the five clades that conform phylum Nematoda, clade I only presents FABPs and now this novel family of LBPs which seems to be clade specific.

## Concluding remarks

Different members of LBPs are present in all parasitic helminths. In some cases, they can be found broadly distributed like FABPs, but in other cases, they are highly specific of a particular group as is the case of HLBP or the clade I from phylum Nematoda, to which P43 and dorylipophorin belong. LBPs are usually found in multigenic families where function might vary even in the same species, as more than one LBP protein might be present and its expression could change through the developmental stages.

Regardless of the enormous amount of information available, a specific function cannot be strictly associated with any of these families of proteins yet. As stated before, LBPs have been proposed to participate in the acquisition and distribution of lipids within tissues of the parasite; and/or in the modulation of the host's immune systems. To date, experiments performed with HLBP and FAR proteins (exclusively found in cestodes and nematodes, respectively) presented strong evidence about their role in the modulation of the host's immune response. On the other hand, FABPs might be involved in the distribution and storage of lipids in nematodes. Whether this is also the case for cestodes remains to be determined. For nemFABPs and NPAs, detailed knowledge about their structure and possible natural ligands is available, but there are no direct functional experiments.

Finally, the HLBPs are the only LBPs with a detailed description of their cargo obtained from their natural environment; this is clearly a missing block of information that needs attention in order to decipher the metabolic routes these proteins are related to.

**Acknowledgements** The authors would like to thank Dr. Marcela Cucher from IMPaM UBA and Dr. Luciana Rodriguez Sawicki from INIBIOLP UNLP for critically reading the manuscript.

**Author contribution** All authors performed the literature search and data analysis. All authors drafted and critically revised the work. All authors read and approved the final manuscript.

**Availability of data and material** Not applicable.

## Declarations

**Additional declarations for articles in life science journals that report the results of studies involving humans and/or animals** Not applicable.

**Ethics approval** Not applicable.

**Consent to participate** All authors agreed to participate.

**Consent for publication** All authors agreed to publish this manuscript.

**Conflicts of interest** The authors declare no competing interests.

## References

- Alvite G, Esteves A (2012) Lipid binding proteins from parasitic platyhelminthes. *Front Physiol* 3. <https://doi.org/10.3389/FPHYS.2012.00363>
- Arend AC, Zaha A, Ayala FJ, Haag KL (2004) The *Echinococcus granulosus* antigen B shows a high degree of genetic variability. *Exp Parasitol* 108(1–2):76–80. <https://doi.org/10.1016/j.exppara.2004.07.009>
- Bancroft AJ, Levy CW, Jowitt TA, Hayes KS, Thompson S, McKenzie EA, Ball MD, Dubaissi E, France AP, Bellina B, Sharpe C, Mironov A, Brown SL, Cook PC, MacDonald AS, Thornton DJ, Grecis RK, MacDonald AS, Thornton DJ, Grecis RK (2019) The major secreted protein of the whipworm parasite tethers to matrix and inhibits interleukin-13 function. *10(1):1–11*. <https://doi.org/10.1038/s41467-019-09996-z>
- Barf T, Lehmann F, Hammer K, Haile S, Axen E, Medina C, Uppenberg J, Svensson S, Rondahl L, Lundbäck T (2009) N-benzylindole carboxylic acids: design and synthesis of potent and selective adipocyte fatty-acid binding protein (a-FABP) inhibitors. *Bioorg Med Chem Lett* 19(6):1745–1748. <https://doi.org/10.1016/j.bmcl.2009.01.084>
- Barletta GP, Franchini G, Corsico B, Fernandez-Alberti S (2019) Fatty acid and retinol-binding protein: unusual protein conformational and cavity changes dictated by ligand fluctuations. *J Chem Inf Model* 59(8):3545–3555. <https://doi.org/10.1021/acs.jcim.9b00364>
- Barrett J (2009) Forty years of helminth biochemistry. *Parasitology* 136(12):1633–1642. <https://doi.org/10.1017/S003118200900568X>
- Barrett J, Saghir N, Timanova A, Clarke K, Brophy PM (1997) Characterisation and properties of an intracellular lipid-binding protein from the tapeworm *Moniezia expansa*. *Eur J Biochem* 250(2):269–275. <https://doi.org/10.1111/j.1432-1033.1997.0269a.x>
- Barrett J (1981) Nutrition and biosynthesis. In: *Biochemistry of Parasitic Helminths*. Macmillan Education UK, pp 149–244. [https://doi.org/10.1007/978-1-349-86119-4\\_4](https://doi.org/10.1007/978-1-349-86119-4_4)
- Basavaraju S, Zhan B, Kennedy MW, Liu Y, Hawdon J, Hotez PJ (2003) Ac-FAR-1, a 20 kDa fatty acid- and retinol-binding protein secreted by adult *Ancylostoma caninum* hookworms: gene transcription pattern, ligand binding properties and structural

- characterisation. *Mol Biochem Parasitol* 126(1):63–71. [https://doi.org/10.1016/S0166-6851\(02\)00253-0](https://doi.org/10.1016/S0166-6851(02)00253-0)
- Becker MM, Kalinna BH, Waine GJ, McManus DP (1994) Gene cloning, overproduction and purification of a functionally active cytoplasmic fatty acid-binding protein (Sj-FABPc) from the human blood fluke *Schistosoma japonicum*. *Gene* 148(2):321–325. [https://doi.org/10.1016/0378-1119\(94\)90706-4](https://doi.org/10.1016/0378-1119(94)90706-4)
- Bélgamo JA, Alberca LN, Pórfido JL, Romero FNC, Rodriguez S, Talevi A, Córscico B, Franchini GR (2020) Application of target repositioning and in silico screening to exploit fatty acid binding proteins (FABPs) from *Echinococcus multilocularis* as possible drug targets. *J Comput Aided Mol Des* 34(12):1275–1288. <https://doi.org/10.1007/s10822-020-00352-8>
- Blaxter M, Koutsovoulos G (2015) The evolution of parasitism in Nematoda. *Parasitology* 142(Suppl 1):S26–S39. <https://doi.org/10.1017/S0031182014000791>
- Bradley JE, Nirmalan N, Kläger SL, Faulkner H, Kennedy MW (2001) River blindness: a role for parasite retinoid-binding proteins in the generation of pathology? *Trends Parasitol* 17(10):471–475. [https://doi.org/10.1016/S1471-4922\(01\)02036-0](https://doi.org/10.1016/S1471-4922(01)02036-0)
- Britton C, Moore J, Gilleard JS, Kennedy MW (1995) Extensive diversity in repeat unit sequences of the cDNA encoding the polypeptide antigen/allergen from the bovine lungworm *Dictyocaulus viviparus*. *Mol Biochem Parasitol* 72(1–2):77–88. [https://doi.org/10.1016/0166-6851\(95\)00088-1](https://doi.org/10.1016/0166-6851(95)00088-1)
- Britton C, Roberts B, Marks ND (2016) Functional genomics tools for *Haemonchus contortus* and lessons from other Helminths. *Adv Parasitol* 93:599–623. <https://doi.org/10.1016/bs.apar.2016.02.017>
- Burbelo PD, Leahy HP, Iadarola MJ, Nutman TB (2009) A four-antigen mixture for rapid assessment of *Onchocerca volvulus* infection. *PLoS Negl Trop Dis* 3(5):e438. <https://doi.org/10.1371/journal.pntd.0000438>
- Cardona AJ, González TM, Álvarez PJ (2010) Otitis bovina por Rhabditis bovis en Córdoba, Colombia. Reporte de dos casos. *Rev MVZ Cordoba* 15(3):2240–2244. <https://doi.org/10.21897/rmvz.311>
- Ceol CJ, Horvitz HR (2004) A new class of *C. elegans* synMuv genes implicates a Tip60/NuA4-like HAT complex as a negative regulator of Ras signaling. *Dev Cell* 6(4):563–576. [https://doi.org/10.1016/S1534-5807\(04\)00065-6](https://doi.org/10.1016/S1534-5807(04)00065-6)
- Chemale G, Haag KL, Ferreira HB, Zaha A (2001) *Echinococcus granulosus* antigen B is encoded by a gene family. *Mol Biochem Parasitol* 116(2):233–237. [https://doi.org/10.1016/S0166-6851\(01\)00316-4](https://doi.org/10.1016/S0166-6851(01)00316-4)
- Cheng X, Xiang Y, Xie H, Xu CL, Xie TF, Zhang C, Li Y (2013) Molecular characterization and functions of fatty acid and retinoid binding protein gene (Ab-far-1) in *Aphelenchoides besseyi*. *PLoS ONE* 8(6):e66011. <https://doi.org/10.1371/journal.pone.0066011>
- Curran SP, Ruvkun G (2007) Lifespan regulation by evolutionarily conserved genes essential for viability. *PLoS Genet* 3(4):0479–0487. <https://doi.org/10.1371/journal.pgen.0030056>
- Dutta TK, Banakar P, Rao U (2015) The status of RNAi-based transgenic research in plant nematology. *Front Microbiol* 0(DEC):760. <https://doi.org/10.3389/FMICB.2014.00760>
- Eiras J, Zhu XQ, Yurlova N, Pedrassani D, Yoshikawa M, Nawa Y (2021) *Diectophyme renale* (Goeze, 1782) (Nematoda, Diectophymidae) parasitic in mammals other than humans: a comprehensive review. *Parasitol Int* 81:102269. <https://doi.org/10.1016/j.parint.2020.102269>
- Esteves A, Ehrlich R (2006) Invertebrate intracellular fatty acid binding proteins. *Comp Biochem Physiol C: Toxicol Pharmacol* 142(3–4):262–274. <https://doi.org/10.1016/j.cbpc.2005.11.006>
- Esteves A, Dallagiovanna B, Ehrlich R (1993) A developmentally regulated gene of *Echinococcus granulosus* codes for a 15.5-kilodalton polypeptide related to fatty acid binding proteins. *Mol Biochem Parasitol* 58(2):215–222. [https://doi.org/10.1016/0166-6851\(93\)90043-W](https://doi.org/10.1016/0166-6851(93)90043-W)
- Estuningsih SE, Smooker PM, Wiedosari E, Widjajanti S, Vaiano S, Partoutomo S, Spithill TW (1997) Evaluation of antigens of *Fasciola gigantica* as vaccines against tropical fasciolosis in cattle. *Int J Parasitol* 27(11):1419–1428. [https://doi.org/10.1016/S0020-7519\(97\)00096-9](https://doi.org/10.1016/S0020-7519(97)00096-9)
- Fairfax KC, Vermeire JJ, Harrison LM, Bungiro RD, Grant W, Husain SZ, Cappello M (2009) Characterisation of a fatty acid and retinol binding protein orthologue from the hookworm *Ancylostoma ceylanicum*. *Int J Parasitol* 39(14):1561–1571. <https://doi.org/10.1016/j.ijpara.2009.06.005>
- Fernández V, Ferreira HB, Fernández C, Zaha A, Nieto A (1996) Molecular characterisation of a novel 8-kDa subunit of *Echinococcus granulosus* antigen B. *Mol Biochem Parasitol* 77(2):247–250. [https://doi.org/10.1016/0166-6851\(96\)02602-3](https://doi.org/10.1016/0166-6851(96)02602-3)
- Frayha GJ, Smyth JD (1983) Lipid metabolism in parasitic helminths. *Adv Parasitol* 22(C):309–387. [https://doi.org/10.1016/S0065-308X\(08\)60465-0](https://doi.org/10.1016/S0065-308X(08)60465-0)
- Frosch P, Hartmann M, Mühlischlegel F, Frosch M (1994) Sequence heterogeneity of the echinococcal antigen B. *Mol Biochem Parasitol* 64(1):171–175. [https://doi.org/10.1016/0166-6851\(94\)90145-7](https://doi.org/10.1016/0166-6851(94)90145-7)
- Furuhashi M, Tuncman G, Görgün CZ, Makowski L, Atsumi G, Vailancourt E, Kono K, Babaev VR, Fazio S, Linton MF, Sulsky R, Robl JA, Parker RA, Hotamisligil GS (2007) Treatment of diabetes and atherosclerosis by inhibiting fatty-acid-binding protein aP2. *Nature* 447(7147):959–965. <https://doi.org/10.1038/nature05844>
- Garofalo A, Kläger SL, Rowlinson MC, Nirmalan N, Klion A, Allen JE, Kennedy MW, Bradley JE (2002) The FAR proteins of filarial nematodes: secretion, glycosylation and lipid binding characteristics. *Mol Biochem Parasitol* 122(2):161–170. [https://doi.org/10.1016/S0166-6851\(02\)00097-X](https://doi.org/10.1016/S0166-6851(02)00097-X)
- Giorello AN, Kennedy MW, Butti MJ, Radman NE, Córscico B, Franchini GR (2017) Identification and characterization of the major pseudocoelomic proteins of the giant kidney worm, *Diectophyme renale*. *Parasites Vectors* 10(1):446. <https://doi.org/10.1186/s13071-017-2388-x>
- González G, Nieto A, Fernández C, Örn A, Wernstedt C, Hellman U (1996) Two different 8 kDa monomers are involved in the oligomeric organization of the native *Echinococcus granulosus* antigen B. *Parasite Immunol* 18(12):587–596. <https://doi.org/10.1046/j.1365-3024.1996.d01-38.x>
- Guidi A, Mansour NR, Paveley RA, Carruthers IM, Besnard J, Hopkins AL, Gilbert IH, Bickle QD (2015) Application of RNAi to genomic drug target validation in schistosomes. *PLoS Negl Trop Dis* 9(5):e0003801. <https://doi.org/10.1371/JOURNAL.PNTD.0003801>
- Haag KL, Ayala FJ, Kamenetzky L, Gutierrez AM, Rosenzvit M (2004) Livestock trade history, geography, and parasite strains: the mitochondrial genetic structure of *Echinococcus granulosus* in Argentina. *J Parasitol* 90(2):234–239. <https://doi.org/10.1645/GE-173R>
- Haunerland NH, Chisholm JM (1990) Fatty acid binding protein in flight muscle of the locust, *Schistocerca gregaria*. *Biochim Biophys Acta (BBA)/Lipids Lipid Metab* 1047(3):233–238. [https://doi.org/10.1016/0005-2760\(90\)90521-X](https://doi.org/10.1016/0005-2760(90)90521-X)
- Haunerland NH, Spener F (2004) Fatty acid-binding proteins - Insights from genetic manipulations. *Prog Lipid Res* 43(4):328–349. <https://doi.org/10.1016/j.plipres.2004.05.001>
- Hewitson JP, Grainger JR, Maizels RM (2009) Helminth immunoregulation: the role of parasite secreted proteins in modulating host immunity. *Mol Biochem Parasitol* 167(1):1–11. <https://doi.org/10.1016/j.molbiopara.2009.04.008>

- Hiepe T, Lucius R, Gottstein B (2006) Allgemeine Parasitologie mit Grundzügen der Immunologie, Diagnostik und Bekämpfung. Schweizer Archiv Für Tierheilkunde 103(10):A-615. <https://doi.org/10.1024/0036-7281.148.6.317a>
- Huang F, Dang Z, Suzuki Y, Horiuchi T, Yagi K, Kouguchi H, Irie T, Kim K, Oku Y (2016) Analysis on gene expression profile in oncospheres and early stage metacystodes from echinococcus multilocularis. PLoS Negl Trop Dis 10(4):e0004634. <https://doi.org/10.1371/JOURNAL.PNTD.0004634>
- Ibáñez-Shimabukuro M, Florencia Rey-Burusco M, Gabrielsen M, Franchini GR, Riboldi-Tunnicliffe A, Roe AJ, Griffiths K, Cooper A, Córscico B, Kennedy MW, Smith BO (2019) Structure and ligand binding of As-p18, an extracellular fatty acid binding protein from the eggs of a parasitic nematode. Biosci Rep 39(7). <https://doi.org/10.1042/BSR20191292>
- Iberkleid I, Vieira P, de Almeida Engler J, Firester K, Spiegel Y, Horowitz SB (2013) Fatty acid-and retinol-binding protein, Mj-FAR-1 induces tomato host susceptibility to root-knot nematodes. PLoS ONE 8(5):e64586. <https://doi.org/10.1371/journal.pone.0064586>
- Iberkleid I, Sela N, Brown Miyara S (2015) Meloidogyne javanica fatty acid- and retinol-binding protein (Mj-FAR-1) regulates expression of lipid-, cell wall-, stress- and phenylpropanoid-related genes during nematode infection of tomato. BMC Genomics 16(1):1–26. <https://doi.org/10.1186/s12864-015-1426-3>
- Jia WZ, Yan HB, Lou ZZ, Ni XW, Liu HX, Li HM, Guo AJ, Fu BQ (2011) Genetic variation of the 8-kDa glycoprotein family from echinococcus granulosus, taenia multiceps and taenia hydatigena. Chin Med J 124(18):2849–2856. <https://doi.org/10.3760/cma.j.issn.0366-6999.2011.18.010>
- Jian X, Shu-Hua X, Hui-Qing Q, Sen L, Hotez P, Bing-Gui S, Hai-Chou X, Tie-Hua L, Bin Z (2003) Necator americanus: maintenance through one hundred generations in golden hamsters (Mesocricetus auratus). II. Morphological development of the adult and its comparison with humans. Exp Parasitol 105(3–4):192–200. <https://doi.org/10.1016/j.exppara.2003.12.016>
- Jordanova R, Groves MR, Kostova E, Woltersdorf C, Liebau E, Tucker PA (2009) Fatty acid- and retinoid-binding proteins have distinct binding pockets for the two types of cargo. J Biol Chem 284(51):35818–35826. <https://doi.org/10.1074/jbc.M109.022731>
- Kamenetzky L, Muzulin PM, Gutierrez AM, Angel SO, Zaha A, Guarnera EA, Rosenzvit MC (2005) High polymorphism in genes encoding antigen B from human infecting strains of Echinococcus granulosus. Parasitology 131(6):805–815. <https://doi.org/10.1017/S0031182005008474>
- Kennedy MW (2011) The polypeptide allergens of nematodes (NPAs) - structure at last, but still mysterious. Exp Parasitol 129(2):81–84. <https://doi.org/10.1016/j.exppara.2011.06.007>
- Kennedy MW, Tomlinson LA, Fraser EM, Christie JF (1990) The specificity of the antibody response to internal antigens of Ascaris: heterogeneity in infected humans, and MHC (H-2) control of the repertoire in mice. Clin Exp Immunol 80(2):219–224. <https://doi.org/10.1111/j.1365-2249.1990.tb05237.x>
- Kennedy MW, Britton C, Price NC, Kelly SM, Cooper A (1995) The DvA-1 polypeptide of the parasitic nematode Dictyocaulus viviparus. A small helix-rich lipid-binding protein. J Biol Chem 270(33):19277–19281. <https://doi.org/10.1074/jbc.270.33.19277>
- Kennedy MW, Garside LH, Goodrick LE, McDermott L, Brass A, Price NC, Kelly SM, Cooper A, Bradley JE (1997) The Ov20 protein of the parasitic nematode Onchocerca volvulus. A structurally novel class of small helix-rich retinol-binding proteins. J Biol Chem 272(47):29442–29448. <https://doi.org/10.1074/jbc.272.47.29442>
- Kimber MJ, McKinney S, McMaster S, Day TA, Fleming CC, Maule AG (2007) flp gene disruption in a parasitic nematode reveals motor dysfunction and unusual neuronal sensitivity to RNA interference. FASEB J 21(4):1233–1243. <https://doi.org/10.1096/fj.06-7343com>
- Kiontke K, Sudhaus W (2006) Ecology of Caenorhabditis species. In: WormBook : the online review of C. elegans biology. pp 1–14. <https://doi.org/10.1895/wormbook.1.37.1>
- Kuang L, Colgrave ML, Bagnall NH, Knox MR, Qian M, Wijffels G (2009) The complexity of the secreted NPA and FAR lipid-binding protein families of Haemonchus contortus revealed by an iterative proteomics-bioinformatics approach. Mol Biochem Parasitol 168(1):84–94. <https://doi.org/10.1016/j.molbiopara.2009.07.001>
- Le X, Wang X, Guan T, Ju Y, Li H (2016) Isolation and characterization of a fatty acid- and retinoid-binding protein from the cereal cyst nematode Heterodera avenae. Exp Parasitol 167:94–102. <https://doi.org/10.1016/j.exppara.2016.05.009>
- Lee E-G, Kim S-H, Bae Y-A, Chung J-Y, Suh M, Na B-K, Kim T-S, Kang I, Ma L, Kong Y (2007) A hydrophobic ligand-binding protein of the Taenia solium metacystode mediates uptake of the host lipid: Implication for the maintenance of parasitic cellular homeostasis. Proteomics 7(21):4016–4030. <https://doi.org/10.1002/PMIC.200700332>
- Li BW, Rush AC, Tan J, Weil GJ (2004) Quantitative analysis of gender-regulated transcripts in the filarial nematode Brugia malayi by real-time RT-PCR. Mol Biochem Parasitol 137(2):329–337. <https://doi.org/10.1016/j.molbiopara.2004.07.002>
- Lightowler MW, Liu D, Haralambous A, Rickard MD (1989) Subunit composition and specificity of the major cyst fluid antigens of Echinococcus granulosus. Mol Biochem Parasitol 37(2):171–182. [https://doi.org/10.1016/0166-6851\(89\)90149-7](https://doi.org/10.1016/0166-6851(89)90149-7)
- Lilley CJ, Davies LJ, Urwin PE (2012) RNA interference in plant parasitic nematodes: a summary of the current status. Parasitology 139(5):630–640. <https://doi.org/10.1017/S0031182011002071>
- Liu RZ, Li X, Godbout R (2008) A novel fatty acid-binding protein (FABP) gene resulting from tandem gene duplication in mammals: transcription in rat retina and testis. Genomics 92(6):436–445. <https://doi.org/10.1016/j.ygeno.2008.08.003>
- Logan J, Pearson MS, Manda SS, Choi Y-J, Field M, Eichenberger RM, Mulvenna J, Nagaraj SH, Fujiwara RT, Gazzinelli-Guimaraes P, Bueno L, Mati V, Bethony JM, Mitreva M, Sotillo J, Loukas A (2020) Comprehensive analysis of the secreted proteome of adult Necator americanus hookworms. PLoS Negl Trop Dis 14(5):1–30. <https://doi.org/10.1371/JOURNAL.PNTD.0008237>
- MacDonald K, Buxton S, Kimber MJ, Day TA, Robertson AP, Ribeiro P (2014) Functional characterization of a novel family of acetylcholine-gated chloride channels in Schistosoma mansoni. PLoS Pathog 10(6):e1004181. <https://doi.org/10.1371/journal.ppat.1004181>
- Marcelino AMC, Smock RG, Gierasch LM (2006) Evolutionary coupling of structural and functional sequence information in the intracellular lipid-binding protein family. Proteins: Struct Funct Genet 63(2):373–384. <https://doi.org/10.1002/prot.20860>
- McDermott L, Kennedy MW, McManus DP, Bradley JE, Cooper A, Storch J (2002) How helminth lipid-binding proteins offload their ligands to membranes: differential mechanisms of fatty acid transfer by the ABA-1 Polypeptide Allergen and Ov-FAR-1 Proteins of Nematodes and Sj-FABPc of Schistosomes. Biochemistry 41(21):6706–6713. <https://doi.org/10.1021/bi0159635>
- Meenan NAG, Ball G, Bromek K, Uhrin D, Cooper A, Kennedy MW, Smith BO (2011) Solution structure of a repeated unit of the ABA-1 nematode polypeptide allergen of Ascaris reveals a novel fold and two discrete lipid-binding sites. PLoS Negl Trop Dis 5(4):e1040. <https://doi.org/10.1371/journal.pntd.0001040>
- Mei B, Kennedy MW, Beauchamp J, Komuniecki PR, Komuniecki R (1997) Secretion of a novel, developmentally regulated fatty acid-binding protein into the perivitelline fluid of the parasitic



- nematode, *Ascaris suum*. *J Biol Chem* 272(15):9933–9941. <https://doi.org/10.1074/jbc.272.15.9933>
- Michalski ML, Weil GJ (1999) Gender-specific gene expression in *Brugia malayi*. *Mol Biochem Parasitol* 104(2):247–257. [https://doi.org/10.1016/S0166-6851\(99\)00149-8](https://doi.org/10.1016/S0166-6851(99)00149-8)
- Michalski ML, Monsey JD, Cistola DP, Weil GJ (2002) An embryo-associated fatty acid-binding protein in the filarial nematode *Brugia malayi*. *Mol Biochem Parasitol* 124(1–2):1–10. [https://doi.org/10.1016/S0166-6851\(02\)00081-6](https://doi.org/10.1016/S0166-6851(02)00081-6)
- Moore J, McDermott L, Price NC, Kelly SM, Cooper A, Kennedy MW (1999) Sequence-divergent units of the ABA-1 polypeptide array of the nematode *Ascaris suum* have similar fatty-acid- and retinol-binding properties but different binding-site environments. *Biochem J* 340(1):337–343. <https://doi.org/10.1042/bj3400337>
- Morphew RM, Wilkinson TJ, MacKintosh N, Jahndel V, Paterson S, McVeigh P, Abbas Abidi SM, Saifullah K, Raman M, Ravikumar G, Lacourse J, Maule A, Brophy PM (2016) Exploring and expanding the fatty-acid-binding protein superfamily in *Fasciola* species. *J Proteome Res* 15(9):3308–3321. <https://doi.org/10.1021/acs.jproteome.6b00331>
- Moser D, Tendler M, Griffiths G, Klinkert MQ (1991) A 14-kDa *Schistosoma mansoni* polypeptide is homologous to a gene family of fatty acid binding proteins. *J Biol Chem* 266(13):8447–8454. [https://doi.org/10.1016/S0021-9258\(18\)92995-9](https://doi.org/10.1016/S0021-9258(18)92995-9)
- Muzulin PM, Kamenetzky L, Gutierrez AM, Guarnera EA, Rosenzvit MC (2008) *Echinococcus granulosus* antigen B gene family: further studies of strain polymorphism at the genomic and transcriptional levels. *Exp Parasitol* 118(2):156–164. <https://doi.org/10.1016/j.exppara.2007.07.004>
- Nirmalan N, Cordeiro NJV, Kläger SL, Bradley JE, Allen JE (1999) Comparative analysis of glycosylated and nonglycosylated filarial homologues of the 20-kilodalton retinol binding protein from *Onchocerca volvulus* (Ov20). *Infect Immun* 67(12):6329–6334. <https://doi.org/10.1128/iai.67.12.6329-6334.1999>
- Obal G, Ramos AL, Silva V, Lima A, Batthyany C, Bessio MI, Ferreira F, Salinas G, Ferreira AM (2012) Characterisation of the native lipid moiety of *echinococcus granulosus* antigen B. *PLoS Negl Trop Dis* 6(5):e1642. <https://doi.org/10.1371/journal.pntd.0001642>
- Oriol R, Williams JF, Pérez Esandi MV, Oriol C (1971) Purification of lipoprotein antigens of *Echinococcus granulosus* from sheep hydatid fluid. *Am J Trop Med Hyg* 20(4):569–574. <https://doi.org/10.4269/ajtmh.1971.20.569>
- Paras KL, Miller L, Verocai GG (2018) Ectopic infection by *Diocotylome renale* in a dog from Georgia, USA, and a review of cases of ectopic diocotylomosis in companion animals in the Americas. *Vet Parasitol: Reg Stud Rep* 14:111–116. <https://doi.org/10.1016/j.vprsr.2018.09.008>
- Parks SC, Nguyen C, Nasrolahi S, Juncas D, Lu D, Ramaswamy R, Dhillon H, Buchman A, Akbari OS, Yamanaka N, Boulanger MJ, Dillman AR (2021) Parasitic nematode fatty acid- and retinol-binding proteins compromise host immunity by interfering with host lipid signaling pathways. *BioRxiv*, 2021.03.25.436866. <https://doi.org/10.1101/2021.03.25.436866>
- Pérez MG, Spiliotis M, Rego N, Macchiaroli N, Kamenetzky L, Holroyd N, Cucher MA, Brehm K, Rosenzvit MC (2019) Deciphering the role of miR-71 in *echinococcus multilocularis* early development in vitro. *PLoS Negl Trop Dis* 13(12):1–21. <https://doi.org/10.1371/JOURNAL.PNTD.0007932>
- Phani V, Shivakumara TN, Davies KG, Rao U (2017) Meloidogyne incognita fatty acid- and retinol- binding protein (Mi-FAR-1) affects nematode infection of plant roots and the attachment of pasteuria penetrans endospores. *Front Microbiol* 8(NOV). <https://doi.org/10.3389/FMICB.2017.02122>
- Plenefisch J, Xiao H, Mei B, Geng J, Komuniecki PR, Komuniecki R (2000) Secretion of a novel class of iFABPs in nematodes: coordinate use of the *Ascaris*/*Caenorhabditis* model systems. *Mol Biochem Parasitol* 105(2):223–236. [https://doi.org/10.1016/S0166-6851\(99\)00179-6](https://doi.org/10.1016/S0166-6851(99)00179-6)
- Poole CB, Hornstra LJ, Benner JS, Fink JR, McReynolds LA (1996) Carboxy-terminal sequence divergence and processing of the polypeptide antigen from *Dirofilaria immitis*. *Mol Biochem Parasitol* 82(1):51–65. [https://doi.org/10.1016/0166-6851\(96\)02721-1](https://doi.org/10.1016/0166-6851(96)02721-1)
- Pórfido JL, Herz M, Kiss F, Kamenetzky L, Brehm K, Rosenzvit MC, Córscico B, Franchini GR (2020) Fatty acid-binding proteins in *Echinococcus* spp.: the family has grown. *Parasitol Res* 119(4):1401–1408. <https://doi.org/10.1007/s00436-020-06631-5>
- Prior A, Jones JT, Blok VC, Beauchamp J, McDermott L, Cooper A, Kennedy MW (2001) A surface-associated retinol- and fatty acid-binding protein (GP-FAR-1) from the potato cyst nematode *Globodera pallida*: lipid binding activities, structural analysis and expression pattern. *Biochem J* 356(2):387–394. <https://doi.org/10.1042/0264-6021:3560387>
- Radoslavov G, Jordanova R, Teofanova D, Georgieva K, Hristov P, Salomone-Stagni M, Liebau E, Bankov I (2010) A novel secretory poly-cysteine and histidine-tailed metalloprotein (Ts-PCHTP) from *Trichinella spiralis* (Nematoda). *PLoS ONE* 5(10):e13343. <https://doi.org/10.1371/JOURNAL.PONE.0013343>
- Rey Burusco MFI (2014) Estudio biofísico y estructural de Na-FAR-1, miembro de una nueva familia de proteínas de nematodos que unen ácidos grasos y retinol [Universidad Nacional de La Plata]. <https://doi.org/10.35537/10915/34308>
- Rey-Burusco MF, Ibáñez-Shimabukuro M, Gabrielsen M, Franchini GR, Roe AJ, Griffiths K, Zhan B, Cooper A, Kennedy MW, Córscico B, Smith BO (2015) Diversity in the structures and ligand-binding sites of nematode fatty acid and retinol-binding proteins revealed by Na-FAR-1 from *Necator americanus*. *Biochem J* 471(3):403–414. <https://doi.org/10.1042/BJ20150068>
- Rodríguez-Pérez J, Rodríguez-Medina JR, García-Blanco MA, Hillyer GV (1992) *Fasciola hepatica*: molecular cloning, nucleotide sequence, and expression of a gene encoding a polypeptide homologous to a *Schistosoma mansoni* fatty acid-binding protein. *Exp Parasitol* 74(4):400–407. [https://doi.org/10.1016/0014-4894\(92\)90202-L](https://doi.org/10.1016/0014-4894(92)90202-L)
- Saghir N, Conde PJ, Brophy PM, Barrett J (2001) Biochemical characterisation of a hydrophobic ligand binding protein from the tapeworm *Hymenolepis diminuta*. *Int J Parasitol* 31(7):653–660. [https://doi.org/10.1016/S0020-7519\(01\)00135-7](https://doi.org/10.1016/S0020-7519(01)00135-7)
- Sako Y, Nakao M, Ikejima T, Piao XZ, Nakaya K, Ito A (2000) Molecular characterization and diagnostic value of *Taenia solium* low-molecular-weight antigen genes. *J Clin Microbiol* 38(12):4439–4444. <https://doi.org/10.1128/jcm.38.12.4439-4444.2000>
- Sanchez F, March F, Mercader M, Coll P, Muñoz C, Prats G (1991) Immunochemical localization of major hydatid fluid antigens in protoscoleces and cysts of *Echinococcus granulosus* from human origin. *Parasite Immunol* 13(6):583–592. <https://doi.org/10.1111/j.1365-3024.1991.tb00554.x>
- Sanchez F, Garcia J, March F, Cardeñosa N, Coll P, Muñoz C, Auladell C, Prats G (1993) Ultrastructural localization of major hydatid fluid antigens in brood capsules and protoscoleces of *Echinococcus granulosus* of human origin. *Parasite Immunol* 15(8):441–447. <https://doi.org/10.1111/j.1365-3024.1993.tb00629.x>
- Scapin G, Gordon JJ, Sacchettini JC (1992) Refinement of the structure of recombinant rat intestinal fatty acid-binding apoprotein at 1.2-Å resolution. *J Biol Chem* 267(6):4253–4269. [https://doi.org/10.1016/S0021-9258\(19\)50654-8](https://doi.org/10.1016/S0021-9258(19)50654-8)
- Schaap FG, Van der Vusse GJ, Glatz JFC (2002) Evolution of the family of intracellular lipid binding proteins in vertebrates. *Mol Cell Biochem* 239(1):69–77. <https://doi.org/10.1023/A:1020519011939>



- Selkirk ME, Gregory WF, Jenkins RE, Maizels RM (1993) Localization, turnover and conservation of gp 15/400 in different stages of *Brugia malayi*. *Parasitology* 107(4):449–457. <https://doi.org/10.1017/S0031182000067810>
- Silva-Álvarez V, Folle AM, Ramos AL, Zamarreño F, Costabel MD, García-Zepeda E, Salinas G, Córscico B, Ferreira AM (2015) Echinococcus granulosus antigen B: A Hydrophobic Ligand Binding Protein at the host-parasite interface. *Prostaglandins Leukot Essent Fatty Acids* 93:17–23. <https://doi.org/10.1016/j.plefa.2014.09.008>
- Silva-Álvarez V, Folle AM, Ramos AL, Kitano ES, Iwai LK, Corraliza I, Córscico B, Ferreira AM (2016) Echinococcus granulosus antigen B binds to monocytes and macrophages modulating cell response to inflammation. *Parasit Vectors* 9(1):1–17. <https://doi.org/10.1186/S13071-016-1350-7>
- Silva-Álvarez V, Ramos AL, Folle AM, Lagos S, Dee VM, Ferreira AM (2018) Antigen B from Echinococcus granulosus is a novel ligand for C-reactive protein. *Parasit Immunol* 40(9):e12575. <https://doi.org/10.1111/pim.12575>
- Siracusano A, Riganò R, Ortona E, Profumo E, Margutti P, Buttari B, Delunardo F, Teggi A (2008) Immunomodulatory mechanisms during Echinococcus granulosus infection. *Exp Parasitol* 119(4):483–489. <https://doi.org/10.1016/j.exppara.2008.01.016>
- Smathers RL, Petersen DR (2011) The human fatty acid-binding protein family: Evolutionary divergences and functions. *Hum Genomics* 5(3):170–191. <https://doi.org/10.1186/1479-7364-5-3-170>
- Solovyova AS, Meenan N, McDermott L, Garofalo A, Bradley JE, Kennedy MW, Byron O (2003) The polyprotein and FAR lipid binding proteins of nematodes: shape and monomer/dimer states in ligand-free and bound forms. *Eur Biophys J* 32(5):465–476. <https://doi.org/10.1007/s00249-003-0297-8>
- Spiliotis M, Brehm K (2009) Axenic in vitro cultivation of Echinococcus multilocularis metacystode vesicles and the generation of primary cell cultures. *Methods Mol Biol* 470:245–262. [https://doi.org/10.1007/978-1-59745-204-5\\_17](https://doi.org/10.1007/978-1-59745-204-5_17)
- Stevens L, Rooke S, Falzon LC, Machuka EM, Momanyi K, Murungi MK, Njoroge SM, Odinga CO, Ogendo A, Ogola J, Fèvre EM, Blaxter M (2020) The Genome of *Caenorhabditis bovis*. *Curr Biol* 30(6):1023–1031.e4. <https://doi.org/10.1016/j.cub.2020.01.074>
- Storch J, Corsico B (2008) The emerging functions and mechanisms of mammalian fatty acid-binding Proteins. *Annu Rev Nutr* 28:73–95. <https://doi.org/10.1146/ANNUREV.NUTR.27.061406.093710>
- Storch J, McDermott L (2009) Structural and functional analysis of fatty acid-binding proteins. *J Lipid Res* 50(SUPPL.):S126–S131. <https://doi.org/10.1194/jlr.R800084-JLR200>
- Tomlinson LA, Christie JF, Fraser EM, McLaughlin D, McIntosh AE, Kennedy MW (1989) MHC restriction of the antibody repertoire to secretory antigens, and a major allergen, of the nematode parasite *Ascaris*. *J Immunol* 143(7):2349–2356 (<http://www.ncbi.nlm.nih.gov/pubmed/2778320>)
- Tree TIM, Gillespie AJ, Shepley KJ, Blaxter ML, Tuan RS, Bradley JE (1995) Characterisation of an immunodominant glycoprotein antigen of *Onchocerca volvulus* with homologues in other filarial nematodes and *Caenorhabditis elegans*. *Mol Biochem Parasitol* 69(2):185–195. [https://doi.org/10.1016/0166-6851\(94\)00204-Z](https://doi.org/10.1016/0166-6851(94)00204-Z)
- Tritten L, Tam M, Vargas M, Jardim A, Stevenson MM, Keiser J, Geary TG (2017) Excretory/secretory products from the gastrointestinal nematode *Trichuris muris*. *Exp Parasitol* 178:30–36. <https://doi.org/10.1016/j.exppara.2017.05.003>
- Tsai IJ, Zarowiecki M, Holroyd N, Garcarrubio A, Sanchez-Flores A, Brooks KL, Tracey A, Bobes RJ, Frago G, Sciuotto E, Aslett M, Beasley H, Bennett HM, Cai J, Camicia F, Clark R, Cucher M, De Silva N, Day TA, ... Valdes V (2013) The genomes of four tapeworm species reveal adaptations to parasitism. *Nature* 496(7443):57–63. <https://doi.org/10.1038/nature12031>
- Vieira P, Eves-Van Den Akker S, Verma R, Wantoch S, Eisenback JD, Kamo K (2015) The *pratylenchus penetrans* transcriptome as a source for the development of alternative control strategies: Mining for putative genes involved in parasitism and evaluation of in planta RNAi. *PLoS ONE* 10(12):e0144674. <https://doi.org/10.1371/journal.pone.0144674>
- Wang YT, Liu CH, Zhu HL (2016) Fatty acid binding protein (FABP) inhibitors: a patent review (2012–2015). *Expert Opin Ther Pat* 26(7):767–776. <https://doi.org/10.1080/13543776.2016.1182500> Taylor & Francis
- Wang DW, Xu CL, Ding SW, Huang X, Cheng X, Zhang C, Chen C, Xie H (2018) Identification and function of FAR protein family genes from a transcriptome analysis of *Aphelenchoides besseyi*. *Bioinformatics* 34(17):2936–2943. <https://doi.org/10.1093/bioinformatics/bty209>
- Xia Y, Spence HJ, Moore J, Heaney N, McDermott L, Cooper A, Watson DG, Mei B, Komuniecki R, Kennedy MW (2000) The ABA-1 allergen of *Ascaris lumbricoides*: Sequence polymorphism, stage and tissue-specific expression, lipid binding function, and protein biophysical properties. *Parasitology* 120(2):211–224. <https://doi.org/10.1017/S0031182099005363>
- Xu M, Joo HJ, Paik YK (2011) Novel functions of lipid-binding protein 5 in *Caenorhabditis elegans* fat metabolism. *J Biol Chem* 286(32):28111–28118. <https://doi.org/10.1074/jbc.M111.227165>
- Zarlenga DS, Rhoads ML, A1-Yaman FM (1994) A *Taenia crassiceps* cDNA sequence encoding a putative immunodiagnostic antigen for bovine cysticercosis. *Mol Biochem Parasitol* 67(2):215–223. [https://doi.org/10.1016/0166-6851\(94\)00132-4](https://doi.org/10.1016/0166-6851(94)00132-4)
- Zhang W, Li J, Jones MK, Zhang Z, Zhao L, Blair D, McManus DP (2010) The Echinococcus granulosus antigen B gene family comprises at least 10 unique genes in five subclasses which are differentially expressed. *PLoS Negl Trop Dis* 4(8):e784. <https://doi.org/10.1371/journal.pntd.0000784>
- Zhang C, Xie H, Cheng X, Wang DW, Li Y, Xu CL, Huang X (2015) Molecular identification and functional characterization of the fatty acid- and retinoid-binding protein gene Rs-far-1 in the burrowing nematode *Radopholus similis* (Tylenchida: Pratylenchidae). *PLoS ONE* 10(3):e0118414. <https://doi.org/10.1371/journal.pone.0118414>
- Zheng H, Zhang W, Zhang L, Zhang Z, Li J, Lu G, Zhu Y, Wang Y, Huang Y, Liu J, Kang H, Chen J, Wang L, Chen A, Yu S, Gao Z, Jin L, Gu W, Wang Z, ... Wang S (2013a) The genome of the hydatid tapeworm *Echinococcus granulosus*. *Nat Genet* 45(10):1168–1175. <https://doi.org/10.1038/ng.2757>
- Zheng Y, Blair D, Bradley JE (2013b) Phyletic distribution of fatty acid-binding protein genes. *PLoS ONE* 8(10):e77636. <https://doi.org/10.1371/journal.pone.0077636>
- Zhou L, Chen F, Pan H, Ye J, Dong X, Li C, Lin F (2016) Identifying virulence-associated genes using transcriptomic and proteomic association analyses of the plant parasitic nematode *Bursaphelenchus mucronatus*. *Int J Mol Sci* 17(9):1492. <https://doi.org/10.3390/ijms17091492>
- Zimmerman AW, Veerkamp JH (2002) New insights into the structure and function of fatty acid-binding proteins. *Cell Mol Life Sci* 59(7):1096–1116. <https://doi.org/10.1007/S00018-002-8490-Y>