

Project Wildfire

CSPB 4502 – Group 3

Team

- Natalie Dreher
- Ronald Durham
- Grant Fairbairn



Questions Sought to Answer

- Have wildfires become more or less frequent over time?
 - There was no significant correlation between the number of fires and years between 1992 and 2015.
- In this dataset, what is the most common cause of fires?
 - The "Debris Burning" category was the most common cause in this dataset (by a long shot). But that category is quite vague.
- Which cause of fire most commonly leads to the largest fire size?
 - From 1992 to 2015, the data shows that lightning strikes were the identified cause of the largest wildfires.

Data Preparation Work

- We utilized the data set "1.88 Million US Wildfires" for wildfires occurring in the United States between 1992 and 2015.
 - Short, Karen C. 2017. Spatial wildfire occurrence data for the United States, 1992-2015 [FPAFOD20170508]. 4th Edition. Fort Collins, CO: Forest Service Research Data Archive. https://doi.org/10.2737/RDS-2013-0009.4.
- The dataset was massive, with 1.88 million rows of tabulated data, requiring that it be broken into four CSV segments before importing.
- We needed to do some data cleaning as some of the rows were incomplete. We utilized the attributes that were useful in answering our questions and removed excess or ones not applicable to this project.

Tools Used

CODING:

- Jupyter Notebook
- Python

DATA PROCESSING AND REGRESSION:

- Pandas
- Numpy
- Sklearn
- Statsmodels


VISUALIZATION TOOLS:

- Matplotlib
- Pyplot
- Seaborn
- Plotly
- Colorcet
- Holoviews

Techniques Applied


- We utilized visualization tools (Matplotlib and Seaborn) to analyze the data and look for interesting patterns to analyze, and we created a "heatmap" showing the prevalence of fires from the dataset.
- Once we had an idea of what we were looking for, we set out to look for significance in correlations.
- For example, we used the linear regression tools from the Python statsmodel library to analyze whether wildfires were becoming more or less common over time.

Heatmap Visualization



Knowledge Gained

• We did not find a significant trend in the growth or reduction of fires over time. The R-squared value of "Fires" against "Years" was low at 0.036:




OLS Regression Results								
Dep. Variable	Fires		res	R-squared:		0.036		
Model	:	С	LS A	dj. R-so	uared:	-0.	800	
Method	: Least Squares		res	F-statistic:		0.8265		
Date	: Tue	, 29 Nov 20	22 Pro	b (F-sta	atistic):	0.	373	
Time	:	02:55	:59 Lo	og-Like	lihood:	-260	0.00	
No. Observations	:		24		AIC:	52	24.0	
Df Residuals	Df Residuals:		22		BIC: 5		26.4	
Df Model	:		1					
Covariance Type: nonrobust								
	coef	std err		P>I+I	10.0	025	0	9751
Intercept -6.095					-			-
Year 343.	.3443	377.678	0.909	0.373	-439.	911	1126	5.600
Omnibus:	5.718	Durbin-	Watson	: 0	.925			
Prob(Omnibus):	Prob(Omnibus): 0.057 Jarque-Bera		era (JB)	: 3	3.859			
Skew:	0.938	F	rob(JB)	: 0	.145			
Kurtosis:	3.585	C	ond. No	. 5.80e	+05			

Knowledge Gained


• But we did find an interesting result in comparing fire size by cause.


Lightning strikes accounted for the largest average fire (by a wide margin):



Knowledge Gained

 That led us to analyze the average fire size caused by lightning over time, revealing a positive correlation:





Applications

- The largest area of application from studying wildfire data is the potential for learning about prevention and containment.
- Wildfires are a necessary part of the global ecosystem, and lightning strikes are natural causes.
- While lightning strikes cannot be prevented, early detection of strikes in forested areas prone to the largest fires (e.g., the western portion of the United States) could reduce the burn area.

Thank You

https://github.com/rodu4835/CSPB4502_ProjectWildfire

