

Spatial Degradation of Hepatocyte Zonation

Ruler Battery — Per-Set Report

Auto-generated

2026-06-21

Method (read first)

Candidate zonation rulers were built, each scored as a per-cell coordinate ($\text{coord} = \text{mean_z}(\text{PC}) - \text{mean_z}(\text{PP})$, or a learned projection). **Ruler quality is judged ONLY on Paper 2 / healthy-atlas criteria** (8-marker validation gate, healthy PC-PP anticorrelation, split-half reproducibility). The best healthy ruler is frozen, then transferred to the Paper 1 disease cohort, where H1/H2/H3 are TEST results — never used for selection. Sections below are ordered by healthy-ruler quality ($\text{split-half} + \max(0, -\text{anticorr})$). H1 = donor-level zonation collapse; H2 = zonal-slope loss; H3 = de-zonation/plasticity coupling.

Selection decision.

SELECTED set: expanded_curated

Reason: best HEALTHY-ruler quality among primary candidates.

Healthy metrics used (NOT disease collapse):

- validation: 8/8 markers correct
- healthy PC-PP anticorrelation: -0.479 (want strongly negative)
- healthy split-half reproducibility: +0.670
- size: 26 PC + 23 PP

WARNING: selected using Paper 2 / healthy-ruler criteria ONLY, NOT Paper 1 disease effect (disease is t

unsupervised_p2

primary_candidate


Candidate zonation ruler. **Genes:** 5177 PC + 3302 PP.

Validation (8-marker healthy gate) & controls. Markers correct: 8/8 (PASS). Healthy PC-PP anticorrelation = -0.471 (want strongly negative); split-half reproducibility = +0.691; coherence PC/PP = +0.076/+0.052.

H1 (donor-level collapse, TRANSFER). coord-spread vs stage rho = -0.504 (perm p = 0.0005); PC-PP anticorr trend rho = +0.635 (perm p = 0.0005).

H2 (zonal slope-loss, held-out). 100 genes tested; 83% weaken; median trend rho = -0.260.

H3 (de-zonation ↔ plasticity, donor-level). mean rho = -0.009; 23/47 donors >0; sign-test p = 1.



expanded_curated

primary_candidate


paper2_landmark plus curated extra anchors (26+23), de-duplicated. **Genes:** 26 PC + 23 PP.

Validation (8-marker healthy gate) & controls. Markers correct: 8/8 (PASS). Healthy PC-PP anticorrelation = -0.479 (want strongly negative); split-half reproducibility = +0.670; coherence PC/PP = +0.155/+0.065.

H1 (donor-level collapse, TRANSFER). coord-spread vs stage rho = -0.434 (perm p = 0.003); PC-PP anticorr trend rho = +0.612 (perm p = 0.0005).

H2 (zonal slope-loss, held-out). 49 genes tested; 90% weaken; median trend rho = -0.258.

H3 (de-zonation ↔ plasticity, donor-level). mean rho = +0.025; 33/47 donors >0; sign-test p = 0.0079.



selected_frozen

primary_candidate


The auto-frozen primary ruler (a copy of the best signature set by healthy metrics). **Genes:** 26 PC + 23 PP.

Validation (8-marker healthy gate) & controls. Markers correct: 8/8 (PASS). Healthy PC-PP anticorrelation = -0.479 (want strongly negative); split-half reproducibility = +0.670; coherence PC/PP = +0.155/+0.065.

H1 (donor-level collapse, TRANSFER). coord-spread vs stage rho = -0.434 (perm p = 0.003); PC-PP anticorr trend rho = +0.612 (perm p = 0.0005).

H2 (zonal slope-loss, held-out). 49 genes tested; 90% weaken; median trend rho = -0.258.

H3 (de-zonation ↔ plasticity, donor-level). mean rho = +0.025; 33/47 donors >0; sign-test p = 0.0079.



unsupervised_top100

primary_candidate


The 100+100 strongest +/- loading genes of the unsupervised PCA axis. **Genes:** 100 PC + 100 PP.

Validation (8-marker healthy gate) & controls. Markers correct: 8/8 (PASS). Healthy PC-PP anticorrelation = -0.292 (want strongly negative); split-half reproducibility = +0.821; coherence PC/PP = +0.088/+0.056.

H1 (donor-level collapse, TRANSFER). coord-spread vs stage rho = -0.608 (perm p = 0.0005); PC-PP anticorr trend rho = +0.613 (perm p = 0.0005).

H2 (zonal slope-loss, held-out). 200 genes tested; 92% weaken; median trend rho = -0.324.

H3 (de-zonation ↔ plasticity, donor-level). mean rho = +0.031; 36/47 donors >0; sign-test p = 0.00035.



unsupervised_top50

primary_candidate


The 50+50 strongest +/- loading genes of the unsupervised PCA axis, scored as a signature. **Genes:** 50 PC + 50 PP.

Validation (8-marker healthy gate) & controls. Markers correct: 8/8 (PASS). Healthy PC-PP anticorrelation = -0.306 (want strongly negative); split-half reproducibility = +0.785; coherence PC/PP = +0.121/+0.082.

H1 (donor-level collapse, TRANSFER). coord-spread vs stage rho = -0.617 (perm p = 0.0005); PC-PP anticorr trend rho = +0.607 (perm p = 0.0005).

H2 (zonal slope-loss, held-out). 100 genes tested; 96% weaken; median trend rho = -0.372.

H3 (de-zonation ↔ plasticity, donor-level). mean rho = +0.027; 38/47 donors >0; sign-test p = 2.5e-05.



unsupervised

primary_candidate


Label-free PCA porto-central axis learned on Paper 1 HEALTHY cells (sensitivity version). **Genes:** 4585 PC + 3903 PP.

Validation (8-marker healthy gate) & controls. Markers correct: 7/8 (PASS). Healthy PC-PP anticorrelation = -0.306 (want strongly negative); split-half reproducibility = +0.781; coherence PC/PP = +0.121/+0.082.

H1 (donor-level collapse, TRANSFER). coord-spread vs stage rho = -0.562 (perm p = 0.0005); PC-PP anticorr trend rho = +0.607 (perm p = 0.0005).

H2 (zonal slope-loss, held-out). 100 genes tested; 96% weaken; median trend rho = -0.375.

H3 (de-zonation ↔ plasticity, donor-level). mean rho = +0.016; 32/47 donors >0; sign-test p = 0.019.



paper2_landmark

primary_candidate


Paper 2's exact 20+20 hepatocyte landmark genes (Hepatocyte-{PC,PP}-LM.csv), the genes Paper 2 itself uses to assign per-cell snRNA zonation (eta method). **Genes:** 20 PC + 20 PP.

Validation (8-marker healthy gate) & controls. Markers correct: 8/8 (PASS). Healthy PC-PP anticorrelation = -0.455 (want strongly negative); split-half reproducibility = +0.631; coherence PC/PP = +0.156/+0.062.

H1 (donor-level collapse, TRANSFER). coord-spread vs stage rho = -0.380 (perm p = 0.0085); PC-PP anticorr trend rho = +0.608 (perm p = 0.0005).

H2 (zonal slope-loss, held-out). 40 genes tested; 78% weaken; median trend rho = -0.229.

H3 (de-zonation $\leftrightarrow$ plasticity, donor-level). mean rho = +0.024; 33/47 donors >0; sign-test p = 0.0079.



paper2_top50

primary_candidate


Top 50+50 genes by —center-of-mass— among qj0.05 zonated genes (supplementary table 8). **Genes:** 50 PC + 50 PP.

Validation (8-marker healthy gate) & controls. Markers correct: 8/8 (PASS). Healthy PC-PP anticorrelation = -0.404 (want strongly negative); split-half reproducibility = +0.573; coherence PC/PP = +0.051/+0.034.

H1 (donor-level collapse, TRANSFER). coord-spread vs stage rho = -0.316 (perm p = 0.035); PC-PP anticorr trend rho = +0.564 (perm p = 0.0005).

H2 (zonal slope-loss, held-out). 88 genes tested; 73% weaken; median trend rho = -0.204.

H3 (de-zonation ↔ plasticity, donor-level). mean rho = +0.010; 33/47 donors >0; sign-test p = 0.0079.




Top 100+100 zonated genes by —center-of-mass—. **Genes:** 100 PC + 100 PP.

Validation (8-marker healthy gate) & controls. Markers correct: 8/8 (PASS). Healthy PC-PP anticorrelation = -0.273 (want strongly negative); split-half reproducibility = +0.563; coherence PC/PP = +0.028/+0.020.

H1 (donor-level collapse, TRANSFER). coord-spread vs stage rho = -0.284 (perm p = 0.06); PC-PP anticorr trend rho = +0.423 (perm p = 0.003).

H2 (zonal slope-loss, held-out). 186 genes tested; 59% weaken; median trend rho = -0.066.

H3 (de-zonation ↔ plasticity, donor-level). mean rho = +0.011; 28/47 donors >0; sign-test p = 0.24.



core_curated

interpretability_only


13+8 hand-curated, highly interpretable canonical anchors (GLUL, CYP2E1, ASS1, HAL, ...). **Genes:** 13 PC + 8 PP.

Validation (8-marker healthy gate) & controls. Markers correct: 8/8 (PASS). Healthy PC-PP anticorrelation = -0.278 (want strongly negative); split-half reproducibility = +0.537; coherence PC/PP = +0.185/+0.104.

H1 (donor-level collapse, TRANSFER). coord-spread vs stage rho = -0.541 (perm p = 0.0005); PC-PP anticorr trend rho = +0.562 (perm p = 0.0005).

H2 (zonal slope-loss, held-out). 21 genes tested; 100% weaken; median trend rho = -0.339.

H3 (de-zonation $\leftrightarrow$ plasticity, donor-level). mean rho = +0.033; 36/47 donors >0; sign-test p = 0.00035.



unsupervised_lasso

robustness


Lasso (L1) sparse ruler distilling the Paper-2-trained PCA axis (clean/external training). **Genes:** 126 PC + 149 PP.

Validation (8-marker healthy gate) & controls. Markers correct: 8/8 (PASS). Healthy PC-PP anticorrelation = -0.184 (want strongly negative); split-half reproducibility = +0.629; coherence PC/PP = +0.022/+0.020.

H1 (donor-level collapse, TRANSFER). coord-spread vs stage rho = -0.505 (perm p = 0.0005); PC-PP anticorr trend rho = +0.602 (perm p = 0.0005).

H2 (zonal slope-loss, held-out). 275 genes tested; 69% weaken; median trend rho = -0.144.

H3 (de-zonation ↔ plasticity, donor-level). mean rho = -0.017; 22/47 donors >0; sign-test p = 0.77.



unsupervised_elasticnet

robustness


Elastic-net (L1+L2) sparse ruler distilling the Paper-2-trained PCA axis (clean). **Genes:** 325 PC + 281 PP.

Validation (8-marker healthy gate) & controls. Markers correct: 8/8 (PASS). Healthy PC-PP anticorrelation = -0.068 (want strongly negative); split-half reproducibility = +0.689; coherence PC/PP = +0.016/+0.014.

H1 (donor-level collapse, TRANSFER). coord-spread vs stage rho = -0.492 (perm p = 0.0005); PC-PP anticorr trend rho = +0.483 (perm p = 0.0025).

H2 (zonal slope-loss, held-out). 606 genes tested; 67% weaken; median trend rho = -0.121.

H3 (de-zonation ↔ plasticity, donor-level). mean rho = -0.020; 21/47 donors >0; sign-test p = 0.56.




Top 250+250 zonated genes by —center-of-mass—. **Genes:** 250 PC + 250 PP.

Validation (8-marker healthy gate) & controls. Markers correct: 8/8 (PASS). Healthy PC–PP anticorrelation = +0.041 (want strongly negative); split-half reproducibility = +0.554; coherence PC/PP = +0.015/+0.011.

H1 (donor-level collapse, TRANSFER). coord-spread vs stage rho = -0.265 (perm p = 0.067); PC–PP anticorr trend rho = +0.250 (perm p = 0.081).

H2 (zonal slope-loss, held-out). 483 genes tested; 53% weaken; median trend rho = -0.011.

H3 (de-zonation ↔ plasticity, donor-level). mean rho = +0.006; 26/47 donors >0; sign-test p = 0.56.



paper2_full

exploratory


All significant zonated hepatocyte genes ($q \leq 0.05$, $\max\text{-expr} \leq 1e-4$), split by COM sign (1624+447). **Genes:** 1624 PC + 447 PP.

Validation (8-marker healthy gate) & controls. Markers correct: 8/8 (PASS). Healthy PC-PP anticorrelation = +0.556 (want strongly negative); split-half reproducibility = +0.484; coherence PC/PP = +0.005/+0.009.

H1 (donor-level collapse, TRANSFER). coord-spread vs stage $\rho = -0.042$ (perm $p = 0.77$); PC-PP anticorr trend $\rho = -0.040$ (perm $p = 0.79$).

H2 (zonal slope-loss, held-out). 2035 genes tested; 41% weaken; median trend $\rho = +0.037$.

H3 (de-zonation $\leftrightarrow$ plasticity, donor-level). mean $\rho = +0.000$; 25/47 donors >0 ; sign-test $p = 0.77$.



full

exploratory


Legacy transcriptome-wide unweighted signature mean (main pipeline). **Genes:** 0 PC + 0 PP.

Validation (8-marker healthy gate) & controls. Markers correct: 8/8 (PASS). Healthy PC-PP anticorrelation = +0.532 (want strongly negative); split-half reproducibility = +0.464; coherence PC/PP = +0.008/+0.010.

H1 (donor-level collapse, TRANSFER). coord-spread vs stage rho = +0.021 (perm p = 0.88); PC-PP anticorr trend rho = +0.007 (perm p = 0.97).

H2 (zonal slope-loss, held-out). 1637 genes tested; 42% weaken; median trend rho = +0.034.

H3 (de-zonation ↔ plasticity, donor-level). mean rho = +0.002; 26/47 donors >0; sign-test p = 0.56.



supervised

exploratory

Whole-transcriptome multinomial logistic regression trained on Paper 2 zone labels (landmarks excluded).
Genes: 4504 PC + 3935 PP.

Validation (8-marker healthy gate) & controls. Markers correct: 8/8 (PASS). Healthy PC-PP anticorrelation = -0.005 (want strongly negative); split-half reproducibility = +0.452; coherence PC/PP = +0.064/+0.014.

H1 (donor-level collapse, TRANSFER). coord-spread vs stage rho = +0.122 (perm p = 0.4); PC-PP anticorr trend rho = +0.569 (perm p = 0.0005).

H2 (zonal slope-loss, held-out). 100 genes tested; 87% weaken; median trend rho = -0.169.

H3 (de-zonation ↔ plasticity, donor-level). mean rho = +0.003; 27/47 donors >0; sign-test p = 0.38.

