

Experimental Design

Note 3-1

Model Adequacy Checking

Keunbaik Lee

Sungkyunkwan University

Model checking and diagnostics I

- **Checking assumptions** is important

- Have we fit the right model?
- Normality
- Independence
- Constant variance

$$y_{ij} = (\bar{y}_{..} + (\bar{y}_{i.} - \bar{y}_{..})) + (y_{ij} - \bar{y}_{i.})$$

$$y_{ij} = \hat{y}_{ij} + e_{ij}$$

$$\text{observed} = \text{predicted} + \text{residual}$$

- Note that the predicted response at treatment i is $\hat{y}_{ij} = \bar{y}_{i.}$
- Diagnostics use predicted responses and residuals.

Model checking and diagnostics II

- Normality
 - Histogram of residuals
 - Normal probability plot / QQ plot
 - Shapiro-Wilk Test
- Constant Variance
 - Plot $\hat{\epsilon}_{ij}$ vs \hat{y}_{ij} (residual plot)
 - Bartlett's or Levene's Test
- Independence
 - Plot $\hat{\epsilon}_{ij}$ vs time/space
 - Plot $\hat{\epsilon}_{ij}$ vs variable of interest
- Outliers

Normality Checking in the ANOVA

- Examination of residuals

$$\begin{aligned}e_{ij} &= y_{ij} - \hat{y}_{ij} \\ &= y_{ij} - \bar{y}_{i.}\end{aligned}$$

- Residual plots are very useful - e.g., Q-Q plot
- Shapiro-Wilk, Kolmogorov-Smirnov, Anderson-Darling Tests

Outliers Checking

- Use standardized residuals to check if there is outliers

$$d_{ij} = \frac{e_{ij}}{\sqrt{MSE}}$$

- > 3 or < -3 is a potential outlier
- Be careful for removing outliers

Constant variance checking I

- In some experiments, error variance (σ_i^2) depends on the mean response

$$E(y_{ij}) = \mu_i = \mu + \tau_i$$

So the constant variance assumption is violated.

- Size of error (residual) depends on mean response (predicted value)
- Residual plot
 - Plot $\hat{\epsilon}_{ij}$ vs \hat{y}_{ij}
 - Is the range constant for different levels of \hat{y}_{ij}
 - More formal tests: Bartlett's Test, Modified Levene's Test.
- Modified Levene's Test
 - For each fixed i , calculate the median m_i of $y_{i1}, y_{i2}, \dots, y_{in_i}$.

Constant variance checking II

- Compute the absolute deviation of observation from sample median:

$$d_{ij} = |y_{ij} - m_i|$$

for $i = 1, 2, \dots, a$ and $j = 1, 2, \dots, n_i$.

- Apply ANOVA to the deviations: d_{ij}
- Use the usual ANOVA F -statistic for testing
 $H_0 : \sigma_1^2 = \dots = \sigma_a^2$

See Tensile.SAS.

Non-constant Variance: Impact and Remedy I

- Why concern?
 - Comparison of treatments depends on MSE
 - Incorrect intervals and comparison results
- Variance-Stabilizing Transformations
 - Common transformations

$$\sqrt{x}, \log(x), 1/x, \arcsin(\sqrt{x}), \text{ and } 1/\sqrt{x}$$

- Box-Cox transformations
 - approximate the relationship $\sigma_i = \theta\mu_i^\beta$, then the transformation is $X^{1-\beta}$
 - use maximum likelihood principle
- Ideas for finding proper transformations

Non-constant Variance: Impact and Remedy II

- Consider response Y with mean $E(Y) = \mu$ and variance $\text{var}(Y) = \sigma^2$.
- That σ^2 depends on μ leads to nonconsistent variances for different μ .
- Let f be a transformation and $\tilde{Y} = f(Y)$. What is the mean and variance of \tilde{Y} ?
- Approximate $f(Y)$ by a linear function (Delta Method):

$$f(Y) \approx f(\mu) + (Y - \mu)f'(\mu)$$

Then

$$\begin{aligned}\text{Mean: } \tilde{\mu} &= E(\tilde{Y}) = E(f(Y)) \approx E(f(\mu)) + E((Y - \mu)f'(\mu)) \\ &= f(\mu)\end{aligned}$$

$$\text{Variance: } \tilde{\sigma}^2 = \text{var}(\tilde{Y}) \approx [f'(\mu)]^2 \text{var}(Y) = [f'(\mu)]^2 \sigma^2$$

Non-constant Variance: Impact and Remedy III

- f is a good transformation if $\tilde{\sigma}^2$ does not depend on $\tilde{\mu}$ anymore. So, \tilde{Y} has constant variance for different $f(\mu)$.
- Transformations
 - Suppose σ^2 is a function of μ , that is $\sigma^2 = g(\mu)$
 - Want to find transformation f such that $\tilde{Y} = f(Y)$ has constant variance: $\text{var}(\tilde{Y})$ does not depend on μ .
 - Have shown $\text{var}(\tilde{Y}) \approx [f'(\mu)]^2 \sigma^2 \approx [f'(\mu)]^2 g(\mu)$
 - Want to choose f such that $[f'(\mu)]^2 g(\mu) \approx c$

Distribution	Variance	Transformation
Poisson	$g(\mu) = \mu$	$f(\mu) = \int \frac{1}{\sqrt{\mu}} d\mu \longrightarrow f(X) = \sqrt{X}$
Binomial	$g(\mu) = \mu(1 - \mu)$	$f(\mu) = \int \frac{1}{\sqrt{\mu(1-\mu)}} d\mu \longrightarrow f(X) = \arcsin(\sqrt{X})$
Box-Cox	$g(\mu) = \mu^{2\beta}$	$f(\mu) = \int \mu^{-\beta} d\mu \longrightarrow f(X) = X^{1-\beta}$
Box-Cox	$g(\mu) = \mu^2$	$f(\mu) = \int \frac{1}{\mu} d\mu \longrightarrow f(X) = \log X$

Non-constant Variance: Impact and Remedy IV

- Identify Box-Cox Transformation using Data: Approximate Method

- From the previous slide, if $\sigma_i = \theta \mu_i^\beta$, the transformation is

$$f(Y) = \begin{cases} Y^{1-\beta}, & \beta \neq 1; \\ \log Y, & \beta = 1. \end{cases}$$

So it is crucial to estimate β based on data y_{ij} , $i = 1, \dots, a$.

- We have $\log \sigma_i = \log \theta + \beta \log \mu_i$.
- Let s_i and $\bar{y}_{i\cdot}$ be the sample standard deviations and means. Because $\hat{\sigma}_i = s_i$ and $\hat{\mu}_i = \bar{y}_{i\cdot}$, approximately,

$$\log s_i = \text{constant} + \beta \log \bar{y}_{i\cdot},$$

where $i = 1, \dots, a$.

- We can plot $\log s_i$ against $\log \bar{y}_{i\cdot}$, fit a straight line and use the slope to estimate β .

Non-constant Variance: Impact and Remedy V

- Identify Box-Cox Transformation: Formal Method
 - For a fixed λ , perform analysis of variance on

$$y_{ij}(\lambda) = \begin{cases} \frac{y_{ij}^\lambda - 1}{\lambda \dot{y}^{\lambda-1}}, & \lambda \neq 0; \\ \dot{y} \log y_{ij}, & \lambda = 0, \end{cases}$$

where $\dot{y} = \prod_{i=1}^a \prod_{j=1}^{n_i} y_{ij}^{1/N}$.

Notice that we cannot select the value of λ by directly comparing the SSEs from the ANOVA on y^λ because for each value of λ , the SSEs are measured on different scales.

- Step 1 generates a transformed data $y_{ij}(\lambda)$. Apply ANOVA to the new data and obtain SS_E . Because SS_E depends on λ , it is denoted by $SS_E(\lambda)$.
- Repeat 1 and 2 for various λ in an interval, e.g., $[-2, 2]$, and record $SS_E(\lambda)$

Non-constant Variance: Impact and Remedy VI

- Find λ_0 which minimizes $SS_E(\lambda)$ and pick up a meaningful λ in the neighborhood of λ_0 . Denote it again by λ .
- Now the selected transformation is:

$$f(y_{ij}) = \begin{cases} y_{ij}^{\lambda_0}, & \text{if } \lambda_0 \neq 0; \\ \log y_{ij}, & \text{if } \lambda_0 = 0. \end{cases}$$

See Transformation.SAS.

Nonparametric methods for ANOVA I

H_0 : a treatments are equal vs H_a : at least one not equal.
(But normality assumption is unsatisfied)

■ Kruskal-Wallis Test

- Rank the observations y_{ij} in ascending order
- Replace each observation by its rank R_{ij} (assign average for tied observations)
- Test statistic

$$H = \frac{1}{S^2} \left[\sum_{i=1}^a \frac{R_{i\cdot}^2}{n_i} - \frac{N(N+1)^2}{4} \right] \approx \chi_{a-1}^2$$

where $S^2 = \frac{1}{N-1} \left[\sum_{i=1}^a \sum_{j=1}^{n_i} R_{ij}^2 - \frac{N(N+1)^2}{4} \right]$

Nonparametric methods for ANOVA II

- Decision Rule: reject H_0 if $H > \chi^2_{\alpha, a-1}$.

See Nonparametric.SAS.