

A treatise on phenomenological models of surface second-harmonic generation from crystalline surfaces

Bernardo S. Mendoza and Sean M. Anderson

December 18, 2015

Contents

1	SHG yield per Sipe, Moss, and van Driel [1]	2
1.1	\mathcal{R}_{pP}	2
1.2	\mathcal{R}_{pS}	4
1.3	\mathcal{R}_{sP}	5
1.4	Summary	6
2	SHG yield per Mizrahi and Sipe [2]	7
2.1	\mathcal{R}_{pP}	7
2.2	\mathcal{R}_{pS}	7
2.3	\mathcal{R}_{sP}	7
2.4	Summary	7
3	SHG yield per Mendoza [3]	7
3.1	\mathcal{R}_{pP}	7
3.2	\mathcal{R}_{pS}	7
3.3	\mathcal{R}_{sP}	7
3.4	Summary	7

1 SHG yield per Sipe, Moss, and van Driel [1]

In this treatment, they define the following for all polarizations;

$$\begin{aligned} f_s &= \frac{\kappa}{n\tilde{\omega}} = \frac{\kappa}{\sqrt{\epsilon(\omega)}\tilde{\omega}}, \\ f_c &= \frac{w}{n\tilde{\omega}} = \frac{w}{\sqrt{\epsilon(\omega)}\tilde{\omega}}, \\ f_s^2 + f_c^2 &= 1, \end{aligned} \tag{1}$$

where

$$\begin{aligned} \kappa &= \tilde{\omega} \sin \theta, \\ w_0 &= \sqrt{\tilde{\omega} - \kappa^2} = \tilde{\omega} \cos \theta, \end{aligned} \tag{2}$$

$$w = \sqrt{\tilde{\omega}\epsilon(\omega) - \kappa^2} = \tilde{\omega}k_z(\omega). \tag{3}$$

From this point on, all capital letters and symbols indicate evaluation at 2ω . Common to all three polarization cases studied here, we require the nonzero components for the (111) face for crystals with C_{3v} symmetry,

$$\begin{aligned} \delta_{11} &= \chi^{xxx} = -\chi^{xyy} = -\chi^{yyx}, \\ \delta_{15} &= \chi^{xxz} = \chi^{yyz}, \\ \delta_{31} &= \chi^{zxx} = \chi^{zyy}, \\ \delta_{33} &= \chi^{zzz}. \end{aligned} \tag{4}$$

Lastly, the remaining quantities that will be needed for all three cases are

$$\begin{aligned} A_p &= \frac{4\pi\tilde{\Omega}\sqrt{\epsilon(2\omega)}}{W_0\epsilon(2\omega) + W}, \\ A_s &= \frac{4\pi\tilde{\Omega}}{W_0 + W}. \end{aligned} \tag{5}$$

1.1 \mathcal{R}_{pP}

For the (111) face ($m = 3$), we have

$$\frac{E^{(2\omega)}(\parallel, \parallel)}{E_p^2 A_p} = a_{\parallel, \parallel} + c_{\parallel, \parallel}^{(3)} \cos 3\phi. \tag{6}$$

We extract these coefficients from Table V, noting that $\Gamma = \gamma = 0$ as we are only interested in the surface contribution,

$$\begin{aligned} a_{\parallel, \parallel} &= i\tilde{\Omega}F_s\epsilon(2\omega)\delta_{31} + i\tilde{\Omega}\epsilon(2\omega)F_sf_s^2(\delta_{33} - \delta_{31}) - 2i\tilde{\Omega}f_sf_cF_c\delta_{15}, \\ c_{\parallel, \parallel}^{(3)} &= -i\tilde{\Omega}F_cf_c^2\delta_{11}. \end{aligned}$$

We substitute these in Eq. (6),

$$\begin{aligned} \frac{E^{(2\omega)}(\parallel, \parallel)}{E_p^2 A_p} &= i\tilde{\Omega} F_s \epsilon(2\omega) \delta_{31} + i\tilde{\Omega} \epsilon(2\omega) F_s f_s^2 (\delta_{33} - \delta_{31}) \\ &\quad - 2i\tilde{\Omega} f_s f_c F_c \delta_{15} - i\tilde{\Omega} F_c f_c^2 \delta_{11} \cos 3\phi \end{aligned}$$

and reduce (omitting the (\parallel, \parallel) notation),

$$\begin{aligned} \frac{E^{(2\omega)}}{E_p^2} &= A_p i\tilde{\Omega} [F_s \epsilon(2\omega) (\delta_{31} + f_s^2 (\delta_{33} - \delta_{31})) - f_c F_c (2f_s \delta_{15} + f_c \delta_{11} \cos 3\phi)] \\ &= A_p i\tilde{\Omega} [F_s \epsilon(2\omega) (f_s^2 \delta_{33} + (1 - f_s^2) \delta_{31}) - f_c F_c (2f_s \delta_{15} + f_c \delta_{11} \cos 3\phi)] \\ &= A_p i\tilde{\Omega} [F_s \epsilon(2\omega) (f_s^2 \delta_{33} + f_c^2 \delta_{31}) - f_c F_c (2f_s \delta_{15} + f_c \delta_{11} \cos 3\phi)]. \end{aligned}$$

As every term has an $f_i^2 F_i$, we can factor out the common

$$\frac{1}{\tilde{\omega}^2 \tilde{\Omega} \epsilon(\omega) \sqrt{\epsilon(2\omega)}}$$

factor after substituting the appropriate terms from Eq. (1),

$$\begin{aligned} \frac{E^{(2\omega)}}{E_p^2} &= \frac{A_p i}{\epsilon(\omega) \sqrt{\epsilon(2\omega)} \tilde{\omega}^2} [K \epsilon(2\omega) (\kappa^2 \delta_{33} + w^2 \delta_{31}) - wW (2\kappa \delta_{15} + w \delta_{11} \cos 3\phi)] \\ &= \frac{A_p i \tilde{\Omega}}{\epsilon(\omega) \sqrt{\epsilon(2\omega)}} [\sin \theta \epsilon(2\omega) (\sin^2 \theta \delta_{33} + k_z^2(\omega) \delta_{31}) \\ &\quad - k_z(\omega) k_z(2\omega) (2 \sin \theta \delta_{15} + k_z(\omega) \delta_{11} \cos 3\phi)] \\ &= \frac{A_p i \tilde{\Omega}}{\epsilon(\omega) \sqrt{\epsilon(2\omega)}} [\sin \theta \epsilon(2\omega) (\sin^2 \theta \chi^{zzz} + k_z^2(\omega) \chi^{zzx}) \\ &\quad - k_z(\omega) k_z(2\omega) (2 \sin \theta \chi^{xxz} + k_z(\omega) \chi^{xxx} \cos 3\phi)]. \end{aligned}$$

We substitute Eq. (5) to complete the expression,

$$\begin{aligned} \frac{E^{(2\omega)}}{E_p^2} &= \frac{4i\pi \tilde{\Omega}^2}{\epsilon(\omega) (W_0 \epsilon(2\omega) + W)} [\dots] \\ &= \frac{4i\pi \tilde{\Omega}}{\epsilon(\omega) (\epsilon(2\omega) \cos \theta + k_z(2\omega))} [\dots] \\ &= \frac{4i\pi \tilde{\omega}}{\cos \theta} \frac{1}{\epsilon(\omega)} \frac{2 \cos \theta}{\epsilon(2\omega) \cos \theta + k_z(2\omega)} [\dots]. \end{aligned}$$

However, our interest lies in \mathcal{R}_{pP} which is calculated as

$$\mathcal{R}_{pP} = \frac{I_p(2\omega)}{I_p^2(\omega)} = \frac{2\pi}{c} \left| \frac{E^{(2\omega)}(\parallel, \parallel)}{E_p^2} \right|^2,$$

and we can finally complete the expression,

$$\begin{aligned}
\mathcal{R}_{pP} &= \frac{2\pi}{c} \left| \frac{4i\pi\tilde{\omega}}{\cos\theta} \frac{1}{\epsilon(\omega)} \frac{2\cos\theta}{\epsilon(2\omega)\cos\theta + k_z(2\omega)} r_{pP} \right|^2 \\
&= \frac{32\pi^3\tilde{\omega}^2}{c\cos^2\theta} |t_p(\omega)T_p(2\omega)r_{pP}|^2 \\
&= \frac{32\pi^3\omega^2}{c^3\cos^2\theta} |t_p(\omega)T_p(2\omega)r_{pP}|^2,
\end{aligned} \tag{7}$$

where

$$\begin{aligned}
t_p(\omega) &= \frac{1}{\epsilon(\omega)}, \\
T_p(2\omega) &= \frac{2\cos\theta}{\epsilon(2\omega)\cos\theta + k_z(2\omega)}, \\
r_{pP} &= \sin\theta\epsilon(2\omega)(\sin^2\theta\chi^{zzz} + k_z^2(\omega)\chi^{zxx}) \\
&\quad - k_z(\omega)k_z(2\omega)(2\sin\theta\chi^{xxz} + k_z(\omega)\chi^{xxx}\cos 3\phi).
\end{aligned}$$

1.2 \mathcal{R}_{pS}

We follow the same procedure as above. For the (111) face ($m = 3$),

$$\frac{E^{(2\omega)}(\parallel, \perp)}{E_p^2 A_s} = b_{\parallel, \perp}^{(3)} \sin 3\phi, \tag{8}$$

and we extract the relevant coefficient from Table V with $\Gamma = \gamma = 0$,

$$b_{\parallel, \perp}^{(3)} = i\tilde{\Omega}f_c^2\delta_{11}.$$

Substituting this coefficient and Eq. (5) into Eq. (8),

$$\begin{aligned}
\frac{E^{(2\omega)}(\parallel, \perp)}{E_p^2} &= A_s i\tilde{\Omega}f_c^2\delta_{11} \sin 3\phi \\
&= \frac{A_s i\tilde{\Omega}}{\tilde{\omega}^2\epsilon(\omega)} \omega^2 \delta_{11} \sin 3\phi \\
&= \frac{A_s i\tilde{\Omega}}{\epsilon(\omega)} k_z^2(\omega) \delta_{11} \sin 3\phi \\
&= \frac{A_s i\tilde{\Omega}}{\epsilon(\omega)} k_z^2(\omega) \chi^{xxx} \sin 3\phi \\
&= \frac{4i\pi\tilde{\Omega}^2}{W_0 + W} \frac{1}{\epsilon(\omega)} k_z^2(\omega) \chi^{xxx} \sin 3\phi \\
&= 4i\pi\tilde{\Omega} \frac{1}{\epsilon(\omega)} \frac{1}{\cos\theta + k_z(2\omega)} k_z^2(\omega) \chi^{xxx} \sin 3\phi \\
&= \frac{4i\pi\omega}{c\cos\theta} \frac{1}{\epsilon(\omega)} \frac{2\cos\theta}{\cos\theta + k_z(2\omega)} k_z^2(\omega) \chi^{xxx} \sin 3\phi
\end{aligned}$$

As before, we must calculate

$$\mathcal{R}_{pS} = \frac{2\pi}{c} \left| \frac{E^{(2\omega)}(\parallel, \perp)}{E_s^2} \right|^2,$$

to obtain the final expression,

$$\begin{aligned} \mathcal{R}_{pS} &= \frac{2\pi}{c} \left| \frac{4i\pi\omega}{c \cos \theta} \frac{1}{\epsilon(\omega)} \frac{2 \cos \theta}{\cos \theta + k_z(2\omega)} k_z^2(\omega) \chi^{xxx} \sin 3\phi \right|^2 \\ &= \frac{32\pi^3 \omega^2}{c^3 \cos^2 \theta} \left| \frac{1}{\epsilon(\omega)} \frac{2 \cos \theta}{\cos \theta + k_z(2\omega)} k_z^2(\omega) \chi^{xxx} \sin 3\phi \right|^2 \\ &= \frac{32\pi^3 \omega^2}{c^3 \cos^2 \theta} |t_p(\omega) T_s(2\omega) k_z^2(\omega) r_{pS}|^2, \end{aligned} \quad (9)$$

where

$$\begin{aligned} t_p(\omega) &= \frac{1}{\epsilon(\omega)}, \\ T_s(2\omega) &= \frac{2 \cos \theta}{\cos \theta + k_z(2\omega)}, \\ r_{pS} &= k_z^2(\omega) \chi^{xxx} \sin 3\phi. \end{aligned}$$

1.3 \mathcal{R}_{sP}

We follow the same procedure as above for the final polarization case. For the (111) face ($m = 3$),

$$\frac{E^{(2\omega)}(\perp, \parallel)}{E_s^2 A_p} = a_{\perp, \parallel} + c_{\perp, \parallel}^{(3)} \cos 3\phi, \quad (10)$$

and we extract the relevant coefficients from Table V with $\Gamma = \gamma = 0$,

$$\begin{aligned} a_{\perp, \parallel} &= i\tilde{\Omega} F_s \epsilon(2\omega) \delta_{31}, \\ c_{\perp, \parallel}^{(3)} &= i\tilde{\Omega} F_c \delta_{11}. \end{aligned}$$

Substituting this coefficient and Eq. (5) into Eq. (10),

$$\begin{aligned}
\frac{E^{(2\omega)}(\perp, \parallel)}{E_s^2} &= A_p(i\tilde{\Omega}F_s\epsilon(2\omega)\delta_{31} + i\tilde{\Omega}F_c\delta_{11}\cos 3\phi) \\
&= A_pi\tilde{\Omega}(F_s\epsilon(2\omega)\delta_{31} + F_c\delta_{11}\cos 3\phi) \\
&= \frac{A_pi\tilde{\Omega}}{\sqrt{\epsilon(2\omega)}}(\sin\theta\epsilon(2\omega)\delta_{31} + k_z(2\omega)\delta_{11}\cos 3\phi) \\
&= \frac{A_pi\tilde{\Omega}}{\sqrt{\epsilon(2\omega)}}(\sin\theta\epsilon(2\omega)\chi^{zxx} + k_z(2\omega)\chi^{xxx}\cos 3\phi) \\
&= \frac{4i\pi\tilde{\Omega}^2}{W_0\epsilon(2\omega) + W}(\sin\theta\epsilon(2\omega)\chi^{zxx} + k_z(2\omega)\chi^{xxx}\cos 3\phi) \\
&= \frac{4i\pi\tilde{\Omega}}{\epsilon(2\omega)\cos\theta + k_z(2\omega)}(\sin\theta\epsilon(2\omega)\chi^{zxx} + k_z(2\omega)\chi^{xxx}\cos 3\phi) \\
&= \frac{4i\pi\omega}{c\cos\theta}\frac{2\cos\theta}{\epsilon(2\omega)\cos\theta + k_z(2\omega)}(\sin\theta\epsilon(2\omega)\chi^{zxx} + k_z(2\omega)\chi^{xxx}\cos 3\phi).
\end{aligned}$$

And we finally obtain \mathcal{R}_{sP} ,

$$\begin{aligned}
\mathcal{R}_{sP} &= \frac{2\pi}{c} \left| \frac{E^{(2\omega)}(\perp, \parallel)}{E_s^2} \right|^2 \\
&= \frac{2\pi}{c} \left| \frac{4i\pi\omega}{c\cos\theta}\frac{2\cos\theta}{\epsilon(2\omega)\cos\theta + k_z(2\omega)}(\sin\theta\epsilon(2\omega)\chi^{zxx} + k_z(2\omega)\chi^{xxx}\cos 3\phi) \right|^2 \\
&= \frac{32\pi^3\omega^2}{c^3\cos^2\theta} \left| \frac{2\cos\theta}{\epsilon(2\omega)\cos\theta + k_z(2\omega)}(\sin\theta\epsilon(2\omega)\chi^{zxx} + k_z(2\omega)\chi^{xxx}\cos 3\phi) \right|^2 \\
&= \frac{32\pi^3\omega^2}{c^3\cos^2\theta} |t_s(\omega)T_p(2\omega)r_{sP}|^2, \tag{11}
\end{aligned}$$

where

$$\begin{aligned}
t_s(\omega) &= 1, \\
T_p(2\omega) &= \frac{2\cos\theta}{\epsilon(2\omega)\cos\theta + k_z(2\omega)}, \\
r_{sP} &= \sin\theta\epsilon(2\omega)\chi^{zxx} + k_z(2\omega)\chi^{xxx}\cos 3\phi.
\end{aligned}$$

1.4 Summary

We unify the final expressions for the SHG yield, Eqs. (7), (9), and (11), as

$$\mathcal{R}_iF = \frac{32\pi^3\omega^2}{c^3\cos^2\theta} |t_i(\omega)T_F(2\omega)r_{iF}|^2. \tag{12}$$

The necessary factors are summarized in Table 1.

iF	$t_i(\omega)$	$T_F(2\omega)$	r_{iF}
pP	$\frac{1}{\epsilon(\omega)}$	$\frac{2 \cos \theta}{\epsilon(2\omega) \cos \theta + k_z(2\omega)}$	$\sin \theta \epsilon(2\omega) (\sin^2 \theta \chi^{zzz} + k_z^2(\omega) \chi^{zxx}) - k_z(\omega) k_z(2\omega) (2 \sin \theta \chi^{xxz} + k_z(\omega) \chi^{xxx} \cos 3\phi)$
pS	$\frac{1}{\epsilon(\omega)}$	$\frac{2 \cos \theta}{\cos \theta + k_z(2\omega)}$	$k_z^2(\omega) \chi^{xxx} \sin 3\phi$
sP	1	$\frac{2 \cos \theta}{\epsilon(2\omega) \cos \theta + k_z(2\omega)}$	$\sin \theta \epsilon(2\omega) \chi^{zxx} + k_z(2\omega) \chi^{xxx} \cos 3\phi$

Table 1: The necessary factors for Eq. (12) for each polarization case.

2 SHG yield per Mizrahi and Sipe [2]

2.1 \mathcal{R}_{pP}

2.2 \mathcal{R}_{pS}

2.3 \mathcal{R}_{sP}

2.4 Summary

3 SHG yield per Mendoza [3]

3.1 \mathcal{R}_{pP}

3.2 \mathcal{R}_{pS}

3.3 \mathcal{R}_{sP}

3.4 Summary

References

- [1] J. E. Sipe, D. J. Moss, and H. M. van Driel. Phenomenological theory of optical second- and third-harmonic generation from cubic centrosymmetric crystals. *Phys. Rev. B*, 35(3):1129–1141, January 1987.
- [2] V. Mizrahi and J. E. Sipe. Phenomenological treatment of surface second-harmonic generation. *J. Opt. Soc. Am. B*, 5(3):660–667, 1988.
- [3] B. S. Mendoza. Longitudinal Gauge Theory of Second Harmonic Generation at Semiconductor Surfaces. In *Epioptics-7: Proceedings of the 24th Course of the International School of Solid State Physics: Erice, Italy, 20-26 July 2002*, volume 23, page 52, 2004.