

# Expressions for arbitrary rotation on $\chi(-2\omega; \omega, \omega)$

Sean M. Anderson<sup>\*</sup> and Bernardo S. Mendoza  
*Centro de Investigaciones en Óptica, A.C., León 37150, Mexico*

Yujin Cho  
*Department of Physics, University of Texas at Austin, Austin, Texas 78712, USA*

(Dated: May 10, 2018)

We develop explicit expressions for rotating the nonlinear susceptibility tensor  $\chi(-2\omega; \omega, \omega)$  through an arbitrary angle  $\gamma$ . Each component in the rotated frame of reference is a combination of different components from the non-rotated system; setting the angle of rotation to  $\pi/2$  makes both systems equivalent.

To take the components of  $\chi(-2\omega; \omega, \omega)$  from the crystallographic frame to the lab frame, we can simply apply a standard rotational matrix,

$$R = \begin{pmatrix} R_{Xx} & R_{Xy} & R_{Xz} \\ R_{Yx} & R_{Yy} & R_{Yz} \\ R_{Zx} & R_{Zy} & R_{Zz} \end{pmatrix} = \begin{pmatrix} \sin \gamma & -\cos \gamma & 0 \\ \cos \gamma & \sin \gamma & 0 \\ 0 & 0 & 1 \end{pmatrix},$$

such that

$$\chi^{IJK} = \sum_{ijk} R_{Ii} R_{Jj} R_{Kk} \chi^{ijk},$$

where  $I, J$ , and  $K$  ( $i, j, k$ ) cycle through  $X, Y$ , or  $Z$  ( $x, y, z$ ). Fig. 1 depicts this rotation over any arbitrary angle  $\gamma$ . Since we only consider a rotation in the  $xy$ -plane along  $\gamma$ , the  $z$  and  $Z$  axes are the same.




FIG. 1: The translation from the non-rotated  $xyz$  coordinates to the rotated  $XYZ$  system.

Therefore, our  $\chi^{IJK}$  components in terms of the original  $ijk$  coordinate system are

$$\begin{aligned}
\chi^{XXX} &= \sin^3 \gamma \chi^{xxx} + \sin \gamma \cos^2 \gamma \chi^{xyy} - 2 \sin^2 \gamma \cos \gamma \chi^{xxy} \\
&\quad - \sin^2 \gamma \cos \gamma \chi^{yxx} - \cos^3 \gamma \chi^{yyy} + 2 \sin \gamma \cos^2 \gamma \chi^{yyx}, \\
\chi^{XYY} &= \sin \gamma \cos^2 \gamma \chi^{xxx} + \sin^3 \gamma \chi^{xyy} + 2 \sin^2 \gamma \cos \gamma \chi^{xxy} \\
&\quad - \cos^3 \gamma \chi^{yxx} - \sin^2 \gamma \cos \gamma \chi^{yyy} - 2 \sin \gamma \cos^2 \gamma \chi^{yyx}, \\
\chi^{XZZ} &= \sin \gamma \chi^{xzz} - \cos \gamma \chi^{yzz}, \\
\chi^{XYZ} &= \chi^{XZY} = \sin^2 \gamma \chi^{xyz} + \sin \gamma \cos \gamma \chi^{xxz} - \sin \gamma \cos \gamma \chi^{yyz} - \cos^2 \gamma \chi^{yxz}, \\
\chi^{XXZ} &= \chi^{XZX} = -\sin \gamma \cos \gamma \chi^{xyz} + \sin^2 \gamma \chi^{xxz} + \cos^2 \gamma \chi^{yyz} - \sin \gamma \cos \gamma \chi^{yxz}, \\
\chi^{XXY} &= \chi^{XYX} = \sin^2 \gamma \cos \gamma \chi^{xxx} - \sin^2 \gamma \cos \gamma \chi^{xyy} + (\sin^3 \gamma - \sin \gamma \cos^2 \gamma) \chi^{xxy} \\
&\quad - \sin \gamma \cos^2 \gamma \chi^{yxx} + \sin \gamma \cos^2 \gamma \chi^{yyy} + (\cos^3 \gamma - \sin^2 \gamma \cos \gamma) \chi^{yyx},
\end{aligned} \tag{1}$$

for the  $\chi^{XJK}$  components,

$$\begin{aligned}
\chi^{YXX} &= \sin^2 \gamma \cos \gamma \chi^{xxx} + \cos^3 \gamma \chi^{xyy} - 2 \sin \gamma \cos^2 \gamma \chi^{xxy} \\
&\quad + \sin^3 \gamma \chi^{yxx} + \sin \gamma \cos^2 \gamma \chi^{yyy} - 2 \sin^2 \gamma \cos \gamma \chi^{yyx}, \\
\chi^{YYX} &= \cos^3 \gamma \chi^{xxx} + \sin^2 \gamma \cos \gamma \chi^{xyy} + 2 \sin \gamma \cos^2 \gamma \chi^{xxy} \\
&\quad + \sin \gamma \cos^2 \gamma \chi^{yxx} + \sin^3 \gamma \chi^{yyy} + 2 \sin^2 \gamma \cos \gamma \chi^{yyx}, \\
\chi^{YZZ} &= \cos \gamma \chi^{xzz} + \sin \gamma \chi^{yzz}, \\
\chi^{YYZ} &= \chi^{YZY} = \sin \gamma \cos \gamma \chi^{xyz} + \cos^2 \gamma \chi^{xxz} + \sin^2 \gamma \chi^{yyz} + \sin \gamma \cos \gamma \chi^{yxz}, \\
\chi^{YXZ} &= \chi^{YZX} = -\cos^2 \gamma \chi^{xyz} + \sin \gamma \cos \gamma \chi^{xxz} - \sin \gamma \cos \gamma \chi^{yyz} + \sin^2 \gamma \chi^{yxz}, \\
\chi^{YXY} &= \chi^{YYX} = \sin \gamma \cos^2 \gamma \chi^{xxx} - \sin \gamma \cos^2 \gamma \chi^{xyy} - (\cos^3 \gamma - \sin^2 \gamma \cos \gamma) \chi^{xxy} \\
&\quad + \sin^2 \gamma \cos \gamma \chi^{yxx} - \sin^2 \gamma \cos \gamma \chi^{yyy} + (\sin^3 \gamma - \sin \gamma \cos^2 \gamma) \chi^{yyx},
\end{aligned}$$

for the  $\chi^{YJK}$  components, and lastly

$$\begin{aligned}
\chi^{ZXX} &= \sin^2 \gamma \chi^{zxx} + \cos^2 \gamma \chi^{zyy} - 2 \sin \gamma \cos \gamma \chi^{zxy}, \\
\chi^{ZYY} &= \cos^2 \gamma \chi^{zxx} + \sin^2 \gamma \chi^{zyy} + 2 \sin \gamma \cos \gamma \chi^{zxy}, \\
\chi^{ZZZ} &= \chi^{zzz}, \\
\chi^{ZYZ} &= \chi^{ZZY} = \sin \gamma \chi^{zyz} + \cos \gamma \chi^{zxx}, \\
\chi^{ZXX} &= \chi^{ZZX} = -\cos \gamma \chi^{zyz} + \sin \gamma \chi^{zxx}, \\
\chi^{ZXY} &= \chi^{ZYY} = \sin \gamma \cos \gamma \chi^{zxx} - \sin \gamma \cos \gamma \chi^{zyy} - \cos 2\gamma \chi^{zxy},
\end{aligned}$$

for the  $\chi^{ZJK}$  components. Fortunately, the intrinsic permutation symmetry of SHG is also present in the new coordinate system, such that  $\chi^{IJK} = \chi^{IKJ}$ ; therefore, there are only 18 unique components in either system. Setting  $\gamma = \pi/2$  signifies that there is no rotation, and thus  $\chi^{IJK} = \chi^{ijk}$ .




FIG. 2:  $\chi^{XXX}$  for three values of  $\gamma$  calculated for a system with  $C_{3v}$  symmetry.

It should also be clear that the crystal symmetries do **not** follow into the rotated system. For instance, the  $C_{3v}$  symmetry satisfies the following,

$$\begin{aligned}\chi^{xxx} &= -\chi^{xyy} = -\chi^{yxy}, \\ \chi^{yxx} &= \chi^{yyy} = 0.\end{aligned}$$

In the rotated system, the top relationship holds true such that  $\chi^{XXX} = -\chi^{XYX} = -\chi^{YXY}$ . However, we also obtain that

$$\chi^{YYY} = \cos 3\gamma \chi^{xxx},$$

which is not necessarily zero. Fortunately, we can simply apply the crystal symmetry to the non-rotated system before transforming to the rotated system. As an example case, we present  $\chi^{XXX}$  for three values of  $\gamma$  for a system with  $C_{3v}$  symmetry in Fig. 2. The nonzero components in the original coordinates are presented in Fig. 3, multiplied by the appropriate prefactors from Eq. (1). We can see how we recover the component in the original coordinates when  $\gamma = \pi/2$ .

\* sma@cio.mx




FIG. 3: Nonzero components of  $\chi^{ijk}$  multiplied by the appropriate prefactors, for three different values of  $\gamma$ .