List all the techniques or methods to check that vectors are dependent pr independent along with examples

To check whether vectors are linearly independent or dependent, there isn't just one method—you actually have several powerful techniques. I'll explain each in a simple way with examples so you can pick what suits your problem.

🔹 1. Definition Method (Basic Idea)

Concept:

Vectors are dependent if one vector can be written as a combination of others.

Example:

Let

$v_1 = (1, 2), v_2 = (2, 4)$

Check:

$v_2 = 2 \cdot v_1$

👉 So they are dependent

🔹 2. Determinant Method (Square Matrix Only)

Concept:

- If det ≠ 0 → independent
- If det = 0 → dependent

Example:

$v_1 = (1, 2), v_2 = (3, 4)$

Matrix:

$$\begin{bmatrix} 1 & 3 \\ 2 & 4 \end{bmatrix}$$

$$\det \begin{bmatrix} 1 & 3 \\ 2 & 4 \end{bmatrix} = (1)(4) - (3)(2) = -2$$

👉 det ≠ 0 → independent

🔹 3. Row Reduction (Gaussian Elimination)

Concept:

Convert matrix to row echelon form:

- Pivot in every column → independent
- Zero row appears → dependent

Example:

$v_1 = (1, 2, 3), v_2 = (2, 4, 6)$

Matrix:

$$\begin{bmatrix} 1 & 2 & 3 \\ 2 & 4 & 6 \end{bmatrix}$$

Row operation:

$R_2 \rightarrow R_2 - 2R_1 \rightarrow (0, 0, 0)$

👉 Zero row → dependent

🔹 4. Rank Method

Concept:

- Rank = number of vectors → independent
- Rank < number of vectors → dependent

Example:

3 vectors in R^3 but rank = 2

👉 dependent

🔹 5. Augmented System Method (Solve $c_1v_1 + c_2v_2 + \ldots = 0$)

Concept:

Solve:

$$c_1v_1 + c_2v_2 + \ldots = 0$$

- Only trivial solution (all c = 0) → independent
- Non-zero solution → dependent

Example:

$v_1 = (1,2)$, $v_2 = (2,4)$

$$c_1(1,2) + c_2(2,4) = (0,0)$$

Solution:

$c_1 = -2c_2$

👉 Non-zero solution exists → dependent

🔹 6. Geometric Interpretation

Concept:

- Same direction (parallel) → dependent
- Different directions → independent

Example:

(1,2) and (2,4) → same line → dependent
(1,0) and (0,1) → perpendicular → independent

🔹 7. Number of Vectors vs Dimension Rule

Concept:

- If vectors > dimension → always dependent

Example:

4 vectors in $\mathbb{R}^3$

👉 Automatically dependent

🔹 8. Eigenvector Idea (Advanced)

If eigenvectors correspond to different eigenvalues, they are always independent.

🔹 9. Gram Matrix Method

Form matrix $A^T A$:

- det ≠ 0 → independent
- det = 0 → dependent

(Used in higher math / numerical methods)

🔹 10. Cross Product Method (Only in $\mathbb{R}^3$)

Concept:

For 2 vectors:

- Cross product ≠ 0 → independent
- Cross product = 0 → dependent

Example:

$v_1 = (1,0,0)$, $v_2 = (0,1,0)$

👉 $v_1 \times v_2 \neq 0$ → independent

🔥 Quick Summary Table

Method	Independent Condition
Definition	No vector is combination of others
Determinant	det ≠ 0

Method	Independent Condition
Row reduction	No zero row
Rank	Rank = number of vectors
System solution	Only trivial solution
Geometry	Not parallel
Dimension rule	vectors ≤ dimension
Cross product	≠ 0

💡 Which method should YOU use?

- Small 2×2 or 3×3 → Determinant
- Large system → Row reduction / Rank
- Conceptual understanding → Definition / Geometry

If you want, I can now:

✔ ↪ Solve a full 3×3 independence problem step-by-step

✔ Or ↪ give you tricky exam-type questions that students usually get wrong

Also what trivial non trivial solution

Good question—this concept is very important for understanding linear independence.

🔹 What is a *Trivial Solution*?

Definition:

A trivial solution means all constants are zero.

For equation:

$$c_1v_1 + c_2v_2 + \cdots + c_nv_n = 0$$

👉 Trivial solution:

$$c_1 = 0,\ c_2 = 0,\ \ldots,\ c_n = 0$$

Example:

Let

$v_1 = (1,0)$, $v_2 = (0,1)$

Solve:

$$c_1(1,0) + c_2(0,1) = (0,0)$$

This gives:

$c_1 = 0$, $c_2 = 0$

👉 Only trivial solution → vectors are independent

🔹 What is a *Non-Trivial Solution*?

Definition:

A non-trivial solution means at least one constant ≠ 0, but equation still equals zero vector.

Example:

$v_1 = (1,2)$, $v_2 = (2,4)$

Solve:

$$c_1(1,2) + c_2(2,4) = (0,0)$$

This gives:

$$c_1 = -2c_2$$

👉 Choose:

$c_2 = 1 \rightarrow c_1 = -2$

So:

$$-2v_1 + 1v_2 = 0$$

👉 Non-zero values exist → non-trivial solution → vectors are dependent

---

🔥 Key Idea (Very Important)

Type of Solution	Meaning	Result
Only trivial	All c = 0	Independent
Non-trivial exists	Some c ≠ 0	Dependent

🔹 Simple Intuition (Easy Way)

- Trivial = "boring solution" → always exists
- Non-trivial = "interesting solution" → proves dependence

🔹 One-Line Shortcut

👉 If you can make:

$$c_1v_1 + c_2v_2 + \cdots = 0$$

without all c = 0,

then vectors are dependent

If you want, I can give:

✔ ↪ Exam-style MCQs on this

✔ Or ↪ a full 3×3 system where we find trivial/non-trivial step-by-step

In a sense of unique and infinite mamy solution

Great—now you're connecting this to unique vs infinite solutions, which is exactly how exams test this concept.

🔹 Core Equation

We always check:

$$c_1v_1 + c_2v_2 + \cdots + c_nv_n = 0$$

This is a homogeneous system.

🔹 Case I: Unique Solution

$c_1v_1 + c_2v_2 + \cdots + c_nv_n = 0$

When does this have a unique solution?

👉 Only when:

$$c_1 = c_2 = \cdots = c_n = 0$$

👉 That means:

- Only trivial solution exists
- No freedom, no parameters

Conclusion:

✔ Unique solution → vectors are independent

Example:

$v_1 = (1,0)$, $v_2 = (0,1)$

System:

$$c_1(1,0) + c_2(0,1) = (0,0)$$

Solution:

$c_1 = 0$, $c_2 = 0$

👉 Only one solution → independent

🔹 Case 2: Infinite Solutions

When does this happen?

👉 When system has free variables

That means:

- One variable depends on another
- Infinite combinations possible

Example:

$v_1 = (1,2)$, $v_2 = (2,4)$

System:

$$c_1(1,2) + c_2(2,4) = (0,0)$$

You get:

$$c_1 = -2c_2$$

Let:

$c_2 = t \rightarrow c_1 = -2t$

👉 Infinite solutions (t can be anything)

Conclusion:

✔ Infinite solutions → vectors are dependent

🔥 Final Connection (Most Important)

Type of Solution	Meaning	Vectors
Unique solution	Only trivial (all 0)	Independent
Infinite solutions	Non-trivial exists	Dependent

🔹 Why this always happens (Quick Logic)

- This system is homogeneous, so:
  - It always has at least one solution (trivial)
- So only two possibilities:
  - Only trivial → unique
  - More solutions → infinite

👉 Never exactly 2 or 3 solutions (important MCQ point)

🔹 Shortcut to Remember

👉

Independent → Unique solution

Dependent → Infinite solutions

If you want next, I can:

✔ ↪ Show how this links with rank and pivots (very exam important)

✔ Or ↪ solve a 3×3 system step-by-step using row reduction

Free variable example in matrix

Good—this is exactly where many students get confused. Let's make free variables very clear with a proper matrix example.

🔹 What is a Free Variable?

A free variable is a variable that:

- does not have a pivot (leading 1) in its column
- can take any value (parameter)

  👉 That's why it creates infinite solutions

🔹 Example 1 (Simple 2×2 Case)

Consider system:

$$\begin{cases} x + 2y = 0 \\ 2x + 4y = 0 \end{cases}$$

Matrix form:

$$\begin{bmatrix} 1 & 2 \\ 2 & 4 \end{bmatrix}$$

Row reduce:

$R_2 \to R_2 - 2R_1$

$$\begin{bmatrix} 1 & 2 \\ 0 & 0 \end{bmatrix}$$

🔍 Now observe:

- First column → pivot (leading 1) → x is basic variable
- Second column → no pivot → y is free variable

Solution:

From first row:

$$x + 2y = 0 \Rightarrow x = -2y$$

Let:

$$y = t$$

Then:

$$x = -2t$$

Final Answer:

$$(x, y) = (-2t, t)$$

👉 Infinite solutions

👉 y is free variable

🔹 Example 2 (3 Variables – More Clear)

System:

$$\begin{cases} x + y + z = 0 \\ 2x + 2y + 2z = 0 \end{cases}$$

Matrix:

$$\begin{bmatrix} 1 & 1 & 1 \\ 2 & 2 & 2 \end{bmatrix}$$

Row reduce:

$R_2 \to R_2 - 2R_1$

$$\begin{bmatrix} 1 & 1 & 1 \\ 0 & 0 & 0 \end{bmatrix}$$

🔍 Identify variables:

- Column 1 → pivot → x is basic
- Column 2 → no pivot → y is free

- Column 3 → no pivot → z is free

Solution:

$$x + y + z = 0 \Rightarrow x = -y - z$$

Let:

$$y = s, \quad z = t$$

Then:

$$x = -s - t$$

Final Answer:

$$(x, y, z) = (-s - t, s, t)$$

👉 Two free variables → more freedom → infinite solutions

🔥 Key Takeaways

Term	Meaning
Pivot variable	Has leading 1 (depends on system)
Free variable	No pivot, can take any value
Free variable exists	Infinite solutions
No free variable	Unique solution

🔹 Shortcut Trick (Exam Tip)

👉 Count pivots:

- If pivots = number of variables → no free variable → unique solution

- If pivots < variables → free variables exist → infinite solutions

---

If you want next, I can:

✔ Show how to identify free variables directly without full row reduction

✔ Or connect this with rank = number of pivots (very important for exams)



Is this conversation helpful so far?