

Prover-Skeptic Games and Logical Pluralism

Rohan French
University of Groningen

20TH AMSTERDAM COLLOQUIUM
AMSTERDAM • DECEMBER 17, 2015

A Problem for Logical Pluralists

The Built-In Opponent Conception of Deduction

Introducing Prover-Skeptic Games

Prover-Skeptic Games: An Example

Proofs & Explanations

Prover-Skeptic Games for Substructural Logics

*A Problem for Logical
Pluralists*

A characterisation of a pluralist position needs to explain what it is to endorse all of the various consequence relations the pluralist accepts and how they relate to an intuitive notion of logical consequence.

—Keefe, 2014, p.1376

Two Central Questions

Two Central Questions

- ▶ What is it to ‘endorse’ a consequence relation?

Two Central Questions

- ▶ What is it to ‘endorse’ a consequence relation?
- ▶ Why are these consequence relations relations of logical consequence?

*The Built-In Opponent
Conception of Deduction*

Multi-Agent View of Logical Consequence

- ▶ Logical consequence grounded in specialised multi-agent discursive practices.

- ▶ Logical consequence grounded in specialised multi-agent discursive practices.
- ▶ View **proofs** as grounded in semi-adversarial debates between a Prover/Proponent and a Skeptic/Opponent.

- ▶ Logical consequence grounded in specialised multi-agent discursive practices.
- ▶ View **proofs** as grounded in semi-adversarial debates between a Prover/Proponent and a Skeptic/Opponent.
- ▶ Prover seeks to establish the conclusion.

- ▶ Logical consequence grounded in specialised multi-agent discursive practices.
- ▶ View **proofs** as grounded in semi-adversarial debates between a Prover/Proponent and a Skeptic/Opponent.
- ▶ Prover seeks to establish the conclusion.
- ▶ Skeptic forces Prover to make their reasoning explicit.

- ▶ Logical consequence grounded in specialised multi-agent discursive practices.
- ▶ View **proofs** as grounded in semi-adversarial debates between a Prover/Proponent and a Skeptic/Opponent.
- ▶ Prover seeks to establish the conclusion.
- ▶ Skeptic forces Prover to make their reasoning explicit.
- ▶ Deductive proofs are winning strategies for Prover in these debates.

Proponent's job is not only to 'beat Opponent'; she also seeks to persuade Opponent of the truth of the conclusion, if he has granted the truth of the premises. In fact, the goal is not only to show that the conclusion follows from the premises, but also why it does; this corresponds to the idea that deductive arguments ought to have explanatory value.

– Dutilh Novaes, forthcoming

Internalization of Skeptic

- ▶ Proofs are not usually thought of as a multi-agent enterprise.

- ▶ Proofs are not usually thought of as a multi-agent enterprise.
- ▶ This is because Skeptic has become **internalized** to the method of giving deductive proofs.

- ▶ Proofs are not usually thought of as a multi-agent enterprise.
- ▶ This is because Skeptic has become **internalized** to the method of giving deductive proofs.
- ▶ In effect their role is played 'offline' by Prover.

- ▶ Proofs are not usually thought of as a multi-agent enterprise.
- ▶ This is because Skeptic has become **internalized** to the method of giving deductive proofs.
- ▶ In effect their role is played 'offline' by Prover.
- ▶ This provides a bridge between multi-agent dialogical and mono-agent inferential practices.

Introducing Prover-Skeptic Games

*In the library two students, **Penelope** and **Scott** are arguing over the validity of the argument from $A \rightarrow B$ and $B \rightarrow C$ to $A \rightarrow C$. Petunia thinks that this argument is valid and is trying to convince Scott, who is skeptical.*

*In the library two students, **Penelope** and **Scott** are arguing over the validity of the argument from $A \rightarrow B$ and $B \rightarrow C$ to $A \rightarrow C$. Petunia thinks that this argument is valid and is trying to convince Scott, who is skeptical.*

- **Penelope (1):** I reckon that $A \rightarrow C$ follows from $A \rightarrow B$ and $B \rightarrow C$

*In the library two students, **Penelope** and **Scott** are arguing over the validity of the argument from $A \rightarrow B$ and $B \rightarrow C$ to $A \rightarrow C$. Petunia thinks that this argument is valid and is trying to convince Scott, who is skeptical.*

- ▶ **Penelope (1):** I reckon that $A \rightarrow C$ follows from $A \rightarrow B$ and $B \rightarrow C$
- ▶ **Scott (1):** Yeah? Well if that's so then suppose I grant you $A \rightarrow B$ and $B \rightarrow C$ along with A , how are you mean to get C ?

- ▶ **Penelope (2):** If you grant me B I can get C from $B \rightarrow C$ (which you just granted).

- ▶ **Penelope (2):** If you grant me B I can get C from $B \rightarrow C$ (which you just granted).
- ▶ **Scott (2):** But why should I grant you B?

- ▶ **Penelope (2):** If you grant me B I can get C from $B \rightarrow C$ (which you just granted).
- ▶ **Scott (2):** But why should I grant you B?
- ▶ **Penelope (3):** Well if you were to grant me A then I could get B from $A \rightarrow B$ which you granted at the start.

- ▶ **Penelope (2):** If you grant me B I can get C from $B \rightarrow C$ (which you just granted).
- ▶ **Scott (2):** But why should I grant you B?
- ▶ **Penelope (3):** Well if you were to grant me A then I could get B from $A \rightarrow B$ which you granted at the start.
- ▶ **Scott (3):** But why should I grant you A?

- ▶ **Penelope (2):** If you grant me B I can get C from $B \rightarrow C$ (which you just granted).
- ▶ **Scott (2):** But why should I grant you B?
- ▶ **Penelope (3):** Well if you were to grant me A then I could get B from $A \rightarrow B$ which you granted at the start.
- ▶ **Scott (3):** But why should I grant you A?
- ▶ **Penelope (4):** You granted it to me at the start!

- ▶ **Penelope (2):** If you grant me B I can get C from $B \rightarrow C$ (which you just granted).
- ▶ **Scott (2):** But why should I grant you B?
- ▶ **Penelope (3):** Well if you were to grant me A then I could get B from $A \rightarrow B$ which you granted at the start.
- ▶ **Scott (3):** But why should I grant you A?
- ▶ **Penelope (4):** You granted it to me at the start!

- ▶ **Penelope (2):** If you grant me B I can get C from $B \rightarrow C$ (which you just granted).
- ▶ **Scott (2):** But why should I grant you B?
- ▶ **Penelope (3):** Well if you were to grant me A then I could get B from $A \rightarrow B$ which you granted at the start.
- ▶ **Scott (3):** But why should I grant you A?
- ▶ **Penelope (4):** You granted it to me at the start!

Penelope leaves the library triumphantly.

A Natural Deduction System

AXIOMS

$$A \succ A$$

AXIOMS

$$A \succ A$$

INTRODUCTION/ELIMINATION RULES

$$\frac{\Gamma, A \succ B}{\Gamma \succ A \rightarrow B} (\rightarrow I) \qquad \frac{\Gamma \succ A \quad \Delta \succ A \rightarrow B}{\Gamma, \Delta \succ B} (\rightarrow E)$$

AXIOMS

$$A \succ A$$

INTRODUCTION/ELIMINATION RULES

$$\frac{\Gamma, A \succ B}{\Gamma \succ A \rightarrow B} (\rightarrow I) \qquad \frac{\Gamma \succ A \quad \Delta \succ A \rightarrow B}{\Gamma, \Delta \succ B} (\rightarrow E)$$

STRUCTURAL RULES

$$\frac{\Gamma, A, A \succ B}{\Gamma, A \succ B} (W) \qquad \frac{\Gamma \succ B}{\Gamma, A \succ B} (K)$$

A Natural Deduction System: Example & Notation

$$\frac{\frac{A \rightarrow B \succ A \rightarrow B \quad A \succ A}{A \rightarrow B, A \succ B} (\rightarrow E) \quad B \rightarrow C \succ B \rightarrow C}{\frac{A \rightarrow B, B \rightarrow C, A \succ C}{A \rightarrow B, B \rightarrow C \succ A \rightarrow C} (\rightarrow I)} (\rightarrow E)$$

A Natural Deduction System: Example & Notation

$$\frac{\frac{A \rightarrow B \succ A \rightarrow B \quad A \succ A}{A \rightarrow B, A \succ B} (\rightarrow E) \quad \frac{B \rightarrow C \succ B \rightarrow C}{B \rightarrow C, B \rightarrow C} (\rightarrow E)}{\frac{A \rightarrow B, B \rightarrow C, A \succ C}{A \rightarrow B, B \rightarrow C \succ A \rightarrow C} (\rightarrow I)} (\rightarrow E)$$

- Let us write $\Gamma \succ A \Rightarrow_{(I)}^* \Delta \succ B$ iff we can derive $\Delta \succ B$ from $\Gamma \succ A$ using $(\rightarrow I)$.

A Natural Deduction System: Example & Notation

$$\frac{\frac{A \rightarrow B \succ A \rightarrow B \quad A \succ A}{A \rightarrow B, A \succ B} (\rightarrow E) \quad \frac{B \rightarrow C \succ B \rightarrow C}{B \rightarrow C, B \rightarrow C} (\rightarrow E)}{\frac{A \rightarrow B, B \rightarrow C, A \succ C}{A \rightarrow B, B \rightarrow C \succ A \rightarrow C} (\rightarrow I)}$$

- Let us write $\Gamma \succ A \Rightarrow_{(I)}^* \Delta \succ B$ iff we can derive $\Delta \succ B$ from $\Gamma \succ A$ using $(\rightarrow I)$.

A Natural Deduction System: Example & Notation

$$\frac{\frac{A \rightarrow B \succ A \rightarrow B \quad A \succ A}{A \rightarrow B, A \succ B} (\rightarrow E) \quad \frac{B \rightarrow C \succ B \rightarrow C}{B \rightarrow C, B \rightarrow C} (\rightarrow E)}{\frac{A \rightarrow B, B \rightarrow C, A \succ C}{A \rightarrow B, B \rightarrow C \succ A \rightarrow C} (\rightarrow I)} (\rightarrow E)$$

- Let us write $\Gamma \succ A \Rightarrow_{(I)}^* \Delta \succ B$ iff we can derive $\Delta \succ B$ from $\Gamma \succ A$ using $(\rightarrow I)$.
- $[\Gamma_1 \succ A_1, \dots, \Gamma_n \succ A_n] \Rightarrow_{(E)}^* \Delta \succ B$ iff we can derive $\Delta \succ B$ from the $\Gamma_i \succ A_i$ s using $(\rightarrow E)$ and the structural rules.

A Natural Deduction System: Example & Notation

$$\frac{\frac{\frac{A \rightarrow B \succ A \rightarrow B \quad A \succ A}{A \rightarrow B, A \succ B} (\rightarrow E) \quad B \rightarrow C \succ B \rightarrow C}{\frac{A \rightarrow B, B \rightarrow C, A \succ C}{A \rightarrow B, B \rightarrow C \succ A \rightarrow C} (\rightarrow I)} (\rightarrow E)$$

- Let us write $\Gamma \succ A \Rightarrow_{(I)}^* \Delta \succ B$ iff we can derive $\Delta \succ B$ from $\Gamma \succ A$ using $(\rightarrow I)$.
- $[\Gamma_1 \succ A_1, \dots, \Gamma_n \succ A_n] \Rightarrow_{(E)}^* \Delta \succ B$ iff we can derive $\Delta \succ B$ from the $\Gamma_i \succ A_i$ s using $(\rightarrow E)$ and the structural rules.

DEFINITION: A *dialogue* over the sequent $\Gamma \succ A$ is a possibly infinite sequence $\Sigma_1, \mathfrak{s}_1, \Sigma_2, \mathfrak{s}_2, \dots$ where:

DEFINITION: A *dialogue* over the sequent $\Gamma \succ A$ is a possibly infinite sequence $\Sigma_1, \mathfrak{s}_1, \Sigma_2, \mathfrak{s}_2, \dots$ where:

1. $\Sigma_1 = [\Gamma \succ A]$

» *Prover begins by asserting the sequent under discussion.*

DEFINITION: A *dialogue* over the sequent $\Gamma \succ A$ is a possibly infinite sequence $\Sigma_1, s_1, \Sigma_2, s_2, \dots$ where:

1. $\Sigma_1 = [\Gamma \succ A]$
2. $s_i \Rightarrow_{(I)}^* \sigma$, for some non-axiomatic sequent $\sigma \in \Sigma_i$ where s_i is of the form $\Theta \succ p_i$ for some propositional atom p_i .

» *Skeptic challenges an (interesting) Prover assertion*

DEFINITION: A *dialogue* over the sequent $\Gamma \succ A$ is a possibly infinite sequence $\Sigma_1, \mathfrak{s}_1, \Sigma_2, \mathfrak{s}_2, \dots$ where:

1. $\Sigma_1 = [\Gamma \succ A]$
2. $\mathfrak{s}_i \Rightarrow_{(I)}^* \sigma$, for some non-axiomatic sequent $\sigma \in \Sigma_i$ where \mathfrak{s}_i is of the form $\Theta \succ p_i$ for some propositional atom p_i .
3. $\Sigma_{i+1} \Rightarrow_{(E)}^* \mathfrak{s}_i$

» *Prover replies, offering sequents which derive the challenge.*

Prover-Skeptic Games: An Example

Revisiting the Previous Example

Consider the following dialogue over $p \rightarrow q, q \rightarrow r \succ p \rightarrow r$.

P(1) $[p \rightarrow q, q \rightarrow r \succ p \rightarrow r]$

» *Prover Starts*

Revisiting the Previous Example

Consider the following dialogue over $p \rightarrow q, q \rightarrow r \succ p \rightarrow r$.

P(1) $[p \rightarrow q, q \rightarrow r \succ p \rightarrow r]$

S(1) $p \rightarrow q, q \rightarrow r, p \succ r$

» *Skeptic*

*challenges the
sole Prover
assertion.*

Revisiting the Previous Example

Consider the following dialogue over $p \rightarrow q, q \rightarrow r \succ p \rightarrow r$.

P(1) $[p \rightarrow q, q \rightarrow r \succ p \rightarrow r]$

S(1) $p \rightarrow q, q \rightarrow r, p \succ r$

P(2) $[q \rightarrow r \succ q \rightarrow r, p \rightarrow q, p \succ q]$

» *Prover replies, as these sequents derive the challenge sequent.*

Revisiting the Previous Example

Consider the following dialogue over $p \rightarrow q, q \rightarrow r \succ p \rightarrow r$.

P(1) $[p \rightarrow q, q \rightarrow r \succ p \rightarrow r]$

» *Skeptic*

S(1) $p \rightarrow q, q \rightarrow r, p \succ r$

challenges the

P(2) $[q \rightarrow r \succ q \rightarrow r, p \rightarrow q, p \succ q]$

only

S(2) $p \rightarrow q, p \succ q$

non-axiomatic

sequent.

Revisiting the Previous Example

Consider the following dialogue over $p \rightarrow q, q \rightarrow r \succ p \rightarrow r$.

P(1) $[p \rightarrow q, q \rightarrow r \succ p \rightarrow r]$

S(1) $p \rightarrow q, q \rightarrow r, p \succ r$

P(2) $[q \rightarrow r \succ q \rightarrow r, p \rightarrow q, p \succ q]$

S(2) $p \rightarrow q, p \succ q$

P(3) $[p \rightarrow q \succ p \rightarrow q, p \succ p]$

» *Prover replies,
and wins the
dialogue.*

Revisiting the Previous Example

Consider the following dialogue over $p \rightarrow q, q \rightarrow r \succ p \rightarrow r$.

P(1) $[p \rightarrow q, q \rightarrow r \succ p \rightarrow r]$

S(1) $p \rightarrow q, q \rightarrow r, p \succ r$

P(2) $[q \rightarrow r \succ q \rightarrow r, p \rightarrow q, p \succ q]$

S(2) $p \rightarrow q, p \succ q$

P(3) $[p \rightarrow q \succ p \rightarrow q, p \succ p]$

Proofs & Explanations

- ▶ The semi-adversarial debates at the heart of the Built-In Opponent conception of deduction are clearly norm governed.

- ▶ The semi-adversarial debates at the heart of the Built-In Opponent conception of deduction are clearly norm governed.
- ▶ For example: norms of turn taking; not questioning obvious claims.

- ▶ The semi-adversarial debates at the heart of the Built-In Opponent conception of deduction are clearly norm governed.
- ▶ For example: norms of turn taking; not questioning obvious claims.
- ▶ These are constitutive of the practice of engaging in these semi-adversarial debates.

- ▶ Another normative aspect of this practice is that both participants have to agree on what counts as a response to a challenge.

- ▶ Another normative aspect of this practice is that both participants have to agree on what counts as a response to a challenge.
- ▶ Prover moves are answers to the question ‘why does this follow’

- ▶ Another normative aspect of this practice is that both participants have to agree on what counts as a response to a challenge.
- ▶ Prover moves are answers to the question ‘why does this follow’
- ▶ What counts as an answer to a question is interest relative.

- ▶ Another normative aspect of this practice is that both participants have to agree on what counts as a response to a challenge.
- ▶ Prover moves are answers to the question ‘why does this follow’
- ▶ What counts as an answer to a question is interest relative.
- ▶ Sometimes the relevant interests can have logical repercussions.

- ▶ Mathematicians working on the foundations of geometry at the beginning of the 20th century appear to have been concerned not just with *what* axioms are involved in proof, but also the *number of times* they are appealed to in it.

- ▶ Mathematicians working on the foundations of geometry at the beginning of the 20th century appear to have been concerned not just with *what* axioms are involved in proof, but also the *number of times* they are appealed to in it.
- ▶ For example: Hessenberg's 1905 proof that Desargues axiom follows from a threefold use of the Pappus axiom.

- ▶ Mathematicians working on the foundations of geometry at the beginning of the 20th century appear to have been concerned not just with *what* axioms are involved in proof, but also the *number of times* they are appealed to in it.
- ▶ For example: Hessenberg's 1905 proof that Desargues axiom follows from a threefold use of the Pappus axiom.
- ▶ One way of understanding this is as involving a change in what counts as a Prover response in a deductive dialogue.

Prover-Skeptic Games for Substructural Logics

- Let S be a subset of $\{(W), (K)\}$. Then an *S-dialogue* is a dialogue as defined above except where Prover is limited to using the structural rules in S in answering a Skeptic challenge.

- ▶ Let S be a subset of $\{(W), (K)\}$. Then an *S-dialogue* is a dialogue as defined above except where Prover is limited to using the structural rules in S in answering a Skeptic challenge.
- ▶ Winning strategies in *S-dialogues* characterise validity in the implicational logic with only structural rules from S .

- ▶ Let S be a subset of $\{(W), (K)\}$. Then an *S-dialogue* is a dialogue as defined above except where Prover is limited to using the structural rules in S in answering a Skeptic challenge.
- ▶ Winning strategies in S -dialogues characterise validity in the implicational logic with only structural rules from S .
- ▶ E.g. $\{(W)\}$ -dialogues characterise the implicational fragment of the relevant logic R .

- ▶ On the dialogical conception of logic used here, logic is normative for a specialised kind of semi-adversarial debates.

- ▶ On the dialogical conception of logic used here, logic is normative for a specialised kind of semi-adversarial debates.
- ▶ Such debates can be governed by different norms which determine what counts as an acceptable response to a Skeptic challenge.

- ▶ On the dialogical conception of logic used here, logic is normative for a specialised kind of semi-adversarial debates.
- ▶ Such debates can be governed by different norms which determine what counts as an acceptable response to a Skeptic challenge.
- ▶ Differences in these norms results in different logics being correct.

Thank You!