

A HYPERSEQUENT CALCULUS FOR CONTINGENT EXISTENCE

Rohan French
University of Groningen

15TH CONGRESS OF LOGIC, METHODOLOGY AND PHILOSOPHY OF SCIENCE
HELSINKI • AUGUST 2015

MOTIVATION

THE PROBLEM WITH QUANTIFIED MODAL LOGIC

- The most technically natural way of presenting quantified modal logic yields a modal logic in which we can derive the *Barcan formula*.

$$\Diamond \exists x A \rightarrow \exists x \Diamond A$$

THE PROBLEM WITH QUANTIFIED MODAL LOGIC

- The most technically natural way of presenting quantified modal logic yields a modal logic in which we can derive the *Barcan formula*.

$$\Diamond \exists x A \rightarrow \exists x \Diamond A$$

- But the Barcan formula has false instances according to commonsense metaphysics.

“In both areas [quantified tense and modal logic], in fact, we have a choice between a certain amount of awkwardness and a certain amount of superstition.”

–ARTHUR N. PRIOR

Past, Present and Future, p.160

- ▶ Goal here is to navigate between awkwardness and superstition by looking at what's going on in proofs of the Barcan formula.

- ▶ Goal here is to navigate between awkwardness and superstition by looking at what's going on in proofs of the Barcan formula.
- ▶ To do this we will use modal hypersequents in the style of Restall (2012). A *Hypersequent* is a multiset of sequents which we will write as

$$X_1 \vdash Y_1 \quad | \quad \dots \quad | \quad X_n \vdash Y_n$$

- Rules for the propositional connectives are ‘local’ to a sequent component $X_i \vdash Y_i$.

$$\frac{\mathcal{H} \mid X \vdash A, B, Y}{\mathcal{H} \mid X \vdash A \vee B, Y} [\vee R]$$

$$\frac{\mathcal{H} \mid X, A \vdash Y \quad \mathcal{H} \mid X, B \vdash Y}{\mathcal{H} \mid X, A \vee B \vdash Y} [\vee L]$$

- Rules for the propositional connectives are ‘local’ to a sequent component $X_i \vdash Y_i$.

$$\frac{\mathcal{H} \mid X \vdash A, B, Y}{\mathcal{H} \mid X \vdash A \vee B, Y} [\vee R]$$

$$\frac{\mathcal{H} \mid X, A \vdash Y \quad \mathcal{H} \mid X, B \vdash Y}{\mathcal{H} \mid X, A \vee B \vdash Y} [\vee L]$$

- Rules for the modal operators have a more ‘global’ flavour

$$\frac{\mathcal{H} \mid X' \vdash Y', A \quad \mid \quad X \vdash Y}{\mathcal{H} \mid X' \vdash Y' \quad \mid \quad X \vdash Y, \Diamond A} [\Diamond R]$$

$$\frac{\mathcal{H} \mid \vdash A \quad \mid \quad X \vdash Y}{\mathcal{H} \mid X, \Diamond A \vdash Y} [\Diamond L]$$

DERIVATIONS OF THE BARCAN FORMULA

A BARCAN DERIVATION

$$\frac{\frac{\frac{Fa \vdash Fa}{Fa \vdash \quad | \quad \vdash \Diamond Fa} [\Diamond R]}{Fa \vdash \quad | \quad \vdash \exists x \Diamond Fx} [\exists R]}{\frac{\exists x Fx \vdash \quad | \quad \vdash \exists x \Diamond Fx} [\exists L]}{\Diamond \exists x Fx \vdash \exists x \Diamond Fx} [\Diamond L]$$

A BARCAN DERIVATION

$$\frac{\frac{\frac{Fa \vdash Fa}{Fa \vdash \quad | \quad \vdash \Diamond Fa} [\Diamond R]}{Fa \vdash \quad | \quad \vdash \exists x \Diamond Fx} [\exists R]}{\frac{\exists x Fx \vdash \quad | \quad \vdash \exists x \Diamond Fx}{\Diamond \exists x Fx \vdash \exists x \Diamond Fx} [\exists L]} [\Diamond L]$$

- An adequate quantified modal logic needs to tell us not only that the above proof is invalid...

A BARCAN DERIVATION

$$\frac{\frac{\frac{Fa \vdash Fa}{Fa \vdash \quad | \quad \vdash \Diamond Fa} [\Diamond R]}{Fa \vdash \quad | \quad \vdash \exists x \Diamond Fx} [\exists R]}{\frac{\exists x Fx \vdash \quad | \quad \vdash \exists x \Diamond Fx} [\exists L]}{\Diamond \exists x Fx \vdash \exists x \Diamond Fx} [\Diamond L]$$

- An adequate quantified modal logic needs to tell us not only that the above proof is invalid...
- ... but also, by way of justifying the logical awkwardness that may be involved, where it led us astray and why.

- ▶ According to Kripke such proofs are question begging...

KRIPKE'S ANALYSIS OF THE PROBLEM

- ▶ According to Kripke such proofs are question begging...
- ▶ ... involving tacit appeals to principles very similar to those they are meant to be proofs of.

KRIPKE'S ANALYSIS OF THE PROBLEM

- ▶ According to Kripke such proofs are question begging...
- ▶ ... involving tacit appeals to principles very similar to those they are meant to be proofs of.
- ▶ If we make the implicit binding of free-variables in formulas explicit then we can no longer derive the Barcan formula.

KRIPKE'S ANALYSIS OF THE PROBLEM

- Unfortunately Kripke's proposal requires us to use a language without individual constants.

KRIPKE'S ANALYSIS OF THE PROBLEM

- ▶ Unfortunately Kripke's proposal requires us to use a language without individual constants.
- ▶ So his solution only works due to a “deliberate impoverishment of the formal machinery”

KRIPKE'S ANALYSIS OF THE PROBLEM

- ▶ Unfortunately Kripke's proposal requires us to use a language without individual constants.
- ▶ So his solution only works due to a “deliberate impoverishment of the formal machinery”
- ▶ Is this the only way to bring out this insight?

PRIMITIVE SOPE MARKERS

PRIMITIVE SCOPE MARKERS

- ▶ Let us employ variables in both their usual *object denoting* role, as well as a new *scope-indicating* role.

PRIMITIVE SCOPE MARKERS

- ▶ Let us employ variables in both their usual *object denoting* role, as well as a new *scope-indicating* role.
- ▶ So, for example, we can distinguish between statements like
 - ▶ “concerning a , it is possibly F ”

$$a \Diamond Fa$$

PRIMITIVE SCOPE MARKERS

- ▶ Let us employ variables in both their usual *object denoting* role, as well as a new *scope-indicating* role.
- ▶ So, for example, we can distinguish between statements like
 - ▶ “concerning a , it is possibly F ”

$a \Diamond Fa$

and

- ▶ “possibly, concerning a it is F ”

$\Diamond a Fa$

PRIMITIVE SCOPE MARKERS

- ▶ Let us employ variables in both their usual *object denoting* role, as well as a new *scope-indicating* role.
- ▶ So, for example, we can distinguish between statements like
 - ▶ “*concerning* a , it is *possibly* F ”

$$a \Diamond Fa$$

and

- ▶ “*possibly, concerning* a it is F ”

$$\Diamond a Fa$$

- ▶ Where we can only bind a with a quantifier in the first case, but not the second.

- Formulas in the language which we'll be concerned with can be constructed by taking closed sentences of the standard language of modal logic and:
 1. Removing occurrences of ' \exists '
 2. Uniformly substituting individual constants for variables which are not bound by a quantifier.

A SKETCH OF A FORMAL LANGUAGE

- Formulas in the language which we'll be concerned with can be constructed by taking closed sentences of the standard language of modal logic and:
 1. Removing occurrences of \exists
 2. Uniformly substituting individual constants for variables which are not bound by a quantifier.
- $\text{'t} \Diamond \text{Ft'}$, $\text{'}\neg \text{t} \Diamond \exists x \text{Gtx'}$ and $\text{'t} \Diamond x \text{Rtx'}$ are all formulas.

A SKETCH OF A FORMAL LANGUAGE

- ▶ Formulas in the language which we'll be concerned with can be constructed by taking closed sentences of the standard language of modal logic and:
 1. Removing occurrences of \exists
 2. Uniformly substituting individual constants for variables which are not bound by a quantifier.
- ▶ $t \Diamond F t$, $\neg t \Diamond \exists x G t x$ and $t \Diamond x R t x$ are all formulas.
- ▶ $\Diamond F t$ (no scope for t) and $t R t u$ (no scope for u) are not.

PRIMITIVE SCOPE MARKERS

- Semantically the idea is that only terms which have wide scope are required to denote in order for a formula containing those terms to be true.

PRIMITIVE SCOPE MARKERS

- Semantically the idea is that only terms which have wide scope are required to denote in order for a formula containing those terms to be true.
- Our rules for the existential quantifier, then, are the following (where in $[\exists L]$ a must be an *eigenvariable*):

$$\frac{\mathcal{H} \mid X \vdash Y, u_1 \dots u_m u A_u^x}{\mathcal{H} \mid X \vdash Y, u_1 \dots u_m \exists x A} [\exists R]$$

$$\frac{\mathcal{H} \mid X, u_1 \dots u_m a A_a^x \vdash Y}{\mathcal{H} \mid X, u_1 \dots u_m \exists x A \vdash Y} [\exists L]$$

So our rules for the quantifiers only bind variables whose scope-marking occurrence has wide-scope.

OPAQUE & TRANSPARENT RULES

- ▶ With our scope markers we can now draw a distinction between two different kinds of insertion rules for connective.

- ▶ With our scope markers we can now draw a distinction between two different kinds of insertion rules for connective.
- ▶ *Transparent* insertion rules make the terms available to be bound in the premises so available in the conclusion.

- ▶ With our scope markers we can now draw a distinction between two different kinds of insertion rules for connective.
- ▶ *Transparent* insertion rules make the terms available to be bound in the premises so available in the conclusion.
- ▶ Disjunction, for example, has transparent insertion rules

$$\frac{\mathcal{H} \mid X, u_1 \dots u_m A \vdash Y \quad \mathcal{H} \mid X, u_1 \dots u_m B \vdash Y}{\mathcal{H} \mid X, u_1 \dots u_m (A \vee B) \vdash Y} [\vee L]$$

$$\frac{\mathcal{H} \mid X \vdash u_1 \dots u_m A, u_1 \dots u_m B, Y}{\mathcal{H} \mid X \vdash u_1 \dots u_m (A \vee B), Y} [\vee R]$$

- ▶ *Opaque* insertion rules make terms which are available to be bound in the premise unavailable in the conclusion.

- ▶ *Opaque* insertion rules make terms which are available to be bound in the premise unavailable in the conclusion.
- ▶ The rules for the modal operators are opaque in this sense.

$$\frac{\mathcal{H} \mid X \vdash Y \quad \mid \quad t_1 \dots t_n A \vdash}{\mathcal{H} \mid X, \Diamond t_1 \dots t_n A \vdash Y} [\Diamond L]$$

$$\frac{\mathcal{H} \mid X \vdash Y \quad \mid \quad X' \vdash Y', t_1 \dots t_n A}{\mathcal{H} \mid X \vdash Y, \Diamond t_1 \dots t_n A \quad \mid \quad X' \vdash Y'} [\Diamond R]$$

- ▶ *Opaque* insertion rules make terms which are available to be bound in the premise unavailable in the conclusion.
- ▶ The rules for the modal operators are opaque in this sense.

$$\frac{\mathcal{H} \mid X \vdash Y \quad \mid \quad t_1 \dots t_n A \vdash}{\mathcal{H} \mid X, \Diamond t_1 \dots t_n A \vdash Y} [\Diamond L]$$

$$\frac{\mathcal{H} \mid X \vdash Y \quad \mid \quad X' \vdash Y', t_1 \dots t_n A}{\mathcal{H} \mid X \vdash Y, \Diamond t_1 \dots t_n A \quad \mid \quad X' \vdash Y'} [\Diamond R]$$

- ▶ Similarly the rules for negation are also opaque.

- ▶ We don't want to ban quantification into modal contexts altogether, though.

- ▶ We don't want to ban quantification into modal contexts altogether, though.
- ▶ We just want to restrict when such quantification can occur.

- ▶ We don't want to ban quantification into modal contexts altogether, though.
- ▶ We just want to restrict when such quantification can occur.
- ▶ *Exportation Rules* allow for this possibility. The exportation rules for \Diamond are the following, where for $[\Diamond\text{ExpR}]$ we require that there be some formula a $B \in X$.

$$\frac{\mathcal{H} \mid X \vdash Y, \Diamond aA}{\mathcal{H} \mid X \vdash Y, a\Diamond A} [\Diamond\text{ExpR}] \qquad \frac{\mathcal{H} \mid X, \Diamond aA \vdash Y}{\mathcal{H} \mid X, a\Diamond A \vdash Y} [\Diamond\text{ExpL}]$$

UNRESTRICTED EXPORTATION & THE BARCAN FORMULA

- ▶ The Barcan formula is not derivable in the system sketched above.

- ▶ The Barcan formula is not derivable in the system sketched above.
- ▶ But it is derivable in the system we get by replacing the $[\Diamond \text{ExpR}]$ rule with the rule of ‘unrestricted exportation’, which drops the restriction placed on that rule.

- ▶ The Barcan formula is not derivable in the system sketched above.
- ▶ But it is derivable in the system we get by replacing the $[\Diamond \text{ExpR}]$ rule with the rule of ‘unrestricted exportation’, which drops the restriction placed on that rule.

$$\begin{array}{c}
 \frac{aFa \vdash aFa}{aFa \vdash \quad | \quad \vdash \Diamond aFa} [\Diamond R] \\
 \frac{aFa \vdash \quad | \quad \vdash \Diamond aFa}{aFa \vdash \quad | \quad \vdash a\Diamond Fa} [UE] \\
 \frac{aFa \vdash \quad | \quad \vdash a\Diamond Fa}{aFa \vdash \quad | \quad \vdash \exists x \Diamond Fx} [\exists R] \\
 \frac{aFa \vdash \quad | \quad \vdash \exists x \Diamond Fx}{\exists x Fx \vdash \quad | \quad \vdash \exists x \Diamond Fx} [\exists L] \\
 \frac{\exists x Fx \vdash \quad | \quad \vdash \exists x \Diamond Fx}{\Diamond \exists x Fx \vdash \exists x \Diamond Fx} [\Diamond L]
 \end{array}$$

- ▶ This approach gives us a very natural explanation of what goes ‘wrong’ in Barcan derivations—they involve exportation inferences which are hidden in the standard language.

- ▶ This approach gives us a very natural explanation of what goes ‘wrong’ in Barcan derivations—they involve exportation inferences which are hidden in the standard language.
- ▶ Moreover, the only justification for those inferences appears to be question begging (e.g. all terms denote an existent in every possibility).

- ▶ This approach gives us a very natural explanation of what goes ‘wrong’ in Barcan derivations—they involve exportation inferences which are hidden in the standard language.
- ▶ Moreover, the only justification for those inferences appears to be question begging (e.g. all terms denote an existent in every possibility).
- ▶ So we have a system which does away with the unwanted superstition without causing *too* much formal awkwardness.

THANK YOU!