

# Axioms of Probability

## Axioms of Probability

- For any event  $A$ ,

$$P(A) \geq 0$$


## Axioms of Probability


- For any event  $A$ ,

$$P(A) \geq 0$$

- If  $A_1, A_2, A_3, \dots, A_n$  are disjoint events (i.e.,  $A_i \cap A_j = \phi \quad \forall i \neq j$ ) then

$$P(\cup A_i) = \sum_i P(A_i)$$





## Axioms of Probability

- For any event  $A$ ,


$$P(A) \geq 0$$

- If  $A_1, A_2, A_3, \dots, A_n$  are disjoint events (i.e.,  $A_i \cap A_j = \phi \quad \forall i \neq j$ ) then

$$P(\cup A_i) = \sum_i P(A_i)$$


- If  $\Omega$  is the universal set containing all events then

$$P(\Omega) = 1$$




## Random Variable (intuition)

- Suppose a student can get one of 3 possible grades in a course:  $A, B, C$




## Random Variable (intuition)

- Suppose a student can get one of 3 possible grades in a course:  $A, B, C$
- One way of interpreting this is that there are 3 possible events here



## Random Variable (intuition)


- Suppose a student can get one of 3 possible grades in a course:  $A, B, C$
- One way of interpreting this is that there are 3 possible events here
- Another way of looking at this is there is a *random variable*  $G$  which each student to one of the 3 possible values



## Random Variable (intuition)

- Suppose a student can get one of 3 possible grades in a course:  $A, B, C$
- One way of interpreting this is that there are 3 possible events here
- Another way of looking at this is there is a *random variable*  $G$  which each student to one of the 3 possible values
- And we are interested in  $P(G = g)$  where  $g \in \{A, B, C\}$







## Random Variable (intuition)

- Suppose a student can get one of 3 possible grades in a course:  $A, B, C$
- One way of interpreting this is that there are 3 possible events here
- Another way of looking at this is there is a *random variable*  $G$  which each student to one of the 3 possible values
- And we are interested in  $P(G = g)$  where  $g \in \{A, B, C\}$
- Of course, both interpretations are conceptually equivalent

## Random Variable (intuition)


- But the second one (using random variables) is more compact






## Random Variable (intuition)

- But the second one (using random variables) is more compact
- Specially, when there are multiple attributes associated with a student (outcome) - *grade, height, age, etc.*



## Random Variable (intuition)


- But the second one (using random variables) is more compact
- Specially, when there are multiple attributes associated with a student (outcome) - *grade, height, age, etc.*
- We could have one random variable corresponding to each attribute



## Random Variable (intuition)


- But the second one (using random variables) is more compact
- Specially, when there are multiple attributes associated with a student (outcome) - *grade, height, age, etc.*
- We could have one random variable corresponding to each attribute
- And then ask for outcomes (or students) where  $Grade = g$ ,  $Height = h$ ,  $Age = a$  and so on


## Random Variable (formal)



## Random Variable (formal)

- A random variable is a *function* which maps each outcome in  $\Omega$  to a value






## Random Variable (formal)


- A random variable is a *function* which maps each outcome in  $\Omega$  to a value
- In the previous example,  $G$  (or  $f_{grade}$ ) maps each student in  $\Omega$  to a value:  $A$ ,  $B$  or  $C$






## Random Variable (formal)

- A random variable is a *function* which maps each outcome in  $\Omega$  to a value
- In the previous example,  $G$  (or  $f_{\text{grade}}$ ) maps each student in  $\Omega$  to a value:  $A$ ,  $B$  or  $C$
- The event  $\text{Grade} = A$  is a shorthand for the event  $\{\omega \in \Omega : f_{\text{Grade}} = A\}$




## Random Variable (continuous v/s discrete)

- A random variable can either take continuous values (for example, *weight, height*)



## Random Variable (continuous v/s discrete)

- A random variable can either take continuous values (for example, *weight, height*)
- Or discrete values (for example, *grade, nationality*)



## Random Variable (continuous v/s discrete)

- A random variable can either take continuous values (for example, *weight, height*)
- Or discrete values (for example, *grade, nationality*)
- For this discussion we will mainly focus on discrete random variables

## Marginal Distribution

- What do we mean by *marginal distribution* over a random variable ?

## Marginal Distribution

- What do we mean by *marginal distribution* over a random variable ?
- Consider our random variable  $G$  for grades

$G$	$P(G = g)$
A	0.1
B	0.2
C	0.7

## Marginal Distribution

- What do we mean by *marginal distribution* over a random variable ?
- Consider our random variable  $G$  for grades
- Specifying the marginal distribution over  $G$  means specifying

$$P(G = g) \quad \forall g \in A, B, C$$

$G$	$P(G = g)$
A	0.1
B	0.2
C	0.7

## Marginal Distribution

- What do we mean by *marginal distribution* over a random variable ?
- Consider our random variable  $G$  for grades
- Specifying the marginal distribution over  $G$  means specifying

$$P(G = g) \quad \forall g \in A, B, C$$

- We denote this marginal distribution compactly by  $P(G)$



## Joint Distribution

- Consider two random variable  $G$  (grade) and  $I$  (intelligence  $\in \{\mathbf{High}, \mathbf{Low}\}$ )

## Joint Distribution

- Consider two random variable  $G$  (grade) and  $I$  (intelligence  $\in \{\mathbf{H}igh, \mathbf{L}ow\}$ )
- The joint distribution over these two random variables assigns probabilities to all events involving these two random variables

$$P(G = g, I = i) \quad \forall (g, i) \in \{A, B, C\} \times \{H, L\}$$

## Joint Distribution

- Consider two random variable  $G$  (grade) and  $I$  (intelligence  $\in \{\mathbf{H}igh, \mathbf{L}ow\}$ )
- The joint distribution over these two random variables assigns probabilities to all events involving these two random variables

$G$	$I$	$P(G = g, I = i)$
A	High	0.3
A	Low	0.1
B	High	0.15
B	Low	0.15
C	High	0.1
C	Low	0.2

$$P(G = g, I = i) \quad \forall (g, i) \in \{A, B, C\} \times \{H, L\}$$

## Joint Distribution

$G$	$I$	$P(G = g, I = i)$
A	High	0.3
A	Low	0.1
B	High	0.15
B	Low	0.15
C	High	0.1
C	Low	0.2

- Consider two random variable  $G$  (grade) and  $I$  (intelligence  $\in \{\mathbf{H}igh, \mathbf{L}ow\}$ )
- The joint distribution over these two random variables assigns probabilities to all events involving these two random variables

$$P(G = g, I = i) \quad \forall (g, i) \in \{A, B, C\} \times \{H, L\}$$

- We denote this joint distribution compactly by  $P(G, I)$

## Conditional Distribution

- Consider two random variable  $G$  (grade) and  $I$  (intelligence)

$G$	$P(G I = H)$
A	0.6
B	0.3
C	0.1

$G$	$P(G I = L)$
A	0.3
B	0.4
C	0.3

## Conditional Distribution

$G$	$P(G I = H)$
A	0.6
B	0.3
C	0.1

$G$	$P(G I = L)$
A	0.3
B	0.4
C	0.3

- Consider two random variable  $G$  (grade) and  $I$  (intelligence)
- Suppose we are given the value of  $I$  (say,  $I = H$ ) then the conditional distribution  $P(G|I)$  is defined as

$$P(G = g|I = H) = \frac{P(G = g, I = H)}{P(I = H)} \forall g \in \{A, B, C\}$$

## Conditional Distribution

$G$	$P(G I = H)$
A	0.6
B	0.3
C	0.1

$G$	$P(G I = L)$
A	0.3
B	0.4
C	0.3

- Consider two random variable  $G$  (grade) and  $I$  (intelligence)
- Suppose we are given the value of  $I$  (say,  $I = H$ ) then the conditional distribution  $P(G|I)$  is defined as

$$P(G = g|I = H) = \frac{P(G = g, I = H)}{P(I = H)} \forall g \in \{A, B, C\}$$

- More compactly defined as

$$P(G|I) = \frac{P(G, I)}{P(I)}$$

or

$$\underbrace{P(G, I)}_{\text{joint}} = \underbrace{P(G|I)}_{\text{conditional}} * \underbrace{P(I)}_{\text{marginal}}$$

## Joint Distribution ( $n$ random variables)

- The joint distribution of  $n$  random variables assigns probabilities to all events involving the  $n$  random variables,

$X_1$	$\dots$	$X_n$	$P(X_1, X_2, \dots, X_n)$
$\dots$	$\dots$	$\dots$	$\dots$
$\dots$	$\dots$	$\dots$	$\dots$
$\dots$	$\dots$	$\dots$	$\dots$

$$\sum = 1$$



## Joint Distribution ( $n$ random variables)

- The joint distribution of  $n$  random variables assigns probabilities to all events involving the  $n$  random variables,
- In other words it assigns

$$P(X_1 = x_1, X_2 = x_2, \dots, X_n = x_n)$$

*for all possible values that variable  $X_i$  can take*

$X_1$	$\dots$	$X_n$	$P(X_1, X_2, \dots, X_n)$
$\dots$	$\dots$	$\dots$	$\dots$
$\dots$	$\dots$	$\dots$	$\dots$
$\dots$	$\dots$	$\dots$	$\dots$

$$\sum = 1$$

## Joint Distribution ( $n$ random variables)

$X_1$	$\dots$	$X_n$	$P(X_1, X_2, \dots, X_n)$
$\dots$	$\dots$	$\dots$	$\dots$
$\dots$	$\dots$	$\dots$	$\dots$
$\dots$	$\dots$	$\dots$	$\dots$

$$\sum = 1$$

- The joint distribution of  $n$  random variables assigns probabilities to all events involving the  $n$  random variables,
- In other words it assigns

$$P(X_1 = x_1, X_2 = x_2, \dots, X_n = x_n)$$

*for all possible values that variable  $X_i$  can take*

- If each random variable  $X_i$  can take two values then the joint distribution will assign probabilities to the  $2^n$  possible events

## Joint Distribution ( $n$ random variables)

- The joint distribution over two random variables  $X_1$  and  $X_2$  can be written as,

$$P(X_1, X_2) = P(X_2|X_1)P(X_1) = P(X_1|X_2)P(X_2)$$

$X_1$	$\dots$	$X_n$	$P(X_1, X_2, \dots, X_n)$
$\dots$	$\dots$	$\dots$	$\dots$
$\dots$	$\dots$	$\dots$	$\dots$
$\dots$	$\dots$	$\dots$	$\dots$

## Joint Distribution ( $n$ random variables)

$X_1$	$\dots$	$X_n$	$P(X_1, X_2, \dots, X_n)$
$\dots$	$\dots$	$\dots$	$\dots$
$\dots$	$\dots$	$\dots$	$\dots$
$\dots$	$\dots$	$\dots$	$\dots$

- The joint distribution over two random variables  $X_1$  and  $X_2$  can be written as,

$$P(X_1, X_2) = P(X_2|X_1)P(X_1) = P(X_1|X_2)P(X_2)$$

- Similarly for  $n$  random variables

$$P(X_1, X_2, \dots, X_n)$$

## Joint Distribution ( $n$ random variables)

$X_1$	$\dots$	$X_n$	$P(X_1, X_2, \dots, X_n)$
$\dots$	$\dots$	$\dots$	$\dots$
$\dots$	$\dots$	$\dots$	$\dots$
$\dots$	$\dots$	$\dots$	$\dots$

- The joint distribution over two random variables  $X_1$  and  $X_2$  can be written as,

$$P(X_1, X_2) = P(X_2|X_1)P(X_1) = P(X_1|X_2)P(X_2)$$

- Similarly for  $n$  random variables

$$\begin{aligned} P(X_1, X_2, \dots, X_n) \\ = P(X_2, \dots, X_n|X_1)P(X_1) \end{aligned}$$

## Joint Distribution ( $n$ random variables)

$X_1$	$\dots$	$X_n$	$P(X_1, X_2, \dots, X_n)$
$\dots$	$\dots$	$\dots$	$\dots$
$\dots$	$\dots$	$\dots$	$\dots$
$\dots$	$\dots$	$\dots$	$\dots$

- The joint distribution over two random variables  $X_1$  and  $X_2$  can be written as,

$$P(X_1, X_2) = P(X_2|X_1)P(X_1) = P(X_1|X_2)P(X_2)$$

- Similarly for  $n$  random variables

$$\begin{aligned} &P(X_1, X_2, \dots, X_n) \\ &= P(X_2, \dots, X_n|X_1)P(X_1) \\ &= P(X_3, \dots, X_n|X_1, X_2)P(X_2|X_1)P(X_1) \end{aligned}$$

## Joint Distribution ( $n$ random variables)

$X_1$	$\dots$	$X_n$	$P(X_1, X_2, \dots, X_n)$
$\dots$	$\dots$	$\dots$	$\dots$
$\dots$	$\dots$	$\dots$	$\dots$
$\dots$	$\dots$	$\dots$	$\dots$

- The joint distribution over two random variables  $X_1$  and  $X_2$  can be written as,

$$P(X_1, X_2) = P(X_2|X_1)P(X_1) = P(X_1|X_2)P(X_2)$$

- Similarly for  $n$  random variables

$$\begin{aligned} &P(X_1, X_2, \dots, X_n) \\ &= P(X_2, \dots, X_n|X_1)P(X_1) \\ &= P(X_3, \dots, X_n|X_1, X_2)P(X_2|X_1)P(X_1) \\ &= P(X_4, \dots, X_n|X_1, X_2, X_3)P(X_3|X_2, X_1) \\ &\quad P(X_2|X_1)P(X_1) \end{aligned}$$

## Joint Distribution ( $n$ random variables)

$X_1$	$\dots$	$X_n$	$P(X_1, X_2, \dots, X_n)$
$\dots$	$\dots$	$\dots$	$\dots$
$\dots$	$\dots$	$\dots$	$\dots$
$\dots$	$\dots$	$\dots$	$\dots$

- The joint distribution over two random variables  $X_1$  and  $X_2$  can be written as,

$$P(X_1, X_2) = P(X_2|X_1)P(X_1) = P(X_1|X_2)P(X_2)$$

- Similarly for  $n$  random variables

$$\begin{aligned} &P(X_1, X_2, \dots, X_n) \\ &= P(X_2, \dots, X_n|X_1)P(X_1) \\ &= P(X_3, \dots, X_n|X_1, X_2)P(X_2|X_1)P(X_1) \\ &= P(X_4, \dots, X_n|X_1, X_2, X_3)P(X_3|X_2, X_1) \\ &\quad P(X_2|X_1)P(X_1) \\ &= P(X_1) \prod_{i=2}^n P(X_i|X_1^{i-1}) \quad (\text{chain rule}) \end{aligned}$$



## From Joint Distributions to Marginal Distributions

$A$	$B$	$P(A = a, B = b)$
High	High	0.3
High	Low	0.25
Low	High	0.35
Low	Low	0.1

$A$	$P(A = a)$
High	0.55
Low	0.45

$B$	$P(B = a)$
High	0.65
Low	0.35

- Suppose we are given a joint distribution over two random variables  $A, B$

## From Joint Distributions to Marginal Distributions

$A$	$B$	$P(A = a, B = b)$
High	High	0.3
High	Low	0.25
Low	High	0.35
Low	Low	0.1

$A$	$P(A = a)$
High	0.55
Low	0.45

$B$	$P(B = a)$
High	0.65
Low	0.35

- Suppose we are given a joint distribution over two random variables  $A, B$
- The marginal distributions of  $A$  and  $B$  can be computed as

$$P(A = a) = \sum_{\forall b} P(A = a, B = b)$$

$$P(B = b) = \sum_{\forall a} P(A = a, B = b)$$

$A$	$B$	$P(A = a, B = b)$
High	High	0.3
High	Low	0.25
Low	High	0.35
Low	Low	0.1

$A$	$P(A = a)$
High	0.55
Low	0.45

$B$	$P(B = a)$
High	0.65
Low	0.35

## From Joint Distributions to Marginal Distributions

- Suppose we are given a joint distribution over two random variables  $A, B$
- The marginal distributions of  $A$  and  $B$  can be computed as

$$P(A = a) = \sum_{\forall b} P(A = a, B = b)$$

$$P(B = b) = \sum_{\forall a} P(A = a, B = b)$$

- More compactly written as

$$P(A) = \sum_B P(A, B)$$

$$P(B) = \sum_A P(A, B)$$

What if there are  $n$  random variables ?

$A$	$B$	$P(A = a, B = b)$
High	High	0.3
High	Low	0.25
Low	High	0.35
Low	Low	0.1

$A$	$P(A = a)$
High	0.55
Low	0.45

$B$	$P(B = a)$
High	0.65
Low	0.35

- Suppose we are given a joint distribution over  $n$  random variables  $X_1, X_2, \dots, X_n$

## What if there are $n$ random variables ?

$A$	$B$	$P(A = a, B = b)$
High	High	0.3
High	Low	0.25
Low	High	0.35
Low	Low	0.1

$A$	$P(A = a)$
High	0.55
Low	0.45

$B$	$P(B = a)$
High	0.65
Low	0.35

- Suppose we are given a joint distribution over  $n$  random variables  $X_1, X_2, \dots, X_n$
- The marginal distributions over  $X_1$  can be computed as

$$\begin{aligned} P(X_1 = x_1) \\ = \sum_{\forall x_2, x_3, \dots, x_n} P(X_1 = x_1, X_2 = x_2, \dots, X_n = x_n) \end{aligned}$$

A	B	$P(A = a, B = b)$
High	High	0.3
High	Low	0.25
Low	High	0.35
Low	Low	0.1

A	$P(A = a)$
High	0.55
Low	0.45

B	$P(B = a)$
High	0.65
Low	0.35

What if there are  $n$  random variables ?

- Suppose we are given a joint distribution over  $n$  random variables  $X_1, X_2, \dots, X_n$
- The marginal distributions over  $X_1$  can be computed as

$$\begin{aligned}
 P(X_1 = x_1) &= \sum_{\forall x_2, x_3, \dots, x_n} P(X_1 = x_1, X_2 = x_2, \dots, X_n = x_n)
 \end{aligned}$$

- More compactly written as

$$P(X_1) = \sum_{X_2, X_3, \dots, X_n} P(X_1, X_2, \dots, X_n)$$

## Conditional Independence

- Two random variables  $X$  and  $Y$  are said to be independent if

$$P(X|Y) = P(X)$$

## Conditional Independence

- Two random variables  $X$  and  $Y$  are said to be independent if

$$P(X|Y) = P(X)$$

- We denote this as  $X \perp\!\!\!\perp Y$



## Conditional Independence

- Two random variables  $X$  and  $Y$  are said to be independent if

$$P(X|Y) = P(X)$$

- We denote this as  $X \perp\!\!\!\perp Y$
- In other words, knowing the value of  $Y$  does not change our belief about  $X$

## Conditional Independence

- Two random variables  $X$  and  $Y$  are said to be independent if

$$P(X|Y) = P(X)$$

- We denote this as  $X \perp\!\!\!\perp Y$
- In other words, knowing the value of  $Y$  does not change our belief about  $X$
- We would expect *Grade* to be dependent on *Intelligence* but independent of *Weight*

## Conditional Independence

- Recall that by Chain Rule of Probability

$$P(X, Y) = P(X)P(Y|X)$$

- Two random variables  $X$  and  $Y$  are said to be independent if

$$P(X|Y) = P(X)$$

- We denote this as  $X \perp\!\!\!\perp Y$
- In other words, knowing the value of  $Y$  does not change our belief about  $X$
- We would expect *Grade* to be dependent on *Intelligence* but independent of *Weight*

## Conditional Independence

- Recall that by Chain Rule of Probability

$$P(X, Y) = P(X)P(Y|X)$$

- However, if  $X$  and  $Y$  are independent, then

$$P(X, Y) = P(X)P(Y)$$

- Two random variables  $X$  and  $Y$  are said to be independent if

$$P(X|Y) = P(X)$$

- We denote this as  $X \perp\!\!\!\perp Y$
- In other words, knowing the value of  $Y$  does not change our belief about  $X$
- We would expect *Grade* to be dependent on *Intelligence* but independent of *Weight*