

Exercise 6.1

()

Exercise 6.2

(LIF in weak gravitational field)

Exercise 6.2: Local inertial frame in a weak gravitational field

A four-dimensional manifold has coordinates (t, x, y, z) and line element

$$ds^2 = -(1 + 2\phi)dt^2 + (1 - 2\phi)(dx^2 + dy^2 + dz^2), \quad (6)$$

where $|\phi(t, x, y, z)| \ll 1$ everywhere.

(a) At any point P with coordinates (t_0, x_0, y_0, z_0) find a coordinate transformation to a locally inertial coordinate system, to first order in ϕ . (1 pt)

- We have to perform a change of coordinates such that the metric in the new frame, $\bar{g}_{\mu\nu}$, is locally flat around P
- This means that $\bar{g}_{\mu\nu} = \eta_{\mu\nu}$
- The first derivatives of $\bar{g}_{\mu\nu}$ at P must vanish,

$$\bar{g}_{\mu\nu}(x) = \eta_{\mu\nu} + \frac{\partial^2 g_{\mu\nu}}{\partial x^\alpha \partial x^\beta} \Big|_P (x^\alpha - x_P^\alpha)(x^\beta - x_P^\beta) + \mathcal{O}(\Delta x^3) \quad (7)$$

with

$$\bar{g}_{\bar{\mu}\bar{\nu}} = \frac{\partial x^\mu}{\partial x^{\bar{\mu}}} \frac{\partial x^\nu}{\partial x^{\bar{\nu}}} g_{\mu\nu}.$$

We can define

$$\Lambda^\mu_{\bar{\nu}} \equiv \frac{\partial x^\mu}{\partial x^{\bar{\nu}}}.$$

As η is a diagonal metric we can try with a diagonal transformation matrix Λ leading also to a diagonal metric $\bar{g}_{\mu\nu}$ (this is the simplest initial guess). Thus, we have,

$$\begin{aligned} \bar{g}_{00}(x) &= [\Lambda^0_{\bar{0}}(x)]^2 g_{00}(x) = -[1 + 2\phi(x)][\Lambda^0_{\bar{0}}(x)]^2 \\ \bar{g}_{ii}(x) &= [\Lambda^i_{\bar{i}}(x)]^2 g_{ii}(x) = [1 - 2\phi(x)][\Lambda^i_{\bar{i}}(x)]^2. \end{aligned}$$

In order $\bar{g}_{\mu\nu}$ to take the form (7) the transformation matrix must fulfill the following equations:

$$\begin{aligned} -1 &= -[1 + 2\phi(P)][\Lambda^0_{\bar{0}}(P)]^2 \longrightarrow \Lambda^0_{\bar{0}}(P) = 1 - \phi(P) + \mathcal{O}(\phi^2) \\ 1 &= [1 - 2\phi(P)][\Lambda^i_{\bar{i}}(P)]^2 \xrightarrow{\text{Taylor expansion}} \Lambda^i_{\bar{i}}(P) = 1 + \phi(P) + \mathcal{O}(\phi^2) \end{aligned}$$

and

$$\begin{aligned} 0 &= 2 \frac{\partial \Lambda^0_{\bar{0}}}{\partial x^\mu} \Big|_P g_{00}(P) + [\Lambda^0_{\bar{0}}(P)]^2 \frac{\partial g_{00}}{\partial x^\mu} \Big|_P \longrightarrow \frac{\partial \Lambda^0_{\bar{0}}}{\partial x^\mu} \Big|_P = -\partial_\mu \phi \Big|_P + \mathcal{O}(\phi^2) \\ 0 &= 2 \frac{\partial \Lambda^i_{\bar{i}}}{\partial x^\mu} \Big|_P g_{ii}(P) + [\Lambda^i_{\bar{i}}(P)]^2 \frac{\partial g_{ii}}{\partial x^\mu} \Big|_P \longrightarrow \frac{\partial \Lambda^i_{\bar{i}}}{\partial x^\mu} \Big|_P = \partial_\mu \phi \Big|_P + \mathcal{O}(\phi^2). \end{aligned}$$

The elements of the transformation matrix thus read,

$$\begin{aligned} \Lambda^0_{\bar{0}}(x) &= 1 - \phi(P) - \partial_\mu \phi \Big|_P (x^\mu - x_P^\mu) + \mathcal{O}(\phi^2) \\ \Lambda^i_{\bar{i}}(x) &= 1 + \phi(P) + \partial_\mu \phi \Big|_P (x^\mu - x_P^\mu) + \mathcal{O}(\phi^2). \end{aligned}$$

and with the non-diagonal terms equal to zero.

(b) At what rate does such a frame accelerate with respect to the original coordinates, again to first order in ϕ ? Consider that the free-falling observer moves with non-relativistic velocity in the non-inertial system. (1 pt)

The world-line of the free-falling frame measured by an observer located in the non-inertial frame obeys the following geodesic equation,

$$\frac{d^2 x^i}{d\tau^2} = -\Gamma^i_{\mu\nu} \frac{dx^\mu}{d\tau} \frac{dx^\nu}{d\tau} = -\Gamma^i_{00} \left(\frac{dt}{d\tau} \right)^2 - \Gamma^i_{jl} \frac{dx^j}{d\tau} \frac{dx^l}{d\tau} - 2\Gamma^i_{j0} \frac{dx^j}{d\tau} \frac{dt}{d\tau}$$

The Christoffel symbols are of order $\Gamma \sim \mathcal{O}(\phi)$ and since the free-falling body moves with non-relativistic velocity in the non-inertial frame, $d\tau = dt + \mathcal{O}(v, \phi)$. Thus, it is clear that the second term in the right-hand side is of order $\mathcal{O}(3)$ and the last one of order $\mathcal{O}(2)$. The leading contribution comes from the first term and is of order $\mathcal{O}(\phi)$. Hence, $\frac{d^2 x^i}{dt^2} = -\Gamma^i_{00} + \mathcal{O}(\phi^2) = -\partial_i \phi + \mathcal{O}(\phi^2)$

ϕ can be interpreted as the Newtonian gravitational potential.

(c) Compute the elements of the Riemann tensor in the original (non-inertial) frame to first order in ϕ . (1 pt)

The Christoffel symbols are as follows :

$$\begin{aligned}\Gamma^0_{00} &= \partial_0\phi + \mathcal{O}(\phi^2) & \Gamma^i_{00} &= \partial^i\phi + \mathcal{O}(\phi^2) \\ \Gamma^0_{ij} &= -\delta_{ij}\partial_0\phi + \mathcal{O}(\phi^2) & \Gamma^i_{0j} &= -\delta^i_j\partial_0\phi + \mathcal{O}(\phi^2) \\ \Gamma^0_{0i} &= \partial_i\phi + \mathcal{O}(\phi^2) & \Gamma^i_{jl} &= \delta_{jl}\partial^i\phi - \delta^i_j\partial_l\phi - \delta^i_l\partial_j\phi + \mathcal{O}(\phi^2)\end{aligned}$$

Notice that $\partial_i\phi = \partial^i\phi + \mathcal{O}(\phi)$, so partial derivatives with upper and lower indices can be interchanged if we work to first order in ϕ . The components of the Riemann tensor will take the following form if we neglect contributions of order $\mathcal{O}(\phi^2)$ or higher,

$$R^\mu{}_{\nu\alpha\beta} = \partial_\alpha\Gamma^\mu{}_{\nu\beta} - \partial_\beta\Gamma^\mu{}_{\nu\alpha} + \mathcal{O}(\phi^2)$$

Thus, we have,

$$\begin{aligned}R^\mu{}_{\nu 00} &= \mathcal{O}(\phi^2) \\ R^\mu{}_{\nu 0i} &= \partial_0\Gamma^\mu{}_{\nu i} - \partial_i\Gamma^\mu{}_{\nu 0} + \mathcal{O}(\phi^2) \\ R^\mu{}_{\nu ij} &= \partial_i\Gamma^\mu{}_{\nu j} - \partial_j\Gamma^\mu{}_{\nu i} + \mathcal{O}(\phi^2)\end{aligned}$$

Equivalently,

$$\begin{aligned} R^\mu{}_{\nu 00} &= \mathcal{O}(\phi^2) \\ R^0{}_{00i} &= \mathcal{O}(\phi^2) \\ R^0{}_{j0i} &= -\delta_{ij}\partial_0^2\phi - \partial_i\partial_j\phi + \mathcal{O}(\phi^2) \\ R^j{}_{00i} &= -\delta_i^j\partial_0^2\phi - \partial_i\partial^j\phi + \mathcal{O}(\phi^2) \\ R^j{}_{l0i} &= \delta_{li}\partial_0\partial^j\phi - \delta_i^j\partial_0\partial_l\phi + \mathcal{O}(\phi^2) \\ R^0{}_{0ij} &= \mathcal{O}(\phi^2) \\ R^0{}_{lij} &= -\delta_{lj}\partial_i\partial_0\phi + \delta_{li}\partial_j\partial_0\phi + \mathcal{O}(\phi^2) \\ R^l{}_{0ij} &= -\delta_j^l\partial_i\partial_0\phi + \delta_i^l\partial_j\partial_0\phi + \mathcal{O}(\phi^2) \\ R^l{}_{kij} &= \delta_{kj}\partial_i\partial^l\phi - \delta_j^l\partial_i\partial_k\phi - \delta_{ki}\partial_j\partial^l\phi + \delta_i^l\partial_j\partial_k\phi + \mathcal{O}(\phi^2) \end{aligned}$$

Exercise 6.3

(Killing vectors and conserved quantities)

Exercise 6.3: Killing vectors and conserved quantities

(a) Consider the scalar quantity $u^\alpha \xi_\alpha$, with \vec{u} the four-velocity of a free-falling particle. What equation must be fulfilled by $\vec{\xi}$ if this scalar remains constant along the particle's trajectory? (1.5 pt)

If $u^\alpha \xi_\alpha$ is constant along the particle's trajectory it has to satisfy the following equation: $\frac{d}{d\tau} (u^\alpha \xi_\alpha) = 0$,

where τ is the proper time of the particle.

This equation can be rewritten as follows,

$$\frac{du^\alpha}{d\tau} \xi_\alpha + u^\alpha \frac{d\xi_\alpha}{d\tau} = 0 \longrightarrow (u^\beta \nabla_\beta u^\alpha) \xi_\alpha + u^\alpha u^\beta \nabla_\beta \xi_\alpha = 0.$$

A free particle satisfies the geodesic equation $u^\beta \nabla_\beta u^\alpha = 0$, so we have $\longrightarrow u^\alpha u^\beta \nabla_\beta \xi_\alpha = 0$.

Regardless of the 4-velocity of the free-falling particle this equation is automatically fulfilled if $\nabla_\beta \xi_\alpha$ is antisymmetric, i.e. if

$$\boxed{\nabla_\beta \xi_\alpha + \nabla_\alpha \xi_\beta = 0} \quad \text{This is Killing's equation.}$$

(b) Express (6) in spherical coordinates (t, r, θ, φ) . (1 pt)

$$ds^2 = -(1 + 2\phi)dt^2 + (1 - 2\phi)(dx^2 + dy^2 + dz^2)$$

$$x = r \sin(\theta) \cos(\varphi)$$

$$y = r \sin(\theta) \sin(\varphi)$$

$$z = r \cos(\theta).$$

Thus, we have,

$$dx = \sin(\theta) \cos(\varphi)dr + r \cos(\theta) \cos(\varphi)d\theta - r \sin(\theta) \sin(\varphi)d\varphi$$

$$dy = \sin(\theta) \sin(\varphi)dr + r \cos(\theta) \sin(\varphi)d\theta + r \sin(\theta) \cos(\varphi)d\varphi$$

$$dz = \cos(\theta)dr - r \sin(\theta)d\theta.$$

This leads to:

$$ds^2 = -(1 + 2\phi)dt^2 + (1 - 2\phi)[dr^2 + r^2\{d\theta^2 + \sin^2(\theta)d\varphi^2\}]$$

(c) Assume now that the metric under consideration is static and spherically symmetric. Find two linearly independent Killing vectors in this particular spacetime, and prove that p_0 and p_φ are conserved along the trajectory of a free-falling particle.

$$\nabla_\beta \xi_\alpha + \nabla_\alpha \xi_\beta = 0 \longrightarrow \partial_\mu \xi_\nu + \partial_\nu \xi_\mu - 2\Gamma^\alpha_{\mu\nu} \xi_\alpha = 0$$

$$ds^2 = -(1+2\phi)dt^2 + (1-2\phi)[dr^2 + r^2\{d\theta^2 + \sin^2(\theta)d\varphi^2\}]$$

For the spacetime determined by the line element (50) with $\phi = \phi(r)$ there are only nine Christoffel symbols different from zero, namely $\Gamma^\theta_{\theta r}, \Gamma^\theta_{\varphi\varphi}, \Gamma^0_{0r}, \Gamma^\varphi_{\varphi r}, \Gamma^\varphi_{\varphi\theta}, \Gamma^r_{rr}, \Gamma^r_{\theta\theta}, \Gamma^r_{\varphi\varphi}$ and Γ^r_{00} . Each Killing vector has to satisfy the following set of equations:

$$\begin{aligned} \partial_0 \xi_0 - \Gamma^r_{00} \xi_r &= 0 & \partial_0 \xi_\theta + \partial_\theta \xi_0 &= 0 \\ \partial_r \xi_r - \Gamma^r_{rr} \xi_r &= 0 & \partial_0 \xi_\varphi + \partial_\varphi \xi_0 &= 0 \\ \partial_\theta \xi_\theta - \Gamma^r_{\theta\theta} \xi_r &= 0 & \partial_r \xi_\theta + \partial_\theta \xi_r - 2\Gamma^\theta_{r\theta} \xi_\theta &= 0 \\ \partial_\varphi \xi_\varphi - \Gamma^\theta_{\varphi\varphi} \xi_\theta - \Gamma^r_{\varphi\varphi} \xi_r &= 0 & \partial_r \xi_\varphi + \partial_\varphi \xi_r - 2\Gamma^\varphi_{r\varphi} \xi_\varphi &= 0 \\ \partial_0 \xi_r + \partial_r \xi_0 - 2\Gamma^0_{0r} \xi_0 &= 0 & \partial_\theta \xi_\varphi + \partial_\varphi \xi_\theta - 2\Gamma^\varphi_{\varphi\theta} \xi_\varphi &= 0 \end{aligned}$$

Let us see whether there exists a Killing vector with $\xi_r = \xi_\theta = \xi_\varphi = 0$. Using this in the Killing equations we find:

$$\partial_\varphi \xi_0 = \partial_\theta \xi_0 = \partial_0 \xi_0 = 0$$

$$\partial_r \xi_0 - 2\Gamma^0_{0r} \xi_0 = 0, \quad \text{so } \xi_0 = \xi_0(r)$$

Using $\Gamma^0_{0r} = \frac{\partial_r \phi}{1+2\phi}$ we can solve the last equation and obtain:

$$\tilde{d}\xi = A (1 + 2\phi, 0, 0, 0) \longrightarrow \boxed{\vec{\xi} = A (1, 0, 0, 0)} \quad \text{where } A \text{ is an integration constant.}$$

Now let us search for another Killing vector, but with $\xi_r = \xi_\theta = \xi_0 = 0$.

The following equations have to be fulfilled,

$$\begin{aligned} \partial_\varphi \xi_\varphi &= \partial_\theta \xi_r = \partial_0 \xi_\varphi &= 0 \\ \partial_r \xi_\varphi - 2\Gamma^\varphi_{\varphi r} \xi_\varphi &= 0 \\ \partial_\theta \xi_\varphi - 2\Gamma^\varphi_{\varphi \theta} \xi_\varphi &= 0. \end{aligned}$$

In this case $\xi_\varphi = \xi_\varphi(r, \theta)$. As $\Gamma^\varphi_{\varphi r} = 1/r - \partial_r \phi / (1 - 2\phi)$ and $\Gamma^\varphi_{\varphi \theta} = \cot g(\theta)$ we find:

$$\begin{aligned} \partial_r \xi_\varphi - \frac{2}{r} \xi_\varphi &= 0 \\ \partial_\theta \xi_\varphi - 2 \cot g(\theta) \xi_\varphi &= 0. \end{aligned}$$

The solution reads,

$$\tilde{d}\xi = B (0, 0, 0, r^2 \sin^2(\theta)(1 - 2\phi)) \longrightarrow \boxed{\vec{\xi} = B (0, 0, 0, 1)}$$

Any linear combination of the two fields will also satisfy the Killing equation

It is clear that along the trajectory of a free-falling particle,

$$u_0 = C_1 \quad ; \quad u_\varphi = C_2 ,$$

or, equivalently,

$$p_0 = \bar{C}_1 \quad ; \quad p_\varphi = \bar{C}_2 .$$

(d) Show that these conservation laws can be also obtained directly from the geodesic equation. (1 pt)

The geodesic equation reads, $u^\nu \nabla_\nu u_\mu = 0 \longrightarrow u^\nu \partial_\nu u_\mu = \Gamma^\alpha_{\nu\mu} u_\alpha u^\nu$

$$\frac{du_\mu}{d\tau} = \frac{1}{2} g_{\beta\nu,\mu} u^\beta u^\nu \longrightarrow m \frac{dp_\mu}{d\tau} = \frac{1}{2} g_{\beta\nu,\mu} p^\beta p^\nu$$

Since the metric does not depend on time nor the angle φ we find that p_0 and p_φ remain constant along the trajectory of a free-falling particle.