

## 2nd April Shift 1


**Q1** A light wave is propagating with plane wave fronts of the type  $x + y + z = \text{constant}$ . The angle made by the direction of wave propagation with the x-axis is :

- (A)  $\cos^{-1} \left( \frac{1}{\sqrt{3}} \right)$  (B)  $\cos^{-1} \left( \frac{2}{3} \right)$   
 (C)  $\cos^{-1} \left( \frac{1}{3} \right)$  (D)  $\cos^{-1} \left( \sqrt{\frac{2}{3}} \right)$

**Q2** The equation for real gas is given by  $\left( P + \frac{a}{V^2} \right) (V - b) = RT$ , where P, V, T and R are the pressure, volume, temperature and gas constant, respectively. The dimension of  $ab^{-2}$  is equivalent to that of :

- (A) Planck's constant  
 (B) Compressibility  
 (C) Strain  
 (D) Energy density


**Q3** A cord of negligible mass is wound around the rim of a wheel supported by spokes with negligible mass. The mass of wheel is 10 kg and radius is 10 cm and it can freely rotate without any friction. Initially the wheel is at rest. If a steady pull of 20 N is applied on the cord, the angular velocity of the wheel, after the cord is unwound by 1 m, would be :



- (A) 20 rad/s (B) 30 rad/s  
 (C) 10 rad/s (D) 0 rad/s


**Q4** A slanted object AB is placed on one side of convex lens as shown in the diagram. The image is formed on the opposite side. Angle made by

the image with principal axis is :



- (A)  $-\frac{\alpha}{2}$  (B)  $-45^\circ$   
 (C)  $+45^\circ$  (D)  $-\alpha$

**Q5** Consider two infinitely large plane parallel conducting plates as shown below. The plates are uniformly charged with a surface charge density  $+\sigma$  and  $-2\sigma$ . The force experienced by a point charge  $+q$  placed at the mid point between two plates will be:



- (A)  $\frac{\sigma q}{4\epsilon_0}$  (B)  $\frac{3\sigma q}{2\epsilon_0}$   
 (C)  $\frac{3\sigma q}{4\epsilon_0}$  (D)  $\frac{\sigma q}{2\epsilon_0}$

**Q6** A river is flowing from west to east direction with speed of 9 km/h. If a boat capable of moving at a maximum speed of 27 km/h in still water, crosses the river in half a minute, while moving with maximum speed at an angle of  $150^\circ$  to direction of river flow, then the width of the river is :

- (A) 300 m  
 (B) 112.5 m  
 (C) 75 m  
 (D)  $112.5 \times \sqrt{3}$  m


**Q7** A point charge  $+q$  is placed at the origin. A second point charge  $+9q$  is placed at  $(d, 0, 0)$  in Cartesian coordinate system. The point in between them where the electric field vanishes is :

- (A)  $(4d/3, 0, 0)$



- (B)  $(d/4, 0, 0)$   
 (C)  $(3d/4, 0, 0)$   
 (D)  $(d/3, 0, 0)$
- Q8** The battery of a mobile phone is rated as  $4.2\text{ V}$ ,  $5800\text{ mAh}$ . How much energy is stored in it when fully charged?  
 (A)  $43.8\text{ kJ}$  (B)  $48.7\text{ kJ}$   
 (C)  $87.7\text{ kJ}$  (D)  $24.4\text{ kJ}$
- Q9** A particle is subjected two simple harmonic motions as :  
 $x_1 = \sqrt{7} \sin 5t\text{ cm}$  and  
 $x_2 = 2\sqrt{7} \sin (5t + \frac{\pi}{3})\text{ cm}$  where  $x$  is displacement and  $t$  is time in seconds. The maximum acceleration of the particle is  $x \times 10^{-2}\text{ ms}^{-2}$ . The value of  $x$  is :  
 (A) 175 (B)  $25\sqrt{7}$   
 (C)  $5\sqrt{7}$  (D) 125
- Q10** The relationship between the magnetic susceptibility ( $\chi$ ) and the magnetic permeability ( $\mu$ ) is given by :  
 ( $\mu_0$  is the permeability of free space and  $\mu_r$  is relative permeability)  
 (A)  $\chi = \frac{\mu}{\mu_0} - 1$   
 (B)  $\chi = \frac{\mu_r}{\mu_0} + 1$   
 (C)  $\chi = \mu_r + 1$   
 (D)  $\chi = 1 - \frac{\mu}{\mu_0}$
- Q11** A zener diode with  $5\text{ V}$  zener voltage is used to regulate an unregulated de voltage input of  $25\text{ V}$ . For a  $400\Omega$  resistor connected in series, the zener current is found to be 4 times load current. The load current ( $I_L$ ) and load resistance ( $R_L$ ) are :  
 (A)  $I_L = 20\text{ mA}$ ;  $R_L = 250\Omega$   
 (B)  $I_L = 10\text{ A}$ ;  $R_L = 0.5\Omega$   
 (C)  $I_L = 0.02\text{ mA}$ ;  $R_L = 250\Omega$   
 (D)  $I_L = 10\text{ mA}$ ;  $R_L = 500\Omega$
- Q12** In an adiabatic process, which of the following statements is true ?  
 (A) The molar heat capacity is infinite  
 (B) Work done by the gas equals the increase in internal energy  
 (C) The molar heat capacity is zero  
 (D) The internal energy of the gas decreases as the temperature increases


- Q13** A square Lamina OABC of length  $10\text{ cm}$  is pivoted at 'O'. Forces act at Lamina as shown in figure. If Lamina remains stationary, then the magnitude of  $F$  is :



- (A) 20 N (B) 0 (zero)  
 (C) 10 N (D)  $10\sqrt{2}\text{ N}$
- Q14** Let  $B_1$  be the magnitude of magnetic field at center of a circular coil of radius  $R$  carrying current  $I$ . Let  $B_2$  be the magnitude of magnetic field at an axial distance ' $x$ ' from the center. For  $x : R = 3 : 4$ ,  $\frac{B_2}{B_1}$  is :  
 (A) 4 : 5 (B) 16 : 25  
 (C) 64 : 125 (D) 25 : 16
- Q15** Considering Bohr's atomic model for hydrogen atom :  
 (A) the energy of H atom in ground state is same as energy of  $\text{He}^+$  ion in its first excited state.  
 (B) the energy of H atom in ground state is same as that for  $\text{Li}^{++}$  ion in its second excited state.  
 (C) the energy of H atom in its ground state is same as that of  $\text{He}^+$  ion for its ground state.  
 (D) the energy of  $\text{He}^+$  ion in its first excited state is same as that for  $\text{Li}^{++}$  ion in its ground state  
 (A) (B), (D) only  
 (B) (A), (B) only  
 (C) (A), (D) only  
 (D) (A), (C) only
- Q16** Moment of inertia of a rod of mass ' $M$ ' and length ' $L$ ' about an axis passing through its center and normal to its length is ' $\alpha$ '. Now the rod is cut into two equal parts and these parts are joined symmetrically to form a cross shape. Moment of inertia of cross about an axis passing through its center and normal to plane containing cross is :  
 (A)  $\alpha$  (B)  $\alpha/4$   
 (C)  $\alpha/8$  (D)  $\alpha/2$

**Q17**





A spherical surface separates two media of refractive indices 1 and 1.5 as shown in figure.

Distance of the image of an object 'O', is :

(C is the center of curvature of the spherical surface and R is the radius of curvature)

- (A) 0.24 m right to the spherical surface  
 (B) 0.4 m left to the spherical surface  
 (C) 0.24 m left to the spherical surface  
 (D) 0.4 m right to the spherical surface

**Q18 Match List-I with List-II.**

**List-I**

**List-II**


- |                              |                        |
|------------------------------|------------------------|
| (A) Coefficient of viscosity | (I) $[ML^0T^{-3}]$     |
| (B) Intensity of wave        | (II) $[ML^{-2}T^{-2}]$ |
| (C) Pressure gradient        | (III) $[M^{-1}LT^2]$   |
| (D) Compressibility          | (IV) $[ML^{-1}T^{-1}]$ |

Choose the correct answer from the options given below :

- (A) (A)–(I), (B)–(IV), (C)–(III), (D)–(II)  
 (B) (A)–(IV), (B)–(I), (C)–(II), (D)–(III)  
 (C) (A)–(IV), (B)–(II), (C)–(I), (D)–(III)  
 (D) (A)–(II), (B)–(III), (C)–(IV), (D)–(I)

- Q19** A small bob of mass 100 mg and charge  $+10 \mu C$  is connected to an insulating string of length 1 m. It is brought near to an infinitely long nonconducting sheet of charge density ' $\sigma$ ' as shown in figure. If string subtends an angle of  $45^\circ$  with the sheet at equilibrium the charge density of sheet will be :

(Given,  $\epsilon_0 = 8.85 \times 10^{-12} \text{ F/m}$  and acceleration due to gravity,  $g = 10 \text{ m/s}^2$ )



- (A)  $0.885 \text{ nC/m}^2$   
 (B)  $17.7 \text{ nC/m}^2$   
 (C)  $885 \text{ nC/m}^2$   
 (D)  $1.77 \text{ nC/m}^2$

- Q20** A monochromatic light is incident on a metallic plate having work function  $\phi$ . An electron, emitted normally to the plate from a point A with maximum kinetic energy, enters a constant magnetic field, perpendicular to the initial velocity of electron. The electron passes through a curve and hits back the plate at a point B. The distance between A and B is :

(Given : The magnitude of charge of an electron is  $e$  and mass is  $m$ ,  $h$  is Planck's constant and  $c$  is velocity of light. Take the magnetic field exists throughout the path of electron)

- (A)  $\sqrt{2 m \left( \frac{hc}{\lambda} - \phi \right)} / eB$   
 (B)  $\sqrt{m \left( \frac{hc}{\lambda} - \phi \right)} / eB$   
 (C)  $\sqrt{8 m \left( \frac{hc}{\lambda} - \phi \right)} / eB$   
 (D)  $2\sqrt{m \left( \frac{hc}{\lambda} - \phi \right)} / eB$

- Q21** A vessel with square cross-section and height of 6 m is vertically partitioned. A small window of  $100 \text{ cm}^2$  with hinged door is fitted at a depth of 3 m in the partition wall. One part of the vessel is filled completely with water and the other side is filled with the liquid having density  $1.5 \times 10^3 \text{ kg/m}^3$ . What force one needs to apply on the hinged door so that it does not get opened ?

(Acceleration due to gravity =  $10 \text{ m/s}^2$ )

- Q22** A steel wire of length 2 m and Young's modulus  $2.0 \times 10^{11} \text{ Nm}^{-2}$  is stretched by a force. If Poisson ratio and transverse strain for the wire are 0.2 and  $10^{-3}$  respectively, then the elastic potential energy density of the wire is \_\_\_\_\_  $\times 10^5$  (in SI units)

- Q23** If the measured angular separation between the second minimum to the left of the central maximum and the third minimum to the right of the central maximum is  $30^\circ$  in a single slit diffraction pattern recorded using  $628 \text{ nm}$  light, then the width of the slit is \_\_\_\_\_  $\mu\text{m}$ .

- Q24**  $\gamma_A$  is the specific heat ratio of monoatomic gas A having 3 translational degrees of freedom.  $\gamma_B$  is the specific heat ratio of polyatomic gas B having 3 translational, 3 rotational degrees of freedom and 1 vibrational mode. If  $\frac{\gamma_A}{\gamma_B} = \left(1 + \frac{1}{n}\right)$ , then the value of  $n$  is \_\_\_\_\_.



**Q25** A person travelling on a straight line moves with a uniform velocity  $v_1$  for a distance  $x$  and with a uniform velocity  $v_2$  for the next  $\frac{3}{2}x$  distance. The average velocity in this motion is  $\frac{50}{7}$  m/s. If  $v_1$  is 5 m/s then  $v_2 =$  \_\_\_\_ m/s.

**Q26** The largest  $n \in \mathbb{N}$  such that  $3^n$  divides  $50!$  is:

- (A) 21 (B) 22  
(C) 20 (D) 23

**Q27** Let one focus of the hyperbola  $H : \frac{x^2}{a^2} - \frac{y^2}{b^2} = 1$  be at  $(\sqrt{10}, 0)$  and the corresponding directrix be  $x = \frac{9}{\sqrt{10}}$ . If  $e$  and  $l$  respectively are the eccentricity and the length of the latus rectum of  $H$ , then  $9(e^2 + l)$  is equal to:

- (A) 14 (B) 15  
(C) 16 (D) 12

**Q28** The number of sequences of ten terms, whose terms are either 0 or 1 or 2, that contain exactly five 1s and exactly three 2s, is equal to

- (A) 360 (B) 45  
(C) 2520 (D) 1820

**Q29** Let  $f : \mathbb{R} \rightarrow \mathbb{R}$  be a twice differentiable function such that

$$(\sin x \cos y)(f(2x + 2y) - f(2x - 2y)) = (\cos x \sin y)(f(2x + 2y) + f(2x - 2y))$$

for all  $x, y \in \mathbb{R}$ .

If  $f'(0) = \frac{1}{2}$ , then the value of  $24f''\left(\frac{5\pi}{3}\right)$  is:

- (A) 2 (B) -3  
(C) 3 (D) -2

**Q30** Let  $A = \begin{bmatrix} \alpha & -1 \\ 6 & \beta \end{bmatrix}$ ,  $\alpha > 0$ , such that  $\det(A) = 0$  and  $\alpha + \beta = 1$ . If  $I$  denotes  $2 \times 2$  identity matrix, then the matrix  $(I + A)^8$  is:

- (A)  $\begin{bmatrix} 4 & -1 \\ 6 & -1 \end{bmatrix}$   
(B)  $\begin{bmatrix} 257 & -64 \\ 514 & -127 \end{bmatrix}$   
(C)  $\begin{bmatrix} 1025 & -511 \\ 2024 & -1024 \end{bmatrix}$   
(D)  $\begin{bmatrix} 766 & -255 \\ 1530 & -509 \end{bmatrix}$

**Q31** The term independent of  $x$  in the expansion of

$$\left( \frac{(x+1)}{(x^{2/3} + 1 - x^{1/3})} - \frac{(x-1)}{(x - x^{1/2})} \right)^{10}, x > 1 \text{ is:}$$

- (A) 210 (B) 150

- (C) 240 (D) 120

**Q32** If  $\theta \in [-2\pi, 2\pi]$ , then the number of solutions of  $2\sqrt{2} \cos^2 \theta + (2 - \sqrt{6}) \cos \theta - \sqrt{3} = 0$ , is equal to:

- (A) 12 (B) 6  
(C) 8 (D) 10

**Q33** Let  $a_1, a_2, a_3 \dots$  be in an A.P. such that

$$\sum_{k=1}^{12} a_{2k-1} = -\frac{72}{5}a_1, a_1 \neq 0. \text{ If}$$

$$\sum_{k=1}^n a_k = 0, \text{ then } n \text{ is:}$$

- (A) 11 (B) 10  
(C) 18 (D) 17

**Q34** If the function

$$f(x) = 2x^3 - 9ax^2 + 12a^2x + 1, \text{ where } a$$

$> 0$ , attains its local maximum and local minimum values at  $p$  and  $q$ , respectively, such that  $p^2 = q$ , then  $f(3)$  is equal to:

- (A) 55 (B) 10  
(C) 23 (D) 37

**Q35** Let  $z$  be a complex number such that  $|z| = 1$ . If  $\frac{2+k^2z}{k+\bar{z}} = kz, k \in \mathbb{R}$ , then the maximum distance of  $k + ik^2$  from the circle  $|z - (1 + 2i)| = 1$  is:

- (A)  $\sqrt{5} + 1$  (B) 2  
(C) 3 (D)  $\sqrt{3} + 1$

**Q36** If  $\vec{a}$  is nonzero vector such that its projections on the vectors  $2\hat{i} - \hat{j} + 2\hat{k}, \hat{i} + 2\hat{j} - 2\hat{k}$  and  $\hat{k}$  are equal, then a unit vector along  $\vec{a}$  is:

- (A)  $\frac{1}{\sqrt{155}}(-7\hat{i} + 9\hat{j} + 5\hat{k})$   
(B)  $\frac{1}{\sqrt{155}}(-7\hat{i} + 9\hat{j} - 5\hat{k})$   
(C)  $\frac{1}{\sqrt{155}}(7\hat{i} + 9\hat{j} + 5\hat{k})$   
(D)  $\frac{1}{\sqrt{155}}(7\hat{i} + 9\hat{j} - 5\hat{k})$

**Q37** Let  $A$  be the set of all functions  $f : \mathbb{Z} \rightarrow \mathbb{Z}$  and  $R$  be a relation on  $A$  such that

$$R = \{(f, g) : f(0) = g(1) \text{ and } f(1) = g(0)\}.$$

Then  $R$  is:

- (A) Symmetric and transitive but not reflexive  
(B) Symmetric but neither reflexive nor transitive  
(C) Reflexive but neither symmetric nor transitive  
(D) Transitive but neither reflexive nor symmetric

**Q38** For  $\alpha, \beta, \gamma \in \mathbb{R}$ , if  $\lim_{x \rightarrow 0} \frac{x^2 \sin \alpha x + (\gamma - 1)e^{x^2}}{\sin 2x - \beta x} = 3$ , then  $\beta + \gamma - \alpha$  is equal to:



- (A) 7 (B) 4  
(C) 6 (D) -1

**Q39** If the system of linear equations

$$3x + y + \beta z = 3$$

$$2x + \alpha y - z = -3$$

$$x + 2y + z = 4$$

has infinitely many solutions, then the value of  $22\beta - 9\alpha$  is :

- (A) 49 (B) 31  
(C) 43 (D) 37

**Q40** Let  $P_n = \alpha^n + \beta^n, n \in \mathbf{N}$ . If

$P_{10} = 123, P_9 = 76, P_8 = 47$  and  $P_1 = 1$ , then the quadratic equation having roots  $\frac{1}{\alpha}$  and  $\frac{1}{\beta}$  is :

- (A)  $x^2 - x + 1 = 0$  (B)  $x^2 + x - 1 = 0$   
(C)  $x^2 - x - 1 = 0$  (D)  $x^2 + x + 1 = 0$

**Q41** If  $S$  and  $S'$  are the foci of the ellipse

$\frac{x^2}{18} + \frac{y^2}{9} = 1$  and  $P$  be a point on the ellipse, then  $\min(SP, S'P) + \max(SP, S'P)$  is equal to :

- (A)  $3(1 + \sqrt{2})$  (B)  $3(6 + \sqrt{2})$   
(C) 9 (D) 27

**Q42** Let the vertices Q and R of the triangle PQR lie on the line  $\frac{x+3}{5} = \frac{y-1}{2} = \frac{z+4}{3}$ ,  $QR = 5$  and the coordinates of the point P be  $(0, 2, 3)$ . If the area of the triangle PQR is  $\frac{m}{n}$  then :

- (A)  $m - 5\sqrt{21}n = 0$   
(B)  $2m - 5\sqrt{21}n = 0$   
(C)  $5m - 2\sqrt{21}n = 0$   
(D)  $5m - 21\sqrt{2}n = 0$

**Q43** Let ABCD be a tetrahedron such that the edges AB, AC and AD are mutually perpendicular. Let the areas of the triangles ABC, ACD and ADB be 5, 6 and 7 square units respectively. Then the area (in square units) of the  $\Delta BCD$  is equal to :

- (A)  $\sqrt{340}$  (B) 12  
(C)  $\sqrt{110}$  (D)  $7\sqrt{3}$

**Q44** Let  $a \in \mathbf{R}$  and A be a matrix of order  $3 \times 3$  such

$$\text{that } \det(A) = -4 \text{ and } A + I = \begin{bmatrix} 1 & a & 1 \\ 2 & 1 & 0 \\ a & 1 & 2 \end{bmatrix},$$

where  $I$  is the identity matrix of order  $3 \times 3$ .

If  $\det((a+1) \text{adj}((a)A))$  is  $2^m 3^n$ ,  $m, n \in \{0, 1, 2, \dots, 20\}$ , then  $m + n$  is equal to :

- (A) 14 (B) 17  
(C) 15 (D) 16

**Q45** Let the focal chord PQ of the parabola  $y^2 = 4x$  make an angle of  $60^\circ$  with the positive x-axis, where P lies in the first quadrant. If the circle, whose one diameter is PS, S being the focus of the parabola, touches the y-axis at the point  $(0, \alpha)$ , then  $5\alpha^2$  is equal to :

- (A) 15 (B) 25  
(C) 30 (D) 20

**Q46** Let  $[x]$  denote the greatest integer function. If

$$\int_0^{e^3} \left[ \frac{1}{e^{x-1}} \right] dx = \alpha - \log_e 2, \text{ then } \alpha^3 \text{ is equal to}$$

**Q47** Let  $f : \mathbf{R} \rightarrow \mathbf{R}$  be a thrice differentiable odd function satisfying

$$f'(x) \geq 0, f''(x) = f(x), f(0) = 0, f'(0) = 3$$

Then  $9f(\log_e 3)$  is equal to \_\_\_\_\_.


**Q48** If the area of the region

$$\{(x, y) : |4 - x^2| \leq y \leq x^2, y \leq 4, x \geq 0\} \text{ is } \left( \frac{80\sqrt{2}}{\alpha} - \beta \right), \alpha, \beta \in \mathbf{N}, \text{ then } \alpha + \beta \text{ is equal to}$$

**Q49** Three distinct numbers are selected randomly from the set  $\{1, 2, 3, \dots, 40\}$ . If the probability, that the selected numbers are in an increasing G.P. is  $\frac{m}{n}$ ,  $\gcd(m, n) = 1$ , then  $m + n$  is equal to

**Q50** The absolute difference between the squares of the radii of the two circles passing through the point  $(-9, 4)$  and touching the lines  $x + y = 3$  and  $x - y = 3$ , is equal to \_\_\_\_\_.

**Q51** Designate whether each of the following compounds is aromatic or not aromatic



- (A) e, g aromatic and a, b, c, d, f, h not aromatic  
(B) b, e, f, g aromatic and a, c, d, h not aromatic



- (C) a, b, c, d aromatic and e, f, g, h not aromatic  
 (D) a, c, d, e, h aromatic and b, f, g not aromatic

**Q52** An optically active alkyl halide  $C_4H_9Br$  [A] reacts with hot KOH dissolved in ethanol and forms alkene [B] as major product which reacts with bromine to give dibromide [C]. The compound [C] is converted into a gas [D] upon reacting with alcoholic  $NaNH_2$ . During hydration 18 gram of water is added to 1 mole of gas [D] on warming with mercuric sulphate and dilute acid at 333 K to form compound [E]. The IUPAC name of compound [E] is :

- (A) But - 2 - yne (B) Butan - 2 - ol  
 (C) Butan - 2 - one (D) Butan - 1 - al



**Q53** The property/properties that show irregularity in first four elements of group-17 is/are :

- (A) Covalent radius  
 (B) Electron affinity  
 (C) Ionic radius  
 (D) First ionization energy



Choose the **correct** answer from the options given below:

- (A) B and D only (B) A and C only  
 (C) B only (D) A, B, C and D

**Q54** Which of the following graph correctly represents the plots of  $K_H$  at 1 bar gases in water versus temperature ?




(C)



**Q55** According to Bohr's model of hydrogen atom, which of the following statement is **incorrect**?

- (A) Radius of 3<sup>rd</sup> orbit is nine times larger than that of 1<sup>st</sup> orbit.  
 (B) Radius of 8<sup>th</sup> orbit is four times larger than that of 4<sup>th</sup> orbit.  
 (C) Radius of 6<sup>th</sup> orbit is three time larger than that of 4<sup>th</sup> orbit.  
 (D) Radius of 4<sup>th</sup> orbit is four times larger than that of 2<sup>nd</sup> orbit.

**Q56** Thermometer



Two vessels A and B are connected via stopcock. The vessel A is filled with a gas at a certain pressure. The entire assembly is immersed in water and is allowed to come to thermal equilibrium with water. After opening the stopcock the gas from vessel A expands into vessel B and no change in temperature is observed in the thermometer. Which of the following statement is true?

- (A)  $dw \neq 0$   
 (B)  $dq \neq 0$   
 (C)  $dU \neq 0$   
 (D) The pressure in the vessel B before opening the stopcock is zero.

**Q57** A solution is made by mixing one mole of volatile liquid A with 3 moles of volatile liquid B. The





vapour pressure of pure A is 200 mm Hg and that of the solution is 500 mm Hg. The vapour pressure of pure B and the least volatile component of the solution, respectively, are :


- (A) 1400 mmHg, A  
(B) 1400 mmHg, B  
(C) 600 mmHg, B  
(D) 600 mmHg, A

- Q58**  $\text{CaCO}_3(s) + 2\text{HCl}(aq) \rightarrow \text{CaCl}_2(aq) + \text{CO}_2(g) + \text{H}_2\text{O}(l)$   
Consider the above reaction, what mass of  $\text{CaCl}_2$  will be formed if 250 mL of 0.76 M reacts with 1000 g of  $\text{CaCO}_3$  ?  
(Given : Molar mass of Ca, C, O, H and Cl are 40, 12, 16, 1 and 35.5  $\text{g mol}^{-1}$ , respectively)  
(A) 3.908 g (B) 2.636 g  
(C) 10.545 g (D) 5.272 g


- Q59** If equal volumes of  $\text{AB}_2$  and  $\text{XY}$  (both are salts) aqueous solutions are mixed, which of the following combination will give a precipitate of  $\text{AY}_2$  at 300 K ?  
(Given  $K_{sp}$  (at 300 K) for  $\text{AY}_2 = 5.2 \times 10^{-7}$ )  
(A)  $3.6 \times 10^{-3}\text{M AB}_2$ ,  $5.0 \times 10^{-4}\text{M XY}$   
(B)  $2.0 \times 10^{-4}\text{M AB}_2$ ,  $0.8 \times 10^{-3}\text{M XY}$   
(C)  $2.0 \times 10^{-2}\text{M AB}_2$ ,  $2.0 \times 10^{-2}\text{M XY}$   
(D)  $1.5 \times 10^{-4}\text{M AB}_2$ ,  $1.5 \times 10^{-3}\text{M XY}$

- Q60** Among  $\text{SO}_2$ ,  $\text{NF}_3$ ,  $\text{NH}_3$ ,  $\text{XeF}_2$ ,  $\text{ClF}_3$  and  $\text{SF}_4$ , the hybridization of the molecule with non-zero dipole moment and highest number of lone-pairs of electrons on the central atom is  
(A)  $\text{sp}^3$  (B)  $\text{dsp}^2$   
(C)  $\text{sp}^3 \text{d}^2$  (D)  $\text{sp}^3 \text{d}$

- Q61** Given below are two statements :



**Statement (I) :** will react with NaOH and also with Tollen's reagent.



**Statement (II) :** will undergo self aldol condensation very easily.

In the light of the above statements, choose the **most appropriate answer** from the options given below :

- (A) Statement I is incorrect but Statement II is correct  
(B) Statement I is correct but Statement II is incorrect  
(C) Both Statement I and Statement II are incorrect  
(D) Both Statement I and Statement II are correct

- Q62** Identify the correct statement among the following:  
(A) All naturally occurring amino acids except glycine contain one chiral centre.  
(B) All naturally occurring amino acids are optically active.  
(C) Glutamic acid is the only amino acid that contains a  $-\text{COOH}$  group at the side chain.  
(D) Amino acid, cysteine easily undergo dimerization due to the presence of free SH group.

- Q63** The correct order of basic nature on aqueous solution for the bases  $\text{NH}_3$ ,  $\text{H}_2\text{N}-\text{NH}_2$ ,  $\text{CH}_3\text{CH}_2\text{NH}_2$ , and  $(\text{CH}_3\text{CH}_2)_2\text{NH}$ ,  $(\text{CH}_3\text{CH}_2)_3\text{N}$  is  
(A)  $\text{NH}_3 < \text{H}_2\text{N}-\text{NH}_2 < (\text{CH}_3\text{CH}_2)_3\text{N} < \text{CH}_3\text{CH}_2\text{NH}_2 < (\text{CH}_3\text{CH}_2)_2\text{NH}$   
(B)  $\text{NH}_3 < \text{H}_2\text{N}-\text{NH}_2 < \text{CH}_3\text{CH}_2\text{NH}_2 < (\text{CH}_3\text{CH}_2)_2\text{NH} < (\text{CH}_3\text{CH}_2)_3\text{N}$   
(C)  $\text{H}_2\text{N}-\text{NH}_2 < \text{NH}_3 < (\text{CH}_3\text{CH}_2)_3\text{N} < \text{CH}_3\text{CH}_2\text{NH}_2 < (\text{CH}_3\text{CH}_2)_2\text{NH}$   
(D)  $\text{NH}_2-\text{NH}_2 < \text{NH}_3 < \text{CH}_3\text{CH}_2\text{NH}_2 < (\text{CH}_3\text{CH}_2)_3\text{N} < (\text{CH}_3\text{CH}_2)_2\text{NH}$

- Q64** Given below are two statements :

**Statement (I) :** The metallic radius of Al is less than that of Ga.

**Statement (II) :** The ionic radius of  $\text{Al}^{3+}$  is less than that of  $\text{Ga}^{3+}$ .

In the light of the above statements, choose the most appropriate answer from the options given below :

- (A) Both Statement I and Statement II are incorrect  
(B) Statement I is incorrect but Statement II is correct  
(C) Statement I is correct but Statement II is incorrect  
(D) Both Statement I and Statement II are correct



**Q65** Given below are two statements :

**Statement (I) :** In octahedral complexes, when  $\Delta_o < P$  high spin complexes are formed. When  $\Delta_o > P$  low spin complexes are formed.

**Statement (II) :** In tetrahedral complexes because of  $\Delta_t < P$ , low spin complexes are rarely formed.

In the light of the above statements, choose the most appropriate answer from the options given below :

- (A) Statement I is correct but Statement II is incorrect.  
 (B) Both Statement I and Statement II are incorrect  
 (C) Statement I is incorrect but Statement II is correct  
 (D) Both Statement I and Statement II are correct

**Q66** Choose the correct tests with respective observations.

- (A)  $\text{CuSO}_4$  (acidified with acetic acid) +  $\text{K}_4[\text{Fe}(\text{CN})_6] \rightarrow$  Chocolate brown precipitate.  
 (B)  $\text{FeCl}_3 + \text{K}_4[\text{Fe}(\text{CN})_6] \rightarrow$  Prussian blue precipitate.  
 (C)  $\text{ZnCl}_2 + \text{K}_4[\text{Fe}(\text{CN})_6]$ , neutralised with  $\text{NH}_4\text{OH} \rightarrow$  White or bluish white precipitate.  
 (D)  $\text{MgCl}_2 + \text{K}_4[\text{Fe}(\text{CN})_6] \rightarrow$  Blue precipitate.  
 (E)  $\text{BaCl}_2 + \text{K}_4[\text{Fe}(\text{CN})_6]$ , neutralised with  $\text{NaOH} \rightarrow$  White precipitate.

Choose the correct answer from the options given below :

- (A) A, D and E only  
 (B) B, D and E only  
 (C) A, B and C only  
 (D) C, D and E only


**Q67** On complete combustion 1.0 g of an organic compound (X) gave 1.46 g of  $\text{CO}_2$  and 0.567 g of  $\text{H}_2\text{O}$ . The empirical formula mass of compound (X) is \_\_\_\_\_ g.

(Given molar mass in  $\text{g mol}^{-1}$  C : 12, H : 1, O : 16)

- (A) 30 (B) 45  
 (C) 60 (D) 15

**Q68**





Consider the following compound (X)



The most stable and least stable carbon radicals, respectively, produced by homolytic cleavage of corresponding C – H bond are :

- (A) II, IV (B) III, II  
 (C) I, IV (D) II, I

**Q69** Consider the following molecules :



The correct order of rate of hydrolysis is :

- (A)  $r > q > p > s$   
 (B)  $q > p > r > s$   
 (C)  $p > r > q > s$   
 (D)  $p > q > r > s$

**Q70** A molecule with the formula  $\text{AX}_4\text{Y}$  has all its elements from p-block. Element A is rarest, monoatomic, non-radioactive from its group and has the lowest ionization enthalpy value among A, X and Y. Elements X and Y have first and second highest electronegativity values respectively among all the known elements. The shape of the molecule is :

- (A) Square pyramidal  
 (B) Octahedral  
 (C) Pentagonal planar  
 (D) Trigonal bipyramidal


**Q71** A transition metal (M) among Mn, Cr, Co and Fe has the highest standard electrode potential ( $\text{M}^{3+}/\text{M}^{2+}$ ). It forms a metal complex of the type  $[\text{M}(\text{CN})_6]^{4-}$ . The number of electrons






present in the  $e_g$  orbital of the complex is \_\_\_\_\_.

- Q72** Consider the following electrochemical cell at standard condition.



$$E_{\text{cell}} = +0.4 \text{ V}$$


The couple  $\text{QH}_2/\text{Q}$  represents quinhydrone electrode, the half cell reaction is given below



$$\left[ \text{Given : } E_{\text{Ag}^+/\text{Ag}}^{\circ} = +0.8 \text{ V and } \frac{2.303 RT}{F} = 0.06 \text{ V} \right]$$

The  $\text{pK}_b$  value of the ammonium halide salt ( $\text{NH}_4\text{X}$ ) used here is \_\_\_\_\_. (nearest integer)

- Q73** 0.1 mol of the following given antiviral compound (P) will weigh \_\_\_\_\_  $\times 10^{-1}$  g




(Given : molar mass in  $\text{g mol}^{-1}$  H : 1, C : 12, N : 14, O : 16, F : 19, I : 127)

- Q74** Consider the following equilibrium,  
 $\text{CO(g)} + 2\text{H}_2(\text{g}) \rightleftharpoons \text{CH}_3\text{OH(g)}$ , 0.1 mol of CO along with a catalyst is present in a  $2 \text{ dm}^3$  flask maintained at 500 K. Hydrogen is introduced into the flask until the pressure is 5 bar and 0.04 mol of  $\text{CH}_3\text{OH}$  is formed. The  $K_p^{\circ}$  is \_\_\_\_\_  $\times 10^{-3}$  (nearest integer).  
 Given :  $R = 0.08 \text{ dm}^3 \text{ bar K}^{-1} \text{ mol}^{-1}$

Assume only methanol is formed as the product and the system follows ideal gas behaviour.

**Q75**

For the reaction  $\text{A} \rightarrow \text{products}$



The concentration of A at 10 minutes is \_\_\_\_\_  $\times 10^{-3} \text{ mol L}^{-1}$  (nearest integer).

The reaction was started with  $2.5 \text{ mol L}^{-1}$  of A.



Android App | iOS App | PW Website

## Answer Key

Q1 (A)  
Q2 (D)  
Q3 (A)  
Q4 (B)  
Q5 (B)  
Q6 (B)  
Q7 (B)  
Q8 (C)  
Q9 (A)  
Q10 (A)  
Q11 (D)  
Q12 (C)  
Q13 (C)  
Q14 (C)  
Q15 (B)  
Q16 (B)  
Q17 (B)  
Q18 (B)  
Q19 (D)  
Q20 (C)  
Q21 150  
Q22 25  
Q23 6  
Q24 3  
Q25 10  
Q26 (B)  
Q27 (C)  
Q28 (C)  
Q29 (B)  
Q30 (D)  
Q31 (A)  
Q32 (C)  
Q33 (A)

Q34 (D)  
Q35 (A)  
Q36 (C)  
Q37 (B)  
Q38 (A)  
Q39 (B)  
Q40 (B)  
Q41 (D)  
Q42 (B)  
Q43 (C)  
Q44 (D)  
Q45 (A)  
Q46 8  
Q47 36  
Q48 22  
Q49 4949  
Q50 768  
Q51 (D)  
Q52 (C)  
Q53 (A)  
Q54 (D)  
Q55 (C)  
Q56 (D)  
Q57 (D)  
Q58 (C)  
Q59 (C)  
Q60 (D)  
Q61 (B)  
Q62 (D)  
Q63 (D)  
Q64 (B)  
Q65 (D)  
Q66 (C)

[Android App](#)[iOS App](#)[PW Website](#)

Q67 (A)  
Q68 (D)  
Q69 (D)  
Q70 (A)  
Q71 1  
Q72 6  
Q73 372  
Q74 74  
Q75 2435



[Android App](#) | [iOS App](#) | [PW Website](#)



[Android App](#) | [iOS App](#) | [PW Website](#)

# Hints & Solutions

Note: scan the QR code to watch video solution

Q1 Video Solution:



Q2 Video Solution:



Q3 Video Solution:



Q4 Video Solution:



Q5 Video Solution:



Q6 Video Solution:



Q7 Video Solution:



Q8 Text Solution:

Video Solution:



Q9 Video Solution:



Q10 Video Solution:



Q11 Video Solution:



Q12 Video Solution:



[Android App](#)

| [iOS App](#)

| [PW Website](#)



Q13 Video Solution:



Q14 Video Solution:



Q15 Video Solution:



Q16 Video Solution:



Q17 Video Solution:



Q18 Video Solution:



Q19 Video Solution:



Q20 Video Solution:



Q21 Video Solution:



Q22 Video Solution:



Q23 Video Solution:



Q24 Video Solution:



Android App

iOS App

PW Website





Q25 Video Solution:



Q26 Video Solution:



Q27 Video Solution:



Q28 Video Solution:



Q29 Video Solution:



Q30 Video Solution:



Q31 Video Solution:



Q32 Video Solution:



Q33 Video Solution:



Q34 Video Solution:



Q35 Video Solution:



Q36 Video Solution:



[Android App](#)

| [iOS App](#)

| [PW Website](#)



Q37 Video Solution:



Q38 Video Solution:



Q39 Video Solution:



Q40 Video Solution:



Q41 Video Solution:



Q42 Video Solution:



Q43 Video Solution:



Q44 Video Solution:



Q45 Video Solution:



Q46 Video Solution:



Q47 Video Solution:



Q48 Video Solution:



[Android App](#)

| [iOS App](#)

| [PW Website](#)



Q49 Video Solution:



Q50 Video Solution:



Q51 Video Solution:



Q52 Video Solution:



Q53 Video Solution:



Q54 Video Solution:



Q55 Video Solution:



Q56 Video Solution:



Q57 Video Solution:



Q58 Video Solution:



Q59 Video Solution:



Q60 Video Solution:



Android App

iOS App

PW Website



Q61 Video Solution:



Q62 Video Solution:



Q63 Video Solution:



Q64 Video Solution:



Q65 Video Solution:



Q66 Video Solution:



Q67 Video Solution:



Q68 Video Solution:



Q69 Video Solution:



Q70 Video Solution:



Q71 Video Solution:



Q72 Text Solution:

.

Video Solution:



[Android App](#)

| [iOS App](#)

| [PW Website](#)



Q73 Video Solution:



Q74 Video Solution:



Q75 Video Solution:



[Android App](#)

| [iOS App](#)

| [PW Website](#)

