


Ans:- 1(a)

DFA

① DFA represents as Deterministic finite Automata.

② There is only one transition for one input in a node.



③ Transition in DFA is : $\delta \Rightarrow Q \times \Sigma \rightarrow Q$

④ DFA is considered as one machine.


⑤ DFA takes less time for computation but more space.

⑥ Difficult to construct

NFA

① NFA represents as Non-Deterministic finite automata.

② There can be more than one transition of one input in a node.



③ Transition in NFA is : $\delta \Rightarrow Q \times \Sigma \rightarrow 2^Q$

④ NFA can be considered as large no. of small machines working simultaneously.

⑤ NFA takes more time for computation but less space.


⑥ Easy to construct

Ans: 1(b)


Given ; transition table :-

Of Melay Machine

| | $q=0$ | | $q=1$ | | |
|-------|-------|--------|-------|--------|--|
| | state | output | state | output | |
| q_0 | q_1 | 1 | q_2 | 0 | |
| q_2 | q_4 | 1 | q_4 | 1 | |
| q_3 | q_2 | 0 | q_3 | 1 | |
| q_4 | q_3 | 0 | q_4 | 1 | |



Moore machine (Automata)



→ These are two new states are formed q_2' , q_3' .

Transition table is :- (Moore machine)

| states | 0 | 1 | output |
|---------|--------|--------|--------|
| → q_1 | q_1 | q_2 | 1 |
| q_2 | q_4 | q_4 | 0 |
| q_2' | q_2 | q_2 | 1 |
| q_3 | q_2 | q_3 | 1 |
| q_3' | q_2 | q_3 | 0 |
| q_4 | q_3' | q_2' | 1 |

Ans:- 1 (c)

The table of my Hill nerode is:-

→ q_0

| q_1 | q_2 | q_3 | q_4 | q_5 | q_6 | q_7 | q_8 | |
|-------|-----------------------------|--------------|--------------|--------------|--------------|--------------|--------------|-------|
| q_1 | \checkmark | \checkmark | \checkmark | \checkmark | \checkmark | \checkmark | \checkmark | |
| q_2 | \checkmark | \checkmark | \checkmark | \checkmark | \checkmark | \checkmark | \checkmark | |
| q_3 | \checkmark | \checkmark | \checkmark | \checkmark | \checkmark | \checkmark | \checkmark | |
| q_4 | \checkmark | \checkmark | \checkmark | \checkmark | \checkmark | \checkmark | \checkmark | |
| q_5 | \checkmark | \checkmark | \checkmark | \checkmark | \checkmark | \checkmark | \checkmark | |
| q_6 | \checkmark | \checkmark | \checkmark | \checkmark | \checkmark | \checkmark | \checkmark | |
| q_7 | \checkmark | \checkmark | \checkmark | \checkmark | \checkmark | \checkmark | \checkmark | |
| q_8 | \checkmark | \checkmark | \checkmark | \checkmark | \checkmark | \checkmark | \checkmark | |
| q_0 | q_1 | q_2 | q_3 | q_4 | q_5 | q_6 | q_7 | q_8 |

step-1

Mark all final states

step-2

(q_1, q_0) , (q_4, q_0) (q_4, q_1) (q_4, q_2) (q_4, q_3)
 (q_7, q_0) (q_7, q_1) (q_7, q_2) (q_7, q_3) (q_7, q_4) (q_7, q_5)
 (q_7, q_6)
 (q_8, q_0) (q_8, q_1) (q_8, q_2) (q_8, q_3) (q_8, q_4) (q_8, q_5)
 (q_8, q_6) (q_8, q_7)

step-3

find transition for - unmarked state

$$\begin{aligned}
 (q_1, q_0) &\rightarrow (q_0, 0) = q_1 \quad (q_0, 1) = q_4 \\
 (q_1, 0) &= q_2 \quad (q_1, 1) = q_3
 \end{aligned}$$

different marked

$$\begin{aligned}
 (q_4, q_0) &\rightarrow (q_4, 0) = q_5 \quad (q_4, 1) = q_6 \\
 (q_0, 0) &= q_1 \quad (q_0, 1) = q_4
 \end{aligned}$$

same marked

$$\begin{aligned}
 (q_4, q_1) &\rightarrow (q_4, 0) = q_5 \quad (q_4, 1) = q_6 \\
 (q_1, 0) &= q_2 \quad (q_1, 1) = q_3
 \end{aligned}$$

$$\begin{aligned}
 (q_4, q_2) &\rightarrow (q_4, 0) = q_5 \quad (q_4, 1) = q_6 \\
 (q_2, 0) &= q_7 \quad (q_2, 1) = q_8
 \end{aligned}$$

$$\begin{aligned}
 (q_4, q_3) &\rightarrow (q_4, 0) = q_5 \quad (q_4, 1) = q_6 \\
 (q_3, 0) &= q_8 \quad (q_3, 1) = q_7
 \end{aligned}$$

$$\begin{aligned}
 (q_7, q_0) &\rightarrow (q_7, 0) = q_7 \quad (q_7, 1) = q_7 \\
 (q_0, 0) &= q_1 \quad (q_0, 1) = q_4
 \end{aligned}$$

$$\begin{aligned}
 (q_8, q_1) &\rightarrow (q_8, 0) = q_7 \quad (q_8, 1) = q_7 \\
 (q_1, 0) &= q_2 \quad (q_1, 1) = q_3
 \end{aligned}$$

Similarly for all state we find transition.

Step 4


After marking all the states we get

(q_0, q_7, q_8)

(q_1, q_4)

(q_2, q_3, q_5, q_6)

So, Reduced DFA will be




step- 4.

After marking all the states, we get,

$\{q_0\}$, $\{q_1, q_8\}$, $\{q_1, q_4\}$, $\{q_2, q_3, q_5, q_6\}$

So, reduced DFA will be \Rightarrow





| states \ input | 0 | 1 |
|--------------------------|--------------------------|--------------------------|
| $\rightarrow q_0$ | $\{q_1, q_4\}$ | $\{q_1, q_4\}$ |
| $\{q_1, q_4\}$ | $\{q_2, q_3, q_5, q_6\}$ | $\{q_2, q_3, q_5, q_6\}$ |
| $\{q_1, q_8\}$ | $\{q_1, q_8\}$ | $\{q_1, q_8\}$ |
| $\{q_2, q_3, q_5, q_6\}$ | $\{q_1, q_8\}$ | $\{q_1, q_8\}$ |

Ans: 2(b)

$$R.E = (0+1)^* (00+11)^* (0+1)^*$$

$$L = \{ 0011, 000111, 010001, \dots \}$$



Transition table :-


| States \ i/o | 0 | 1 | ε |
|------------------|----------------|----------------|----------------|
| → q ₀ | q ₀ | q ₀ | q ₁ |
| q ₁ | q ₁ | q ₂ | — |
| q ₂ | q ₂ | q ₁ | q ₃ |
| q ₃ | q ₃ | q ₃ | — |

E-closer : $q_0 = q_0, q_1$

— 11 — $q_1 = q_1$

— 11 — $q_2 = q_2, q_3$

— 11 — $q_3 = q_3$



ज्ञानादेव तु कैवल्यम्

Ans- 2(c)

→ To prove any language is non-regular we operate pumping lemma to it

According to pumping lemma:-

Any ~~long~~ string belong to language it can be divided into

three parts as $x y^i z$ where $i \geq 0$ has some pumping lemma P it follows three rules are:-

- (i) $x y^i z \in L \quad \forall i \geq 0, n \geq 1$
- (ii) $y \neq \epsilon, i \geq 0$
- (iii) $|xy| \geq 0$

Let $L = \{a^n b^n \mid n \geq 1\}$

$L = \{ab, aabb, aaabbb, \dots\}$

⇒ step-1

Let take $n = 3$

for:-

$n = 1$

ab

$x = a$

$y = b$

$z = \epsilon$

ab^i

$i = 2$

$abb \notin L$

\therefore ~~it is not~~

\rightarrow for $n = 2$

$aaabbb$

$x = a, y = ab, z = b$

$a(ab)^i b$

$i = 2$

$aaabbbb$

$i = 3$

$aaaaabbbb$

\vdots

for ~~if~~ let take;

$x = a, y = aa, z = bb$

$a(a)^i bb$

$i = 2$

$aaaabb \notin L$

\rightarrow for $n = 3$

$aaaaabbbb$

$x = aa, y = abb, z = b$

$aa(abbb)^i b$

$$i = 2$$

aaabbbbbb $\notin L$

eg

step - 2 $i > 0$

(i) $i = 2$ (ii) $i = 2$ (iii) $i = 2$

Here $i \neq 0$

step 3 $(xy) \neq 0$

(i) ab $\neq 0$

(ii) aab $\neq 0$

(iii) aaabb $\neq 0$

Ans \therefore It is not a regular language

3(a)

(2)

$$\frac{(a+b)^*}{(0+1)^* (00)^+}$$

(1)

$$\frac{(1)^+ (1)^*}{(1)^+ (1)^*}$$

$$\frac{(1)^+ (1)^*}{(1)^+ (1)^*}$$

Ans- 3 & (c)

Given;

$$E \rightarrow E + T / T \quad \text{--- (i)}$$

$$T \rightarrow T * F / F \quad \text{--- (ii)}$$

$$F \rightarrow (E) / a \quad \text{--- (iii)}$$

Step-1

Elimination of null propagation.

Step-2

Elimination of Unit Propagation

$$E \rightarrow T \text{ \& \; } T \rightarrow F$$

&

$$T \rightarrow T * F / F \text{ and } F \rightarrow (E) / a$$

T will be

$$T \rightarrow T * F / (E) / a \quad \text{--- (iv)}$$

for

$$E \rightarrow T$$

put (iv) in (i)

$$E \rightarrow E + T / T * F / (E) / a \quad \text{--- (v)}$$

~~$E \rightarrow E + T / T * F / (E) / a$~~

Step-3 convert it into CNF (partially) .

$$E \rightarrow EC_1T / TC_2F / CEC_3 / a$$

$$C_1 \rightarrow +$$

$$C_2 \rightarrow *$$

$$C_3 \rightarrow]$$

Step-4

Rename

$$E \rightarrow A_1 ; C_1 \rightarrow A_2 ; T \rightarrow A_3 ; C_2 \rightarrow A_4 ; F \rightarrow A_5 ; E \rightarrow A_6 \\ C_3 \rightarrow A_7$$

$$A_1 \rightarrow A_1 A_2 A_3 / A_3 A_4 A_5 / C A_6 A_7 / a$$

Step-5

left Recursion

$$X_i \leftarrow X_j - X_i \rightarrow X_j ; i \leq j$$

$$A_1 \rightarrow A_3 A_4 A_5 / C A_6 A_7 / a / A_1 A_2 A_3$$

$$Z_1 \rightarrow A_2 A_3 Z_1 / A_2 A_3$$

Then,

$$A_1 \rightarrow A_3 A_4 A_5 / C A_6 A_7 / a / Z_1 / A_3 A_4 A_5 Z_1 \\ C A_6 A_7 Z_1 / a Z_1$$

8


$$z_1 \rightarrow + A_3 z_1 / + A_3$$

So the @ GNF is correspond it given

are

$$A_1 \rightarrow A_3 A_4 A_5 / (A_6 A_7 / a) z_1 / A_3 A_4 A_5 z_1 / (A_6 A_7 z_1 / a z_1$$

$$z_1 \rightarrow + A_3 z_1 / + A_3$$



ज्ञानादेन तु कैवल्यम्

Ans: 3(d)

$S \rightarrow 0.B / 1.A$

$A \rightarrow 0.0S / 1.AA$

$B \rightarrow 1.1S / 0.BB$

→ LMD (left most derivative)

$S \xrightarrow{LM} 0.B$

$\xrightarrow{RLM} 0.BB$

$\xrightarrow{LM} 00.BB.B$

$\xrightarrow{LM} 001.S.B.B$

$\xrightarrow{LM} 001.1.A.B.B$

$\xrightarrow{LM} 0011.0.B.B$

$\xrightarrow{LM} 00110.1.S.B$

$\xrightarrow{LM} 001101.0.B$

$\xrightarrow{LM} 0011010.1$

समाप्ति नु कृतम्

→ RMD (Right most derivative)

$S \xrightarrow{RM} 0.B$

→ $0.BB$

→ $0.B1$

→ $00.BB1$

→ $01.S1$

→ $011.A1$


→ $0110.S1$

→ $01101.A1$

→ ~~0110101~~

$S \xrightarrow{RM} 0B$
 $\xrightarrow{RM} \cancel{01S}$
 $\xrightarrow{RM} 00B$
 $\xrightarrow{RM} 001S$
 $\xrightarrow{RM} 0011A$
 $\xrightarrow{RM} 00110S$
 $\xrightarrow{RM} 001101A$
 $\xrightarrow{RM} 0011010B$
 $\xrightarrow{RM} 00110101$

Derivation tree:-



Ans:- (4) (b)



→ Push down Automata

$$L = \{a^n b^{2n}\} \text{ where } n \geq 1$$

$$L = \{abb, aabbbb, \dots\}$$



$$\text{Let } n = 2$$

$$w = a|a|b|b|b|\epsilon$$



wz \downarrow

| | | | | |
|---|---|---|---|---|
| a | a | b | b | b |
|---|---|---|---|---|



ज्ञानदेव तु केशवस्य

Ans- 4(c)

Turing Machine :-

Given;


$$L = \{a^n b^n c^n\} \text{ where } n \geq 0$$

$$L = \{a b c, a a b b c c, \dots\}$$

Let $n=2$

$w =$


| | | | | | | | |
|---|---|---|---|---|---|---|---|
| B | a | a | b | b | c | c | B |
|---|---|---|---|---|---|---|---|



$w =$


| | | | | | | | |
|---|--------------|--------------|---|---|---|---|---|
| B | a | a | b | b | c | c | B |
|---|--------------|--------------|---|---|---|---|---|

$\uparrow \quad \uparrow$




$w =$

| | | | | | | |
|--------------|--------------|--------------|--------------|--------------|--------------|--------------|
| B | a | a | b | c | c | B |
|--------------|--------------|--------------|--------------|--------------|--------------|--------------|



$w =$

| | | | | | | | |
|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|
| B | a | a | b | b | c | c | B |
|--------------|--------------|--------------|--------------|--------------|--------------|--------------|--------------|



| States | Read | write | Direction |
|--------|------|-------|-----------|
|--------|------|-------|-----------|

| | | | |
|-------|---|---|---|
| q_0 | a | x | R |
|-------|---|---|---|

| | | | |
|-------|---|---|---|
| q_0 | a | a | R |
|-------|---|---|---|

| | | | |
|-------|---|---|---|
| q_1 | b | y | R |
|-------|---|---|---|

| | | | |
|-------|---|---|---|
| q_1 | b | b | R |
|-------|---|---|---|

| | | | |
|-------|---|---|---|
| q_2 | c | z | L |
|-------|---|---|---|

| | | | |
|-------|---|---|---|
| q_2 | b | b | L |
|-------|---|---|---|

| | | | |
|-------|---|---|---|
| q_2 | y | y | L |
|-------|---|---|---|

| | | | |
|-------|---|---|---|
| q_2 | a | a | L |
|-------|---|---|---|

| | | | |
|-------|---|---|---|
| q_2 | x | x | L |
|-------|---|---|---|

| | | | |
|-------|---|---|---|
| q_3 | B | B | R |
|-------|---|---|---|

And ~~Transformation~~

~~$S(a, n, R)$~~

Ans:- 5 (a)

PARTIAL FUNCTION:-

→ Partial function is a type of function in which every value of input there can be distinct output.

But for the same value of input there can't be same value of output.

INITIAL FUNCTION:-

There are three types of initial functions.

(i) Zero function :- Every value of input the output is always zero.

Eg:-

$$Z(6) = 0 ; Z(5) = 0 \quad \text{i.e.,} \quad Z(n) = 0$$
$$Z(6.14) = 0$$

(2) Successive function :- for every value of x
the output is successive i.e.,
 $x+1$

$$S(6) = 7$$

$$S(9) = 10 \quad ; \quad \text{i.e.,} \quad S(n) = n+1$$

(3) Projection function :- In projection function value of i the output will be i .

$$U_i^j(x_1, x_2, x_3, x_4)$$

$$\text{Here } j=4$$

$$i=2$$

So,

$$U_2^4(x_1, x_2, x_3, x_4) = x_2$$

ज्ञानादेव तु केवलम्

Eg:-

$$U_2^3(6, 4, 7) = 4$$

Ans:- 5(b)

① $f(x, y) \equiv x * y$

step-1

put $y = 0$.

$$f(x, 0) = x * 0 = 0 = G_1(x)$$

step-2

put $y = y+1$

$$f(x, y+1) = x * (y+1)$$

$$= x * y + x * 1$$

$$= x * y + x$$

step 3

ज्ञानादेव तु केवल्यम्

$$f(x, y, f(x, y+1)) = x * y + x$$

$$= x * U_1^0(f(x), f(y), f(x, y+1))$$

$$= U_1^0(f(x), f(y), f(x, y+1)) +$$

$$U_1^0(f(x, y+1))$$

$\therefore I +$ is primitive recursive fn.

$$\Rightarrow \textcircled{11} \quad f(x, y) = x^y$$

Step-1 Put $x=0$

$$f(x, 0) = x = G(x)$$

Step-2 Put $y = y+1$

$$f(x, y+1) = x^{y+1} \\ = x^y \cdot x = x^y * x$$

Step-3

$$f(x, y, f(x, y+1)) = x^y + x \\ = U_1^1(x, y, f(x, y+1)) \\ = U_1^2(x, y, f(x, y+1)) \rightarrow * \\ = U_1^3(x, y+2)$$

\therefore it is recursive fn.

\Rightarrow Any function can be a primitive recursive function when

it is one of the initial function i.e. zero function, successor function, projection function and

we can obtain it by composition and rearranging.

Ans- 5(c)

→ There are two types of complexity in Turing machine


→ Space complexity & Time complexity
⇒ SPACE COMPLEXITY:-

→ Space complexity is a space occupied in the system by Turing machine to do computation

→ The space or the values of no. of tape cell required to perform computation is space complexity of Turing machine

→ The space complexity is of Turing machine is

$S(n) = 4$



Here it is 4.


→ The space complexity is the amount of space taken by Turing machine for computation.

⇒ TIME COMPLEXITY

→ Time taken by the Turing machine for computation is known as time complexity.

→ In Turing machine the no. of transition or steps of pointer is space complexity.


→ It is $T(n) = O(n)$



ज्ञानादेव तु कैवल्यम्

4(a) NPDA

- ① Non Deterministic push down automata
- ② It is less strong as compared to PDA
- ③ It is not common & for use
- ④ Its time & space complexity is High



ज्ञानादेव तु कैवल्यम्


PDA

- ① Deterministic push down automata
- ② It is more strong.
- ③ The uses of PDA is high in industrial
- ④ The space & time complexity is low ~~com~~ compared to NPDA


Ans - 2(a)

Property of regular grammar


- (1) A ~~is~~ Non-terminal to Nonterminal -
can be done
- (2) There can be not any
duplicate states.




ज्ञानादेव तु कैवल्यम्




ज्ञानादेन तु केवलम्




ज्ञानदेव तु कैवल्यम्




ज्ञानादयः तु कैवल्यम्




ਸਿਮਰਿ ਤੁ ਭਗਤੁ




सामान्य बुद्धि





ज्ञानं त्वं कैवल्यम्






ज्ञानादेव तु कैवल्यम्



सिद्धि विद्या





$A \ B \ B$
 ~~$0 \ S \ B \ B$~~
 ~~$0 \ B$~~
 ~~$A \ B \ B$~~
 0
 $B \ B$
 $1 \ S$
 0

1

1000

1000

